

Determination of the presence of hyaluronic acid in preparations containing amino acids: The molecular weight characterization

A. Bellomaria^a, R. Nepravishta^b, U. Mazzanti^c, M. Marchetti^c, P. Piccioli^c, M. Paci^c

^a Department of Chemical Sciences and Technology, University of Rome Tor Vergata, Via della Ricerca Scientifica 1, 00133 Rome, Italy

^b Cooperativa di Ricerca Finalizzata, Via della Ricerca Scientifica 1, 00133 Rome, Italy

^c Italfarmacia srl, Via Tor Sapienza, 8, Rome, Italy

ARTICLE INFO

Article history:

Received 17 April 2014

Received in revised form 2 July 2014

Accepted 19 July 2014

Available online 29 July 2014

Keywords:

Hyaluronic acid

Separation

DOSY

NMR spectroscopy

ABSTRACT

Several pharmaceutical preparations contain hyaluronic acid in the presence of a large variety of low molecular weight charged molecules like amino acids. In these mixtures, it is particularly difficult to determine the concentration and the molecular weight of the hyaluronic acid fragments. In fact zwitterionic compounds in high concentration behave by masking the hyaluronic acid due to the electrostatic interactions between amino acids and hyaluronic acid. In such conditions the common colorimetric test of the hyaluronic acid determination appears ineffective and in the ¹H NMR spectra the peaks of the polymer disappear completely. By a simple separation procedure the presence of hyaluronic acid was revealed by the DMAB test and ¹H NMR while its average molecular weight in the final product was determined by DOSY NMR spectroscopy alone. The latter determination is very important due to the healthy effects of some sizes of this polymer's fragments.

© 2014 Elsevier B.V. All rights reserved.

1. Introduction

Hyaluronic acid (HYA) is a linear polysaccharide composed of repeating units of D-glucuronic acid and D-N-acetylglucosamine, [→4)-beta-D-GlcA-(1→3)-beta-D-GlcNAc-(1→], extracted from a variety of animal tissues. Owing to its unique physico-chemical properties and relevant biological activities, hyaluronic acid has been used in a wide variety of products, such as food, biomedicine, biomaterials and cosmetics. Due to limited sources and the risk of viral infection the acid is now produced by optimized microbial fermentation, with high purification efficiency, low production

costs and a low risk of cross-species viral infection (). The characterization of the molecular weight appears particularly important since fragments of hyaluronic acids with low-molecular-weight (LMW-HYA) reveal different biological activities (). In fact average molecular weights in the range of 45.2–145 kDa fragments show pronounced free radical scavenging and antioxidant activities compared to those carried out by hyaluronic acid of very high MW (). LMW-HYAs also behave as effective angiogenic factors () and can promote excisional wound healing through enhanced angiogenesis (). Other evidence indicates that LMW-HYAs inhibit colorectal carcinoma growth by decreasing tumor cell proliferation and stimulating immune response (). Their use has been also described as a new adjuvant candidate in the preparation of dendritic cell-based anti-cancer vaccines with potent immune-stimulatory properties (). In fact for this reason it appears important to reduce the high-molecular weight hyaluronic acid in the LMW-HYA and many methods of degradation have been proposed () including the physical method (), the chemical approach () and the enzymatic process ()

Abbreviations: HYA, hyaluronic acid; DMAB, p-dimethylaminobenzaldehyde; DOSY, Diffusion Ordered Spectroscopy; LMW-HYA, low molecular weight hyaluronic acid; ADS, Amino acid Deprived and Sonicated; AD, Amino acid Deprived.

* Corresponding author at: Department of Chemical Sciences and Technology, University of Rome Tor Vergata, Via della Ricerca Scientifica 1, 00133 Rome, Italy. Tel.: +39 0672594410; fax: +39 0672594830.

E-mail addresses: (A. Bellomaria),

(R. Nepravishta),

(U. Mazzanti), (M. Marchetti),

(P. Piccioli), (M. Paci).

¹ Contributed equally to this work.

² Tel.: +39 0672594410; fax: +39 0672594830.

³ Tel.: +39 0672594461.

⁴ Tel.: +39 0645441800.

⁵ Tel.: +39 0672594446; fax: +39 0672594328.

). Fragments have also been produced by electron beam irradiation, gamma ray irradiation, microwave irradiation and thermal treatment (). In pharmaceutical products containing amino acids and other molecules, the HYA content is difficult to detect and quantify. Indeed, the large number of interactions between this polyelectrolyte polymer and the small charged or zwitterionic molecules creates a variety of non-selective physico-chemical interactions resulting in a large structure. The aim of this study was the characterization of HYA molecular weight by using a rapid, robust and simple NMR technique. The detection of HYA was performed by NMR spectroscopy, paralleled by a specific colorimetric test.

By Diffusion Ordered Spectroscopy (DOSY) NMR the MW determination was directly carried out in a rapid experiment. The commercial product AAA, Italy (an subcutaneous administration pharmaceutical form) which contains 0.1% w/v of HYA and several amino acids was analyzed. For the detection of HYA the direct routine DMAB test () and the ^1H NMR spectroscopy were used. Both techniques failed to reveal any presence of HYA in the intact product. Therefore this paper reports the results obtained regarding AAA by adopting a particular partial purification procedure through a nearly complete deprivation of the amino acid content. The procedure adopted allowed the characterization of the presence of HYA by a colorimetric test and ^1H NMR spectroscopy and the determination of the characteristic MW of the polymer by DOSY NMR.

2. Materials and methods

The AAA sample used for the analysis constituted of 0.1% HYA as sodium salt, L-isoleucine, L-leucine, L-lysine, L-proline, L-valine, glycine, L-serine, L-alanine, L-cysteine, sodium bicarbonate placed in sterile 5 mL vials was kindly donated by the producer in Italy. DMAB, sodium tetraborate and HYA at 13, 50, and 208 kDa mean MW were obtained from Sigma Aldrich (USA). The Mini Dialysis Kit at 1000 Da cut-off was purchased from Amersham Biosciences (USA).

The *p*-dimethylaminobenzaldehyde (DMAB) assay is a UV–Vis test for oligosaccharides containing N-acetylglucosamine in the reducing end. In this work we used a slightly modified method as previously described by . For the assay two working solutions were used: tetraborate solution, prepared by dissolving 3 g of sodium tetraborate in 10 mL H_2O at pH 9, and then DMAB solution, prepared by dissolving 0.5 g DMAB in an acid mixture made up of 0.6 mL of HCl 12 N and 4.4 mL of glacial acetic acid. Both working solutions were prepared immediately before use. In order to perform the DMAB assay 0.1 mL of AAA sample was added to 0.1 mL of tetraborate solution in a glass tube vortexed, heated for 5 min and then placed in cold water. At this point 0.8 mL of DMAB solution was added and the sample was incubated for 30 min at 37 °C. The UV–Vis measurements were performed on a Perkin Elmer spectrophotometer Lambda Bio by transferring the sample in a quartz cuvette of 1 cm path-length and the absorbance was measured at $\lambda = 544$ nm.

The amino acid deprivation of the sample was achieved by precipitation of 5 mL of AAA sample with 100 mL methanol which was then removed by evaporation. The precipitate was solubilized in 0.6 mL of H_2O and dialyzed for 48 h using the mini-dialysis kit at 1000 Da cut-off for complete amino acid deprivation. During the dialysis the amino acid deprivation from the sample was assayed by ^1H NMR spectroscopy following the decrease of specific resonances of amino acids until their complete absence and the consequent appearance of characteristic NMR HYA resonances.

The sample sonication was performed in an ultrasound bath operating at 28 kHz frequency for 6 h in order to achieve the fragmentation of the HA.

NMR experiments were performed on a Bruker Avance instrument operating at 700.13 MHz. All the samples were prepared by dissolving them in H_2O containing 10% D_2O for the lock signal and transferred on 5 mm NMR tubes. In order to avoid the water signal at 4.7 ppm the ^1H NMR experiments were acquired using the zgpr pulse sequence with 32 K and 64 scans at 298 K.

The DOSY calibration curve of HYA at different MW was performed in order to correlate the MW of the HYA molecules with their Diffusion coefficient. For the lowest limit of the curve the Diffusion coefficient of water molecules (MW = 18 Da) was taken into consideration. For this three HYA samples were purchased from Sigma Aldrich with a mean MW of respectively 13 kDa, 50 kDa, and 208 kDa. HYA reference samples were dissolved in water and 10% D_2O at 0.1% w/v concentration transferred in 5 mm NMR tubes; for each of them the DOSY spectra at 298 K were acquired. The log Diffusion coefficient (logD) was then correlated to the log MW of the HYA molecules giving an R^2 of 0.99. All the DOSY spectra were performed by using the ledbgppr2s pulse sequence in order to suppress the water signal at 4.7 ppm. During the DOSY experiment 32 mono dimensional spectra were obtained with 64 scans in a linear increasing gradient varying from 5% to 95% with a Δ of 70 ms and a δ of 2 ms. The spectra were then analyzed by using the DOSY module implemented in Bruker software TOPSPIN 3.1.

3. Results

As already reported the presence of amino acids and other molecules resulted both in the complete absence of reactivity in the DMAB test and the absence of the HYA resonance in the NMR spectrum. In fact in the NMR spectrum in only the resonances of the amino acids contained in the sample appear while the resonances due to the HYA are not visible. In fact it is well known that the HYA resonances are located at around 5.2 ppm and between 3.9 and 3.2 ppm in the ^1H NMR spectrum with a well-defined envelope (). Evidently the high concentration of amino acids with respect to HYA may mask the peaks of HYA; moreover, intermolecular interactions which HYA is able to establish with amino acids change to a great extent the tumbling of the macromolecule while the NMR resonances broaden as to be undetectable. It is important to mention that the original AAA sample studied was prepared with HYA (MW of 1.7 MDa) at a concentration of about 0.1% w/v with the addition of amounts of amino acids such as: L-isoleucine, L-leucine, L-lysine chloridrate, L-proline, L-valine, glycine, L-serine, L-alanine, L-cysteine and sodium bicarbonate. This solution was autoclaved for the sterilization with cycles of 20 min at 121 °C and a pressure of 2 bar. The final product was inserted in disposable vials for direct use as a pharmaceutical preparation.

3.1. The analysis of hyaluronic acid by colorimetric DMAB method

To reveal the HYA in the AAA sample, the DMAB assay was performed and the development of a red colored product due to the reaction of the terminal N-acetylglucosamine unit of HYA with the reagent DMAB () followed spectropho-




Fig. 1. NMR spectrum of the AAA sample.

tometrically. In this case also the test was unsuccessful, indicating the presence of negligible amounts of free HYA in the preparation, very far from the declared concentration of 0.1% w/v of HYA in the original product.

Because the presence of HYA in the AAA product was warranted, a purification procedure for the amino acid deprivation of samples was adopted as reported in Section (The scheme is in). Moreover one of the vials of AAA under study was exposed to ultrasound for 6 h in order to investigate the effects of ultrasounds on MW distribution (see below). The samples have been labeled as Amino acid Deprived and Sonicated (ADS) and Amino acid Deprived (AD), respectively.

The results of the DMAB test of the samples obtained are reported in . The absorbance due to the color developed by the reaction is clearly visible. As expected the color, developed in samples at a known mean MW of 13, 50, and 208 kDa (samples b, c, and d in), increases upon decreasing the MW. In fact the reaction involves the reducing terminal of the chain of HYA which, at the same monomer molarity, is more concentrated in low MW samples. Both microdialyzed samples (e and g in) showed a very low absorbance, which increased significantly after many steps of purification from amino acids (f-ADS and h-AD in). Moreover, the sonicated sample (f-ADS in) presented the highest value of absorbance. In fact the longer sonication time leads to a more fragmented HYA. In the NMR spectra appears also evident that in the purified samples the resonances of HYA at around 5.2 ppm and between 3.9 and 3.2 ppm are clearly visible (see h and f). The comparison of NMR spectra of ADS and AD reported in shows a slight difference between sonicated (f) and non-sonicated (h) in their resonances line width. This is in line with the fragmentation induced by sonication on the AADS sample which decreases the resonances line widths according to the lower molecular weight of the fragments. These results obtained after many steps of purification confirm that a large number of amino acid molecules strongly bind to the HYA polymer interfering in the DMAB test and increasing the molecular tumbling and the internal flexibility to broaden the NMR resonances beyond detection.

3.2. The characterization of molecular weight of hyaluronic acid by DOSY NMR

The characterization of the molecular weight of HYA samples has been also reported by other authors using other techniques




Fig. 2. The scheme of the procedure adopted for the amino acids deprivation of the samples of AAA as reported in Section . Letters are used to identify the samples.




Fig. 3. Results of DMAB colorimetric test. (a) AAA sample, (b) 13 kDa HYA sample, (c) 50 kDa HYA sample, (d) 208 kDa HYA sample, (e) sonicated for 6 h sample, (f) ADS sample, (g) Non-sonicated sample and (h) AD sample.




Fig. 4. NMR spectra of the AAA sample (g) ¹H NMR spectrum before amino acid deprivation and (h) ¹H NMR spectrum after deprivation of amino acids without sonication, and (f) ¹H NMR spectrum after deprivation of amino acids and sonication as reported in Section and the procedure of .

such as electrophoresis on SDS–PAGE and agarose gel (

); gel filtration chromatography (

) dynamic light scattering (

). Other colorimetric methods such as carbazole determination of uronic acid (

), high performance capillary electrophoresis (

), and, to a lesser extent, HPLC have also all been

used for quantification. Our approach is based on DOSY-NMR technique, an attractive non-destructive method making it possible to

determine a sample's MW by measuring its Diffusion coefficient, using small quantities; the technique is relatively fast and accurate

when compared to the upper mentioned methods. The DOSY-NMR technique allowed us to characterize the MW of the HYA polymer

in the AAA samples. DOSY-NMR is a powerful tool for structural studies of macromolecules and their interactions making it possible

to measure translational diffusion of dissolved molecules. In fact the value of Diffusion coefficient provides direct information on

translational dynamics, including intermolecular interactions (

), aggregation and conformational changes (

). Moreover, DOSY in the second dimension makes it possible to measure Diffusion

coefficient without a physical separation of mixture components (

). In our case this approach allowed us to determine the MW of the HYA fragments provided

that a calibration curve had been measured. The result of the calibration curve is reported in where in the log–log plot the

molecular Diffusion coefficients are reported as a function of the

MW of the polymers studied. (The polymers of 15, 30, 200 kDa have been used as standards). To extend meaningfully the measurements

of MW of fragments of HYA outside the calibration interval the Diffusion coefficient measured by the resonance of




Fig. 5. Calibration curve of Diffusion coefficient of HYA at different MWs from commercial standards. The value for water molecule observed in the spectra has been inserted.

water molecule has been included due to high linearity behavior of that value. In fact in the calibration curve including water Diffusion coefficient the linearity correlation coefficient was $R^2 = 0.99$. Strictly the hydrodynamic radius should be reported in abscissa instead of MWs, but at a first approximation one can consider the correspondence between the two values. The results of the DOSY experiments performed on the sample AD (left panel) and ADS (right panel) are reported in Fig. 6 where the resonances of the two samples are easily distinguishable at different MWs.

3.3. The effects of sonication on molecular weights of HYA

The DOSY spectra shown in Fig. 6 indicated that the ultrasound treatments led to a fragmentation of the hyaluronic acid in the preparation of AAA samples. In fact in the two samples the average MW of hyaluronic acid appears to be significantly different. In particular in the AD sample, the MW was estimated at about 8000–11000 Da. On the ADS sample the MW was estimated at about 270–500 Da, demonstrating that in our case the sonication produces a fragmentation of HYA polymer. This approach may be considered as complementary with the methods already established for the determination of MW of hyaluronic acid. The results reported here are a demonstration of utility of the DOSY NMR tech-

nique. Further studies for a complete validation of this technique with results obtained by other techniques is planned in the near future.

4. Discussion and conclusion

In conclusion the procedure of purification and verification by DMAB test and by NMR spectroscopy of HYA present in commercial preparations for human use, such as AAA has proved successful. It must be concluded that the presence of large amounts of amino acids is able to completely mask the colorimetric test and to fail the determination of the hyaluronic acid by NMR spectroscopy. The purification procedure adopted led to a complete characterization by ^1H NMR of the HYA present in the samples, removing the interference of other compounds. Moreover the DOSY-NMR spectroscopy allowed us to determine the molecular weight of the HYA fragments present in the product. Furthermore, the sonication in the conditions adopted led to a further increase in fragmentation of the HYA visible from the DOSY spectra. Thus, the sonication of AAA improves the fragmentation process and may find a broad application for obtaining LMW-HYA. Indeed, fragments have been studied and were found to actively stimulate many types of cells and to facilitate the clearance of debris or infectious agents ().

Acknowledgment

We are grateful to Mr. Fabio Bertocchi for NMR technical support during this work.

References




Fig. 6. The DOSY NMR spectra of the AD sample (left) and ADS sample (right). The arrows are for visual guidance.

