

Diffusion of Polymethylene Chain Molecules in Nonpolar Solvents


Bruce A. Kowert*

 Cite This: <https://dx.doi.org/10.1021/acs.jpcb.0c00063> Read Online

ACCESS |

 Metrics & More Article Recommendations

ABSTRACT: The translational diffusion constants, D , for solutes with polymethylene chains are compared with the predictions of a hydrodynamic bead model based on Kirkwood–Riseman theory. The solute–solvent combinations include (a) n -alkanes in n -alkanes; (b) n -alkanes in benzene, toluene, tetralin, decalin, and CCl_4 ; (c) 1-alkenes in n -alkanes; (d) 1-phenylalkanes in n -alkanes; and (e) 1-phenylalkanes in isocetane (2,2,4,4,6,8,8-heptamethylnonane), pristane (2,6,10,14-tetramethylpentadecane), and squalane (2,6,10,15,19,23-hexamethyltetracosane). The bead model gives good overall agreement with an average difference of less than 3% between 207 experimental and calculated diffusion constants that include published data as well as new D values determined for 1-alkenes and 1-phenylalkanes using capillary flow techniques. The calculated values are obtained using chain element (bead) radii that decrease as the solvent viscosity increases. The bead model's results are comparable to those obtained using cylinder and lollipop diffusion for many of the same solute–solvent systems; the three models are compared and discussed. The results for the 1-alkenes and n -alkanes in the n -alkanes are the first from a Kirkwood–Riseman analysis in a homologous series of solvents.



INTRODUCTION

Polymethylene chain molecules such as the n -alkanes are model systems for the study of the hydrodynamic interactions between the frictional elements of polymers. Kirkwood–Riseman theory^{1,2} takes these interactions into account and has been used in analyses of macromolecular transport properties. Paul and Mazo,³ in an early test of the theory, considered chains with up to 40 spherical elements (beads) and used Monte Carlo-generated conformers to calculate the size parameter $\phi(r)$ in the Stokes–Einstein relation for the translational diffusion constant, D , with stick boundary conditions^{4,5}

$$D = k_B T / 6\pi\eta\phi(r) \quad (1)$$

where k_B is Boltzmann's constant, T is the absolute temperature, and η is the viscosity; $\phi(r)$ has units of length and is determined by the number of chain elements, their size, and the hydrodynamic interactions. Paul and Mazo's values of $\phi(r)$, calculated with an element radius $a = 0.77$ Å, gave diffusion constants in agreement with those for n -alkanes with 5–32 carbon atoms in CCl_4 .³ The fit was not as good for n -alkane solutes in other solvents but could be improved using different values of a that were attributed³ to solute–solvent interactions.

In a subsequent series of papers, Garcia de la Torre and co-workers^{6,7} carried out Kirkwood–Riseman-based calculations of $\phi(r)$ for polymethylene chains using an improved conformation-averaging procedure and a more rigorous method of solving the associated hydrodynamic equations. Their results were con-

sistent with the experimental D values for the n -alkanes in CCl_4 and benzene for a chain element radius of 0.6 Å.^{6,7} Other groups^{8–10} have used Kirkwood–Riseman theory to analyze the diffusion of n -alkanes, but the number of solute–solvent systems considered was relatively small. In this paper, the results in refs 6 and 7 are used to determine an average difference of <3% between 207 experimental and calculated diffusion constants that vary by a factor of ~ 100 , as do the solutions' viscosities.

We find an average difference of <3% for 131 diffusion constants of n -alkane solutes in seven n -alkanes,^{11–24} benzene,^{25,26} toluene,¹⁰ tetralin,²⁵ decalin,²⁵ and CCl_4 .^{10,27} Similar analyses give average differences of <2% between 26 experimental and calculated D values of 1-alkenes in five n -alkanes²⁸ and <3% for 50 diffusion constants of 1-phenylalkanes in five n -alkanes, squalane (2,6,10,15,19,23-hexamethyltetracosane), isocetane (2,2,4,4,6,8,8-heptamethylnonane, HPMN), and pristane (2,6,10,14-tetramethylpentadecane).^{29,30} The results for the n -alkanes and 1-alkenes in the n -alkanes are the first from a Kirkwood–Riseman analysis in a homologous series of solvents.

Received: January 3, 2020

Revised: April 9, 2020

The analyses include new D values, determined using capillary flow techniques,^{31,32} for 1-alkenes (1-hexadecene and 1-eicosene in n -hexane, n -octane, n -decane, n -dodecane, n -tetradecane, n -hexadecane, and squalane) and 1-phenylalkanes (ethylbenzene, 1-phenylpentane, 1-phenyloctane, 1-phenylundecane, 1-phenyltetradecane, and 1-phenylheptadecane in n -hexadecane). Five of these 20 diffusion constants could not be compared with calculated values for reasons explained in the “Hydrodynamic Bead Model” section.

Our average difference for the bead model is comparable to those obtained for many of the same solute–solvent systems using the cylinder and lollipop diffusion models.^{29,30} The three models and their results are compared.

EXPERIMENTAL METHODS

Chemicals and Sample Preparation. In this and the following sections, n -C_{*i*}, 1-C_{*i*}, and 1-C_{*i*}Ph are used for particular n -alkanes, 1-alkenes, and 1-phenylalkanes, respectively. The suppliers for our solutes and solvents were (a) Aldrich: 1-C₈Ph (98%), 1-C₁₁Ph (99%), 1-C₁₇Ph (97%), n -C₈ (99+%), n -C₁₄ (99+%), and squalane (99%); (b) Sigma-Aldrich: n -C₆ (≥99%), n -C₁₀ (≥99%), and n -C₁₆ (99%); (c) Alfa Aesar: 1-C₅Ph (96%) and 1-C₁₄Ph (97%); (d) TCI: 1-C₁₆ (>99%) and 1-C₂₀ (>97%); (e) Sigma: n -C₁₂ (99%); and (f) Acros: ethylbenzene (99.8%). All substances were used as received. Solutions were prepared by adding two to five drops of the liquid solutes to 6.0–8.0 mL of the solvents.

Data Acquisition and Analysis. The microcapillary method used to determine the D values was developed by Bello and co-workers.³³ The acquisition of the solutes’ elution profiles was described in refs 31 and 32. The data acquisition system, including the microcapillary (76.5 μm i.d.) and variable wavelength detector, was the same as that in ref 32; wavelengths of 260 and 192 nm were used for the 1-phenylalkanes and 1-alkenes, respectively. The diffusion constants were obtained by comparing the sigmoidal elution profiles with those calculated using Taylor’s theory,³⁴ as described in refs 29, 31, and 32. All profiles were taken at room temperature, which was 22.50 ± 0.50 and 25.00 ± 0.25 °C for the 1-alkenes and 1-phenylalkanes, respectively. Multiple determinations of D were made for each solute–solvent pair; their average values, with uncertainties of ±2.5%, are given in Tables 1 and 2.

Table 1. Diffusion Constants of 1-C₁₆ and 1-C₂₀ in Alkanes at 22.50 ± 0.50 °C

solvent	10 ⁶ D (expt), cm ² s ^{−1}	
	1-C ₁₆	1-C ₂₀
n -C ₆	24.0	20.1
n -C ₈	15.0 ₅	13.0
n -C ₁₀	9.81	8.18
n -C ₁₂	6.61	5.67
n -C ₁₄	4.94	4.04
n -C ₁₆	3.75	3.08 ₅
squalane	0.876	0.667

Solute–Solvent Systems. The solvents for the n -alkane,^{10–27} 1-alkene,²⁸ and 1-phenylalkane^{29,30} solutes are given in Tables 3–5, respectively, as are the temperatures at which they were studied. A temperature near (or equal to) 25 °C was used for all solutions; the solvents’ viscosities^{25,35–39} at those temperatures are given in Tables 6–8 and will be used

Table 2. Diffusion Constants of Alkylbenzenes in n -C₁₆ at 25.0 ± 0.25 °C

solute	10 ⁶ D (expt), cm ² s ^{−1}
ethylbenzene	7.76 ₅
1-C ₃ Ph	5.91
1-C ₈ Ph	4.60
1-C ₁₁ Ph	3.78
1-C ₁₄ Ph	3.24
1-C ₁₇ Ph	2.95

when discussing our results in the sections “Solvent Dependence of σ ” and “Solvent-Dependent Size Parameters”. The n -alkanes in toluene, CCl₄, and benzene (Table 3) and the 1-alkenes in the n -alkanes (Table 4) were studied at additional temperatures; the viscosities at those temperatures^{25,35–39} are not given but were used in our analyses. The diffusion constants in refs 10–27 were reported from two to four significant figures; the average differences between the experimental and calculated D values reported here are given to two significant figures with a third subscripted figure.

RESULTS AND DISCUSSION

Hydrodynamic Bead Model. Garcia de la Torre and co-workers^{6,7} calculated values of $\phi(r) = f_i^*$ averaged over Monte Carlo-generated polymethylene conformers at 25 °C with a 500 cal mol^{−1} energy difference between t and g rotational isomers. Hydrodynamic interactions were taken into account using a modified Oseen tensor; translation–rotation coupling was included as was excluded volume. Stokes law

$$f_i = 6\pi\eta\sigma \quad (2)$$

was used as the friction coefficient for the N spherical chain elements (beads) with a common hydrodynamic radius, σ , and a separation $b = 1.53$ Å between elements. Values of f_i^* were reported as a function of σ/b for $0.0654 \leq \sigma/b \leq 0.490$ and $50 \geq N \geq 5$.^{6,7} We calculated diffusion constants using values of f_i^* interpolated from Table 2 of ref 7 and Figure 3 of ref 6; σ/b was varied to find the smallest average difference between the experimental and calculated D values for a given type of solute in a given solvent. The best values of σ/b and σ were solvent-dependent and are given in Tables 6, 7, and 8 for the n -alkanes, 1-alkenes, and 1-phenylalkanes, respectively; the corresponding average differences are given in Table 9.

The agreement for the n -alkane solutes’ 131 experimental and calculated diffusion constants ($32 \geq N \geq 5$) was generally good with an average difference of 2.81%. The average difference was 2.6% in the n -alkanes, 3.2% in the other hydrocarbon solvents (benzene, toluene, tetralin, and decalin), and 2.2% in CCl₄ (Table 9). Our values of $\sigma = 0.661$ Å for CCl₄ and 0.623 Å for benzene (Table 6) are consistent with $\sigma \sim 0.6$ Å found for both solvents by Garcia de la Torre and co-workers (who did not vary σ/b to the same extent as this analysis).^{6,7}

The n -alkanes’ diffusion constants in toluene,¹⁰ CCl₄,^{10,11,27} and benzene^{25,26} were determined over ranges of 73.0, 43.5, and 2.2 °C, respectively, but temperature-dependent values of σ/b were not needed to produce acceptable agreement with experiment. We found average differences of 2.2% in CCl₄, 1.9% in toluene, and 4.1% in benzene (Table 9) using a single value of σ/b for each solvent (Table 6). Paul and Mazo’s values of $\phi(r)$ were shown to be insensitive to temperature for n -alkane solutes;^{3,8,10} that appears to be the case for f_i^* as well.

Table 3. *n*-Alkane Solutes and Their Solvents

solvent	<i>i</i> for <i>n</i> -C _{<i>i</i>} solutes	<i>T</i> (°C)
<i>n</i> -C ₆ ^a	32, 24, 18, 16, 12, 10, 8, 7, 6, 5	25.0
<i>n</i> -C ₇ ^b	18, 16, 14, 12, 10, 8, 7, 6	25.0
<i>n</i> -C ₈ ^c	32, 24, 16, 14, 12, 8	25.0
<i>n</i> -C ₁₀ ^d	16, 7	25.0
<i>n</i> -C ₁₂ ^e	18, 16, 12, 8, 7, 6	25.0
<i>n</i> -C ₁₄ ^f	8, 7	25.0
<i>n</i> -C ₁₆ ^g	16, 12, 10, 8, 7, 6	25.0
benzene ^h	32, 29, 28, 18, 14, 12, 10, 9, 8, 7, 6, 5	22.2
benzene ⁱ	20, 18, 14, 10, 5	20.0
toluene ^j	26, 22, 18, 14, 10, 6	27.0, 50.0, 75.0, 100.0
tetralin ^h	28, 18, 14, 12, 9, 7, 6, 5	22.2
decalin ^h	28, 18, 14, 12, 10, 9, 8, 7, 6, 5	22.2
CCl ₄ ^k	32, 28, 22, 20, 18, 16, 12, 10, 8, 7, 6, 5	25.0
CCl ₄ ^j	26, 18, 14, 10, 6	20.5, 35.0, 50.0, 64.0

^aThe *D* value for 6 is the average of the values from ref 11; the rest are from ref 12. ^bThe *D* value for 18 is from ref 13; that for 16 is from ref 14; those for 14, 12, and 10 are from ref 15; that for 8 is from ref 16; that for 7 is from ref 17; and that for 6 is the average of those from ref 13. ^cThe *D* value for 8 is the average of the values in refs 18–21; that for 12 is from ref 19; that for 14 is from ref 15; and those for 16, 24, and 32 are from ref 12. ^dThe *D* value for 7 is from ref 15; the *D* value for 16 is from ref 11. ^eThe *D* value for 6 is from 22, that for 7 is from ref 15, those for 8 and 18 are from ref 19, that for 12 is the average of the values in refs 23 and 18, and the value for 16 is from ref 12. ^fThe *D* values for 7 and 8 are from ref 15. ^gThe *D* values for 6, 7, and 8 are from ref 12; those for 10 and 12 are from ref 11; and that for 16 is the average of the values in refs 23 and 24. ^hAll *D* values are from ref 25. ⁱAll *D* values are from ref 26. ^jAll *D* values are from ref 10. ^kThe *D* values for 28 and 32 are from ref 27; all others are from ref 11.

Table 4. 1-Alkene Solutes and Their Solvents

solvent	<i>i</i> for 1-C _{<i>i</i>} solutes ^a	<i>T</i> (°C)
<i>n</i> -C ₈	14, 12, 10, 8	25.0
<i>n</i> -C ₈	20, 16	22.5
<i>n</i> -C ₁₀	14, 12, 10, 8	25.0
<i>n</i> -C ₁₀	20, 16	22.5
<i>n</i> -C ₁₂	14, 12, 10, 8	25.0
<i>n</i> -C ₁₂	20, 16	22.5
<i>n</i> -C ₁₄	14, 12, 10, 8	25.0
<i>n</i> -C ₁₄	20, 16	22.5
<i>n</i> -C ₁₆	20, 16	22.5

^aThe *D* values for 1-C₁₆ and 1-C₂₀ are from this work; the others were calculated using the data in ref 28.

Table 5. 1-Phenylalkane Solutes and Their Solvents

solvent ^a	<i>i</i> for 1-C _{<i>i</i>} Ph solutes ^a	<i>T</i> (°C)
<i>n</i> -C ₉	17, 14, 11, 8, 5	24.0
<i>n</i> -C ₁₀	17, 14, 11, 8, 5	23.5
<i>n</i> -C ₁₂	17, 14, 11, 8, 5	20.5
<i>n</i> -C ₁₅	17, 14, 13, 12, 11, 9, 8, 7, 6, 5	24.5
<i>n</i> -C ₁₆	17, 14, 11, 8, 5	25.0
HPMN	17, 14, 11, 8, 5	21.0
pristane	17, 14, 11, 8, 5	25.5
squalane	17, 14, 13, 12, 11, 9, 8, 7, 6, 5	23.0

^aThe *D* values in *n*-C₁₆ are from this work; those in *n*-C₉, *n*-C₁₀, and *n*-C₁₂ are from ref 29; and those in *n*-C₁₅, HPMN, pristane, and squalane are from ref 30.

While the terminal groups of the 1-alkenes and 1-phenylalkanes are obviously different from the *n*-alkanes' methyl groups, we found an average difference of 1.8% for these solutes' 76 diffusion constants by assuming that the 1-phenylalkanes' phenyl ring and the 1-alkenes' double bond added a single equivalent element to their chains of *n*_c sp³ carbon atoms (*N* = *n*_c + 1); solutes with *n*_c ≥ 5 were considered with 17 ≥ *n*_c ≥ 5 for the 1-phenylalkanes and 18 ≥ *n*_c ≥ 6 for the 1-

Table 6. Values of the Solvents' Viscosities, σ/b , σ , and c_β for *n*-Alkane Solutes

solvent ^a	<i>T</i> ^b (°C)	10 ² η ^b (P)	σ/b	σ^c (Å)	c_β^d
<i>n</i> -C ₆	25.0	0.300 ^e	0.464	0.710	
<i>n</i> -C ₇	25.0	0.385 ^e	0.429	0.656	
<i>n</i> -C ₈	25.0	0.508 ^e	0.380	0.581	
<i>n</i> -C ₁₀	25.0	0.838 ^e	0.274	0.419	
<i>n</i> -C ₁₂	25.0	1.38 ^e	0.196	0.300	
<i>n</i> -C ₁₄	25.0	2.09 ^e	0.163	0.249	
<i>n</i> -C ₁₆	25.0	3.03 ^e	0.135	0.207	
toluene	27.0	0.535 ^f	0.422	0.646	
benzene	22.2	0.632 ^f	0.407	0.623	0.213
tetralin	22.2	2.10 ^g	0.222	0.340	0.121
decalin	22.2	2.64 ^g	0.196	0.300	0.099
CCl ₄	25.0	0.894 ^f	0.432	0.661	0.225

^aSee Table 3 for the list of solutes. ^bIf the *D* values were analyzed at more than one temperature in a given solvent (see Table 3), the temperature closest to (if not at) 25 °C and the corresponding viscosity are given. ^cValues of σ for *b* = 1.53 Å. ^dFrom ref 46 for benzene and CCl₄ at 25 °C, decalin and tetralin at 22.2 °C. ^eFrom ref 35. ^fViscosities for toluene, benzene, and CCl₄ at all temperatures in Table 3 are from ref 36. ^gFrom ref 25.

Table 7. Values of the Solvents' Viscosities, σ/b , and σ for 1-Alkene Solutes

solvent ^a	<i>T</i> (°C)	10 ² η ^b (P)	σ/b	σ^c (Å)
<i>n</i> -C ₈	25.0	0.508	0.419	0.641
<i>n</i> -C ₁₀	25.0	0.838	0.295	0.451
<i>n</i> -C ₁₂	25.0	1.38	0.221	0.338
<i>n</i> -C ₁₄	25.0	2.09	0.173	0.265
<i>n</i> -C ₁₆	25.0	3.03	0.147	0.225

^aSee Table 4 for the list of solutes. ^bThe viscosities at 25 °C are given, although some of the *D* values were obtained at 22.5 °C (see Table 4); the viscosities for both temperatures are from ref 35. ^cValues of σ for *b* = 1.53 Å.

Table 8. Values of the Solvents' Viscosities, σ/b , and σ for 1-Phenylalkane Solutes

solvent ^a	<i>T</i> (°C)	10 ² η (P)	σ/b	σ^b (Å)	1-C _i Ph/ <i>n</i> -C _i ^c	1-C _i Ph/1-C _i ^d
<i>n</i> -C ₉	24.0	0.674 ^e	0.459	0.702		
<i>n</i> -C ₁₀	23.5	0.857 ^e	0.453	0.693	1.65	1.54
<i>n</i> -C ₁₂	20.5	1.50 ^e	0.353	0.540	1.80	1.60
<i>n</i> -C ₁₅	24.5	2.57 ^e	0.217	0.332		
<i>n</i> -C ₁₆	25.0	3.03 ^e	0.216	0.330 _s	1.60	1.47
HPMN	21.0	3.44 ^f	0.234	0.358		
pristane	25.5	6.61 ^g	0.114	0.174		
squalane	23.0	30.3 ^h	0.0717	0.110		

^aSee Table 5 for the list of solutes. ^bValues of σ for $b = 1.53$ Å. ^c $\sigma(1\text{-C}_i\text{Ph})/\sigma(n\text{-C}_i)$. ^d $\sigma(1\text{-C}_i\text{Ph})/\sigma(1\text{-C}_i)$. ^eFrom ref 35. ^fFrom ref 37. ^gFrom ref 38. ^hFrom ref 39.

Table 9. Values of Average Percentage Difference

solutes	no. of solutes ^a	solvents	avg % diff
<i>n</i> -C _i	40	<i>n</i> -C _i	2.6 ₀
<i>n</i> -C _i	8	tetralin	3.7 ₉
<i>n</i> -C _i	10	decalin	4.2 ₄
<i>n</i> -C _i	17	benzene	4.1 ₆
<i>n</i> -C _i	24	toluene	1.9 ₇
<i>n</i> -C _i	59	te, de, bz, to ^b	3.2 ₃
<i>n</i> -C _i	32	CCl ₄	2.2 ₉
<i>n</i> -C _i	131	total	2.8 ₁
1-C _i	26	<i>n</i> -C _i	1.5 ₀
1-C _i Ph	30	<i>n</i> -C _i	1.7 ₀
1-C _i Ph	20	H, P, S ^c	2.6 ₆
1-C _i Ph	50	total	2.0 ₈
1-C _i , 1-C _i Ph	total	76	1.8 ₈
all solutes	207	all solvents	2.4 ₇

^aFrom Tables 3–5 for *n*-C_i, 1-C_i, and 1-C_iPh, respectively. ^bTetralin, decalin, benzene, and toluene. ^cHPMN, pristane, and squalane.

alkenes. The average differences were 2.0₈% for the 1-phenylalkanes in the *n*-alkanes, HPMN, pristane, and squalane and 1.5₀% for the 1-alkenes in the *n*-alkanes (Table 9); the 1-alkenes were studied at two temperatures (Table 4), and as for the *n*-alkanes, a single value of σ/b gave satisfactory agreement in each solvent (Table 7).

This degree of agreement would seem to indicate that, for this model and these solutes, the differences in the shapes and sizes of their terminal groups and the attendant modifications of the conformational averaging are not as important as the number of elements and their nonpolar character (as well as that of the solvents). The 2.4₇% average difference between the 207 experimental and calculated diffusion constants for the *n*-alkanes, 1-alkenes, and 1-phenylalkanes (Figure 1 and Table 9) indicates the model's usefulness for solutions in which the *D* values and viscosities vary from $5.57 \times 10^{-5} \text{ cm}^2 \text{ s}^{-1}$ for *n*-C₆ in toluene¹⁰ at 100 °C ($\eta = 0.280 \text{ cP}$ ³⁶) to $5.68 \times 10^{-7} \text{ cm}^2 \text{ s}^{-1}$ for 1-C₁₇Ph in squalane³⁰ at 23 °C ($\eta = 30.3 \text{ cP}$ ³⁹). It would be interesting to see what differences are found when calculations are carried out for alkyl chains with terminal phenyl and vinyl groups.

The values of σ/b for our fits determine the strength of the hydrodynamic interactions between the beads, which is given by⁴⁰

$$\lambda = f_i / 8\pi\eta b \quad (3)$$

which becomes

$$\lambda = (3/4)(\sigma/b) \quad (4)$$




Figure 1. Comparison of the experimental diffusion constants (in $\text{cm}^2 \text{ s}^{-1}$) for *n*-alkane, 1-alkene, and 1-phenylalkane solutes with the predictions of a hydrodynamic bead model based on Kirkwood–Riseman theory.

when eq 2 is used for f_i . Zwanzig, Kiefer, and Weiss, in an analysis of Kirkwood–Riseman theory,⁴⁰ said that “interesting values of λ can range from 0.10 to 10”. While $\lambda = 0.10$ was not stated to be an absolute lower limit, it gives $\sigma/b = 0.133$, which is greater than the values for the 1-phenylalkanes in pristane (0.114) and squalane (0.0717) and is close to that for the *n*-alkanes in *n*-C₁₆ (0.135).

A reviewer questioned whether it was appropriate to use a hydrodynamic model for these small values of σ/b and addressed the issue by calculating the self-diffusion constants for *n*-C₈, *n*-C₁₂, and *n*-C₁₆. These are solvents for which our *n*-alkane solutes, including their self-diffusion constants, had values of σ/b that decreased from 0.380 for *n*-C₈ to 0.196 for *n*-C₁₂ to 0.135 for *n*-C₁₆. The reviewer's *D* values for the three *n*-alkanes were obtained from simulations in boxes of varying length. The diffusion constants for each of the *n*-alkanes increased as the box length increased. The *D* values in each box for a given *n*-alkane became equal; however, if Yeh and Hummer's hydrodynamically based correction to infinite box size⁴¹ was applied, inferring that the molecules' diffusion can be described by classical hydrodynamics and the Kirkwood–Riseman equation can be used for our solvents with higher viscosities and smaller values of σ/b .

Calculations of f_i^* for more values of σ/b than those given in refs 6 and 7 also would be helpful because the minimum average differences could not be determined for the 1-alkenes in *n*-C₆ and squalane. In *n*-C₆, values of σ/b greater than the upper limit of 0.490 are required for the 1-C_i with $i = 8, 10, 12$, and 14 from ref 28 and $i = 16$ and 20 from Table 1. In squalane, values less than the lower limit of 0.0654 are needed for the 1-C_i with $i = 14$ from ref 31 and $i = 16$ and 20 from Table 1, although additional simulations would seem necessary to check if values of $\sigma/b <$

0.0654 can be used. The D values for 1- C_{16} , 1- C_{20} , and ethylbenzene (with $n_c = 2$) were reported, since they may be useful in other analyses.

Solvent Dependence of σ . Tables 6–8 give our values of σ and, as explained in the “Solute–Solvent Systems” section, the solvents’ viscosities at the temperatures close to (if not at) 25 °C. The values of σ for a given type of solute decrease as the viscosity of a given type of solvent increases; for example, $\sigma = 0.710$, 0.419, and 0.207 Å for the n -alkanes in n - C_6 , n - C_{10} , and n - C_{16} , respectively. The values of σ also are small compared to 2.00 Å, the common van der Waals radius of the methyl and methylene groups;⁴² the largest is $\sigma = 0.710$ Å for the n -alkanes in n - C_6 , and the smallest is 0.110 Å for the 1-phenylalkanes in squalane.

The results for the n -alkane and 1-alkene solutes in the n -alkanes are, to our knowledge, the first from a Kirkwood–Riseman-based study in a homologous series of solvents; the values of σ can be fitted to

$$\sigma = A_\sigma / \eta^p \quad (5)$$

where η is in cP at 25 °C (Table 6). The fits, shown in Figure 2, give $A_\sigma = 0.377_5$ Å, $p = 0.559$, and $R^2 = 0.995$ for the n -alkane




Figure 2. Values of σ for the n -alkane and 1-alkene solutes in the n -alkane solvents vs the solvents’ viscosities. The fit is to eq 5.

solutes and $A_\sigma = 0.417$ Å, $p = 0.588$, and $R^2 = 0.995$ for the 1-alkenes; the uncertainties in σ are $\pm 3.5\%$. The 1-phenylalkanes also are solutes in a homologous series of the n - C_i ($i = 9, 10, 12, 15, 16$), but the ranges of σ and η are narrower than those for the n -alkanes and 1-alkenes. The n -alkanes in benzene, toluene, decalin, and tetralin are not in a homologous series nor are the 1-phenylalkanes in HPMN, pristane, and squalane, but as seen in Figure 3, where all three types of solutes in the hydrocarbon solvents are included, the trend is obvious; σ decreases by a factor of ~ 6.5 , while the viscosities increase by a factor of ~ 100 .

Stepito and co-workers⁹ found a parameter similar to σ in their analysis of the diffusion constants of n -alkanes in four of the solvents considered here: benzene and CCl_4 at 25.0 °C and decalin and tetralin at 22.2 °C. Kirkwood–Riseman calculations were carried out using the rotational-isomeric-state model⁴³ and a segmental friction constant for the solute suggested by Freire and Horta^{44–46}

$$f_i = 6\pi\eta c_\beta \beta \quad (6)$$




Figure 3. Comparison of the values of σ that give the best agreement between experimental and calculated D values and the solvents’ viscosities for n -alkanes in n -alkanes; 1-alkenes in n -alkanes; n -alkanes in benzene, toluene, tetralin, and decalin; 1-phenylalkanes in n -alkanes; and 1-phenylalkanes in HPMN, pristane, and squalane.

where $\beta = (\langle r^2 \rangle / n)^{1/2}$ is the effective bond length of a polymethylene chain with n bonds and a root-mean-square end-to-end distance $\langle r^2 \rangle^{1/2}$; $c_\beta \beta$ is the effective segment radius. The experimental diffusion constants for the n -alkanes could be fitted with solvent-dependent values of c_β that (with the exception of CCl_4 and benzene) decreased as the viscosity increased;^{45,46} they are given in Table 6. Figure 4 shows a




Figure 4. Comparison of the values of c_β from ref 46 with our values of σ from Table 6.

reasonably good correlation between c_β and σ ; our values of σ for CCl_4 and benzene have the same reversed viscosity dependence as c_β (Table 6). There is a difference between Stepito’s approach and ours; the segmental friction constant in eq 4 depends on chain length (through β),^{44–46} while Garcia de la Torre’s, eq 2, does not.⁹ Note that Stepito’s^{9,45,46} parameters were c_b and b ; we have used c_β and β to avoid confusion with our $b = 1.53$ Å.

COMPARISON OF MODELS

Cylinder and Lollipop Models. The diffusion constants for many of the solute–solvent systems considered here also have

been compared with the predictions of the cylinder and lollipop models.³⁰ The solute size parameters from those analyses likewise decreased as the solvents' viscosities increased and were small compared to the van der Waals radius of the methyl and methylene groups.⁴² The cylinder model⁴⁷ assumed the solute to be a rigid rod divided into elements with diameter d and length l . The experimental diffusion constants were compared with values calculated as a function of l/d . For each value of l/d , the value of d that gave the best agreement with experiment for a given type of solute in a given solvent was determined. A value of $l/d = 1.57$ gave the lowest average difference of 3.3% between 150 experimental and calculated D values.³⁰ The values of d generally decreased as the viscosity increased for n -alkanes in n -alkanes; 1-alkenes in n -alkanes; 1-phenylalkanes in n -alkanes; and 1-phenylalkanes in HPMN, pristane, and squalane.³⁰

The diffusion constants of 1-phenylalkanes in n -alkanes, HPMN, pristane, and squalane also were analyzed using the lollipop model.⁴⁸ The phenyl group is the candy, a sphere with radius σ_1 , and the alkyl chain is the rigid handle, a linear array of spheres with radius σ_2 . For each value of σ_1/σ_2 , the value of σ_2 that gave the best agreement with experiment for a given type of solute in a given solvent was determined. A value of $\sigma_1/\sigma_2 = 2.17$ gave the lowest average difference of 3.7% between 85 experimental and calculated D values.³⁰ The values of σ_2 that gave the best agreement in the n -alkanes decreased as the viscosity increased, as did those in in HPMN, pristane, and squalane.³⁰

The average differences for the bead, cylinder, and lollipop models are comparable, although that for the bead model, which is more characteristic of the solutes, is slightly smaller and was obtained using more D values. The bead model also varied only one size parameter, σ (b remained fixed), while the cylinder model varied l/d and d and the lollipop model varied σ_1/σ_2 and σ_2 . One of the conclusions from the cylinder and lollipop analyses, however, was that the reasonably good agreement indicated that the alkyl chains of the n -alkanes, 1-alkenes, and 1-phenylalkanes were relatively extended.^{29,30} This was consistent with Monte Carlo and molecular dynamics (MD) simulations as well as experimental results;^{49–52} the time-averaged shapes calculated for the n -C_{*i*} with $i = 8, 12, 16$, and 20 were⁵⁰ “basically rodlike” with “little change in the length between dynamic time steps”. The bead model was intended for n -alkane solutes, and its success with them indicates the values of f_i^* are consistent with an extended shape;⁷ this may be a reason the model works as well as it does for the 1-alkenes and 1-phenylalkanes.

Solvent-Dependent Size Parameters. The solvent-dependent values of σ , like the solutes' size parameters for the cylinder and lollipop models,³⁰ are indicative of solute–solvent interactions. Zwanzig and Harrison⁵³ pointed out that these effective hydrodynamic sizes are a measure of the coupling of the solute motion to the solvent flow; the weaker the coupling, the smaller the value of σ . Two trends in our values of σ , which are most clearly seen in the n -alkane solvents, are consistent with this explanation. The first is the strength of the solutes' interactions with a common solvent. It would be expected to decrease in the order 1-phenylalkanes > 1-alkenes > n -alkanes because of the differences in their structures. Table 8 shows this to be the case in n -C₁₀, n -C₁₂, and n -C₁₆; the values of σ for the 1-phenylalkanes are 47–60% larger than those for the 1-alkenes and 60–80% larger than those for the n -alkanes.

The second trend is the decrease in σ as the viscosity increases. Solutes moving through the n -alkanes' relatively extended regions have relatively weak interactions with them.⁵⁰

The lengths of these regions increase as their chain lengths (and viscosities) increase, leading to longer intervals of weak interactions and smaller values of σ (Table 6). As seen in Figure 3, the correlation is evident although not as strong for the other solute–solvent groupings. Molecular dynamics simulations would be helpful in gaining a more detailed understanding of the relation between the values of σ and the intermolecular forces in these systems.

Others, including Paul and Mazo,³ have discussed solvent-dependent size parameters in terms of solute–solvent interactions. Stepto and co-workers interpreted their values of c_β in terms of the solutes' interactions with the actual structure of the solvents as opposed to their assumed continuous nature.^{9,45,46} Garcia de la Torre et al.⁷ pointed out that both local structure (short-range) and excluded volume (long-range) interactions contribute to the hydrodynamic properties of real chains.

SUMMARY AND CONCLUSIONS

The diffusion constants of 1-C₁₆ and 1-C₂₀ in a series of n -alkanes and several 1-phenylalkanes in n -C₁₆ have been determined at room temperature using capillary flow techniques. They are among the 207 D values that were compared with the predictions of a hydrodynamic bead model^{6,7} based on Kirkwood–Riseman theory.^{1,2} The solutes and solvents, whose D values and viscosities both vary by a factor of ~ 100 , include (a) n -alkanes in n -alkanes; (b) n -alkanes in benzene, toluene, tetralin, decalin, and CCl₄; (c) 1-alkenes in n -alkanes; (d) 1-phenylalkanes in n -alkanes; and (e) 1-phenylalkanes in methyl-substituted alkanes. The agreement is generally good; the average difference between the experimental and calculated D values is 2.4%. This result is comparable to that for many of the same solute–solvent systems using the cylinder and lollipop diffusion models.^{29,30}

The diffusion constants were calculated as a function of the bead radius, which was found to decrease as the solvent viscosity increased; this correlation of solute size parameters with viscosity was also found for many of the same solute–solvent combinations using the cylinder and lollipop models. The solvent dependence of the size parameters is attributed to solute–solvent interactions. The results for the 1-alkenes and n -alkanes in the n -alkanes are the first from a Kirkwood–Riseman analysis in a homologous series of solvents.

AUTHOR INFORMATION

Corresponding Author

Bruce A. Kowert – Department of Chemistry, Saint Louis University, St. Louis, Missouri 63103, United States;
orcid.org/0000-0002-4030-8670; Phone: 314-977-2837;
Email: bruce.kowert@slu.edu; Fax: 314-977-2521

Complete contact information is available at:
<https://pubs.acs.org/10.1021/acs.jpcb.0c00063>

Notes

The author declares no competing financial interest.

ACKNOWLEDGMENTS

The Department of Chemistry, Saint Louis University, has supported this research. The data acquisition system and detector were purchased with grants to Dr. Barry Hogan from Research Corporation and the donors of the Petroleum Research Fund, administered by the American Chemical

Society. We thank Dr. Michael Lewis for the 1-C₃Ph and 1-C₈Ph. We thank the referee for the helpful comments and calculations regarding the small values of σ/b .

REFERENCES

- (1) Kirkwood, J. G.; Riseman, J. The Intrinsic Viscosities and Diffusion Constants of Flexible Macromolecules in Solution. *J. Chem. Phys.* **1948**, *16*, 565–573.
- (2) Kirkwood, J. The General Theory of Reversible Processes in Solutions of Macromolecules. *J. Polym. Sci.* **1954**, *12*, 1–14.
- (3) Paul, E.; Mazo, R. M. Calculations of the Diffusion Coefficients of *n*-Alkanes. *J. Chem. Phys.* **1968**, *48*, 1405–1407.
- (4) Tyrrell, H. J. V.; Harris, K. R. *Diffusion in Liquids*; Butterworths: London, 1984.
- (5) Longuet-Higgins, H. C. Transport Processes in Fluids. *Il Nuovo Cim.* **1955**, *1* (Ser. 10, Suppl. 2), 140–155.
- (6) Garcia de la Torre, J.; Freire, J. J. Intrinsic Viscosities and Translational Diffusion Coefficients of *n*-Alkanes in Solution. *Macromolecules* **1982**, *15*, 155–159.
- (7) Garcia de la Torre, J.; Lopez Martinez, M. C.; Tirado, M. M.; Freire, J. J. Monte Carlo Study of Hydrodynamic Properties of Flexible Linear Chains: Analysis of Several Approximate Methods. *Macromolecules* **1984**, *17*, 2715–2722.
- (8) Dewan, R. K.; Van Holde, K. E. Role of Hydrodynamic Interaction in the Diffusion of *n*-Alkanes in Carbon Tetrachloride. *J. Chem. Phys.* **1963**, *39*, 1820–1824.
- (9) Mokrys, I. J.; Rigby, D.; Stepto, R. F. T. The Diffusion of Short Polymer Chains in Dilute Solution. *Ber. Bunsenges. Phys. Chem.* **1979**, *83*, 446–454.
- (10) Chen, H.-C.; Chen, S.-H. Taylor Dispersion Measurement of the Diffusivities of Polymethylenes in Dilute Solution. *Chem. Eng. Sci.* **1985**, *40*, 521–526.
- (11) Suarez-Iglesias, O.; Medina, I.; Pizarro, C.; Bueno, J. L. Modeling of Tracer Diffusion in Liquids when Solute-Solvent Interactions Are Present. *Fluid Phase Equilib.* **2007**, *253*, 155–164.
- (12) Hayduk, W.; Ioakimidis, S. Liquid Diffusivities in Normal Paraffin Solutions. *J. Chem. Eng. Data* **1976**, *21*, 255–260.
- (13) Winkelmann, J. Diffusion Coefficient of Octadecane in Heptane. In *Diffusion in Gases, Liquids and Electrolytes*. *Physical Chemistry*; Lechner, M. D., Ed.; Springer: Berlin, Heidelberg, 2018; Vol. 15B2.
- (14) Bidlack, D. L.; Anderson, D. K. Mutual Diffusion in Nonideal, Nonassociating Liquid Systems. *J. Phys. Chem.* **1964**, *68*, 3790–3794.
- (15) Lo, H. Y. Diffusion Coefficients in Binary Liquid *n*-Alkane Systems. *J. Chem. Eng. Data* **1974**, *19*, 236–241.
- (16) Matthews, M. A.; Akgerman, A. Diffusion Coefficients for Binary Alkane Mixtures to 573 K and 3.5 MPa. *AIChE J.* **1987**, *33*, 881–885.
- (17) Ertl, H.; Dullien, F. A. L. Self-Diffusion and Viscosity of Some Liquids as a Function of Temperature. *AIChE J.* **1973**, *19*, 1215–1223.
- (18) Iwahashi, M.; Yamaguchi, Y.; Ogura, Y.; Suzuki, M. Dynamical Structures of Normal Alkanes, Alcohols, and Fatty Acids in the Liquid State as Determined by Viscosity, Self-Diffusion Coefficient, Infrared Spectra, and ¹³C NMR Spin-Lattice Relaxation Time Measurements. *Bull. Chem. Soc. Jpn.* **1990**, *63*, 2154–2158.
- (19) Van Geet, A. L.; Adamson, A. W. Diffusion in Liquid Hydrocarbon Mixtures. *J. Phys. Chem.* **1964**, *68*, 238–246.
- (20) McCall, D. W.; Douglass, D. C.; Anderson, E. W. Self-Diffusion Studies by Means of Nuclear Magnetic Resonance Spin-Echo Techniques. *Ber. Bunsenges. Phys. Chem.* **1963**, *67*, 336–340.
- (21) Tofts, P. S.; Lloyd, D.; Clark, C. A.; Barker, G. J.; Parker, G. J. M.; McConville, P.; Baldock, C.; Pope, J. M. Test Liquids for Quantitative MRI Measurements of Self-Diffusion Coefficient In Vivo. *Magn. Reson. Med.* **2000**, *43*, 368–374.
- (22) Shieh, J. C.; Lyons, P. A. Transport Properties of Liquid *n*-Alkanes. *J. Phys. Chem.* **1969**, *73*, 3258–3264.
- (23) Tofts, P. S.; Lloyd, D.; Clark, C. A.; Barker, G. J.; Parker, G. J. M.; McConville, P.; Baldock, C.; Pope, J. M. Test Liquids for Quantitative MRI Measurements of Self-Diffusion Coefficient In Vivo. *Magn. Reson. Med.* **2000**, *43*, 368–374.
- (24) Dymond, J. H.; Harris, K. R. The Temperature and Density Dependence of the Self-Diffusion Coefficient of *n*-Hexadecane. *Mol. Phys.* **1992**, *75*, 461–466.
- (25) Varoqui, R.; Daune, M.; Freund, L. Configuration and Mechanism of Diffusion of *n*-Paraffins in Dilute Solution. *J. Chim. Phys. Phys.-Chim. Biol.* **1961**, *58*, 394–401.
- (26) Rossi, C.; Bianchi, E.; Rossi, A. Measurement of the Diffusion in Benzene. *J. Chim. Phys. Phys.-Chim. Biol.* **1958**, *55*, 97–101.
- (27) Hayduk, W.; Buckley, W. D. Effect of Molecular Size and Shape on Diffusivity in Dilute Liquid Solutions. *Chem. Eng. Sci.* **1972**, *27*, 1997–2003.
- (28) Kowert, B. A.; Turner, R. M., II; Caldwell, C. V. C. Diffusion of 1-Alkenes and Cyclohexene in Alkane Solvents. *Chem. Phys.* **2008**, *344*, 114–120.
- (29) Kowert, B. A.; Register, P. M. Diffusion of Benzene and Alkylbenzenes in *n*-Alkanes. *J. Phys. Chem. B* **2015**, *119*, 12931–12937.
- (30) Kowert, B. A. Diffusion of Benzene and Alkylbenzenes in Nonpolar Solvents. *J. Phys. Chem. B* **2018**, *122*, 1940–1947.
- (31) Kowert, B. A.; Watson, M. B. Diffusion of Organic Solutes in Squalane. *J. Phys. Chem. B* **2011**, *115*, 9687–9694.
- (32) Kowert, B. A.; Watson, M. B.; Dang, N. C. Diffusion of Squalene in *n*-Alkanes and Squalane. *J. Phys. Chem. B* **2014**, *118*, 2157–2163.
- (33) Bello, M. S.; Rezzonico, R.; Righetti, P. G. Use of Taylor-Aris Dispersion for Measurement of a Solute Diffusion Coefficient in Thin Capillaries. *Science* **1994**, *266*, 773–776.
- (34) Taylor, G. The Dispersion of Matter in Turbulent Flow through a Pipe. *Proc. Royal Soc. London, Ser. A* **1954**, *223*, 446–468 and references therein..
- (35) Viswanath, D. S.; Natarajan, G. *Data Book on the Viscosity of Liquids*; Hemisphere Publishing: New York, 1989.
- (36) Yaws, C. L. *Handbook of Viscosity*; Gulf Publishing Co.: Houston, TX, 1995; Vols. 1, 2.
- (37) Prak, D. J. C.; Truelove, P. C.; Cowart, J. S. Density, Viscosity, Speed of Sound, Surface Tension, and Flash Point of *n*-Hexadecane and 2,2,4,4,6,8,8-Heptamethylnonane and of Algal-Based Hydrotreated Renewable Diesel. *J. Chem. Eng. Data* **2013**, *58*, 920–926.
- (38) Et-Tahir, A.; Boned, C.; Lagourette, B.; Xans, P. Determination of the Viscosity of Various Hydrocarbons and Mixtures of Hydrocarbons Versus Temperature and Pressure. *Int. J. Thermophys.* **1995**, *16*, 1309–1334.
- (39) Barlow, A. J.; Erginsav, A.; Lamb, J. Viscoelastic Relaxation of Supercooled Liquids. II. *Proc. Royal Soc. London, Ser. A* **1967**, *298*, 481–494.
- (40) Zwanzig, R.; Kiefer, J.; Weiss, G. H. On the Validity of the Kirkwood-Riseman Theory. *Proc. Natl. Acad. Sci. U. S. A.* **1968**, *60*, 381–386.
- (41) Yeh, I.-C.; Hummer, G. System-Size Dependence of Diffusion Coefficients and Viscosities from Molecular Dynamics Simulations with Periodic Boundary Conditions. *J. Phys. Chem. B* **2004**, *108*, 15873–15879.
- (42) Pauling, L. *The Nature of the Chemical Bond*, 3rd ed.; Cornell University Press: Ithaca, NY, 1960; pp 257–264.
- (43) Flory, P. J. *Statistical Mechanics of Chain Molecules*; Interscience: New York, 1969.
- (44) Freire, J. J.; Horta, A. Mean Reciprocal Distances of Short Polymethylene Chains. Calculation of the Translational Diffusion Coefficient of *n*-Alkanes. *J. Chem. Phys.* **1976**, *65*, 4049–4054.
- (45) Edwards, C. J. C.; Kaye, A.; Stepto, R. F. T. Translation Diffusion of Polymer Chains in Dilute Solutions. *Macromolecules* **1984**, *17*, 773–782.
- (46) Stepto, R. F. T. Interpretation of the Effects of Segmental Friction on the Diffusion Coefficients of Polymer Chains. *Makromol. Chem.* **1989**, *190*, 549–557.
- (47) Hansen, S. Translational Friction Coefficients for Cylinders of Arbitrary Axial Ratios Estimated by Monte Carlo Simulation. *J. Chem. Phys.* **2004**, *121*, 9111–9115.
- (48) Garcia de la Torre, J.; Bloomfield, V. A. Hydrodynamic Properties of Macromolecular Complexes. I. Translation. *Biopolymers* **1977**, *16*, 1747–1763.

(49) Thomas, L. L.; Christakis, T. J.; Jorgensen, W. L. Conformation of Alkanes in the Gas Phase and Pure Liquids. *J. Phys. Chem. B* **2006**, *110*, 21198–21204.

(50) Zhang, Y.; Venable, R. M.; Pastor, R. W. Molecular Dynamics Simulations of Neat Alkanes: The Viscosity Dependence of Rotational Relaxation. *J. Phys. Chem.* **1996**, *100*, 2652–2660.

(51) Unruh, T.; Smuda, C.; Busch, S.; Neuhaus, J.; Petry, W. Diffusive Motions in Liquid Medium-Chain *n*-Alkanes as Seen by Quasielastic Time-of-Flight Neutron Spectroscopy. *J. Chem. Phys.* **2008**, *129*, 121106.

(52) Morhenn, H.; Busch, S.; Unruh, T. Chain Dynamics in a Hexadecane Melt as Seen by Neutron Scattering and Identified by Molecular Dynamics Simulations. *J. Phys.: Condens. Matter* **2012**, *24*, 375108.

(53) Zwanzig, R.; Harrison, A. K. Modifications of the Stokes-Einstein Formula. *J. Chem. Phys.* **1985**, *83*, 5861–5862.