

Week 1: Euclidean Vector Spaces

Exercise 1

- The Cartesian product $A_1 \times \cdots \times A_n$ of sets A_1, \dots, A_n is defined as
 - $A_1 \times \cdots \times A_n := A_1 \setminus (A_2 \cup \dots \cup A_n)$
 - $A_1 \times \cdots \times A_n := \{(a_1, \dots, a_n) \mid a_1 \in A_1, \dots, a_n \in A_n\}$
 - $A_1 \times \cdots \times A_n := \{a_1 \cdot a_2 \cdot \dots \cdot a_n \mid a_1 \in A_1, \dots, a_n \in A_n\}$
- Let $n \in \mathbb{N} \setminus \{0\}$. Then \mathbb{R}^n contains
 - n real numbers
 - n -tuples of real numbers
 - n -tuples of vectors
- Which of the following sets together with standard addition and scalar multiplication does not form a real vector space?
 - The set of integers \mathbb{Z}
 - The set of complex numbers \mathbb{C}
 - The set of real-valued, continuous functions $\{f: \mathbb{R}^n \rightarrow \mathbb{R} \mid f \text{ continuous}\}$
- Scalar multiplication in a real vector space $V = (\mathcal{V}, +, \cdot)$ is given by a mapping
 - $\mathcal{V} \times \mathcal{V} \rightarrow \mathbb{R}$
 - $\mathbb{R} \times \mathcal{V} \rightarrow \mathcal{V}$
 - $\mathbb{R} \times \mathcal{V} \rightarrow \mathbb{R}$
- How many vector subspaces does \mathbb{R}^2 have?
 - two: $\{\mathbf{0}\}$ and \mathbb{R}^2
 - four: $\{\mathbf{0}\}, \mathbb{R} \times \{0\}, \{0\} \times \mathbb{R}$ (the axes), \mathbb{R}^2
 - infinitely many
- Which of the following subsets of \mathbb{R}^2 is not a vector subspace?
 - $\{\mathbf{0}\}$
 - $\{\mathbf{x} \in \mathbb{R}^2 \mid x_1 = 2x_2\}$
 - $\{\mathbf{x} \in \mathbb{R}^2 \mid x_1 = x_2 + 1\}$
- For which of the following objects does it make sense to say that they are “linearly independent”?
 - Elements of a vector space $\mathbf{v}_1, \dots, \mathbf{v}_n \in \mathcal{V}$
 - Real numbers $\lambda_1, \dots, \lambda_n \in \mathbb{R}$
 - The linear combination $\lambda_1 \mathbf{v}_1 + \dots + \lambda_n \mathbf{v}_n$
- Which of the following vectors form a basis of \mathbb{R}^2 ?
 - $\begin{bmatrix} 2 \\ 4 \end{bmatrix}, \begin{bmatrix} 2 \\ 3 \end{bmatrix}$
 - $\begin{bmatrix} 1 \\ 0 \end{bmatrix}, \begin{bmatrix} -1 \\ 0 \end{bmatrix}$
 - $\begin{bmatrix} 1 \\ 5 \end{bmatrix}, \begin{bmatrix} 3 \\ -6 \end{bmatrix}, \begin{bmatrix} 2 \\ 3 \end{bmatrix}$
- What is the scalar product of the vectors $\begin{bmatrix} 1 \\ -2 \\ 0 \end{bmatrix}$ and $\begin{bmatrix} 3 \\ 0 \\ 3 \end{bmatrix}$?
 - 3
 - 5
 - 7
- Which of the following statements is false?
 - For any Euclidean vector space $(V, \langle \cdot, \cdot \rangle)$ the function $\|\cdot\|$ defined by $\|\mathbf{v}\| := \sqrt{\langle \mathbf{v}, \mathbf{v} \rangle}$ ($\mathbf{v} \in \mathcal{V}$) satisfies the properties of a norm.
 - For any Euclidean vector space $(V, \langle \cdot, \cdot \rangle)$ it holds that $\langle \mathbf{v}, \mathbf{w} \rangle \geq 0$ for all $\mathbf{v}, \mathbf{w} \in \mathcal{V}$.
 - Let M be a subspace of an Euclidean vector space $(V, \langle \cdot, \cdot \rangle)$. Then
$$\{\mathbf{v} \in \mathcal{V} \mid \forall \mathbf{u} \in M: \mathbf{v} \perp \mathbf{u}\}$$
is also subspace of V .

Exercise 2

Consider $(\mathbb{R} \setminus \{-1\}, \star)$, where $a \star b := ab + a + b$ with $a, b \in \mathbb{R} \setminus \{-1\}$.

1. Show that $(\mathbb{R} \setminus \{-1\}, \star)$ is an Abelian group.
2. Solve $3 \star x \star x = 15$ in the Abelian group $(\mathbb{R} \setminus \{-1\}, \star)$.
3. Is $(\mathbb{R} \setminus \{-1\}, +, \star)$ a field? Justify!

Exercise 3

Show: If $\mathbf{v}_1, \dots, \mathbf{v}_n$ form an orthonormal basis of a Euclidean vector space $(V, \langle \cdot, \cdot \rangle)$, the following holds for all $\mathbf{x} \in V$:

$$\mathbf{x} = \sum_{i=1}^n \langle \mathbf{x}, \mathbf{v}_i \rangle \mathbf{v}_i$$

Hint: Establish first that \mathbf{x} can be expressed as $\mathbf{x} = \lambda_1 \mathbf{v}_1 + \dots + \lambda_n \mathbf{v}_n$. Then show that $\langle \mathbf{x}, \mathbf{v}_i \rangle = \lambda_i$ for all $1 \leq i \leq n$.

Exercise 4

Let $(V, \langle \cdot, \cdot \rangle)$ be a Euclidean vector space and U an r -dimensional vector subspace $U \subseteq V$ with orthonormal basis $\mathbf{u}_1, \dots, \mathbf{u}_r$. The orthogonal projection of a vector $\mathbf{v} \in V$ onto U is given by

$$p(\mathbf{v}) := \sum_{i=1}^r \langle \mathbf{v}, \mathbf{u}_i \rangle \mathbf{u}_i$$

1. Compute the orthogonal projection of the vector $(25, 0)^\top$ onto the subspace spanned by the vector $(3, 4)^\top$. Visualize the subspace and the projection in a drawing with both vectors.
2. Let $\mathbf{x} \in V$ and $\lambda \in \mathbb{R}$. Show:

$$p(\lambda \mathbf{x}) = \lambda p(\mathbf{x}).$$

3. Let $\mathbf{x}, \mathbf{y} \in V$. Show:

$$p(\mathbf{x} + \mathbf{y}) = p(\mathbf{x}) + p(\mathbf{y}).$$

Exercise 5

Let $\langle \cdot, \cdot \rangle$ be the standard scalar product on \mathbb{R}^n .

1. Show that the mapping

$$\begin{aligned} k: \mathbb{R}^2 \times \mathbb{R}^2 &\rightarrow \mathbb{R} \\ (\mathbf{x}, \mathbf{y}) &\mapsto \langle \mathbf{x}, \mathbf{y} \rangle^2 \end{aligned}$$

does not define a scalar product on \mathbb{R}^2 .

2. Consider the mapping $\Phi: \mathbb{R}^2 \rightarrow \mathbb{R}^3$ with

$$\begin{bmatrix} x_1 \\ x_2 \end{bmatrix} \mapsto \begin{bmatrix} x_1^2 \\ x_2^2 \\ \sqrt{2} \cdot x_1 x_2 \end{bmatrix}.$$

Show:

$$\langle \Phi(\mathbf{x}), \Phi(\mathbf{y}) \rangle = k(\mathbf{x}, \mathbf{y})$$

Exercise 6

The 1-norm is often used for finding sparse solutions to an optimization problem (vectors or matrices with many entries equal to zero). This will be demonstrated in the following exercise:

We are looking for a vector $\mathbf{w} = (x, y)^\top \in \mathbb{R}^2$, which solves the optimization problem

$$\max_{\mathbf{w}} f(\mathbf{w}) \quad \text{s.t.} \quad \|\mathbf{w}\| = 1$$

Consider $f(x, y) = \frac{1}{2}x + y$ and compare the solutions to this optimization problem for the 1- and the 2-norm.

1. Draw the set of all points on the x - y -plane that have 2-norm equal to 1 (i.e. the ℓ_2 unit circle).

$$C_2 := \left\{ \mathbf{w} \in \mathbb{R}^2 \mid \|\mathbf{w}\|_2 = \sqrt{x^2 + y^2} = 1 \right\}.$$

2. Draw the set of all points on the x - y -plane that have 1-norm equal to 1 (i.e. the ℓ_1 unit circle).

$$C_1 := \{\mathbf{w} \in \mathbb{R}^2 \mid \|\mathbf{w}\|_1 = |x| + |y| = 1\}.$$

3. Draw contour lines $c = f(x, y)$ for $c = 0.5$, $c = 1$ and $c = 1.1$.

4. Where in your drawing does the solution to the optimization problem lie for the 1-norm? Where does it lie for the 2-norm?

Exercise 7

Given training data (red and blue) a new data point (green) should be classified using the k -NN algorithm. Which label does the new data point receive for $k \in 1, 2, 5$?

