




Data Science
Machine Learning
~Somyanush


Simple Linear RegressionSupervised Machine Learning

$$y = mx + c$$




$$h_0(x) = \theta_0 + \theta_1 x$$

~~If New θ_0 = Intercept
if $x=0$ then $h_0(x) = \theta_0$~~



~~If New θ_1 = Slope or coefficient~~




Errors = $y - \hat{y}$ [where \hat{y} = predicted value of y by model]

Now our aim is to minimise the sum of all errors

$$\min \left[\sum (y - \hat{y})^2 \right]$$

$$\hat{y} = \text{predicted}$$



Machine Learning → Notes Part 2

Cost function

Gradient Descent → An algorithm to minimize a function by optimizing its parameters

Let's say there is a Science Test Max marks = 50

Friend asks you to guess his marks?

guess1 → 45 ?


friend → Too far

Guess2 → 40 ?

friend → Still far

guess3 → 37

friend → Very close



Imp → In GD we start with random guess & then slowly move to the right answer.

How fast (slow) we'll converge to the answer is determined by learning rate


$$\text{New value} = \text{Old value} - \text{Stepsize}$$

where StepSize = Learning rate × Slope

Eg → A simple square function

$$f(x) = x^2$$

Now as per GD this function should be minimised.



New step 1) Random guess (-2, 3) ($x \& y$)

Step 2 → If $x = -2$ then $f(x) = x^2$ &

$$\text{Slope of function: } \frac{d}{dx} f(x) = 2x = 2(-2) = -4 \checkmark$$

Step 3 If $x = -1$ then $2x$ ie slope = -2 ✓

Hence we are getting close to zero ie moving in correct direction

Step 4 → If $x = +2$ then $2x$ ie slope = 4 which is beyond zero so we're in wrong direction now

Step 4 → Now $\boxed{\text{new value} = \text{old value} - \text{slope}}_{\times LR}$

& if a function has multiple parameters unlike $f(x) = x^2$, ie function having more than one parameter.

Eg Cost funcⁿ for regression =

$$J(m, c) = \sum_{i=1}^n (y_i - (mx_i + c))^2$$

Step 5 → Now there are two params in b.c.

Step 6 → Again random guess let's say
 $c=0, m=1$

x	y
1	2
3	4

← Training Data

Step 7 → Now cost function for linear regn
 $= [y - (mx + c)]^2$

Plugging the values for training data (1, 2) & (3, 4)

$$J(m, c) = [2 - (c + mx_1)]^2 + [4 - (c + mx_2)]^2$$

partial derivative

$$\frac{\partial J}{\partial c} = -2[2 - (c + m \cdot 1)] - 2[4 - (c + m \cdot 3)]$$

for $c=0$ & $m=1$

$$= -2[2 - 1] - 2[4 - 3]$$

$$= -4$$

Step 8 Now because i differentiated wrt c

so my

$$\begin{aligned} \text{new } c &= \text{old } c - LR \times (-4) && \text{slope} \\ &= 0 - (0.001) \times (-4) \\ &= 0.004 \end{aligned}$$

Step 9 Similar to above now we'll differentiate wrt m & new value of m will be found

Step 10 → New new values will be taken for consideration & entire process will get repeated

Step 11 → When algo will stop?

Ans → When there won't be much improvement in cost for values of c & m

Step 12 → Choosing learning Rate! Try to be moderate

Machine Learning Notes → 3


Multiple Linear Regression

House pricing dataset

No. of Rooms	Size of house	Location	Price
1	1800	x	100000 \$
2	1900	y	200000 \$
3	2000	z	1000000 \$


$$y = m_1 x_1 + m_2 x_2 + m_3 x_3 + \dots + m_n x_n + c$$

Example curve, multiple I/P features,



Performance Metrics used in Linear Regression


$$1) R^2 \text{ squared} = 1 - \frac{SS_{\text{Residual}}}{SS_{\text{Total}}}$$



classmate
Page

$$\text{SS Residual} = \text{Sum of Square Residual} = \sum_{i=1}^n (y_i - \hat{y}_i)^2$$

$$\text{SS Total (Sum of square total)} = \sum_{i=1}^n (y_i - \bar{y}_i)^2$$



$$\therefore R^{\text{square}} = 1 - \frac{\sum_{i=1}^n (y_i - \hat{y}_i)^2}{\sum_{i=1}^n (y_i - \bar{y}_i)^2}$$

$$1 - \frac{\text{Small Number}}{\text{Big Number}} \Rightarrow 1 - \text{Small Number} \approx 1$$

0.70 → 70% Accuracy
0.85 → 85% " & so on

2) Adjusted R squared \leftrightarrow Extra feature (Owner DOB)

$$1 - \frac{(1-R^2)(N-1)}{N-p-1}$$

N = no. of data points

p = no. of independent features

dataset

No. of Rooms	Size	Location	Owner DOB	Price
—	—	—	—	—

$R^2 \uparrow$
 $\text{Adj } R^2 \downarrow$

} via penalty through
denominator.

$$\text{Ex } p=2 \quad R^2 = 80\% \quad \text{Adj } R^2 = 76\%$$

$$p=3 \quad R^2 = 85\% \quad \text{Adj } R^2 = 79\%$$

(Penalty of
non correlated feature)