
a
contrast changes are changes of spread
observed contrast change (σ)
contrast change predicted by the spread change (σ)
one population (slope 1)
two types, Δ = 2 (slope 0.58)
SNAPSI zonal (post hoc)
SNAPSI full 3-D (post hoc)
SNAPSI free, natural (registered)
b
law (filled) vs shift only (open)
SNAPSI zonal-mean
SNAPSI SH warming
SNAPSI full 3-D
CESM2, January 2021
SNAPSI free, natural
pooled, five sets
24 regional cold classes
human forcing, 3 models
temperature, 3 models
precipitation, 4 models
replication: temperature
replication: precipitation
forced contrast minus null contrast (σ)
c
elasticity: one population 1; forced mixing of types ≥ 2 s.d. apart below 0.6
SNAPSI zonal (36) (post hoc)
SNAPSI free (registered)
human forcing, 3 models (registered)
expected under the exact law (post hoc)
Δ = 4
Δ = 3
two types, Δ = 2
one population
elasticity of the contrast with respect to the spread
d
share of the variance that fixed-type mixing requires but is absent
ρ (0 = fixed-type mixing)
replication (CESM2, MPI-GE; registered)
temperature
precipitation
not rejected after Holm (open)
one translated population (1)
fixed response types (0)
forced change of the mean, lowest to highest forcing (s.d. of the lowest)