



Synthesis, Characterization and Evaluation of Quaternary Compounds of 4-(N-Substituted)-3-Pyridyl-5-Mercapto-S-Triazole with Secondary Amines

Anil Manikrao*¹, Prafulla M Sabale², Prashant Prajapati²

¹Food and drug administration, 1st floor Dr. B. R. Ambedkar Market, Jalgaon-425 001

²Department of Pharma Chemistry, Parul Institute of Pharmacy, Limda-391 760,Vadodara, Gujarat.

Abstract In the present study, the interaction of 4-(N-substituted)-3-pyridyl-5-mercapto-s-triazole with secondary amines was explored. The Isonicotinic acid hydrazide was converted into the corresponding potassium dithiocarbazinate, by reacting with carbon disulphide in alkaline medium which undergoes ring closure reaction after further treatment with aqueous potassium hydroxide to give 5-pyridyl-2-mercapto-1,3,4-oxadiazole. 4-(N-pyridylcarboxamido)-3- pyridyl-5-mercapto-s-triazole and 3-pyridyl-4-amino-5-mercapto-s-triazole were obtained in one pot reaction by heating equimolar quantities of oxadiazole with isonicotinic acid hydrazide and hydrazine hydrate respectively. Condensation of triazole with secondary amines in anhydrous dimethyl sulfoxide results in the formation of corresponding quaternary salts. The synthesized compounds were confirmed by IR, ¹NMR spectra and elemental analysis. All the compounds were screened for their preliminary *in-vitro* antibacterial and antifungal activity.

Keywords: 1,3,4-Oxadiazole, Quaternary Compound, Morpholine, Pyrrolidine.

Received on 31-12-2012

Modified on 15-01-2013

Accepted on 10-02-2013

INTRODUCTION

The treatment of infectious diseases still remains an important and challenging problem because of a combination of factors including emerging infectious diseases and the increasing number of multi-drug resistant microbial pathogens. In spite of a large number of antibiotics and chemotherapeutics available for medical use, at the same time the emergence of old and new antibiotic resistance created in the last decades revealed a substantial medical need for new classes of antimicrobial agents. There is perceived need for the discovery of new compounds endowed with antimicrobial activity, possibly

*Address for correspondence:

Dr. Anil Manikrao,
Food and drug administration,
1st floor Dr. B. R. Ambedkar Market,
Jalgaon-425 001, Maharashtra
E-mail: anilmanikrao@gmail.com
Contact: 9373556025

acting through mechanisms of action, which are distinct from those of well-known classes of antibacterial agents to which many clinically relevant pathogens are now resistant. Though the various molecules were designed and synthesized for this aim, it was demonstrated that the mercapto- and thione- substituted 1,2,4-triazoles and their derivatives could be considered as possible antimicrobial agents. The substituted triazoles are heterocyclic compounds, which serve both as biomimetic and reactive pharmacophores and many are key elements with potential biological activities such as analgesic¹, anti-inflammatory²⁻⁵, antimicrobial⁶⁻¹¹, tuberculostatic¹²⁻¹³, and can be used as fungicides¹⁴ and antitumour agents¹⁵⁻¹⁶.

In this study, the interaction of 4-(N-substituted)-3-pyridyl-5-mercapto-s-triazole with secondary amines, pyrrolidine (**a**) and morpholine (**b**) was explored. The Isonicotinic acid hydrazide (INH) (**1**) was converted in to the corresponding potassium dithiocarbazinate, by reacting with carbon disulphide in alkaline medium which

undergoes ring closure reaction after further treatment with aqueous potassium hydroxide to give 5-pyridyl-2-mercapto-1,3,4-oxadiazole (2). 4-(N-pyridyl carboxamido)-3-pyridyl-5-mercapto-s-triazole (3) and 3-pyridyl-4-amino-5-mercapto-s-triazoles (4) were obtained in one pot reaction by heating equimolar quantities of oxadiazole with INH and hydrazine hydrate respectively, according to Reid and Heindel procedure with slight modification¹⁷. Condensation of both triazole (3 & 4) with secondary amines (a & b) when heated at 80°C in anhydrous dimethyl sulfoxide (DMSO) results in the formation of corresponding 4-(N-pyridylcarboxamido)-3-pyridyl-s-triazole-5-thiolate (3a & 3b) and 3-pyridyl-4-amino-s-triazole-5-thiolate (4a & 4b) as a quaternary salts as shown in Scheme-I. This technique of conversion of oxadiazole to triazole is similar to well-known transformation of furans to pyrroles with nucleophilic amines. In this oxadiazole was intermediate in the hydrazinolysis of dithiocarbazinate to triazole. The identity of those was confirmed by elemental analysis and spectral data. All the novel compounds were


evaluated for their preliminary *in-vitro* antibacterial and antifungal activity.

EXPERIMENTAL

GENERAL

Melting points were taken using microprocessor based melting point apparatus (Veego make) containing liquid paraffin and are uncorrected. Thin layer chromatography was used to monitor the progress of the reaction on silica gel-G precoated plates by using chloroform and ethyl acetate (1:1) as the eluent and observed in UV light. IR spectra in KBr were recorded on Shimadzu-8400 FTIR spectrophotometer, ¹H NMR Spectra were recorded on Bruker spectrophotometer(400MHz) in DMSO-d₆/CDCl₃ using trimethylsilane as an internal standard (chemical shifts are expressed in δ, ppm). Isonicotinic acid hydrazide and all reagents were used of analytical grade purchased from commercial suppliers without further purification.

Scheme for the synthesis of Quaternary Compounds of 4-(N-Substituted)-3-Pyridyl-5-Mercapto-S-Triazole



Scheme-I

Preparation of 4-(N-pyridyl carboxamido)-3-pyridyl-5-mercapto-s-triazole (3)

To a solution of Potassium hydroxide (0.015mol) in absolute ethanol, 5-pyridyl-2-mercapto-1, 3, 4-oxadiazole (0.01mol) and carbon disulphide (0.01mol) were added and mixture was stirred for 24 hrs. The solid separated out, to this Isonicotinic acid hydrazide (0.01mol) was added and refluxed at 140⁰ for 8 hrs. The reaction mixture was cooled and diluted with water. On acidification with hydrochloric acid the solid obtained was filtered, washed with water and recrystallized from aqueous ethanol¹⁷ yield 84%, m.p.290-294 ⁰C.

Preparation of 3-pyridyl-4-amino-5-mercapto-s-triazole

A solution of 30 mmole of 5-pyridyl-2-mercapto-1, 3, 4-oxadiazole, 20ml. of water, and 30 mmole 95% hydrazine was refluxed for 4 hours, diluted with 200ml. of cold water, acidified by the drop wise addition of concentrated hydrochloric acid, and filtered. The solid was washed with

a minimum of cold water, and recrystallized from aqueous ethanol¹⁷, yield 62%, m.p. 206 -210 ⁰C.

Synthesis of morpholine-4-ium 4-(N-pyridylcarboxamido)-3-pyridyl-s-triazole-5-thiolate (3b)

30 mmol of 4-(N-pyridylcarboxamido)-3-pyridyl-5-mercapto-s-triazole (3), 30 mmol of Morpholine were added to 20 ml of anhydrous dimethyl sulfoxide and the reaction mixture was stirred vigorously for 12 hr at room temperature under nitrogen atmosphere. Dimethyl sulfoxide was partially recovered under reduced pressure, diluted with 200 ml cold water. Sticky residue obtained which was purified by column chromatography using neutral alumina (25g), using chloroform: methanol (8:2) as an eluent and recrystallised from methanol to obtained desired compound. Yield -32%, m.p. 167⁰C. The preparation of the remaining compounds was carried out as above described procedure. Physicochemical and spectral data of titled compounds (3a-3b & 4a-4b) are shown in Table 1 and 2.

Table: 1. Physicochemical data of Synthesized Compounds (3a-b & 4a-b)

Compound	Molecular Formula	Molecular Weight	%Yield	Melting point °C	R _f *
3a	C ₁₇ H ₁₉ N ₇ OS	369.44	29	281-84	0.17
3b	C ₁₇ H ₁₉ N ₇ O ₂ S	385.13	32	292-95	0.25
4a	C ₁₁ H ₁₆ N ₆ S	264.12	34	248-50	0.24
4b	C ₁₁ H ₁₆ N ₆ OS	280.11	37	262-65	0.19

* Mobile Phase - Chloroform: Methanol (8:2)

Table: 2. Spectral data of Synthesized Compounds (3a-b & 4a-b)

Comp	IR (KBr)cm ⁻¹	¹ H NMR (CDCl ₃)	Elemental analysis (%) Calculated (Found)
3a	3090 (Ar C-H),2490(+NH ₂) 1670(C=ONH),1629(C=N) 1261(C-N)	2.1(2H,m,NH ⁺ ₂),2.8(4H,m,CH ₂ NCH ₂),1.6(4H,m,2CH ₂), 7.5-7.9(4H,dd,3-H,5-H), 8.5-9.06(4H,dd,2-H,6-H) & 11.6 (1H,s,NH)	C, 55.27; H, 5.18; N, 26.54 (C,55.82; H,5.07; N, 26.19)
3b	3090(ArC-H), 2487(+NH ₂) 1670(C=ONH),1629(C=N) 1261(C-N)1114(C-O-C)	2.36(2H,m,NH ⁺ ₂),2.8(4H,m,CH ₂ NCH ₂),3.6(4H,m,CH ₂ OCH ₂),7.6-8.0 (4H,dd,3-H,5-H),8.7-9.0 (4H,dd,2-H,6-H) & 12.6 (1H,s,NH)	C, 52.97; H, 4.97; N, 25.44 (C,53.057;H,4.91; N, 25.18)
4a	3298 (NH ₂), 3045 (Ar C-H), 2484(+NH ₂),1629(C=N), 1270(C-N)	2.14(2H,m,NH ⁺ ₂),2.7(4H,m,CH ₂ NCH ₂),1.6(4H,m,2CH ₂),7.7(2H,d,3-H,5-H),8.7 (2H,d,2-H,6-H)&5.9 (2H,s,NH ₂)	C, 49.98; H, 6.10; N, 31.79 (C,50.12; H,5.98; N, 31.64)
4b	3295(NH ₂),2923(ArC-H), 2510(+NH ₂),1621(C=N), 1255(C-N),1118(C-O-C)	2.3(2H,m,NH ⁺ ₂),2.9(4H,m,CH ₂ NCH ₂),3.8(4H,m,CH ₂ OCH ₂), 7.7(2H,d,3-H,5-H), 8.6(2H,d,2-H,6-H) & 5.9 (2H,s,NH ₂)	C, 47.13; H, 5.75; N, 29.98 (C,47.80; H,5.45; N, 30.14)

RESULTS AND DISCUSSION:

Condensation of 4-(N-pyridylcarboxamido)-3-pyridyl-5-mercapto-s-triazole and 3-pyridyl-4-amino-5-mercapto-s-triazole with secondary amines when heated at 60°C in anhydrous dimethyl sulfoxide results in the formation of corresponding 4-(N-pyridylcarboxamido)-3-pyridyl-s-triazole-5-thiolate and 3-pyridyl-4-amino-s-triazole-5-thiolate as a quaternary salts. The identity of those was confirmed by IR, ¹H NMR and elemental analysis data. From the IR and ¹H NMR spectral data, the possibility of the formation of the corresponding thiosemicarbazide is excluded by the absence of the absorption band at 1520 and 1320 cm⁻¹(ascribed to C=S) in the IR spectra and there are no signals of C(S) NH (at 7.8-8.8 ppm) group in the data of ¹H NMR spectra of compounds synthesized. From the antifungal screening it was found that the compounds showed significant activity compared to standard Griseofulvin. However no conclusive structural activity relationship has emerged from the antifungal screening.

Antibacterial screening revealed that the compounds exhibit weak to moderate activity.

BIOLOGICAL EVALUATION ANTIBACTERIAL ACTIVITY

Studies on the antibacterial activity of the synthesized compounds (**3a-3b** & **4a-4b**) have been screened using cup-plate agar diffusion method¹⁸ against four pathogenic organisms, viz., *Staphylococcus aureus* (G⁺) (ATCC No. 25923), *Klebsiella pneumoniae* (G⁻) (ATCC No. 700603), *Escherichia coli* (G⁻) (ATCC No. 87064) and *Pseudomonas aeruginosa* (G⁻) (ATCC No. 17933) by measuring the inhibition zone in mm at two concentrations (100 and 150 µg/ml).

Streptomycin (100 and 150 µg/ml) was used as a standard and was also screened under similar conditions for comparison. The results of the antibacterial studies are shown in **Table-3**.

Table: 3. Antibacterial Activity of Synthesized Compounds (3a-b & 4a-b)

Compound	<i>Staphylococcus aureus</i>		<i>Klebsiella pnemoniae</i>		<i>Escherichia coli</i>		<i>Pseudomonas aeruginosa</i>	
	100 µg/ml	150 µg/ml	100 µg/ml	150 µg/ml	100 µg/ml	150 µg/ml	100 µg/ml	150 µg/ml
3a	09	12	10	16	08	11	12	15
3b	10	13	12	17	10	12	14	18
4a	08	10	11	15	08	11	13	16
4b	09	11	12	16	10	13	14	18
Streptomycin	15	18	16	21	13	16	18	22

Zone of inhibition expressed in mm

ANTIFUNGAL ACTIVITY

The antifungal activity studies of the novel triazole derivatives(**3a-3b** & **4a-4b**) have been screened using cup-plate agar diffusion method¹⁸ against the four fungi *Aspergillus flavus*, *A.fumigatus*, *penicillium* and

Trichophyton by expressing the zone of inhibition in mm at two concentrations (100 and 150 µg/ml). Griseofulvin (100 and 150 µg/ml) was used as a standard and was also screened under similar conditions for comparison. The results of the antifungal studies are shown in **Table-4**.

Table: 4. Antifungal Activity of Synthesized Compounds (3a-b & 4a-b)

Compound	<i>Aspergillus flavus</i>		<i>Aspergillus fumigates</i>		<i>Penicillium</i>		<i>Trichophyton</i>	
	100 µg/ml	150 µg/ml	100 µg/ml	150 µg/ml	100 µg/ml	150 µg/ml	100 µg/ml	150 µg/ml
3a	10	13	10	14	07	10	10	12
3b	11	14	12	16	08	11	11	14
4a	11	13	11	15	08	11	10	12
4b	11	13	13	17	09	10	11	12
Griseofulvin	12	15	13	17	11	14	13	16

Zone of inhibition expressed in mm

CONCLUSION:

Morpholine and/or pyrrolidine formed quaternary salts when heated with 4-(N-pyridyl carboxamido)-3-pyridyl-5-mercapto-s-triazole (**3**) and/or 3-pyridyl-4-amino-5-mercapto-s-triazole (**4**) at 80°C in DMSO. The quaternary salts showed significant antifungal activity in primary screening against *Aspergillus flavus*, *A. fumigatus* and moderate activity against *penicillium* and *Trichophyton*. Antibacterial screening revealed that the quaternary salts exhibit weak to moderate activity against *Staphylococcus aureus*, *Klebsiella pneumoniae*, *Escherichia coli* and *Pseudomonas aeruginosa*.

ACKNOWLEDGEMENT:

The authors are thankful to SAIF, Punjab University, Chandigarh for NMR spectral analysis and Indian Institute of Technology, Pawai, Mumbai for elemental analysis of the new compounds. We would like to thanks Dr. Devanshu J Patel, Managing Trustee Parul Trust for providing necessary infrastructure and Dr. Rajesh K. S. Principal, Parul Institute of Pharmacy, Limda, Vadodara for offering precious suggestions.

REFERENCES:

1. Tozkoparan B, Küpeli E, Yeşilada E, Ertan M. Preparation of 5-aryl-3-alkylthio-1,2,4-triazoles and corresponding sulfones with Antiinflammatory–analgesic activity. *Bioorg Med Chem*. 2007;15:1808–14.

2. Palaska E, Sahin G, Kelicen P, Durlu NT, Altinok G. Synthesis and anti-inflammatory activity of 1-acylthiosemicarbazides, 1,3,4-oxadiazoles, 1,3,4-thiadiazoles and 1,2,4-triazole-3-thiones. *Farmaco*. 2002;57:101-7.
3. Kumar H, Javed SA, Khan SA, Amir M. 1,3,4-Oxadiazole/thiadiazole and 1,2,4-triazole derivatives of biphenyl-4-yloxy acetic acid: synthesis and preliminary evaluation of biological properties. *Eur J Med Chem*. 2008;43:2688-98.
4. Labanauskas L, Udrenaite E, Gaidelis P, Brukstus A. Synthesis of 5-(2-,3- and 4-methoxyphenyl)-4H-1,2,4-triazole-3-thiol derivatives exhibiting anti-inflammatory activity. *Farmaco*. 2004;59:255-59.
5. Abd Alla MS, Hegab MI, Abo Taleb NA, Hasabelnaby SM, Goudah A. Synthesis and anti-inflammatory evaluation of some condensed [4-(3,4-dimethylphenyl)-1(2H)-oxo-phthalazin-2-yl]acetic acid hydrazide. *Eur J Med Chem*. 2010;45:1267-77.
6. Ulusoy N, Gürsoy A, Otük G. timicrobial activity of some 1,2,4-triazole-3-mercaptoacetic acid derivatives. *Farmaco*. 2001;56:947-52.
7. Eswaran S, Adhikari AV, Shetty NS. Synthesis and antimicrobial activities of novel quinoline derivatives carrying 1,2,4-triazole moiety. *Eur J Med Chem*. 2009;44:4637-47.
8. Bayrak H, Demirbas A, Demirbas N, Karaoglu SA. Synthesis of some new 1,2,4-triazoles starting from isonicotinic acid hydrazide and evaluation of their antimicrobial activities. *Eur J Med Chem*. 2009;44:4362-66.

9. Padmavathi V, Sudhakar Reddy G, Padmaja A, Kondaiah P, Ali-Shazia. Synthesis, antimicrobial and cytotoxic activities of 1,3,4-oxadiazoles, 1,3,4-thiadiazoles and 1,2,4-triazoles. *Eur J Med Chem.* 2009;44:2106-12.
10. Muralikrishna A, Venkatesh BC, Padmavathi V, Padmaja A, Kondaiah P, Krishna NS. Synthesis, antimicrobial and cytotoxic activities of sulfone linked bis heterocycles. *Eur J Med Chem.* 2012;54:605-14.
11. Bayrak H, Demirbas A, Karaoglu SA, Demirbas N. Synthesis of some new 1,2,4-triazoles, their Mannich and Schiff bases and evaluation of their antimicrobial activities. *Eur J Med Chem.* 2009;44:1057-66.
12. Klimesova V, Zahajska L, Waisser K, Kaustova J, Möllmann U. Synthesis and antimycobacterial activity of 1,2,4-triazole 3-benzylsulfanyl derivatives. *Farmaco.* 2004;59:279-88.
13. Mallikarjuna BP, Sastry BS, Suresh Kumar GV, Rajendraprasad Y, Chandrashekar SM, Sathisha K. Synthesis of new 4-isopropylthiazole hydrazide analogs and some derived clubbed triazole, oxadiazole ring systems--a novel class of potential antibacterial, antifungal and antitubercular agents. *Eur J Med Chem.* 2009;44:4739-46.
14. Collin X, Sauleau A, Coulon J. 1,2,4-Triazole mercapto and aminonitriles as potent antifungal agents. *Bioorg Med Chem Lett.* 2003;13:2601-5.
15. Bhat KS, Poojary B, Prasad DJ, Naik P, Holla BS. Synthesis and antitumor activity studies of some new fused 1,2,4-triazole derivatives carrying 2,4-dichloro-5-fluorophenyl moiety. *Eur J Med Chem.* 2009;44:5066-70.
16. Al-Soud YA, Al-Dweri MN, Al-Masoudi NA. Synthesis, antitumor and antiviral properties of some 1,2,4-triazole derivatives. *Farmaco.* 2004;59:775-83.
17. Jack RR, Ned DH. Improved Synthesis of 5-substituted-4-amino-3-mercapto(4H)-1,2,4-triazoles. *J. Hetero Chem.* 1976; 13:925-26.
18. Seelay HW, Van Demark PJ. *Microbes in Action, A laboratory manual in microbiology* 2nd Edⁿ, 1975, page no. 55.