Aesculus parviflora
Bottlebrush Buckeye

Family: Sapindaceae

Orangerie: 20 | Beltline N_S: 25 | Maintenance: 77 | **TOTAL 122**



HABIT



OTHER



OTHER



FLOWER

QUICK FACTS

Plant Type: Shrub

Evergreen or deciduous
Deciduous

Growth & Size
Rate: Not stated
Size: - 6–12 ft tall × 8–15 ft wide.
- Spreads by suckering to form co

Sun

Water

Soil
- Moist, rich, well-drained soil. - Prefers acidic soil

Zone 8a
Suitable for Atlanta Zone 8a. Native to the southeastern United States (AL, GA, FL, SC). Hardy Zones 4–8/9. Prefers moist, rich woodland conditions and shade to part shade. Not stated for specific Atlanta urban performance beyond general southeastern native status and hardiness

PLANTING NOTES

Establishment
- Plant in moist, rich, well-drained soil in shade to part shade. Water until established. Pruning usually unnecessary. - Can be rejuvenated by severe pruning if needed

ORIGIN AND CHARACTERISTICS

Origin
- Native to southeastern United States (Alabama, Georgia, Florida, South Carolina)

Form
- Dense, mounded, multi-stemmed deciduous shrub. Suckering habit

Foliage
- Palmate leaves with 5–7 leaflets.
Dark green in summer; yellow in autumn

Flowers and fruit
- Erect cylindrical panicles (to 12 in) of tubular white flowers with red anthers and pinkish filaments. Mid-summer bloom. - Glossy pear-shaped nuts (buckeyes) in husks; infrequently produced in northern parts of range.
Poisonous if ingested

Bark
- Not stated

HEALTH

Health
- No serious insect or disease problems stated. Foliage relatively disease-free. - Medium severity poison characteristics; fruit poisonous if ingested

Symptoms
None stated