
Sorbus keissleri Rehder

Keissler's Whitebeam, Mao Xu Hua Qiu


Mirror Garden: 1 **TOTAL 1**



HABIT



OTHER



LEAF



OTHER

QUICK FACTS

Plant Type: Deciduous tree or large shrub

Evergreen or deciduous
Potentially suitable for Atlanta Zone 8a (hardiness estimated Zones 6–9 from limited data). Heat tolerance not well documented. Requires good drainage. Rare in cultivation.
Open items for local performanc

Growth & Size
Rate: Not stated
Size: 7–15 m (23–50 ft) tall.
Medium-sized

Sun

Water

Soil
Moist but well-drained soils. Adaptable to various soil types

Zone 8a
Potentially suitable for Atlanta Zone 8a (hardiness estimated Zones 6–9 from limited data). Heat tolerance not well documented
Requires good drainage. Rare in cultivation.
Open items for local performance

PLANTING NOTES

Planting
None stated

ORIGIN AND CHARACTERISTICS

Origin
Native to China (Guangxi, Guizhou, Hubei, Hunan, Jiangxi, Sichuan, Xizang, Yunnan). Dense forests at 1200–1900 m

Form
Tree or shrub with terete branchlets that are initially tomentose, becoming glabrous. Prominently lenticellate

Foliage
Simple leaves, obovate or oblong-obovate, 7–11 × 3.5–6 cm. Pale green beneath, green above. Tomentose when young, becoming glabrate. Crenulate margins. 8–10 pairs of lateral veins

Flowers and fruit
White flowers, 7–9 mm diameter, in dense terminal compound corymbs 4–5 cm across. May–June. Pedicels and rachis tomentose; Brownish green, ovoid or flattened, 9–12 mm diameter.
2- or 3-loculed. August–September

Bark
Not stated

HEALTH

Health
No serious problems stated in available . General Sorbus susceptibilities (fireblight, etc.) may apply.
Not stated for this species

Symptoms
None stated