
Atlanta Botanical Garden - Expansion

Rudbeckia subtomentosa Pursh 'Henry Eilers'

Henry Eilers Sweet Coneflower, Quilled Sweet Coneflower

Fountain Garden: 14 **TOTAL 14**




OTHER



OTHER



OTHER



FLOWER

QUICK FACTS

Plant Type: Herbaceous perennial

Evergreen or deciduous
Well adapted to Atlanta Zone 8a. Heat and humidity tolerant. Long bloom period into fall. Excellent for borders, pollinator gardens, and cutting. Low maintenance; deadhead for extended bloom o leave seed heads for birds. Reliable performer

Growth & Size
Rate: Not stated
Size: 3–6 ft (0.9–1.8 m) tall; 2–3 ft (0.6–0.9 m) wide.
Upright, clumping, sometimes branching

Sun

Water

Soil
Average, well-drained soils. Heat and humidity tolerant. Good air circulation helps prevent mildew

Zone 8a
Well adapted to Atlanta Zone 8a. Heat and humidity tolerant. Long bloom period into fall. Excellent for borders, pollinator gardens, and cutting. Low maintenance; deadhead for extended bloom or leave seed heads for birds. Reliable performer

PLANTING NOTES

Planting
None stated

ORIGIN AND CHARACTERISTICS

Origin
Selection of a species native to the central United States. Found wild in Illinois; introduced by Larry Lowman

Form
Clump-forming with stiff, upright, leafy stems.
Basal and stem foliage

Foliage
Gray-green, toothed leaves (lower leaves often 3-lobed), downy/softly hairy below. Mild sweet/anise or vanilla scent when crushed

Flowers and fruit
Unique quilled (rolled) yellow ray florets forming a starburst or tubular effect around a dome-shaped dark brown center. Clusters on strong stems. Midsummer to frost (July–September). Excellent cut flowers. Attracts butterflies and birds

Bark
Not stated

HEALTH

Health
Generally pest-free and low maintenance. Deer resistant. No serious problems stated. Occasional powdery mildew in poor air circulation

Symptoms
None stated