REMARK profile: TCGA HPV− programme scores

Step	Patients
Xena GDC TCGA−HNSC samples	566
primary tumour (−01) samples	520
patients (one primary tumour sample each)	520
HPV− (cBioPortal SUBTYPE)	415
HPV− with a TCGA−CDR HNSC record	415
of these, not redacted	415
analysed for PFI (time > 0, event known)	415
analysed for OS (time > 0, event known)	415

Endpoint	Patients	Events	Median follow−up (months)	Stage I−II / III / IV / unknown	Site oral / larynx / other
PFI	415	168	31.1	81 / 67 / 225 / 42	277 / 106 / 32
OS	415	191	35.0	81 / 67 / 225 / 42	277 / 106 / 32

Set	Endpoint	Univariable HR (95% CI)	Multivariable HR (95% CI)	PH test p (score)
Buffa 2010 hypoxia	OS	1.22 (1.05−1.42)	1.20 (1.03−1.39)	0.48
Gavish MP1 G2/M	OS	1.20 (1.04−1.40)	1.20 (1.04−1.39)	0.87
Gavish MP2 G1/S	OS	1.14 (0.98−1.32)	1.16 (1.00−1.35)	0.3
Puram 2017 p−EMT	OS	1.07 (0.93−1.24)	1.06 (0.91−1.24)	0.13
Toustrup 2011 hypoxia	OS	1.15 (0.99−1.34)	1.13 (0.98−1.31)	0.83
Buffa 2010 hypoxia	PFI	1.23 (1.05−1.44)	1.22 (1.04−1.43)	0.11
Gavish MP1 G2/M	PFI	1.22 (1.05−1.42)	1.21 (1.04−1.41)	0.49
Gavish MP2 G1/S	PFI	1.18 (1.01−1.37)	1.19 (1.02−1.39)	0.62
Puram 2017 p−EMT	PFI	1.05 (0.90−1.22)	1.02 (0.86−1.19)	0.028
Toustrup 2011 hypoxia	PFI	1.13 (0.97−1.32)	1.11 (0.95−1.30)	0.19

Cut−off rule: median of each score in the analysed patients (pre−specified; no optimal cut−point search).
Missing covariates: stage missing or [Discrepancy] coded as 'unknown'; site and age complete.
Endpoints: TCGA−CDR PFI (primary) and OS (secondary); scores standardised (HR per SD).