

a
A1 All paired patients (n = 22)
518 higher in tumour | 237 higher in normal squamous
HMSD
HPV16−E1
HOXA10
EFEMP2
HPV16−E7
HSD17B2
GPC3
HPV16−E6
MAGEA4
CYP26A1
IL33
SNCA
AC098973.2
FMO2
HNMT
KLK5
SLC38A5
−log10 FDR
log2 fold change
b
A2 HPV+ patients (n = 17)
439 higher in tumour | 146 higher in normal squamous
HSD17B2
HMSD
HPV16−E1
SMARCD3
HPV16−E7
ZBED2
EFEMP2
OVOS2
HPV16−E6
NEFM
EREG
TAF7L
HPV16−E5
NDN
CYP26A1
KLK5
−log10 FDR
log2 fold change
c
A3 HPV− patients (n = 5)
31 higher in tumour | 29 higher in normal squa
PAX9
SYT8
MAGEA4
HOXA10
PHYHD1
NRN1
MGST1
THSD4
HOXC10
AC098973.2
CHL1
SELENOP
PNCK
FMO2
FOXD1
IGFL2−AS1
−log10 FDR
log2 fold change
d
B HPV+ vs HPV− tumour cells (31 vs 39 patients, study−adjusted)
400 higher in HPV+ | 452 higher in HPV−
SMC1B
SYNGR3
ZFR2
MIR9−3HG
KRTDAP
SYCP2
STAG3
MEI1
EREG
MMP2
HTRA1
VCAM1
KLHL35
IGFBP6
MMP1
CDA
SIM2
CXCL14
INHBA
NNMT
CDKN2A
−log10 FDR
log2 fold change
e
DE genes per comparison
right = higher in tumour / HPV+
left = higher in normal squamous / HPV−
A1 all
237
518
A2 HPV+
146
439
A3 HPV−
29
31
A4 HPV+ vs HPV− change
0
1
B HPV+ vs HPV−
452
400
Genes (FDR < 0.05, |log2FC| >= 1)