

$$A = 240, 46 = B \quad \gcd(240, 46)$$

$$\therefore \text{extended_euclid}(240, 46) = ?$$

	Index i	Quotient q_i	Reminder r	x_i x_{i-2} $= x_{i-2} - q_i x_{i-1}$	y_i y_{i-2} $= y_{i-2} - q_i y_{i-1}$
240/46	0			1 x_2	0 y_2
	1			0 x_1	1 y_1
240, 46	2	$240/46 = 5$	10	$x_2 - q_2 x_1$ $1 - 5 \times 0$ $x = 1$	$y_2 - q_2 y_1$ $0 - 5 \times 1$ $y = -5$
46, 10	3	$46/10 = 4$	6	$x_1 - q_3 x_2$ $0 - 1 \times 1$ $= -1$	$y_1 - q_3 y_2$ $1 - 5 \times (-5)$ $= 26$
10, 6	4	$10/6 = 1$	4	$x_2 - q_4 x_3$ $1 - 4 \times (-1)$ $= 5$	$y_2 - q_4 y_3$ $-5 - 4 \times 26$ $= -109$
6, 4	5	$6/4 = 1$	2	$x_3 - q_5 x_4$ $-1 - 5 \times 1$ $= -6$	$y_3 - q_5 y_4$ $26 - 109 \times 1$ $= -83$
4, 2	6	$4/2 = 2$	0	$x_4 - q_6 x_5$ $5 - 18$ $= -13$	$y_4 - q_6 y_5$ $-109 - 83 \times 2$ $= -275$

when
 $r=0$
stops

Pseudocode

extended_euclid (int A, int B, int *X, int *Y)

{

int x1 = 0, x2 = 1;

int y1 = 1, y2 = 0;

int r, q, r1, r2;

$$\frac{x_i}{x} = \frac{x_{i-2}}{x_2} - \frac{q(x_{i-1})}{x_1}$$

for (r2 = A, r1 = B; r1 != 0;

∴ q = r2 / r1;

r = r2 % r1;

∴ x = x2 - (q * x1);

y = y2 - (y1 * q);

r2 = r1,
r1 = r,
x2 = x1,
x1 = x,
y2 = y1,
y1 = y)

*x = x2;

*y = y2;

∴ return r2;

@x @y = gcd(a, b)

Modular multiplicative inverse using extend euclid

$\therefore A \times \equiv 1 \pmod{m}$ $\therefore x$ is modular Inverse
of A modulo m

$\therefore Ax + my = 1$ \therefore Hence $x \rightarrow mI$ of a

\therefore suppose we have to find mI of $\textcircled{3}$ modulo $\textcircled{5}$

$\therefore 5x + 3y = 1$ \therefore Hence value of y is our mI (ans)

A n_2	B n_1	Reminder, r r	Quotient Q	y_2 0	y_1 1	y $y = y_2 - Q \times y_1$
5	$\textcircled{3}$	$5 \div 3 =$ $\textcircled{2}$	$5/3 = 1$	0	$\textcircled{1}$	$= 0 - 1 \times 1$
3	2	$3 \div 2 = 1$	$3/2 = 1$	1	-1	$= 1 - (-1) \times 1$ $= 2$
2	1	$2 \div 1 = 0$	$2/1 = 2$	-1	2	$= -1 \div 2 \times 2$ $= -5$
1	0	\times	\times	$\textcircled{2}$	-5	.

\downarrow
 $r_1 = 0$
loop breaks

\downarrow
 value
 of y_2
 is
 our
 ans
 mI

$3x + 5y = 1$ \therefore Here x is our
m of $\frac{3}{A}$ modulo $\frac{5}{B}$

A	B	R	Q	x_2	x_1	X
3	5	$3/5 = 3$	$3/5 = 0$	1	0	$= 1 - 0$ $= 1$
5	3	2	1	0	1	

now it has become the same solution of

$$5x + 3y = 1$$

so better do with

$$5x + 3y = 1$$

and find value y

Chinese Remainder theorem (weak form)

$$x \equiv a_1 \pmod{m_1}$$

$$x \equiv a_2 \pmod{m_2}$$

\vdots

$$x \equiv a_n \pmod{m_n}$$

\therefore Find value of x

Rule:-

$$m_1, m_2, \dots, m_n$$

all pairwise coprime

$$\gcd(m_1, m_2) = \gcd(m_2, m_n)$$

$$= \gcd(m_1, m_n) = \dots$$

$$= 1$$

Proof

Suppose,

$$x \equiv a_1 \pmod{m_1}$$

$$x \equiv a_2 \pmod{m_2}$$

\therefore we know m_1, m_2 pairwise coprime

\therefore from Bezout's Identity,

$$m_1 p + m_2 q = 1 \quad \text{--- (1)}$$

$$(Ax + By = 1)$$

using extended euclid, we can find p, q

\therefore now we can say,

$$x = (a_1 m_2 q + a_2 m_1 p) \pmod{m_1 m_2}$$

ଏହି ଭାଗটি existing ହୁଏ, ଏହାର ଉଦାହରଣ କିଛି (or actually)

ଏହି ହୁଏ (ଅର୍ଥ)

RoCal-D
Calcium 500 mg & Vitamin-D₃ 200 IU

$$x \equiv a_1 \pmod{m_1}$$

$$x \equiv a_2 \pmod{m_2}$$

Reef-D
Calcium (Ester-C) 500 mg & Vitamin-D₃ 250 IU

କିମ୍ବା



$$\begin{aligned}
 x &= a_1 m_2 q + a_2 m_1 p \\
 &= a_1 (1 - m_1 p) + a_2 m_1 p \quad \text{--- From (1)} \\
 &= a_1 - a_1 m_1 p + a_2 m_1 p
 \end{aligned}$$

$$x = a_1 + ((a_2 - a_1)p)m_1$$

$$\therefore x \equiv a_1 \pmod{m_1}$$

same way $x \equiv a_2 \pmod{m_2}$ (Proved)

$$\# \quad x = (a_1 m_2 q + a_2 m_1 p) \pmod{m_1 m_2}$$

why mod with $m_1 m_2$?

→ to find the (smallest solution and unique) among infinite solutions

How to sure solution is unique?

suppose we have two solution of x is (x_1, x_2)

$$x_1 \equiv a \pmod{m_1}$$

$$x_2 \equiv a \pmod{m_1}$$

we can say $\therefore x_1 \equiv x_2 \pmod{m_1}$

$$\therefore x_1 - x_2 \equiv 0 \pmod{m_1} \quad | \quad (x_1 - x_2) \text{ is divisible by } m_1$$

$$\text{same} \quad x_1 - x_2 \equiv 0 \pmod{m_2} \quad | \quad \parallel \quad \parallel \quad \parallel \quad \parallel \quad m_2$$

if $(x_1 - x_2)$ is divisibly by both m_1 and m_2
and m_1, m_2 coprime

that means $x_1 - x_2$ is also divisible by $m_1 \times m_2$

$$\therefore x_1 - x_2 \equiv 0 \pmod{m_1 m_2}$$

$$\therefore x_1 \equiv x_2 \pmod{m_1 m_2}$$

hence there exist only one solution

$\therefore x$ is unique to modulo $m_1 m_2$

(proved)

PseudoCode

pair<int, int> chineseReminderTheorem
(vector<int> A, vector<int> m)

<
if (A.size() != m.size())
{ Invalid Input }

∴ int n = A.size();

∴ int a1 = A[0] ✓

∴ int m1 = m[0] ✓

for (int i = 1; i < n; i++) <

int a2 = A[i], m2 = m[i]; ✓

∴ int p, q;

extendedEuclid(m1, m2, &p, &q);

∴ int x = (a1 * q * m2 + a2 * p * m1) % m1 * m2

∴ a1 = x ✓

∴ m1 = m1 * m2 ✓

}

∴ if (a1 < 0) a1 = a1 + m1

∴ return { a1, m1 }

Chinese Remainder Theorem - Strong

$$x \equiv a_1 \pmod{m_1}$$

$$x \equiv a_2 \pmod{m_2}$$

$$\vdots$$
$$x \equiv a_n \pmod{m_n}$$

m_1, m_2, \dots, m_n
not coprime

(*) How to sure there a solution exists?

$$\therefore \gcd(m_1, m_2) = g$$

we know,

$$x \equiv a_1 \pmod{m_1}$$

$$x - a_1 \equiv 0 \pmod{m_1} \quad | \quad m_1 \text{ divide } x - a_1$$

divisor
of m_1
like (g) also divides $x - a_1$ (*)

$$\therefore x - a_1 \equiv 0 \pmod{g}$$

same $x - a_2 \equiv 0 \pmod{g}$

$$\therefore x - a_1 \equiv x - a_2 \pmod{g}$$

$$\therefore a_1 \equiv a_2 \pmod{g}$$

if ~~sta~~ satisfy this
condition then there exist
a solution of x



RoCal-D
Calcium 500 mg & Vitamin D₃ 200 IU

Reef-D
Calcium 500 mg & Vitamin D₃ 200 IU

Now solution

\therefore we know

$$\text{GCD}(m_1, m_2) = g$$

Bézout's Identity

$$\therefore m_1 p + m_2 q = g$$

$$\therefore \frac{m_1}{g} p + \frac{m_2}{g} q = 1 \quad \text{--- (1)}$$

using ext-gcd we can find p, q

$$\therefore X = a_1 \frac{m_2}{g} q + a_2 \frac{m_1}{g} p$$

now same stykproof

$$\therefore x = a_1 \left(1 - \frac{m_1}{g} p\right) + a_2 \frac{m_1}{g} p \quad \text{--- (1)}$$

$$= a_1 - a_1 \frac{m_1}{g} p + a_2 \frac{m_1}{g} p$$

$$= a_1 + \frac{(a_2 - a_1) m_1}{g} p$$

$$\therefore x \equiv a_1 \pmod{m_1}$$

(same)

$$\therefore x \equiv a_2 \pmod{m_2}$$

proved

$$X = \left(a_1 \frac{m_2}{g} a + a_2 \frac{m_1}{g} p \right) \bmod \text{lcm}(m_1, m_2)$$

Pseudocode

same like weak form

just some changes

same

$\therefore \text{int } g = \text{gcd}(m_1, m_2)$

$\therefore \text{if } (a_1 \% g \neq a_2 \% g) \rightarrow \text{no solution}$

$\text{int } p, q$

$\text{ext-gcd}(m_1/g, m_2/g, \&p, \&q)$

$\therefore \text{int } \text{mods} = m_1 / g * m_2$

$$X = \left(a_1 q \frac{m_2}{g} + a_2 p \frac{m_1}{g} \right) \bmod \text{mods}$$

$a_1 = X$

$\text{if } (a_1 < 0) \ a_1 += \text{mod}$

$m_1 = \text{mod}$