

Abstract Measure Theory

Uncultured Tramp

September 13, 2022

Contents

1	Measure Algebras	1
1.1	Subject	1
1.1.1	Definition and Basic Property	1
1.1.2	Measure Algebras Generated by Measure Spaces	4
1.1.3	Stone Representation Theorem	5
1.1.4	Ideals	6
1.1.5	Measure Properties	8
1.1.6	Connections with other Boolean Properties	11
1.1.7	Subspace Measures and Indefinite Integrals	13
1.1.8	Simple Products	14
1.1.9	Strictly Localizable Spaces	16
1.1.10	Subalgebras	17
1.1.11	Localization	19
1.1.12	Stone Spaces	20
1.1.13	Purely Infinite Elements	22
1.2	Topology	24
1.2.1	Subject	24
1.3	Category	25
1.4	Radon-Nikodym Parallels	25
2	Maharam's Theory	25
3	Abstract Ergodic Theory	25
4	Measurable Algebras	25

1 Measure Algebras

1.1 Subject

1.1.1 Definition and Basic Property

MeasureAlgebra :: ? $\sum A : \sigma\text{-DedekindComplete} . A \rightarrow \mathbb{R}_+$

$(A, \mu) : \text{MeasureAlgebra} \iff \forall a \in A . \mu(a) = 0 \iff a = 0 \ \&$

$$\& \forall a : \text{PairwiseDisjointElements}(\mathbb{N}, A) . \mu \left(\bigvee_{n=1}^{\infty} a_n \right) = \sum_{n=1}^{\infty} \mu(a_n)$$

MeasureMonotonicity :: $\forall (A, \mu) : \text{MeasureAlgebra} . \forall a, b \in A . a \leq b \Rightarrow \mu(a) \leq \mu(b)$

Proof =

Write $\mu(b) = \mu(a) + \mu(b \setminus a) \geq \mu(a)$.

□

MeasureStrictMonotonicity :: $\forall (A, \mu) : \text{MeasureAlgebra} . \forall a, b \in A . a > b \Rightarrow \mu(a) > \mu(b)$

Proof =

Definition of measure algebra implies that $\mu(b \setminus a) > 0$.

Write $\mu(b) = \mu(a) + \mu(b \setminus a) > \mu(a)$.

□

MinkovskyIneq :: $\forall (A, \mu) : \text{MeasureAlgebra} . \forall a, b \in A . \mu(a \vee b) \leq \mu(a) + \mu(b)$

Proof =

Write $\mu(a) + \mu(b) = \mu(a \setminus ab) + \mu(ab) + \mu(b \setminus ab) + \mu(ab) \geq \mu(a \setminus ab) + \mu(ab) + \mu(b \setminus ab) = \mu(a \vee b)$.

□

MonotonicSupremumAsLimit :: $\forall (A, \mu) : \text{MeasureAlgebra} . \forall a : \mathbb{N} \uparrow A . \mu \left(\bigvee_{n=1}^{\infty} a_n \right) = \lim_{n \rightarrow \infty} \mu(a_n)$

Proof =

Construct disjoint sequence $b_n = a_n \setminus \bigvee_{k=1}^{n-1} a_k$.

Then by construction $\mu \left(\bigvee_{n=1}^{\infty} a_n \right) = \mu \left(\bigvee_{n=1}^{\infty} b_n \right) = \sum_{n=1}^{\infty} \mu(b_n) = \lim_{n \rightarrow \infty} \sum_{k=1}^n \mu(b_k) = \lim_{n \rightarrow \infty} \mu \left(\bigvee_{k=1}^n b_k \right) = \lim_{n \rightarrow \infty} \mu(a_n)$.

□

SupremumIneq :: $\forall (A, \mu) : \text{MeasureAlgebra} . \forall a : \mathbb{N} \rightarrow A . \mu \left(\bigvee_{n=1}^{\infty} a_n \right) \leq \sum_{n=1}^{\infty} \mu(a_n)$

Proof =

Construct increasing sequence $b_n = \bigvee_{k=1}^n a_n$.

Then by construction $\mu \left(\bigvee_{n=1}^{\infty} a_n \right) = \mu \left(\bigvee_{n=1}^{\infty} b_n \right) = \lim_{n \rightarrow \infty} \mu(b_n) = \lim_{n \rightarrow \infty} \mu \left(\bigvee_{k=1}^n a_k \right) \leq \lim_{n \rightarrow \infty} \sum_{k=1}^n \mu(a_k) = \sum_{n=1}^{\infty} \mu(a_n)$.

□

MonotonicInfimumAsLimit ::

$:: \forall (A, \mu) : \text{MeasureAlgebra} . \forall a : \mathbb{N} \downarrow A . \forall \aleph : \inf_{n \in \mathbb{N}} \mu(a_n) < \infty . \mu \left(\bigwedge_{n=1}^{\infty} a_n \right) = \lim_{n \rightarrow \infty} \mu(a_n)$

Proof =

Without loss of generality assume that $\mu(a_1) < \infty$.

Then construe the increasing sequence $b_n = a_1 \setminus a_n$.

Then $\mu(a_1) - \mu \left(\bigwedge_{n=1}^{\infty} a_n \right) = \mu \left(a_1 \setminus \bigwedge_{n=1}^{\infty} a_n \right) = \mu \left(\bigvee_{n=1}^{\infty} a_1 \setminus a_n \right) = \mu \left(\bigvee_{n=1}^{\infty} b_n \right) = \lim_{n \rightarrow \infty} \mu(b_n) =$
 $= \lim_{n \rightarrow \infty} \mu(a_1 \setminus a_n) = \lim_{n \rightarrow \infty} \mu(a_1) - \mu(a_n) = \mu(a_1) - \lim_{n \rightarrow \infty} \mu(a_n)$.

So basic algebraic manipulations $\mu \left(\bigwedge_{n=1}^{\infty} a_n \right) = \lim_{n \rightarrow \infty} \mu(a_n)$.

□

SupremumExistence ::

$:: \forall (A, \mu) : \text{MeasureAlgebra} . \forall C : \text{UpwardsDirected}(A) . \forall \aleph : \sup_{c \in C} \mu(c) < \infty . \exists a \in A : a = \sup C$

Proof =

1 Assume $\gamma = \sup_{c \in C} \mu(c)$.

2 Then there exists a sequence of $a : \mathbb{N} \rightarrow C$ such that $\mu(a_n) \geq \gamma - 2^{-n}$.

3 As C is upwards closed, it is possible to find $c : \mathbb{N} \rightarrow C$ such that $c_{n+1} \geq a_n \vee c_n$.

4 Then c is monotonic-nondecreasing and so it has $\mu \left(\bigvee_{n=1}^{\infty} c_n \right) = \lim_{n \rightarrow \infty} \mu(c_n) = \gamma$.

4.1 Note that $\gamma \geq \mu(c_n) \geq \gamma - 2^{-n}$.

5 let $d = \bigvee_{n=1}^{\infty} c_n$.

6 $d \geq f$ for every $f \in C$.

6.1 Assume this is false.

6.2 Then $f \setminus d \neq 0$ and so $\alpha = \mu(f \setminus d) > 0$.

6.3 Then there exists n such that $\gamma - \mu(c_n) < \alpha$.

6.4 As C is upwards directed there is $g \in C$ such that $g \geq f \vee c_n$.

6.5 But $\mu(g) \geq \mu(f \vee c_n) = \mu(c_n) + \mu(f \setminus c_n) \geq \mu(c_n) + \mu(f \setminus d) > \gamma$ which is impossible.

7 If there is another f with the property (6), then $d = \bigvee_{n=1}^{\infty} c_n \leq f$ as $c_n \leq f$ for each $n \in \mathbb{N}$.

□

UpperContinuity ::

$$:: \forall (A, \mu) : \text{MeasureAlgebra} . \forall C : \text{UpwardsDirected}(A) . \forall \mathbb{N} : \exists a \in A : a = \sup C . \sup_{c \in C} \mu(c) = \mu(\sup C)$$

Proof =

Case $\sup_{c \in C} \mu(c) = \infty$ is trivial.

Finite case follows from the cconstruction in the previous theorem.

□

DisjointUpperContinuity ::

$$:: \forall (A, \mu) : \text{MeasureAlgebra} . \forall C : \text{PairwiseDisjointElements}(A) . \forall \mathbb{N} : \exists a \in A : a = \sup C . \\ . \mu(\sup C) = \sum_{c \in C} \mu(c)$$

Proof =

Construct a new set $D = \left\{ \bigvee_{n=1}^{\infty} c_k \mid c : \mathbb{N} \rightarrow C \right\}$.

Then D is upwards directed and $\sup C = \sup D$.

$$\text{But this is evedent that } \mu(\sup D) = \sup_{d \in D} \mu(d) = \sup_{c: \mathbb{N} \rightarrow C} \mu\left(\bigvee_{n=1}^{\infty} c_n\right) = \sup_{n \in \mathbb{N}, c: \{1, \dots, n\} \rightarrow C} \sum_{k=1}^n \mu(c_k) = \sum_{c \in C} \mu(c).$$

□

InfimumExistance ::

$$:: \forall (A, \mu) : \text{MeasureAlgebra} . \forall C : \text{DownwaedDirected}(A) . \forall \mathbb{N} : \inf_{c \in C} \mu(c) < \infty . \exists a \in A : a = \inf C$$

Proof =

1 There exists some $a \in C$ such that $\mu(a) < \infty$.

2 Construct another set $D = a \setminus C$.

3 Then D is upwards directed and $\sup_{d \in D} \mu(d) \leq \mu(a) < \infty$.

4 So there is $d = \sup d$.

5 Define $f = a \setminus d$.

6 $f \leq c$ for any $c \in C$ as $a \setminus f \geq a \setminus c$.

7 if some g has property (6) then $a \setminus g \geq d$ and so $g \leq f$.

□

LowerContinuity ::

$$:: \forall (A, \mu) : \text{MeasureAlgebra} . \forall C : \text{DownwardsDirected}(A) . \forall \mathbb{N} : \exists a \in A : a = \inf C . \\ . \forall \sqsupset : \inf_{c \in C} \mu(c) < \infty . \inf_{c \in C} \mu(c) = \mu(\inf C)$$

Proof =

Use the construction in the previous theorem.

□

1.1.2 Measure Algebras Generated by Measure Spaces

measureAlgebra :: MEAS \rightarrow MeasureAlgebra

$$\text{measureAlgebra}(X, \Sigma, \mu) = (A_\mu, \bar{\mu}) := \left(\frac{\Sigma}{\Sigma \cap \mathcal{N}_\mu}, [E] \mapsto \mu(E) \right)$$

This is obviously well defined as $[E] = [F]$ iff $\mu(E \triangle F) = 0$.

canonicalProjection :: $\forall (X, \Sigma, \mu) \in \text{MEAS} . \sigma\text{-BOOL}(\Sigma, A_\mu)$

$$\text{canonicalProjection}(E) = \pi_\mu(E) := [E]$$

1 The algebraic properities are obvious as $\Sigma \cap \mathcal{N}_\mu$ is an ideal.

2 In order to prove sigma-continuity assume $E : \mathbb{N} \rightarrow \Sigma$.

2.1 Let $Z : \mathbb{N} \rightarrow \Sigma \cap \mathcal{N}_\mu$.

$$2.2 \text{ Then } F_Z = \bigvee_{n=1}^{\infty} (E_n \triangle Z_n) = \left(\bigvee_{n=1}^{\infty} E_n \right) \triangle \left(\bigvee_{n=1}^{\infty} Z_n \right).$$

$$2.3 \text{ Note that } \mu \left(\bigvee_{n=1}^{\infty} Z_n \right) \leq \sum_{n=1}^{\infty} \mu(Z_n) = 0.$$

$$2.4 \text{ So } \bigvee_{n=1}^{\infty} Z_n \in \Sigma \cap \mathcal{N}_\mu \text{ as } \mu \geq 0.$$

$$2.5 \text{ Thus } [F_Z] = \left[\bigcap_{n=1}^{\infty} E_n \right] \text{ for any selection of } Z.$$

$$2.6 \text{ This means that } \pi_\mu \left(\bigcap_{n=1}^{\infty} E_n \right) = \bigvee_{n=1}^{\infty} \pi_\mu(E_n).$$

□

MeasureAlgebraMonotonicity :: $\forall (X, \Sigma, \mu) \in \text{MEAS} . \forall T \subset_\sigma \Sigma . \pi_\mu(T) \subset_\sigma A_\mu$

Proof =

1 Clearly $B = \pi_\mu(T) \subset A_\mu$.

2 Also as T is σ -algebra and $\pi - \mu$ is a σ -continuous homomorphism B is again.

□

MeasureAlgebraInverseMonotonicity :: $\forall (X, \Sigma, \mu) \in \text{MEAS} . \forall B \subset_\sigma A_\mu . \pi_\mu^{-1}(B) \subset_\sigma \Sigma$

Proof =

1 Clearly $T = \pi_\mu^{-1}(B) \subset \Sigma$.

2 Assume F is a set constructed by applying σ -algebra operations to setes $E_1, E_2, \dots \in T$.

3 Then $\pi_\mu(F)$ can be constructed by applying same operations to $\pi(E_1), \pi(E_2), \dots$

4 This implies that $\pi_\mu(F) \in B$ and reciprorary $F \in T$.

5 Thus T is a σ -algebra.

□

1.1.3 Stone Representation Theorem

StoneRepresentationTheorem :: $\forall (A, \mu) : \text{MeasureAlgebra} . \exists (X, \Sigma, \nu) \in \text{MEAS} . (A, \mu) = (A_\nu, \bar{\nu})$

Proof =

1 By Loomis-Sikorski representation there exists a set X with a sigma-algebra Σ and

sigma-ideal I such that $\frac{\Sigma}{I} \cong_{\text{BOOL}} A$.

2 Then there is a canonical projection $\pi_I \in \text{BOOL}(\Sigma, A)$.

3 Define $\nu = \pi_I \mu$.

4 ν is measure on Σ .

4.1 $\nu(\emptyset) = \mu(0) = 0$.

4.2 Assume E is a disjoint sequence in Σ .

4.3 Then $\pi_I(E_n)\pi_I(E_m) = \pi_I(E_n \cap E_m) = \pi_I(\emptyset) = 0$, so $\pi_I(E)$ is disjoint in A .

4.4 Thus, $\nu\left(\bigcup_{n=1}^{\infty} E_n\right) = \pi_I \mu\left(\bigcup_{n=1}^{\infty} E_n\right) = \mu\left(\bigvee_{n=1}^{\infty} \pi_I(E_n)\right) = \sum_{n=1}^{\infty} \pi_I \mu(E_n) = \sum_{n=1}^{\infty} \nu(E_n)$.

5 Also by consytuction $\mathcal{N}_\nu \cap \Sigma = I$, so $(A, \mu) = (A_\nu, \bar{\nu})$.

□

spaceOfStone :: $\text{MeasureAlgebra} \rightarrow \text{MEAS}$

SpaceOfStone $(A, \mu) = (Z_A, \dot{\Sigma}_\mu, \dot{\mu}) := \text{StoneRepresentationTheorem}(A, \mu)$

1.1.4 Ideals

PrincipleIdealRestriction :: $\forall (A, \mu) : \text{MeasureAlgebra} . \forall a \in A . \text{MeasureAlgebra}((a), \mu|_{(a)})$

Proof =

This is obvious.

□

measureQuotient ::

$:: \forall (A, \mu) : \text{MeasureAlgebra} . \forall I : \text{Ideal}(A) . \forall [a] \in \frac{A}{I} . \exists \gamma \in \mathbb{R}_{++}^{\infty} . \gamma = \min\{\mu(b) | b \in A, \pi_I(b) = [a]\}$

Proof =

1 $\gamma = \inf\{\mu(b) | b \in A, \pi_I(b) = [a]\}$ exists as a set is bounded by below by 0.

2 If $\gamma = \infty$ then the result is obvious.

3 Otherwise there is a decreasing sequence $b : \mathbb{N} \rightarrow A$ such that $\pi_I(b_n) = [a]$ for any n and $\lim_{n \rightarrow \infty} \mu(b_n) = \gamma$.

4 Then $c = \bigwedge_{n=1}^{\infty} b_n$ is such that $\mu(c) = \gamma$ and $\pi_I(c) = a$.

4.1 Clearly $\pi_I\left(\bigwedge_{n=1}^{\infty} b_n\right) = \bigwedge_{n=1}^{\infty} \pi_I(b_n) = \bigwedge_{n=1}^{\infty} [a] = [a]$.

5 So the infimum is attained.

□

measureQuotient :: $\prod (A, \mu) : \text{MeasureAlgebra} . \prod I : \text{Ideal}(A) . \frac{A}{I} \rightarrow \mathbb{R}_{++}$

measureQuotient (a) = $\mu_I(a) := \min\{\mu(b) | b \in A, \pi_I(b) = a\}$

finiteElementsIdeal :: $\prod (A, \mu) : \text{MeasureAlgebra} . \text{Ideal}(A)$

finiteElementsIdeal () = $A^f := \{a \in A | \mu(a) < \infty\}$

MeasureIdealQuotient :: $\forall (A, \mu) : \text{MeasureAlgebra} . \forall I : \text{Ideal}(A) . \text{MeasureAlgebra} \left(\frac{A}{I}, \mu_I \right)$

Proof =

1 Clearly $\mu_I(0) = 0$.

2 Assume that $[a] \neq 0$.

2.1 Then there exists $b \in A$ such that $\pi_I(a) = [a]$ and $\mu(b) = \mu_I[a]$.

2.2 As $[a] \neq 0$, then $b \neq 0$, and henceforth $\mu(b) \neq 0$.

2.3 Thus, $\mu_I[a] \neq 0$.

3 Assume $[a] : \mathbb{N} \rightarrow \frac{A}{I}$ is disjoint.

3.1 It is possible to select representatives b_n for each $[a_n]$ such that $\mu(b_n) = \mu_I[a_n]$.

3.2 Then $b_n b_m \in I$ if $n \neq m$.

3.3 Construct a new sequence $c_n = b_n + \sum_{k=1}^{n-1} b_n b_k$ is a disjoint representative sequence for $[a_n]$.

3.3.1 In fact $c = b$.

3.4 $\bigvee_{n=1}^{\infty} c_n$ is the minimal representative of $\bigvee_{n=1}^{\infty} [a_n]$.

3.4.1 Assume d is a representative for $\bigvee_{n=1}^{\infty} a_n$.

3.4.2 If $\mu(d) < \mu \left(\bigvee_{n=1}^{\infty} c_n \right)$ then we may construct $c_n \wedge d$ which is smaller then c .

3.4.3 But this is a contradiction.

3.5 So $\mu_I \left(\bigvee_{n=1}^{\infty} [a_n] \right) = \mu \left(\bigvee_{n=1}^{\infty} c_n \right) = \sum_{n=1}^{\infty} \mu(c_n) = \sum_{n=1}^{\infty} \mu_I[a_n]$.

□

1.1.5 Measure Properties

ProbabilityAlgebra :: ?MeasureAlgebra

$(A, \pi) : \text{ProbabilityAlgebra} \iff \pi(e) = 1$

FiniteMeasureAlgebra :: ?MeasureAlgebra

$(A, \mu) : \text{FiniteMeasureAlgebra} \iff \mu(e) < \infty$

σ -FiniteMeasureAlgebra :: ?MeasureAlgebra

$(A, \mu) : \sigma\text{-FiniteMeasureAlgebra} \iff \exists a : \mathbb{N} \rightarrow A . \forall n \in \mathbb{N} . \mu(a_n) < \infty \ \& \ \bigvee_{n=1}^{\infty} a_n = e$

SemifiniteMeasureAlgebra :: ?MeasureAlgebra

$(A, \mu) : \text{SemifiniteMeasureAlgebra} \iff \forall a \in A . \mu(a) = \infty \Rightarrow \exists b \in A . b < a \ \& \ 0 < \mu(b) < \infty$

LocalizableMeasureAlgebra := OrderDedekindComplete & SemifiniteMeasureAlgebra : Type;

ProbabilityConstruction :: $\forall (X, \Sigma, \mu) \in \text{MEAS} . \text{Probability}(X, \Sigma, \mu) \iff \text{ProbabilityAlgebra}(A_\mu, \bar{\mu})$

Proof =

This is obvious.

□

FiniteConstruction :: $\forall (X, \Sigma, \mu) \in \text{MEAS} . \text{Finite}(X, \Sigma, \mu) \iff \text{FiniteMeasureAlgebra}(A_\mu, \bar{\mu})$

Proof =

This is obvious.

□

SigmaFiniteConstruction :: $\forall (X, \Sigma, \mu) \in \text{MEAS} . \sigma\text{-Finite}(X, \Sigma, \mu) \iff \sigma\text{-FiniteMeasureAlgebra}(A_\mu, \bar{\mu})$

Proof =

This is obvious.

□

SemifiniteConstruction ::

$\forall (X, \Sigma, \mu) \in \text{MEAS} . \text{Semifinite}(X, \Sigma, \mu) \iff \text{SemifiniteMeasureAlgebra}(A_\mu, \bar{\mu})$

Proof =

This is obvious.

□

LocalizableConstruction ::

$\forall (X, \Sigma, \mu) \in \text{MEAS} . \text{Localizable}(X, \Sigma, \mu) \iff \text{LocalizableMeasureAlgebra}(A_\mu, \bar{\mu})$

Proof =

This is obvious.

□

AtomInConstruction ::

$$:: \forall (X, \Sigma, \mu) \in \text{MEAS} . \forall E \in \Sigma . E \in \text{Atom}(X, \Sigma, \mu) \iff [E] \in \text{Atom}(A_\mu, \bar{\mu})$$

Proof =

This is obvious.

□

AtomlessConstruction ::

$$:: \forall (X, \Sigma, \mu) \in \text{MEAS} . \forall E \in \Sigma . E \in \text{Atomless}(X, \Sigma, \mu) \iff [E] \in \text{Atomless}(A_\mu, \bar{\mu})$$

Proof =

This is obvious.

□

PurelyAtomicConstruction ::

$$:: \forall (X, \Sigma, \mu) \in \text{MEAS} . \forall E \in \Sigma . E \in \text{PurelyAtomic}(X, \Sigma, \mu) \iff [E] \in \text{PurelyAtomic}(A_\mu, \bar{\mu})$$

Proof =

This is obvious.

□

FinitenessPropertiesIerarchy ::

$$\begin{aligned} &:: \forall (A, \mu) : \text{MeasureAlgebra} . \text{PobabilityAlgebra}(A, \mu) \Rightarrow \text{FiniteMeasureAlgebra}(A, \mu) \Rightarrow \\ &\Rightarrow \sigma\text{-FiniteMeasureAlgebra}(A, \mu) \Rightarrow \text{LocalizableMeasureAlgebra}(A, \mu) \Rightarrow \text{Semifinite}(A, \mu) \end{aligned}$$

Proof =

1 Most implications here are obvious expect the one deriving Localizability from σ -finiteness.

2 So assume that (A, μ) is σ -finite .

2.1 Then the corresponding Stone space $(ZA, \Sigma_\mu, \bar{\mu})$ is σ -finite.

2.2 But then $(ZA, \Sigma_\mu, \bar{\mu})$ is localizable .

2.3 So (A, μ) is also localizable.

□

MeasureAlgebraOfCompletion :: $\forall (X, \Sigma, \mu) \in \text{MEAS} . A_\mu \cong_{\text{BOOL}} A_{\hat{\mu}}$

Proof =

This is basically follows from definitions.

□

MeasureAlgebraOfLocallyDeterminedCompletion ::

$$\begin{aligned} &:: \forall (X, \Sigma, \mu) \in \text{MEAS} . \exists A_\mu \xrightarrow{\phi} A_{\bar{\mu}} : \text{BOOL} . \forall a \in A_{\bar{\mu}} . \hat{\mu}(a) < \infty \Rightarrow \exists b \in A_\mu . \phi(b) = a \ \& \\ &\ \& \forall b \in A_\mu . \hat{\mu}(b) < \infty \Rightarrow \hat{\mu}(\phi(b)) = \hat{\mu}(b) \end{aligned}$$

Proof =

...

□

localDeterminationMorphism :: $\prod (X, \Sigma, \mu) \in \text{MEAS} . \text{BOOL}(A_\mu, A_{\bar{\mu}})$

localDeterminationMorphism () = $\phi_\mu := \text{MeasureAlgebraOfLocallyDeterminedCompletion}$

localDeterminationMorhismInjectivity ::

$$:: \forall (X, \Sigma, \mu) \in \text{MEAS} . \text{Semifinite}(X, \Sigma, \mu) \iff \text{Injective}(A_\mu, A_{\bar{\mu}}, \phi_\mu)$$

Proof =

...

□

localDeterminationMorhismBijectivity ::

$$:: \forall (X, \Sigma, \mu) \in \text{MEAS} . \text{Localizable}(X, \Sigma, \mu) \iff \text{Bijective}(A_\mu, A_{\bar{\mu}}, \phi_\mu)$$

Proof =

...

□

SemifinitenessCriterion :: $\forall (A, \mu) : \text{MeasureAlgebra} .$

$$. \text{SemifiniteMeasureAlgebra}(A, \mu) \iff \exists P : \text{PartitionOfUnity}(A) . \forall p \in P . \mu(p) < \infty$$

Proof =

1 (\Rightarrow) assume first that (A, μ) is semifinite.

1.1 Then A^f is order dense in A .

1.2 By order density theorem there is a desired partition of unity.

2 (\Leftarrow) Let P be the partition of unity.

2.1 Assume $a \in A$ is such that $\mu(a) = \infty$.

2.2 Then there exists $p \in P$ such that $pa \neq 0$.

2.3 Note that this means that $\mu(pa) > 0$.

2.4 Also it is clear that $\mu(pa) \leq \mu(p) < \infty$.

□

SemifiniteneSupElementExpression ::

$$:: \forall (A, \mu) : \text{SemifiniteMeasureAlgebra}(A, \mu) . \forall a \in A . a = \bigvee \{b \in A : b \leq a, \mu(b) < \infty\}$$

Proof =

This follows from the previous theorem.

□

SemifiniteneSupMeasureComputation ::

$$:: \forall (A, \mu) : \text{SemifiniteMeasureAlgebra}(A, \mu) . \forall a \in A . \mu(a) = \bigvee \{\mu(b) \in A : b \leq a, \mu(b) < \infty\}$$

Proof =

This follows from the previous theorem.

□

1.1.6 Connections with other Boolean Properties

SemifiniteIsWeaklyDistributive ::

:: $\forall (A, \mu) : \text{SemifiniteMeasureAlgebra}(A, \mu) . (\sigma, \infty)\text{-WeaklyDistributive}(A, \mu)$

Proof =

- 1 Assume $X : \mathbb{N} \rightarrow 2^A$ is a sequence of downwards selected sets with $\inf X_n = 0$ for every $n \in \mathbb{N}$.
 - 2 Let $C = \{a \in A : \forall n \in \mathbb{N} . \exists x \in X_n . a \geq x\}$.
 - 3 Assume $d \in A$ is such that $d \neq 0$.
 - 4 Then there is an element $d' \leq d$ such that $0 < \mu(d') < \mu(d)$.
 - 5 $\inf_{x \in X} d'x = 0$ for each $n \in \mathbb{N}$.
 - 6 Select a sequence $x : \prod_{n=1}^{\infty} X_n$ such that $\mu(d'x_n) \leq 2^{-n-2}\mu(d')$.
 - 7 Define $c = \sup_{n=1}^{\infty} a_n \in C$.
 - 8 Then $\mu(d'c) \leq \sum_{n=0}^{\infty} \mu(cx_n) < \mu(d')$.
 - 9 This means that $d \not\leq c$.
 - 10 And as d was arbitrary $\inf C = 0$.
-

SemifiniteIffCCC :: $\forall (A, \mu) : \text{SemifiniteMeasureAlgebra}(A, \mu) .$

$. \sigma\text{-FiniteMeasureAlgebra}(A, \mu) \iff \text{WithCountableChainCondition}(A)$

Proof =

- 1 (\Leftarrow) assume that A has ccc.
 - 1.1 Then there is a partition of unity P in A consisting of finite elements as A is semifinite.
 - 1.2 But as A has ccc P must be atmost countable.
 - 1.3 This proves that A is σ -finite.
 - 2 (\Rightarrow) assume that (A, μ) is σ -finite .
 - 2.1 Then there exists a countable partition of unity P of A with finite elements.
 - 2.2 If A is not ccc, then there exists an uncountable refinement Q of A with finite elements.
 - 2.3 Then by pigeonhole principle there exists $p \in P$
 - such that set $Q' = \{q \in Q : q \subset p\}$ such that Q' is uncountable.
 - 2.4 as for $\mu(q) > 0$ for any $q \in Q'$ by pigeonhole principle there exists some $n \in \mathbb{Z}$
 - such that there are an infinite number of $q \in Q'$ with $\mu(q) \in [2^{-n-1}, 2^{-n}]$.
 - 2.5 So $\mu(p) \geq \sum_{q \in Q'} \mu(q) = \infty$, but this is a contradiction.
-

SemifiniteIffProbabilityRenormalizationExists ::

$$\begin{aligned} &:: \forall (A, \mu) : \text{SemifiniteMeasureAlgebra}(A, \mu) . A \neq \{0\} \Rightarrow \\ &\Rightarrow \exists \pi : A \rightarrow \mathbb{R}_+^\infty . \text{ProbabilityAlgebra}(A, \pi) \end{aligned}$$

Proof =

- 1 Corresponding Stone space is σ -finite.
 - 2 So there exists a proper renormalization of $\bar{\mu}$ to a probability π with the same sets of measure zero.
 - 3 Then the measure algebra of (ZA, π) is a probability algebra and $A_\pi \cong_{\text{BOOL}} A$.
-

1.1.7 Subspace Measures and Indefinite Integrals

MeasurableEnvelopePrincipleIdealIsomorphism ::

$$:: \forall (X, \Sigma, \mu) \in \text{MEAS} . \forall Y \subset X . \forall E : \text{MeasurableEnvelope}(X, \Sigma, \mu, Y) . (A_{\mu|Y}, \widehat{\mu|Y}) \cong_{\text{MA}} \left(([E]), \hat{\mu}|_{([E])} \right)$$

Proof =

This result is technically convoluted but actually is pretty intuitive.

□

PrincipleIdealIsomorphism ::

$$:: \forall (X, \Sigma, \mu) \in \text{MEAS} . \forall E \in \Sigma . (A_{\mu|E}, \widehat{\mu|E}) \cong_{\text{MA}} \left(([E]), \hat{\mu}|_{([E])} \right)$$

Proof =

A straightforward application of a previous theorem.

□

ThickEquivalence ::

$$:: \forall (X, \Sigma, \mu) \in \text{MEAS} . \forall Y : \text{Thick}(X, \Sigma, \mu) . (A_{\mu|Y}, \widehat{\mu|Y}) \cong_{\text{MA}} (X, \hat{\mu})$$

Proof =

A straightforward application of a previous theorem.

□

IndefiniteIntegralPrincipleIdealIsomorphism ::

$$:: \forall (X, \Sigma, \mu) \in \text{MEAS} . \forall f \in \mathcal{I}_+(X, \Sigma, \mu) . \exists E \in \Sigma . A_{f \, d\mu} \cong_{\text{BOOL}} ([E])$$

Proof =

We may assume that $\text{supp } f$ has a measurable envelope E .

Then the result is obvious as $\mathcal{N}_\mu \subset \mathcal{N}_{f \, d\mu}$.

□

1.1.8 Simple Products

`simpleProduct` :: $\prod_{I \in \text{SET}} (I \rightarrow \text{MeasureAlgebra}) \rightarrow \text{MeasureAlgebra}$

$$\text{simpleProduct}(A, \mu) = \prod_{i \in I} (A_i, \mu_i) := \left(\prod_{i \in I} A_i, \sum_{i \in I} \mu_i \right)$$

Obviously $\sum_{i \in I} \mu_i(0) = \sum_{i \in I} 0 = 0$.

Also assume $a : \mathbb{N} \rightarrow \prod_{i \in I} A_i$ is disjoint.

$$\text{Then } \sum_{i \in I} \mu_i \left(\bigvee_{n=1}^{\infty} a_n \right) = \sum_{i \in I} \sum_{n=1}^{\infty} \mu_i(a_{n,i}) = \sum_{n=1}^{\infty} \sum_{i \in I} \mu_i(a_{n,i}) = \sum_{n=1}^{\infty} \sum_{i \in I} \mu_i(a_n).$$

□

`PrincipleIdealsInMeasureAlgebras` ::

$$:: \forall I \in \text{SET} . \forall (A, \mu) : I \rightarrow \text{MeasureAlgebra} . (A_i, \mu_i) \cong_{\text{MA}} \left((e_i), \left(\sum_{i \in I} \mu_i \right)_{|(e_i)} \right)$$

`Proof` =

This is pretty obvious.

□

`SimpleProductCoproductCorrespondance` ::

$$:: \forall I \in \text{SET} . \forall (X, \Sigma, \mu) : I \rightarrow \text{MEAS} . \prod_{i \in I} (A_{\mu_i}, \hat{\mu}_i) \cong \text{measureAlgebra} \prod_{i \in I} (X_i, \Sigma_i, \mu_i)$$

`Proof` =

Obvious by Stone Theory.

□

`SimpleProductOfSemifinite` ::

$$:: \forall I \in \text{SET} . \forall (A, \mu) : I \rightarrow \text{SemifiniteMeasureAlgebra} . \text{SemifiniteMeasureAlgebra} \left(\prod_{i \in I} (A, \mu) \right)$$

`Proof` =

Assume a has infinite measure in (A, μ) .

Then there exists $i \in I$ such that $a_i \neq 0$.

As (A_i, μ_i) is semifinite there is $b \leq a_i$ such that $0 < \mu_i(b) < \infty$.

Then $be_i \leq a$ and $0 < \sum_{j \in I} \mu_j(be_i) = \mu_i(b) < \infty$.

□

SimpleProductOfLocalizable ::

$$:: \forall I \in \mathbf{SET} . \forall (A, \mu) : I \rightarrow \mathbf{LocalizableMeasureAlgebra} . \mathbf{LocalizableMeasureAlgebra} \left(\prod_{i \in I} (A, \mu) \right)$$

Proof =

Let J be a set and $a : J \rightarrow \prod_{i \in I} (A_i, \mu_i)$.

Then $\sup_{j \in J} a_j = (\sup_{j \in J} a_{j,i})_{i \in I}$.

□

PoUProductRepresentation ::

$$:: \forall (A, \mu) : \mathbf{MeasureAlgebra} . \forall (e_n)_{n=1}^{\infty} : \mathbf{PartitionOfUnity}(A) . (A, \mu) \cong_{\mathbf{MA}} \prod_{n=1}^{\infty} ((e_n), \mu|_{(e_n)})$$

Proof =

This is pretty obvious.

□

PoUProductRepresentation ::

$$:: \forall (A, \mu) : \mathbf{LocalizableMeasureAlgebra} . \exists I \in \mathbf{SET} . \exists (B, \nu) : I \rightarrow \mathbf{FiniteMeasureAlgebra} . \\ . (A, \mu) \cong_{\mathbf{MA}} \prod_{i \in I} (B_i, \nu_i)$$

Proof =

It is possible to select a partition of unity P of A consisting of finite elements.

Then by previous theorem $(A, \mu) \cong \prod_{p \in P} ((p), \mu|_{(p)})$.

And each $((p), \mu|_{(p)})$ are obviously finite.

□

LocalizableMeasureAlgebrasHasLocallyDeterminedRepresentations ::

$$:: \forall (A, \mu) : \mathbf{LocalizableMeasureAlgebra} . \exists (X, \Sigma, \nu) : \mathbf{LocallyDetermined} . (A, \mu) \cong_{\mathbf{MA}} (A_\nu, \hat{\nu})$$

Proof =

Represent $(A, \mu) \cong_{\mathbf{MA}} \prod_{i \in I} (B_i, \nu_i)$.

Then Stone's spaces $\mathbf{Z} B_i$ correspond to finite measure spaces.

And Stone's space of product correspond to a disjoint union of $\mathbf{Z} B_i$.

But such spaces are trivially locally determined.

□

1.1.9 Strictly Localizable Spaces

StrictlyLocalizableSpacePoU ::

$$:: \forall (X, \Sigma, \mu) : \text{StrictlyLocalizable} . \forall P : \text{PartitionOfUnity}(A_\mu) .$$

$$. \exists E : P \rightarrow \Sigma . \forall p \in P . [E_p] = p \ \& \ \text{Decomposition}(X, \Sigma, \mu, \text{Im } E)$$

Proof =
...
□

1.1.10 Subalgebras

SubalgebraMeasureAlgebra :: $\forall(A, \mu) : \text{MeasureAlgebra} . \forall B \subset_\sigma A . \text{MeasureAlgebra}(B, \mu|_B)$

Proof =

This is obvious.

□

SubalgebraFinifteMeasureAlgebra ::

$:: \forall(A, \mu) : \text{FiniteMeasureAlgebra} . \forall B \subset_\sigma A . \text{FiniteMeasureAlgebra}(B, \mu|_B)$

Proof =

This is obvious.

□

SigmaFiniteSubalgebraMeasureAlgebra ::

$:: \forall(A, \mu) : \sigma\text{-FiniteMeasureAlgebra} . \forall B \subset_\sigma A .$

$. \text{SemifiniteMeasureAlgebra}(B, \mu|_B) \Rightarrow \sigma\text{-FiniteMeasureAlgebra}(B, \mu|_B)$

Proof =

1 The set B^f is order-dense in B .

2 But then B^f is also order-dense in A .

3 Select a finite-measured countable partition of unity P in A .

4 If B is not σ -finite, then there is a subordinate uncountable partition of unity Q .

5 Then there would exist a uncountable refinement of P subordinate to Q .

6 Then P must contain an infinite element, but this is impossible!.

7 So Q must be countable, and so $(B, \mu|_B)$ must be countable.

□

FinifteMeasureAlgebraBySubalgebra ::

$:: \forall(A, \mu) : \text{MeasureAlgebra} . \forall B \subset_\sigma A . \text{FiniteMeasureAlgebra}(B, \mu|_B) \Rightarrow \text{FiniteMeasureAlgebra}(A, \mu)$

Proof =

This is obvious.

□

ProbabilityAlgebraBySubalgebra ::

$:: \forall(A, \mu) : \text{MeasureAlgebra} . \forall B \subset_\sigma A .$

$. \text{ProbabilityAlgebra}(B, \mu|_B) \Rightarrow \text{ProbabilityAlgebra}(A, \mu)$

Proof =

This is obvious.

□

$\text{SigmaFiniteAlgebraBySubalgebra} ::$
 $:: \forall (A, \mu) : \text{MeasureAlgebra} . \forall B \subset_\sigma A .$
 $. \sigma\text{-Finite}(B, \mu|_B) \Rightarrow \sigma\text{-Finite}(A, \mu)$
 $\text{Proof} =$
 This is obvious.
 \square

1.1.11 Localization

MeasureAlgebraCompletion ::

$$\begin{aligned} &:: \forall (A, \mu) : \text{SemifiniteMeasureAlgebra} . \exists ! \hat{\mu} : \tau(A) \rightarrow \mathbb{R}_{++}^{\infty} . \\ & . \hat{\mu}|_A = \mu \ \& \ \text{LocalizableMeasureAlgebra}(\tau(A), \hat{\mu}) \end{aligned}$$

Proof =

- 1 Define $\hat{\mu}(t) = \sup\{\mu(a) \mid a \in A, a \leq t\}$.
 - 2 As A is order dense in $\tau(A)$, it holds that $\hat{\mu}(a) = 0 \iff a = 0$ for any $a \in \tau(A)$.
 - 3 If $t : \mathbb{N} \rightarrow \tau(A)$ is disjoint then $\hat{\mu}\left(\bigvee_{n=1}^{\infty} t_n\right) = \sum_{n=1}^{\infty} \hat{\mu}(t_n)$.
 - 3.1 Write $S = \{a \in A : \exists c : \mathbb{N} \rightarrow A . a = \lim_{n \rightarrow \infty} c_n \ \& \ c \leq t\}$.
 - 3.2 Then there is $s = \sup S \in \tau(A)$.
 - 3.3 We write $\hat{\mu}(s) = \sup_{c \leq t} \mu\left(\bigvee_{n=1}^{\infty} c_n\right) = \sup_{c \leq t} \sum_{n=1}^{\infty} \mu(c_n) = \sum_{n=1}^{\infty} \sup_{c \leq t_n} \mu(c) = \sum_{n=1}^{\infty} \hat{\mu}(t_n)$.
 - 4 Obviously $(\tau(A), \hat{\mu})$ is semifinite and order-complete, and hence Localizable.
-

localization :: **SemifiniteMeasureAlgebra** \rightarrow **LocalizableMeasureAlgebra**

$$\text{localization}(A, \mu) = \left(\tau(A), \tau(\mu)\right) := \text{MeasureAlgebraCompletion}$$

LocalizationFiniteEmbedding ::

$$:: \forall (A, \mu) : \text{SemifiniteMeasureAlgebra} . \iota_{\tau}(A^f) = \tau^f(A)$$

Proof =

- 1 Assume $t \in \tau(A)$ such that $\hat{\mu}(t) < \infty$.
 - 2 Note, $\hat{\mu}(t) = \sup_{a \leq t} \mu(a)$.
 - 3 So we may select an increasing $a : \mathbb{N} \rightarrow A$ such that $\lim_{n \rightarrow \infty} \mu(a_n) = \hat{\mu}(t)$.
 - 4 Then $b = \bigvee_{n=1}^{\infty} a_n \in A$ and $\hat{\mu}(b) = \mu(b) = \hat{\mu}(t)$.
 - 5 So $\mu(t \setminus b) = 0$, and so $t = b \in A$ as clearly $b < t$.
-

1.1.12 Stone Spaces

LocalizableMeasureAlgebraHasStrictlyLocalizableStoneSpace ::

$:: \forall (A, \mu) : \text{LocalizableMeasureAlgebra} . \text{StrictlyLocalizable}(\mathbb{Z} A, \Sigma_\mu, \bar{\mu})$

Proof =

- 1 We already proved that $\bar{\mu}$ is locally determined.
- 2 As (A, μ) is semifinite there is a partition of unity P consisting of finite elements.
- 3 Use Stone representation $S_A(P)$ to construct a corresponding set in $\mathbb{Z} A$.
- 4 Assume $E \in \Sigma_\mu$ such that $\bar{\mu}(E) > 0$.
- 5 By definition of Stone's Space there is a clopen set $F \in \mathbb{Z} A$ such that $E \triangle F$ is meager.
- 6 And there is a Stone representation $a \in A$ such that $F = S_A(a)$.
- 7 Then $\mu(a) = \nu(S_A(a)) = \nu(E) > 0$.
- 8 So, there exists $p \in P$ such that $ap \neq 0$.
- 9 This means that $\nu(E \cap S_A(p)) > 0$.
- 10 As E was arbitrary this means that $S_A(P)$ provides a strict localization for $\bar{\mu}$.

□

MeagerSetsAreNowhereDense ::

$:: \forall (A, \mu) : \text{SemifiniteMeasureAlgebra} . \forall M \in \mathbf{MGR}(\mathbb{Z} A) . \text{NowhereDense}(\mathbb{Z} A, M)$

Proof =

- 1 As it was shown A is (σ, ∞) -WeaklyDistributive boolean algebra.
- 2 And this is a property of (σ, ∞) -WeaklyDistributive boolean algebra.

□

StoneSpaceMeasurableExpression ::

$:: \forall (A, \mu) : \text{SemifiniteMeasureAlgebra} . \forall E \in \Sigma_\mu .$
 $. \exists U : \text{Clopen}(\mathbb{Z} A) . \exists F : \text{NowhereDense}(\mathbb{Z} A) . E = U \cap F$

Proof =

- 1 This is clear from the previous theorem.

□

StoneSpaceMeasureComputation ::

$:: \forall (A, \mu) : \text{SemifiniteMeasureAlgebra} . \forall E \in \Sigma_\mu .$
 $. \bar{\mu}(E) = \sup \left\{ \mu(U) \mid U : \text{Clopen}(\mathbb{Z} A), U \subset E \right\}$

Proof =

- 1 This is clear from the previous theorem.

□

StoneSpaceCLDIsStrictlyLocalizable ::

$:: \forall (A, \mu) : \text{SemifiniteMeasureAlgebra} . \text{StrictlyLocalizable}(\mathbb{Z} A, \bar{\Sigma}_\mu, \bar{\bar{\mu}})$

Proof =

...

□

StoneSpaceCLDZeroSets ::

$:: \forall (A, \mu) : \text{SemifiniteMeasureAlgebra} . \mathcal{N}_{\bar{\mu}} = \mathcal{N}_{\bar{\mu}}$

Proof =

...

□

FiniteStoneSpaceMeasureComputation ::

$:: \forall (A, \mu) : \text{FiniteMeasureAlgebra} . \forall E \in \Sigma_{\mu} .$

$\bar{\mu}(E) = \inf \left\{ \mu(U) \mid U : \text{Clopen}(Z A), E \subset U \right\}$

Proof =

1 This is clear from the previous theorem.

□

1.1.13 Purely Infinite Elements

`purelyInfiniteElements` :: $\prod (A, \mu) : \text{MeasureAlgebra} . \sigma\text{-Ideal}(A)$

`purelyInfiniteElements` () = $I_\infty(\mu := \{a \in A : \forall b \in A . b \leq a \ \& \ \mu(b) < \infty \Rightarrow b = 0\})$

`semifiniteMeasure` :: $\prod (A, \mu) : \text{MeasureAlgebra} . \frac{A}{I_\infty(\mu)} \rightarrow \mathbb{R}_+^\infty$

`semifiniteMeasure` ([a]) = $\mu_{\text{sf}} := \sup\{\mu(b) | b \in A : b \leq a \ \& \ \mu(b) < \infty\}$

If [a] = [b], then $a \triangle b \in I_\infty(\mu)$.

So μ_{sf} is well-defined.

`SemifiniteMeasureIsMeasure` ::

$:: \forall (A, \mu) : \text{MeasureAlgebra} . \text{SemifiniteMeasureAlgebra} \left(\frac{A}{I}, \mu_{\text{sf}} \right)$

Proof =

1 If $\mu_{\text{sf}}[a] = 0$, then clearly $a \in I_\infty$.

2 Assume $[a] : \mathbb{N} \rightarrow A$ is disjoint.

2.1 Then $a_n a_m \in I_\infty$ if $n \neq m$.

2.2 Select increasing $b : \mathbb{N} \rightarrow A^f$ such that $b_n \leq \bigvee_{k=1}^\infty a_k$ and $\lim_{n \rightarrow \infty} \mu(b_n) = \mu_{\text{sf}} \left[\bigvee_{k=1}^\infty a_k \right] = \mu_{\text{sf}} \bigvee_{k=1}^\infty [a_k]$.

2.3 By (2.1) we may assert that $a b_n$ is disjoint and then $\bigvee_{k=1}^\infty a_k b_n = b_n$ for any $n \in \mathbb{N}$.

2.4 So $\mu(b) = \sum_{k=1}^\infty \mu(a_k b_n)$.

2.5 By taking limits and using monotonic convergence theorem

$$\sum_{k=1}^\infty \mu_{\text{sf}}[a_k] = \sum_{k=1}^\infty \lim_{n \rightarrow \infty} \mu(a_k b_n) = \lim_{n \rightarrow \infty} \mu(b_n) = \mu_{\text{sf}} \bigvee_{k=1}^\infty [a_k].$$

3 Clearly $\mu_{\text{sf}}[a] < \mu(a)$.

3.1 If $\mu_{\text{sf}}[a] = \infty$, then $a \notin I_\infty$.

3.2 So it is possible to select $b \in A$ such that $b \leq a$ and $0 < \mu(b) \leq a$.

3.3 $0 < \mu_{\text{sf}}[b] \leq \mu(b) < \infty$.

3.4 This proves that $\left(\frac{A}{I}, \mu_{\text{sf}} \right)$ is semifinite.

□

1.2 Topology

1.2.1 Subject

`measureAlgebraAsTopologicalSpace` :: `MeasureAlgebra` → `TOP`
`measureAlgebraAsTopologicalSpace` $((A, \mu)) = (A, \mu) :=$
 $:= \left(A, \mathcal{W}(A^f \times A^f, \mathbb{R}, \Lambda a \in A^f . \Lambda b \in A^f . \Lambda c \in A . \mu(ac + ab)) \right)$

`measureAlgebraAsUniformSpace` :: `MeasureAlgebra` → `UNI`
`measureAlgebraAsUniformSpace` $((A, \mu)) = (A, \mu) :=$
 $:= \left(A, \mathcal{I}(A^f \times A^f, \mathbb{R}, \Lambda a \in A^f . \Lambda b \in A^f . \Lambda c \in A . \mu(ac \triangle ab)) \right)$

`metricOfFrechetNikodym` :: $\prod (A, \mu) : \text{MeasureAlgebra} . \text{Metric}(A^f)$
`metricOfFrechetNikodym` $() = \rho_\mu := \Lambda a, b \in A^f . \mu(a \triangle b)$

`BooleanOperationsAreUniformlyContinuous` ::
:: $\forall (A, \mu) : \text{MeasureAlgebra} . (*), (\setminus), (\vee), (\wedge) \in \text{UNI}(A \times A, A)$

`Proof` =

- 1 Let \circ stay for any binary operation above.
 - 2 Select $c, d \in A$.
 - 3 Then $\mu(a(c \circ d) + b) \leq \mu(a(c \vee d) + b) \leq \mu(ac + d) + \mu(ad + b)$.
 - 4 So μ is bounded by the sum of uniform functions and is uniformly continuous.
-

`FiniteElementsAreDense` ::
:: $\forall (A, \mu) : \text{MeasureAlgebra} . \text{Dense}(A, A^f)$

`Proof` =

- 1 Select $c \in A$.
 - 2 Then c has a base of neighborhoods of form $U = \{u \in A : \mu(au + ac) \leq r\}$ with $a \in A^f, r \in \mathbb{R}_{++}$.
 - 3 But then $ac \in U$ and $ac \in A^f$.
-

`FiniteMeasureAlgebraHasUniformlyContinuousMeasure` ::
 $\forall (A, \mu) : \text{FiniteMeasureAlgebra} . \mu \in \text{UNI}(A, \mathbb{R}_{++})$

`Proof` =

This is pretty obvious as $\mu = \rho_\mu(0, a)$.

□

`FiniteMeasureAlgebraHasUniformlyContinuousMeasure` ::
 $\forall (A, \mu) : \text{FiniteMeasureAlgebra} . \mu \in \text{UNI}(A, \mathbb{R}_{++})$

`Proof` =

This is pretty obvious as $\mu = \rho_\mu(0, a)$.

□

`SemifinitMeasureAlgebraHasLowerSemicontinuousMeasure` ::

$\forall (A, \mu) : \text{SemifiniteMeasureAlgebra} . \mu \in \text{LowerSemicontinuous}(A, \mathbb{R}_{++}^{\infty})$

`Proof` =

This is pretty obvious as $\mu = \rho_{\mu}(0, a)$.

□

1.3 Category

1.4 Radon-Nikodym Parallels

2 Maharam's Theory

3 Abstract Ergodic Theory

4 Measurable Algebras

Sources:

1. D. H. Fremlin — Measure Theory (32,33,34) 2016