

# Measure.Know

Uncultured Tramp

December 18, 2016

$$\int_A d\lambda$$

# Contents

<b>1</b>	<b>Basic measure theory</b>	<b>3</b>
1.1	Limits of Sets . . . . .	3
1.2	Fields and Measures . . . . .	7
1.3	Measure Extension . . . . .	16
1.4	Lebesgue-Stieltjes Measures and Distributions on the Real Line . . . . .	26
1.5	Lebesgue-Stieltjes Measures in Multivariable Context [!]	34
1.6	Categorical Viewpoint: Boolean Algebras [!]	35
<b>2</b>	<b>Lebesgue Integration</b>	<b>36</b>
2.1	Measurable Functions . . . . .	36
2.2	Integration: Definition and Basic Results . . . . .	44
2.3	Convergence of Integrals . . . . .	48
2.4	Lebesgue VS Riemann [!]	52
2.5	Categorical Viewpoint: Integral As a Functor [!]	53
<b>3</b>	<b>Radon-Nikodym Theory</b>	<b>55</b>
3.1	Jordan-Hahn Decomposition . . . . .	55
3.2	Radon-Nikodym Theorem . . . . .	58
3.3	Absolutely Continuous Functions . . . . .	62
3.4	Differentiation of Measures . . . . .	66
3.5	Categorical Viewpoint: Developing Borel-Null [!]	72
<b>4</b>	<b>Convergence in Measure</b>	<b>73</b>
4.1	Measure Topology . . . . .	73
4.2	Comparison with other modes of convergence . . . . .	74
<b>5</b>	<b>Products over Measure spaces</b>	<b>78</b>
5.1	Product Measure Theorem . . . . .	78
5.2	Fubini Theorem . . . . .	82
5.3	Iterated Integrals . . . . .	85
5.4	Infinite Products . . . . .	87
<b>6</b>	<b>Convergence of Measures</b>	<b>92</b>
6.1	Duals of Measures . . . . .	92
6.2	Extending Linear Functionals . . . . .	95
6.3	Vague Convergence over Compacts . . . . .	97

# 1 Basic measure theory

## 1.1 Limits of Sets

$\text{IncreasingSetSeq} :: ?\mathbb{N} \rightarrow \text{Set}$

$A : \text{IncreasingSetSeq} \iff A \uparrow \iff \forall n \in \mathbb{N} . A_n \subset A_{n+1}$

$\text{IncreasingTo} :: ?\text{IncreasingSetSeq} \times \text{Set}$

$(A, \alpha) : \text{IncreasingTo} \iff A \uparrow \alpha \iff \alpha = \bigcup_{i=1}^{\infty} A_i$

$\text{DecreasingSetSeq} :: ?\mathbb{N} \rightarrow \text{Set}$

$A : \text{DecreasingSetSeq} \iff A \downarrow \iff \forall n \in \mathbb{N} . A_{n+1} \subset A_n$

$\text{IncreasingTo} :: ?\text{IncreasingSetSeq} \times \text{Set}$

$(A, \alpha) : \text{DecreasingTo} \iff A \downarrow \alpha \iff \alpha = \bigcap_{i=1}^{\infty} A_i$

$\text{ComplimentLimit1} :: \forall A \uparrow \alpha . A^c \downarrow \alpha^c$

$\text{ComplimentLimit2} :: \forall A \downarrow \alpha . A^c \uparrow \alpha^c$

$\text{ToDisjoint} :: \forall A : \mathbb{N} \rightarrow ?\Omega . \exists A' : \text{Disjoint}(\Omega, \mathbb{N}) : \bigcup_{i=1}^{\infty} A_i = \bigcup_{i=1}^{\infty} A'_i$

$\text{ToDisjoint}(A) = \Lambda n \in \mathbb{N} . \bigcap_{i=1}^{n-1} A_i^c \cap A_n$

$\text{toDisjoint} :: \forall A : \uparrow_{\Omega} . \exists A' : \text{Disjoint}(\Omega, \mathbb{N}) : \bigcup_{i=1}^{\infty} A_i = \bigcup_{i=1}^{\infty} A'_i$

$\text{toDisjoint}(A) = \Lambda n \in \mathbb{N} . \text{if } n = 1 \text{ then } A_1 \text{ else } (A_n \setminus A_{n-1})$

$\limsup :: (\mathbb{N} \rightarrow ?\Omega) \rightarrow ?\Omega$

$\limsup A = \bigcap_{n=1}^{\infty} \bigcup_{k=n}^{\infty} A_k$

$\liminf :: (\mathbb{N} \rightarrow ?\Omega) \rightarrow ?\Omega$

$$\liminf A = \bigcup_{n=1}^{\infty} \bigcap_{k=n}^{\infty} A_k$$

**LimSupComplement** ::  $\forall A : \mathbb{N} \rightarrow ?\Omega . (\limsup A)^c = \liminf A^c$

**Scatch** :

$$\begin{aligned} (\limsup A)^c &= \left( \bigcap_{n=1}^{\infty} \bigcup_{k=n}^{\infty} A_k \right)^c = \bigcup_{n=1}^{\infty} \left( \bigcup_{k=n}^{\infty} A_k \right)^c = \bigcup_{n=1}^{\infty} \bigcap_{k=n}^{\infty} A_k^c = \\ &= \liminf A^c \end{aligned}$$

**LimInfComplement** ::  $\forall A : \mathbb{N} \rightarrow ?\Omega . (\liminf A)^c = \limsup A^c$

**Scatch** :

$$\begin{aligned} (\liminf A)^c &= \left( \bigcup_{n=1}^{\infty} \bigcap_{k=n}^{\infty} A_k \right)^c = \bigcap_{n=1}^{\infty} \left( \bigcap_{k=n}^{\infty} A_k \right)^c = \bigcap_{n=1}^{\infty} \bigcup_{k=n}^{\infty} A_k^c = \\ &= \limsup A^c \end{aligned}$$

**InfSubsetSup** ::  $\forall A : \mathbb{N} \rightarrow ?\Omega . \liminf A \subset \limsup A$

**Proof** =

**Assume**  $A : \mathbb{N} \rightarrow ?\Omega$ ,

**Assume**  $n \in \mathbb{N}$ ,

**Assume**  $m \in \mathbb{N}$ ,

**Assume**  $a \in \bigcap_{k=m}^{\infty} A_k$ ,

$$a \in \bigcap_{k=m}^{\infty} A_k \rightsquigarrow a \in A_{n+m} \rightsquigarrow a \in \bigcup_{k=n}^{\infty} A_k;$$

$$\bigcap_{k=m}^{\infty} A_k \subset \bigcup_{k=n}^{\infty} A_k;$$

$$\bigcup_{m=1}^{\infty} \bigcap_{k=m}^{\infty} A_k \subset \bigcup_{k=n}^{\infty} A_k \rightsquigarrow \liminf A \subset \text{set} \bigcup_{k=n}^{\infty} A_k;$$

$$\liminf A \subset \bigcap_{n=1}^{\infty} \bigcup_{k=n}^{\infty} A_k \rightsquigarrow \liminf A \subset \limsup A; \square$$

**IncreasingLimit** ::  $\forall A \uparrow \alpha . \liminf A = \limsup A = \alpha$

**Proof** =

**Assume**  $A \uparrow \alpha$ ,

**Assume**  $a \in \alpha$ ,

$(A \uparrow \alpha, a \in \alpha) \rightsquigarrow \exists N \in \mathbb{N} : \forall n > N . a \in A_n$  **Extract**,

$\forall n > N . a \in A_n \rightsquigarrow \forall n > N . a \in \bigcap_{k=n}^{\infty} A_k \rightsquigarrow a \in \liminf A$ ,

$\forall n > N . a \in A_n \rightsquigarrow \forall n \in \mathbb{N} . a \in \bigcup_{k=n}^{\infty} A_k \rightsquigarrow a \in \limsup A$ ;

$\alpha \subset \liminf A, \alpha \subset \limsup A$  **as** (1),

$A \uparrow \alpha \iff \alpha = \bigcup_{n=1}^{\infty} A_n \rightsquigarrow \liminf A, \limsup A \subset \alpha$  **as** (2),

$(1, 2) \rightsquigarrow \liminf A = \limsup A = \alpha; \square$

**DecreasingLimit** ::  $\forall A \downarrow \alpha . \liminf A = \limsup A = \alpha$

**Proof** =

**Assume**  $A \downarrow \alpha$ ,

**Assume**  $a \in \limsup A$ ,

$a \in \limsup A \rightarrow \forall n \in \mathbb{N} . a \in \bigcup_{k=n}^{\infty} A_k$  **as** (2)

**Assume**  $n \in \mathbb{N}$ ,

$(2)(n) \rightsquigarrow a \in \bigcup_{k=n}^{\infty} A_k \rightsquigarrow \exists m \in \mathbb{N} : m \geq n : a \in A_m$  **Extract**,

$A \downarrow . a \in A_m \rightsquigarrow \forall k \in \mathbb{N} : k \leq m . a \in A_k \rightsquigarrow_n a \in A_n$ ;

$\forall n \in \mathbb{N} . a \in A_n \rightsquigarrow a \in \alpha$ ;

$\limsup A \subset \alpha$  **as** (2),

$\alpha \subset \liminf A, (2) \rightsquigarrow \liminf A = \limsup A = \alpha; \square$

**SetLimit** ::  $(\mathbb{N} \rightarrow ??\Omega) \rightarrow ??\Omega$   
 $\alpha : \text{SetLimit}(A) \iff \liminf A = \limsup A = \alpha$

**Example 1.**

$A := \Lambda n \in \mathbb{N} . \text{if } n : \text{Odd then } (-1/n, 1] \text{ else } (-1, 1/n)$   
 $\limsup A = (-1, 1],$   
 $\liminf A = \{0\},$

**Example 2.**

$A := \Lambda n \in \mathbb{N} . \mathbb{B}^2 \left( ((-1)^n/n, 0), 1 \right)$   
 $\limsup A = \overline{\mathbb{B}}^2(0, 1) \setminus \{(0, 1), (0, -1)\}$   
 $\liminf A = \mathbb{B}^2(0, 1)$

**Example 3.**

$\limsup x = X$   
 $A := \Lambda n \in \mathbb{N} . (-\infty, x_n]$   
 $(-\infty, X) \subset \limsup A \subset (-\infty, X]$   
 $\liminf y = Y$   
 $B := \Lambda n \in \mathbb{N} . (-\infty, y_n]$   
 $(-\infty, Y) \subset \liminf A \subset (-\infty, Y]$

**Example 4.**

$a < b < c < d$   
 $A := \Lambda n \in \mathbb{N} . \text{if } n : \text{Odd then } (a, b) \text{ else } (c, d)$   
 $\liminf A = \emptyset$   
 $\limsup A = (a, b) \cup (c, d)$

## 1.2 Fields and Measures

$\text{Algebra} :: \prod \Omega : \text{Set} . ???\Omega$

$\mathcal{F} : \text{Algebra} \iff \Omega \in \mathcal{F} \wedge \forall A, B \in \mathcal{F} . A^c \in \mathcal{F} \wedge A \cup B \in \mathcal{F}$

$\sigma\text{-Algebra} :: ?\text{Algebra}(\Omega)$

$\mathcal{F} : \sigma\text{-Algebra} \iff \forall A : \mathbb{N} \rightarrow \mathcal{F} . \bigcup_{i=1}^{\infty} A_i \in \mathcal{F}$

$\text{GenSigmaAlgebra} :: ???\Omega \rightarrow \sigma\text{-Algebra}(\Omega)$

$\text{GenSigmaAlgebra}(A) = \sigma_{\Omega}(A) = \bigcap \{ \mathcal{F} : \sigma\text{-Algebra}(\Omega) : A \subset \mathcal{F} \}$

$\text{AlgebraContraction} :: \forall \mathcal{A} : ???\Omega . \forall A \in ?\Omega . [\sigma_{\Omega}(\mathcal{A})] \cap A = \sigma_A([\mathcal{A}] \cap A)$

**Proof** =

**Assume**  $\mathcal{A} : ???\Omega$ ,

**Assume**  $A : ?\Omega$ ,

$[\sigma_{\Omega}(\mathcal{A})] \cap A : \sigma\text{-Algebra}(A) , [\mathcal{A}] \cap A \subset [\sigma_{\Omega}(\mathcal{A})] \cap A \rightsquigarrow$

$\rightsquigarrow \sigma_A([\mathcal{A}] \cap A) \subset [\sigma_{\Omega}(\mathcal{A})] \cap A$  **as** (1),

$G := \{ B \in \sigma_{\Omega}(\mathcal{A}) : B \cap A \in \sigma_A(\mathcal{A} \cap A) \}$

$G : \sigma\text{-Algebra}(\Omega) \rightsquigarrow G = \sigma_{\Omega}(\mathcal{A}) \rightsquigarrow [\sigma_{\Omega}(\mathcal{A})] \cap A \subset \sigma_A([\mathcal{A}] \cap A)$  **as** (2),

$(1, 2) \rightsquigarrow [\mathcal{A}] \cap A = [\sigma_{\Omega}(\mathcal{A})] \cap A ; \square$

$\text{Measure} :: \prod \mathcal{F} : \sigma\text{-Algebra}(\Omega) . ?\mathcal{F} \rightarrow \mathbb{R}_+^{\infty}$

$\mu : \text{Measure} \iff \forall A : \text{Disjoint}(\Omega, \mathbb{N}) \& \mathcal{F} . \mu \left( \bigcup_{n=1}^{\infty} A_n \right) = \sum_{i=1}^n \mu(A_i)$

$\text{Probability} :: ?\text{Measure}(\Omega, \mathcal{F})$

$\mathbb{P} : \text{Probability} \iff \mathbb{P}(\Omega) = 1$

$\text{MeasureSpace} := \sum \Omega : \text{Set} . \mathcal{F} : \sigma\text{-Algebra}(\Omega) . \text{Measure}(\Omega, \mathcal{F})$

$\text{ProbbilitySpace} := \sum \Omega : \text{Set} . \mathcal{F} : \sigma\text{-Algebra}(\Omega) . \text{Probability}(\Omega, \mathcal{F})$

$\text{SetFunction} :: \prod \mathcal{F} : \text{Algebra}(\Omega) . ?\mathcal{F} \rightarrow \mathbb{R}^{\infty}$

$f : \text{SetFunction} \iff \{-\infty, \infty\} \not\subset \text{Im } f \wedge \exists A \in \mathcal{F} : f(A) \in \mathbb{R}$

**Charge** :: ?SetFunction( $\Omega, \mathcal{F}$ )

$f : \text{Charge} \iff \forall (A, B) : \text{DisjointPair}(\Omega) . f(A \cup B) = f(A) + f(B)$

**CountablyAdditive** ::  $\prod \mathcal{F} : \sigma\text{-Algebra}(\Omega) . ?\text{SetFunction}(\Omega, \mathcal{F})$

$f : \text{CountablyAdditive} \iff \forall A : \text{Disjoint}(\Omega, \mathbb{N}) \& \mathcal{F} . \mu \left( \bigcup_{n=1}^{\infty} A_n \right) = \sum_{i=1}^n \mu(A_i)$

**ConcentratedOn** ::  $\prod \mathcal{F} : \sigma\text{-Algebra}(\Omega) . \mathcal{F} \rightarrow ?\text{Measure}(\Omega, \mathcal{F})$

$\mu : \text{ConcentratedOn}(A) \iff \mu(A^c) = 0$

**EmptyIsZero** ::  $\forall f : \text{Charge}(\Omega, \mathcal{F}) . f(\emptyset) = 0$

**Scatch** :

$f : \text{SetFunction}(\Omega, \mathcal{F}) \rightsquigarrow \exists A \in \mathcal{F} : f(A) \in \mathbb{R} \text{ Extract}$

$f(A) = f(A) + f(\emptyset) \rightsquigarrow f(\emptyset) = 0 \square$

**UnionDecomposition** ::  $\forall f : \text{Charge}(\Omega, \mathcal{F}) . \forall A, B \in \mathcal{F} .$

$. f(A \cup B) = f(A) + f(B) - f(A \cap B)$

**Scatch** :

$$\begin{aligned} f(A \cup B) &= f(A \cap B^c) + f(A \cap B) + f(B \cap A^c) = \\ &= \left( f(A \cap B^c) + f(A \cap B) \right) + \left( f(B \cap A^c) + f(A \cup B) \right) - f(A \cap B) = \\ &= f(A) + f(B) - f(A \cap B) \square \end{aligned}$$

**MeasureInequality** ::  $\forall \mu : \text{Measure}(\Omega, \mathcal{F}) . \forall A : \mathbb{N} \rightarrow \mathcal{F} . \mu \left( \bigcup_{n=1}^{\infty} A_n \right) \leq \sum_{n=1}^{\infty} \mu(A_n)$

**Finite** :: ?SetFunction( $\Omega, \mathcal{F}$ )

$f : \text{Finite} \iff \text{Im } f \subset \mathbb{R}$

**$\sigma$ -Finite** :: ?SetFunction( $\Omega, \mathcal{F}$ )

$f : \sigma\text{-Finite} \iff \exists A : \mathbb{N} \rightarrow \mathcal{F} : \bigcup_{n=1}^{\infty} A_n = \Omega \wedge \forall n \in \mathbb{N} . f(A_n) \in \mathbb{R}$



**MeasureUpperContinuity** ::  $\forall \mathcal{F} : \sigma\text{-Algebra}(\Omega) . \forall \mu : \text{CountablyAdditive}(\mathcal{F}) .$   
 $. \forall A \uparrow \alpha . \lim_{n \rightarrow \infty} \mu(A_n) = \mu(\alpha)$

**Proof** =

**Assume**  $\mathcal{F} : \sigma\text{-Algebra}(\Omega),$

**Assume**  $\mu : \text{CountablyAdditive}(\mathcal{F}),$

**Assume**  $A \uparrow \alpha,$

$A' := \text{toDisjoint}(A),$

$$\mu(\alpha) = \mu\left(\bigcup_{n=1}^{\infty} A'_n\right) = \sum_{n=1}^{\infty} \mu(A'_n)$$

**Assume**  $n \in \mathbb{N},$

$$\sum_{k=1}^n \mu(A'_k) = \mu(A_1) + \sum_{k=2}^n \mu(A_k \setminus A_{k-1}) = \mu(A_n);$$

$$\forall n \in \mathbb{N} . \sum_{k=1}^n \mu(A'_k) = \mu(A_n) \text{ as (1)}$$

**FromDef** **Serial** (1)  $\leadsto \lim_{n \rightarrow \infty} \mu(A_n) = \mu(\alpha); ; ; \square$

**MeasureLowerContinuity** ::  $\forall \mathcal{F} : \sigma\text{-Algebra}(\Omega) . \forall \mu : \text{CountablyAdditive}(\mathcal{F}) .$   
 $. \forall A \downarrow \alpha . \lim_{n \rightarrow \infty} \mu(A_n) = \mu(\alpha)$

**Proof** =

**Assume**  $\mathcal{F} : \sigma\text{-Algebra}(\Omega),$

**Assume**  $\mu : \text{CountablyAdditive}(\mathcal{F}),$

**Assume**  $A \downarrow \alpha,$

$B := A_1 \setminus A,$

$$B \uparrow A_1 \setminus \alpha \leadsto \lim_{n \rightarrow \infty} \mu(A_1 \setminus A_n) = \mu(A_1 \setminus \alpha) = \mu(A_1) - \mu(\alpha) \text{ as (1),}$$

$$\lim_{n \rightarrow \infty} \mu(A_1 \setminus A_n) = \lim_{n \rightarrow \infty} \mu(A_1) - \mu(A_n) = \mu(A_1) - \lim_{n \rightarrow \infty} \mu(A_n) \text{ as (2),}$$

$$(1, 2) \rightarrow \mu(\alpha) = \lim_{n \rightarrow \infty} \mu(A_n); ; ; \square$$

**UpperContinuous** ::  $\prod \mathcal{F} : \sigma\text{-Algebra}(\Omega) . ?\text{Charge}(\mathcal{F})$

$$\mu : \text{UpperContinuous} \iff \forall A \uparrow \alpha . \lim_{n \rightarrow \infty} \mu(A_n) = \mu(\alpha)$$

**LowerContinuous** ::  $\prod \mathcal{F} : \sigma\text{-Algebra}(\Omega) . ?\text{Charge}(\mathcal{F})$

$$\mu : \text{LowerContinuous} \iff \forall A \downarrow \alpha . \lim_{n \rightarrow \infty} \mu(A_n) = \mu(\alpha)$$

**CountablyAdditivityMark1** ::  $\forall \mathcal{F} : \sigma\text{-Algebra}(\Omega) . \forall \mu : \text{UpperContinuous}(\mathcal{F}) .$   
 $. \mu : \text{CountablyAdditive}(\mathcal{F})$

**Proof** =

**Assume**  $\mathcal{F} : \sigma\text{-Algebra}(\Omega) ,$

**Assume**  $\mu : \text{UpperContinuous}(\mathcal{F}) ,$

**Assume**  $A : \text{Disjoint}(\Omega, \mathbb{N}) : \text{Im}(A) \subset \mathcal{F} ,$

$B := \Lambda n \in \mathbb{N} . \bigcup_{k=1}^n A_k ,$

$B \uparrow \bigcup_{n=1}^{\infty} A_n ,$

$\mu : \text{UpperContinuous}(\mathcal{F}) \rightsquigarrow \lim_{n \rightarrow \infty} \mu(B_n) = \mu \left( \bigcup_{n=1}^{\infty} A_n \right) \text{ as } (1) ,$

**Assume**  $n \in \mathbb{N} ,$

$\mu : \text{UpperContinuous}(\mathcal{F}) \rightsquigarrow \mu : \text{Charge}(\mathcal{F}) ,$

$\text{Charge}(\mathcal{F})(\mu)(A_1, \dots, A_n) \rightsquigarrow \mu(B_n) = \sum_{k=1}^n \mu(A_k) ;$

$\forall n \in \mathbb{N} . \mu(B_n) = \sum_{k=1}^n \mu(A_k) \text{ as } (2) ,$

$\text{Seria}(2) \rightsquigarrow \lim_{n \rightarrow \infty} \mu(B_n) = \sum_{n=1}^{\infty} \mu(A_n) \rightsquigarrow_{(1)} \sum_{n=1}^{\infty} \mu(A_n) = \mu \left( \bigcup_{n=1}^{\infty} A_n \right) ;$

$\mu : \text{CountablyAdditive}(\mathcal{F}) ; ; \square$

**CountablyAdditivityMark1** ::  $\forall \mathcal{F} : \sigma\text{-Algebra}(\Omega) . \forall \mu : \text{LowerContinuous}(\mathcal{F}) .$   
 $. \mu : \text{CountablyAdditive}(\mathcal{F})$

**Proof** =

**Assume**  $\mathcal{F} : \sigma\text{-Algebra}(\Omega) ,$

**Assume**  $\mu : \text{UpperContinuous}(\mathcal{F}) ,$

**Assume**  $A : \text{Disjoint}(\Omega, \mathbb{N}) : \text{Im}(A) \subset \mathcal{F} ,$

$\alpha := \bigcup_{n=1}^{\infty} A_n ,$

$B := \Lambda n \in \mathbb{N} . \alpha \setminus \bigcup_{k=1}^n A_k ,$

$B \downarrow \emptyset \rightsquigarrow \lim_{n \rightarrow \infty} \mu(B_n) = 0 ,$

**Assume**  $n \in \mathbb{N} ,$

$$\begin{aligned}\mu(\alpha) &= \mu(B_n) + \sum_{k=1}^n \mu(A_n) \rightsquigarrow \sum_{k=1}^n \mu(A_n) = \mu(\alpha) - \mu(B_n); \\ \sum_{n=1}^{\infty} \mu(A_n) &= \lim_{n \rightarrow \infty} \mu(\alpha) - \mu(B_n) = \mu(\alpha) - \lim_{n \rightarrow \infty} \mu(B_n) = \mu(\alpha); \\ \mu &: \text{CountablyAdditive}(\mathcal{F});; \square\end{aligned}$$

Example 1

$$\begin{aligned}\text{Assume } \Omega &: \text{Infinite\&Countable}, \\ \mathcal{F} &:= 2^\Omega : \sigma\text{-Algebra}(\Omega), \\ \mu &:= \Lambda A \in \mathcal{F} . \text{ if } A : \text{Finite} \text{ then } 0 \text{ else } \infty : \text{Measure}(A), \\ \text{Assume } n &\in \mathbb{N}, \\ \text{Assume } A &: \text{DisjointElems}(\mathcal{F}, n), \\ \text{Assume } \exists k \in n &: A_k : \text{Infinite}, \\ \mu \left( \bigcup_{k=1}^n A_k \right) &= \infty = \sum_{k=1}^n \mu(A_k); \\ \text{Assume } \forall k \in n &. A_k : \text{Finite} \\ \mu \left( \bigcup_{k=1}^n A_k \right) &= 0 = \sum_{k=1}^n \mu(A_k);; \\ \mu &: \text{Charge}(\mathcal{F}), \\ \Omega &: \text{Infinite\&Countable} \rightsquigarrow \exists \omega : \mathbb{N} \leftrightarrow \Omega \text{ Extract}, \\ \mu(\Omega) = \infty \neq 0 &= \sum_{n=1}^{\infty} 0 = \sum_{n=1}^{\infty} \mu(\{\omega_n\}) \rightsquigarrow \mu ! \text{CountablyAdditive}(\mathcal{F}).\end{aligned}$$

Example 2

$$\begin{aligned}\text{Assume } \Omega &: \text{Infinite}, \\ \mu &:= \Lambda A \in 2^\Omega . \#A \\ \Omega &: \text{Infinite} \rightsquigarrow \exists Z : ?\Omega : Z : \text{Infinite\&Countable} \text{ Extract}, \\ Z &: \text{Infinite\&Countable} \rightsquigarrow \exists z : \mathbb{N} \leftrightarrow \Omega \text{ Extract}, \\ A &:= \Lambda n \in \mathbb{N} . Z \setminus \bigcup_{k=1}^n \{z_n\}, \\ A &\downarrow \emptyset, \\ \text{Assume } n &\in \mathbb{N}, \\ \mu(A_n) &= \infty; \\ \forall n \in \mathbb{N} &. \mu(A_n) = \infty, \\ \lim_{n \rightarrow \infty} \mu(A_n) &= \infty, \\ \mu &! \text{CountablyAdditive}(\mathcal{F}).\end{aligned}$$

Example 3

Assume  $\Omega : \text{Infinite\&Countable}$ ,

$\mathcal{F} := \left\{ A : ?\Omega : \#A < \infty \vee \#A^c < \infty \right\} : \text{Algebra}(\Omega)$ ,

$\mu := \Lambda A \in \mathcal{F} . \text{if } A : \text{Finite} \text{ then } 0 \text{ else } 1 : \mathcal{F} \rightarrow \mathbb{R}^{\infty}$ ,

Assume  $n \in \mathbb{N}$ ,

Assume  $A : \text{DisjointElems}(\mathcal{F}, n)$ ,

Assume Alternative  $\exists k \in n : A_k : \text{Infinite Extract}$ ,

Assume  $i \in n : i \neq k$ ,

$A : \text{DisjointElems}(\mathcal{F}, n), i \neq k \rightsquigarrow A_i \subset A_k^c \rightsquigarrow_{j(\mathcal{F})}$

$\rightsquigarrow_{j(\mathcal{F})} A_i : \text{Finite};$

$\forall i \in n : i \neq k . A_i : \text{Finite} \rightsquigarrow_{j(\mu)} \sum_{i=1}^n \mu(A_i) = 1 = \mu \left( \bigcup_{i=1}^n A_i \right);$

Close Alternative  $\forall k \in n . A_k : \text{Finite}$ ,

$\sum_{i=1}^n \mu(A_i) = 0 = \mu \left( \bigcup_{i=1}^n A_i \right);;$

$\mu : \text{Charge}(\mathcal{F})$ ,

... (as in Ex. 1)

$\mu ! \text{CountablyAdditive}(\mathcal{F})$ .

Example 4

$\mathcal{F} := \left\{ \bigcup_{k=1}^n I_i \mid n \in \mathbb{N}, I : \text{DisjointElem}(\text{RightSemiclosed}(\mathbb{R}), n) \right\}$

def  $\mu : \mathcal{F} \rightarrow \mathbb{R}^{\infty}$

$\mu(-\infty, a] = a$ ,

$\mu(a, b] = a - b$ ,

$\mu(b, \infty) = -b$ ,

$\mu(\mathbb{R}) = 0$ ,

$\mu \left( \bigcup_{i=1}^n I_i \right) = \sum_{i=1}^n \mu(I_i) \text{ Having } I : \text{DisjointElem}(\text{RightSemiclosed}(\mathbb{R}), n)$

Check  $(-\infty, n)$ .

**UT1** ::  $\forall \mu : \text{Charge}(\mathcal{F}) : \mu \geq 0 . \forall A : \text{DisjointElems}(\mathcal{F}, \mathbb{N}) : \bigcup_{n=1}^{\infty} A_n \in \mathcal{F} .$

$$\cdot \mu \left( \bigcup_{n=1}^{\infty} A \right) \geq \sum_{n=1}^{\infty} \mu(A)$$

**Proof** =

**Assume**  $\mu : \text{Charge}(\mathcal{F}) ; \mu \geq 0,$

**Assume**  $A : \text{DisjointElems}(\mathcal{F}, \mathbb{N}) : \bigcup_{n=1}^{\infty} A_n \in \mathcal{F},$

$$\alpha := \bigcup_{n=1}^{\infty} A_n \in \mathcal{F},$$

**Assume**  $n \in \mathbb{N},$

$$a_n := \mu \left( \alpha \setminus \bigcup_{i=1}^n A_n \right),$$

$$\mu(\alpha) = a_n + \sum_{k=1}^n \mu(A_n) \geq \sum_{k=1}^n \mu(A_n);$$

$$\mu(\alpha) \geq \sum_{k=1}^{\infty} \mu(A_n) \square$$

**UT2** ::  $\forall f : \Omega \rightarrow \Omega' . \forall \mathcal{B} : ??\Omega' . \sigma(f^{-1}(\mathcal{B})) = f^{-1}\sigma(\mathcal{B})$

**Proof** =

$$f : \Omega \rightarrow \Omega',$$

$$\mathcal{B} : ??\Omega',$$

**Assume**  $A \in f^{-1}\sigma(\mathcal{B}),$

$A \in f^{-1}\sigma(\mathcal{B}) \rightsquigarrow \exists B \in \sigma(\mathcal{B}) : A = f^{-1}B$  **Extract**,

$$\sigma(B) : \sigma\text{-Algebra}(\Omega') \rightsquigarrow B^{\complement} \in \sigma(B) \rightsquigarrow f^{-1}(B^{\complement}) = A^{\complement} \in f^{-1}\sigma(\mathcal{B});$$

$$(1) : \forall A \in f^{-1}\sigma(\mathcal{B}) . A^{\complement} \in f^{-1}\sigma(\mathcal{B}),$$

**Assume**  $A : \mathbb{N} \rightarrow f^{-1}\sigma(\mathcal{B}),$

$$B := f(A) : \mathbb{N} \rightarrow \sigma(\mathcal{B}),$$

$$\sigma(B) : \sigma\text{-Algebra}(\Omega') \rightsquigarrow \bigcup_{n=1}^{\infty} B_n \in \sigma(B) \rightsquigarrow f^{-1} \bigcup_{n=1}^{\infty} B_n = \bigcup_{n=1}^{\infty} A_n \in f^{-1}\sigma(\mathcal{B});$$

$$(2) : \forall A : \mathbb{N} \rightarrow f^{-1}\sigma(\mathcal{B}) . \bigcup_{n=1}^{\infty} A_n \in f^{-1}\sigma(\mathcal{B})$$

$$(1, 2) \rightsquigarrow f^{-1}\sigma(\mathcal{B}) : \sigma\text{-Algebra}(\Omega) \text{ as } (3),$$

$$\partial\sigma \rightsquigarrow \mathcal{B} \subset \sigma(\mathcal{B}) \rightsquigarrow (4) : f^{-1}\mathcal{B} \subset f^{-1}\sigma(\mathcal{B}),$$

$$\partial\sigma(3, 4) \rightsquigarrow (5) : \sigma(f^{-1}\mathcal{B}) \subset f^{-1}\sigma(\mathcal{B}),$$

$$G := \{B \in \sigma(\mathcal{B}) : f^{-1}(B) \in \sigma(f^{-1}(\mathcal{B}))\} : ??\Omega,$$

$$\text{Assume } B \in \mathcal{B},$$

$$f^{-1}(B) \in f^{-1}(\mathcal{B}) \rightsquigarrow_{\partial\sigma} f^{-1}(B) \in \sigma(f^{-1}(\mathcal{B})) \rightsquigarrow_{jG} B \in G;$$

$$(6) : B \subset G,$$

$$\text{Assume } B \in G,$$

$$B \in G \rightsquigarrow_{jG} f^{-1}(B) \in \sigma(f^{-1}(\mathcal{B})),$$

$$\partial\sigma \rightsquigarrow \sigma(f^{-1}(\mathcal{B})) : \sigma\text{-Algebra}(\Omega) \rightsquigarrow$$

$$\rightsquigarrow (f^{-1}(B))^{\complement} = f^{-1}(B^{\complement}) \in \sigma(f^{-1}(\mathcal{B})) \rightsquigarrow B^{\complement} \in f^{-1}(\mathcal{B});$$

$$(7) : \forall B \in G . B^{\complement} \in G,$$

$$\text{Assume } B : \mathbb{N} \rightarrow G,$$

$$\partial G \rightsquigarrow f^{-1}(B) : \mathbb{N} \rightarrow \sigma(f^{-1}(\mathcal{B})),$$

$$\partial\sigma \rightsquigarrow \sigma(f^{-1}(\mathcal{B})) : \sigma\text{-Algebra}(\Omega) \rightsquigarrow$$

$$\rightsquigarrow \bigcup_{n=1}^{\infty} f^{-1}(B_n) = f^{-1}\left(\bigcup_{n=1}^{\infty} B_n\right) \in \sigma(f^{-1}(\mathcal{B})) \rightsquigarrow \bigcup_{n=1}^{\infty} B_n \in G;$$

$$(8) : \forall B : \mathbb{N} \rightarrow G . \bigcup_{n=1}^{\infty} B_n \in G,$$

$$(7, 8) \rightsquigarrow G : \sigma\text{-Algebra}(\Omega') \text{ as } (9),$$

$$\partial\sigma(6, 9) \rightsquigarrow (10) \rightsquigarrow \sigma(\mathcal{B}) \subset G,$$

$$\partial G \rightsquigarrow (11) : G \subset \sigma(\mathcal{B}),$$

$$(10, 11) \rightsquigarrow G = \sigma(\mathcal{B}) \rightsquigarrow (12) : f^{-1} \sigma \mathcal{B} \subset \sigma f^{-1} \mathcal{B},$$

$$(5, 12) \rightsquigarrow f^{-1} \sigma \mathcal{B} = \sigma f^{-1} \mathcal{B}; ; \square$$

$$\text{UT3} :: \forall \mu : \text{FiniteMeasure}(\Omega, \mathcal{F}) . \forall A : \text{DisjountElems}(\mathcal{F}, X) : \forall x \in X . A_x > 0 . \#X \leq \aleph_0$$

$$\text{Proof} =$$

$$\text{Assume } \mu : \text{FiniteMeasure}(\Omega, \mathcal{F}),$$

$$\text{Assume } A : \text{DisjountElems}(\mathcal{F}, X) : \forall x \in X . A_x > 0,$$

$$b := \Lambda n \in \mathbb{N} . 1/n,$$

$$\text{Assume } n \in \mathbb{N},$$

$$Y_n := \{x \in X : \mu(A_x) \geq b_n\},$$

$$\text{Assume } a : \#Y_n \geq \aleph_0,$$

$$a \rightsquigarrow \exists y : \mathbb{N} \hookrightarrow Y_n \text{ Extract},$$

$$(1) : \mu\left(\bigcup_{i=1}^{\infty} A_{y_i}\right) = \sum_{i=1}^{\infty} \mu(A_{y_i}) \geq \sum_{i=1}^{\infty} b_n = \infty,$$

$$\mu : \text{FiniteMeasure}(\Omega, \mathcal{F}) \rightsquigarrow \mu\left(\bigcup_{i=1}^{\infty} A_{y_i}\right) < \infty \rightsquigarrow_{(1)} \perp;$$

$$(1) : \#Y_n \leq \aleph_0;$$

$$Y : \mathbb{N} \rightarrow ?X,$$

$$(1) : \forall n \in \mathbb{N} . \#Y_n < \aleph_0,$$

$$\mathfrak{d}(A, Y) \rightsquigarrow Y \upharpoonright X \rightsquigarrow (2) : X = \bigcup_{n=1}^{\infty} Y_n,$$

$$(1, 2, \#\mathbb{N} = \aleph_0) \rightsquigarrow \#X \leq \aleph_0; ; \square$$

## 1.3 Measure Extension

**PreBorel** :: Algebra

$$\mathbf{PreBorel} := \left\{ \bigcup_{i=1}^n A_i \mid n \in \mathbb{N}, A : n \rightarrow \mathbf{Semiclosed}(\mathbb{R}) \right\}$$

**PreBorelPreMeasure** :: ?CountablyAdditive (PreBorel)

$$\mu : \mathbf{PreBorelPreMeasure} \iff \exists F : \mathbf{Increase\&RightContinuous}(\mathbb{R}) : \\ : \mu(a, b] = F(b) - F(a)$$

**Extentible** :: ?CountablyAdditive( $\Omega, \mathcal{F}$ ) &  $\mathcal{F} \rightarrow \mathbb{R}_+$

$$\mu : \mathbf{Extentible} \iff \forall A : \mathbf{DisjointElems}(\mathcal{F}, \mathbb{N}) : \bigcup_{n=1}^{\infty} A_n \in \mathcal{F} . \mu \left( \bigcup_{n=1}^{\infty} A_n \right) = \sum_{n=1}^{\infty} \mu(A_n)$$

**FExtentible** :: ?Extentible( $\Omega, \mathcal{F}$ )

$$\mu : \mathbf{FExtentible} \iff \forall A \in \mathcal{F} . \mu(A) < \infty$$

**MonotonicityI** ::  $\forall P : \mathbf{FExtentible}(\mathcal{F}, \Omega) .$

$$. \forall A \uparrow \alpha, B \uparrow \beta : \text{Im } A \cup \text{Im } B \subset \mathcal{F} : \alpha \subset \beta . \lim_{n \rightarrow \infty} P(A_n) = \lim_{n \rightarrow \infty} P(B_n)$$

**Proof** =

**Assume**  $P : \mathbf{FinitelyAdditive}(\mathcal{F}, \Omega) : \text{Im } P \subset [0, 1] : P(\Omega) = 1,$

**Assume**  $\forall A \uparrow \alpha, B \uparrow \beta : \text{Im } A \cup \text{Im } B \subset \mathcal{F} : \alpha \subset \beta . \lim_{n \rightarrow \infty} P(A_n) = \lim_{n \rightarrow \infty} P(B_n),$

**Assume**  $n, m \in \mathbb{N} : n < m,$

$$A_n \subset A_m, P \geq 0 \rightsquigarrow P(A_n) \leq P(A_m),$$

$$B_n \subset B_m, P \geq 0 \rightsquigarrow P(B_n) \leq P(B_m);$$

$$P(A), P(B) : \mathbf{Nondecreasing}(\mathbb{N}, [0, 1]),$$

$$P \leq 1 \rightsquigarrow P(A), P(B) : \mathbf{Bounded}(\mathbb{N}, [0, 1]),$$

**BoundedConvergence**  $\rightsquigarrow P(A), P(B) : \mathbf{Convergent},$

**Assume**  $n \in \mathbb{N},$

$$C := A_n \cap A',$$

$$C \uparrow A_n,$$

$$A_n \in \mathcal{F}, A : \mathbf{Extentible}(\Omega, \mathcal{F}) \rightsquigarrow \lim_{m \rightarrow \infty} \mu(C_n) = \mu(A_n),$$

**Assume**  $m \in \mathbb{N},$

$$C_n \subset A'_m \rightsquigarrow \mu(C_n) \leq \mu(A'_m);$$

$$\lim_{m \rightarrow \infty} C_m \leq \lim_{m \rightarrow \infty} A'_m \rightsquigarrow \mu(A_n) \leq \lim_{m \rightarrow \infty} \mu(A'_m);$$

$$\lim_{m \rightarrow \infty} \mu(A_n) \leq \lim_{m \rightarrow \infty} \mu(A'_m); \square$$



**SetOfUnions** ::  $??\Omega \rightarrow ??\Omega$

$$\mathbf{SetOfUnions}(S) = U(S) = \left\{ \bigcup_{n=1}^{\infty} A_n \mid A : \mathbb{N} \rightarrow S \right\}$$

**Premeasure** ::  $\prod \mathcal{F} : \mathbf{Algebra}(\Omega) . \prod c \in \mathbb{R}_{++} . ?U(\mathcal{F}) \rightarrow [0, c]$

$$\begin{aligned} P : \mathbf{Premeasure} &\iff P|_{\mathcal{F}} : \mathbf{Extendible}(\mathcal{F}, \Omega) : P(\Omega) = c, \\ \forall A, B \in U(\mathcal{F}) . P(A \cup B) &= P(A) + P(B) - P(A \cap B), \\ \forall A, B \in U(\mathcal{F}) : A \subset B . P(A) &\leq P(B), \\ \forall A : \mathbb{N} \rightarrow U(\mathcal{F}) : A \uparrow \alpha . \alpha \in S(\mathcal{F}) \wedge \lim_{n \rightarrow \infty} P(A_n) &= P(\alpha). \end{aligned}$$

**ExtensionI** ::  $\forall \mu : \mathbf{FExtendible}(\Omega, \mathcal{F}) : \mu(\Omega) = 1 . \exists P : \mathbf{Premeasure}(\Omega, \mathcal{F}) : P|_{\mathcal{F}} = \mu$

**Proof** =

**Assume**  $\mu : \mathbf{Extendible}(\Omega, \mathcal{F})$ ,

$$P := \Lambda A \in U(\mathcal{F}) . \text{if } A \in \mathcal{F} \text{ then } \mu(A) \text{ else } \lim_{n \rightarrow \infty} \mu \left( \bigcup_{i=1}^n \partial U(A)_i \right),$$

$$\mu(\Omega) = 1 \rightsquigarrow (1) : P(\Omega) = 1,$$

**Assume**  $A, B \in U(\mathcal{F})$ ,

$$a := \Lambda n \in \mathbb{N} . \bigcup_{i=1}^n \partial U(A)_i : \mathbb{N} \rightarrow \mathcal{F} : a \uparrow A,$$

$$b := \Lambda n \in \mathbb{N} . \bigcup_{i=1}^n \partial U(B)_i : \mathbb{N} \rightarrow \mathcal{F} : b \uparrow B,$$

$$a \cap b \uparrow A \cap B,$$

$$a \cup b \uparrow A \cup B,$$

**Assume**  $n \in \mathbb{N}$ ,

$$(2) : P(a_n \cup b_n) = P(a_n) + P(b_n) - P(a_n \cap b_n);$$

$$(2) : P(A \cup B) = P(A) + P(B) - P(A);$$

$$(2) : \forall A, B \in U(\mathcal{F}) . P(A \cup B) = P(A) + P(B) - P(A),$$

**MonotonicityI**( $\mu$ ) = (3) :  $\forall A, B \in U(\mathcal{F}) : A \subset B . P(A) \leq P(B)$ ,

**Assume**  $A : \mathbb{N} \rightarrow U(\mathcal{F}) : A \uparrow \alpha$ ,

**Assume**  $n \in \mathbb{N}$ ,

$$a^n := \Lambda m \in \mathbb{N} . \bigcup_{i=1}^m \partial U(A_n)_i : \mathbb{N} \rightarrow \mathcal{F} : a^n \uparrow A_n;$$

$$a : \mathbb{N} \rightarrow \mathbb{N} \rightarrow \mathcal{F},$$

$$a' := \Lambda n \in \mathbb{N} . \bigcup_{i=1}^n a_n^i : \mathbb{N} \rightarrow \mathcal{F},$$

$$a' \uparrow \alpha \rightsquigarrow \alpha \in U(\mathcal{F}),$$

**Assume**  $n, m \in \mathbb{N} : n \leq m$ ,

$$a_m^n \subset a'_m \rightsquigarrow P(a_m^n) \leq P(a'_m),$$

$$a'_m \subset A_m \rightsquigarrow P(a'_m) \leq P(A_m) \rightsquigarrow P(a_m^n) \leq P(A_m);$$

$$(4) : \forall n \in \mathbb{N} : \lim_{m \rightarrow \infty} P(a_m^n) \leq \lim_{m \rightarrow \infty} P(a'_m) \leq \lim_{m \rightarrow \infty} P(A_m),$$

$$(4) \rightsquigarrow \forall n \in \mathbb{N} . P(A_n) \leq \lim_{m \rightarrow \infty} P(a'_n) \leq \lim_{m \rightarrow \infty} P(A_m) \rightsquigarrow$$

$$\rightsquigarrow \lim_{n \rightarrow \infty} P(A_n) \leq \lim_{m \rightarrow \infty} P(a'_n) \leq \lim_{m \rightarrow \infty} P(A_m) \rightsquigarrow$$

$$\rightsquigarrow (4) : \lim_{m \rightarrow \infty} P(A_m) = \lim_{n \rightarrow \infty} P(a'_n),$$

$$a' \uparrow \alpha, a' : \mathbb{N} \rightarrow \mathcal{F} \rightsquigarrow \lim_{n \rightarrow \infty} P(a'_n) = P(\alpha) \rightsquigarrow \lim_{n \rightarrow \infty} P(A_n) = P(\alpha);$$

$$(4) : \forall A : \mathbb{N} \rightarrow U(\mathcal{F}) : A \uparrow \alpha . \alpha \in S(\mathcal{F}) \wedge \lim_{n \rightarrow \infty} P(A_n) = P(\alpha),$$

$$(1, 2, 3, 4) \rightsquigarrow P : \text{Premeasure}(\Omega, \mathcal{F}); \square$$

$$\text{OuterMeasure} :: ? \left( ?\Omega \rightarrow \mathbb{R}_+^\infty \right)$$

$$\mu : \text{OuterMeasure} \iff \mu(\emptyset) = 0,$$

$$\forall A, B \in ?\Omega : A \subset B . \mu(A) \leq \mu(B),$$

$$\forall A : \mathbb{N} \rightarrow ?\Omega . \mu \left( \bigcup_{n=1}^{\infty} A_n \right) \leq \sum_{n=1}^{\infty} \mu(A_n)$$

$$\text{OuterExtension} :: \prod \mu : \text{Premeasure}(\Omega, \mathcal{F}) . ?(?\Omega \rightarrow \mathbb{R}^\infty)$$

$$P : \text{OuterExtension} \iff P|_{U(\mathcal{F})} = \mu,$$

$$\forall A, B \in ?\Omega . P(A \cup B) + P(A \cap B) \leq P(A) + P(B),$$

$$\forall A, B \in ?\Omega : A \subset B . P(A) \leq P(B),$$

$$\forall A \uparrow_\Omega \alpha : \lim_{n \rightarrow \infty} P(A_n) = P(\alpha),$$

**ExtensionII** ::  $\forall \mu : \text{Premeasure}(\Omega, \mathcal{F}) . \exists P : \text{OuterExtension}(\mu)$

**Proof** =

**Assume**  $\mu : \text{Premeasure}(\Omega, \mathcal{F})$ ,

$P := \Lambda A \in ?\Omega . \inf\{\mu(S) \mid S \in U(\mathcal{F}) : A \subset S\}$ ,

**Assume**  $A, B \in 2^\Omega$ ,

$\exists P \rightsquigarrow \exists a : \mathbb{N} \rightarrow U(\mathcal{F}) : \forall n \in \mathbb{N} . A \subset a_n : \mu(a_n) \downarrow P(A)$  **Extract**,

$\exists P \rightsquigarrow \exists b : \mathbb{N} \rightarrow U(\mathcal{F}) : \forall n \in \mathbb{N} . B \subset b_n : \mu(b_n) \downarrow P(B)$  **Extract**,

**Assume**  $n \in \mathbb{N}$ ,

$A \subset a_n, B \subset b_n \rightsquigarrow A \cap B \subset a_n \cap b_n, A \cup B \subset a_n \cup b_n$ ,

$\mu(a_n) + \mu(b_n) = \mu(a_n \cup b_n) + \mu(a_n \cap b_n)$ ;

$P(A \cap B) + P(A \cup B) \leq \lim_{n \rightarrow \infty} \mu(a_n \cup b_n) + \mu(a_n \cap b_n) = \lim_{n \rightarrow \infty} \mu(a_n) + \mu(b_n) =$   
 $= P(A) + P(B)$ ;

(1) :  $\forall A, B \in ?\Omega . P(A \cup B) + P(A \cap B) \leq P(A) + P(B)$ ,

**Assume**  $A : A \uparrow_\Omega \alpha$ ,

**Assume**  $\epsilon \in \mathbb{R}_{++}$ ,

**Assume**  $n \in \mathbb{N}$ ,

$\exists P \rightsquigarrow \exists B \in U(\mathcal{F}) : \mu(B) \leq P(A_n) + \epsilon/n!$  **Extract as**  $B_n$ ;

$\beta := \bigcup_{n=1}^{\infty} B_n \in U(\mathcal{F})$ ,

**Assume**  $n \in \mathbb{N}$ ,

**Assume**  $P\left(\bigcup_{k=1}^n B_k\right) \leq P(A_n) + \sum_{k=1}^n \epsilon/k!$ ,

$P\left(\bigcup_{k=1}^{n+1} B_k\right) = P\left(\bigcup_{k=1}^n B_k \cup B_{n+1}\right) = P\left(\bigcup_{k=1}^n B_k\right) + P(B_{n+1}) - P\left(\bigcup_{k=1}^n B_k \cap B_{n+1}\right)$

$A_k \subset \bigcup_{k=1}^n B_k \cap B_{k+1} \rightsquigarrow P\left(\bigcup_{k=1}^n B_k \cap B_{k+1}\right) \geq P(A_n)$ ,

$P\left(\bigcup_{k=1}^{n+1} B_k\right) \leq P(A_n) + P(A_{n+1}) + \sum_{k=1}^{n+1} \epsilon/n! - P(A_n) = P(A_{n+1}) + \sum_{k=1}^{n+1} \epsilon/n! ;$

**Induction**  $\rightsquigarrow P(\alpha) \leq P(\beta) \leq \lim_{n \rightarrow \infty} P(A_{n+1}) + \epsilon\epsilon$ ;

$\lim_{n \rightarrow \infty} P(A_n) \leq P(\alpha) \leq \lim_{n \rightarrow \infty} P(A_n) \rightsquigarrow P(\alpha) = \lim_{n \rightarrow \infty} P(A_n)$ ;

(2) :  $\forall A \uparrow_\Omega \alpha : \lim_{n \rightarrow \infty} P(A_n) = P(\alpha)$ ,

(1, 2)  $\rightsquigarrow P : \text{OuterExtension}(\mu)$ ;  $\square$

**OuterTHM** ::  $\forall \mu : \text{Premeasure}(\Omega, \mathcal{F}) . \forall P : \text{OuterExtension}(\mu) . P : \text{OuterMeasure}(\Omega)$

**Proof** =

**Assume**  $\mu : \text{Premeasure}(\Omega, \mathcal{F})$ ,

**Assume**  $P : \text{OuterExtension}(\mu)$ ,

**Assume**  $A : \mathbb{N} \rightarrow ?\Omega$ ,

**Assume**  $n \in \mathbb{N}$ ,

$$P\left(\bigcup_{k=1}^n A_k\right) \leq P\left(\bigcup_{k=1}^n A_k\right) + \sum_{k=1}^n P\left(A_k \cap \bigcup_{i=k+1}^n A_i\right) \leq \sum_{k=1}^n P(A_k);$$

$$P\left(\bigcup_{k=1}^{\infty} A_k\right) = \lim_{n \rightarrow \infty} P\left(\bigcup_{k=1}^n A_k\right) \leq \lim_{n \rightarrow \infty} \sum_{k=1}^n P(A_k) = \sum_{k=1}^{\infty} P(A_k);$$

$P : \text{OuterMeasure}(\Omega); \square$

**relativeAlgebra** ::  $\text{OuterMeasure}(\Omega) \rightarrow ??\Omega$

$$\text{relativeAlgebra}(P) = H(P) := \left\{ A \in \Omega : P(A) + P(A^c) = P(\Omega) \right\}$$

**extension** ::  $\text{FExtendible}(\Omega, \mathcal{F}) \rightarrow \text{OuterMeasure}(\Omega) \& \text{OuterExtension}(\Omega, \mathcal{F})$

**extension**( $\mu$ ) =  $P_\mu := \text{OuterTHM ExtensionII ExtensionI } \mu$

**ExtensionIII** ::  $\forall \mu : \text{FExtendible}(\Omega, \mathcal{F}) . H(P_\mu) : \sigma\text{-Algebra}(\Omega) : \mathcal{F} \subset H(P_\mu)$

**Proof** =

**Assume**  $\mu : \text{Premeasure}(\Omega, \mathcal{F})$ ,

$P := P_\mu$ ,

**Assume**  $A \in U(\mathcal{F})$ ,

$a := \mathfrak{D}(U)(\mathcal{F}, A) : \mathbb{N} \rightarrow \mathcal{F} : a \uparrow A$ ,

**Assume**  $n \in \mathbb{N}$ ,

$$a_n \subset A \rightsquigarrow A^c \subset a_n^c \rightsquigarrow P(A^c) \leq P(a_n^c)$$

$$P(a_n) + P(A^c) \leq P(a_n) + P(a_n^c) = P(\Omega);$$

$$(1) : P(A) + P(A^c) \leq \lim_{n \rightarrow \infty} P(a_n) + P(A^c) \leq P(\Omega),$$

$$(2) : P(A) + P(A^c) \geq P(A \cup A^c) + P(A \cap A^c) = P(\Omega) + P(\emptyset) = P(\Omega),$$

$$(1, 2) \rightsquigarrow P(A) + P(A^c) = P(\Omega) \rightsquigarrow A \in H(P);$$

$$(1) : U(\mathcal{F}) \subset H(P);$$

**Assume**  $A, B \in H(P)$ ,

**ExtensionII**  $\leadsto (2) : P(A \cap B) + P(A \cup B) \leq P(A) + P(B)$ ,

**ExtensionII**  $\leadsto (3) : P((A \cap B)^c) + P((A \cup B)^c) \leq P(A^c) + P(B^c)$ ,

$(2, 3) \leadsto (4) : P((A \cap B)^c) + P((A \cup B)^c) + P(A \cap B) + P(A \cup B)$   
 $\leq P(A) + P(B) + P(A^c) + P(B^c) = 2P(\Omega)$ ,

$(5) : P(\Omega) \leq P((A \cup B)^c) + P(A \cup B)$ ,

$(6) : P(\Omega) \leq P((A \cap B)^c) + P(A \cap B)$ ,

$(4, 5, 6) \leadsto P(\Omega) = P((A \cup B)^c) + P(A \cup B), P(\Omega) = P((A \cap B)^c) + P(A \cap B) \leadsto$   
 $\leadsto A \cap B, A \cup B \in H(P)$ ;

$H(P) : (\mathbf{Algebra})(\Omega)$ ,

**Assume**  $A : \mathbb{N} \rightarrow H(P) : A \uparrow \alpha$ ,

**Assume**  $\epsilon \in \mathbb{R}_{++}$ ,

**ExtensionII**  $\leadsto \lim_{n \rightarrow \infty} P(A_n) = P(\alpha) \leadsto \exists n \in \mathbb{N} . P(\alpha) \leq P(A_n) + \epsilon$  **Extract**,

$A_n \subset \alpha \leadsto \alpha^c \subset A_n^c \leadsto P(\alpha^c) \leq P(A_n^c)$ ,

$P(\Omega) \leq P(\alpha^c) + P(\alpha) \leq P(A_n^c) + P(A_n) + \epsilon = P(\Omega) + \epsilon$ ;

$P(\alpha^c) + P(\alpha) = P(\Omega) \leadsto P(\alpha) \in H(P)$ ;

$H(P) : \sigma\text{-Algebra}(\Omega)$ ;  $\square$

**extensionToSpace** :  $\mathbf{FExtendible}(\Omega, \mathcal{F}) \rightarrow \mathbf{MeasureSpace}$

**extensionToSpace**( $\mu$ ) =  $\mu^\circ := (\Omega, \sigma(\mathcal{F}), P_{\mu|\sigma(\mathcal{F})})$

**Complete** ::  $? \mathbf{MeasureSpace}$

$(\Omega, \mathcal{F}, \mu) : \mathbf{Complete} \iff \forall A \in \mathcal{F} : \mu(A) = 0 . \forall S \subset A . S \in \mathcal{F}$

**algCompletion** ::  $\mathbf{Measure}(\Omega, \mathcal{F}) \rightarrow \sigma\text{-Algebra}(\Omega)$

**algCompletion**( $\mu$ ) =  $C(\mu) = [\mathcal{F}] \cup [\{A \subset \Omega | \exists F \in \mathcal{F} : \mu(F) = 0 : A \subset F\}]$

**completion** ::  $\mathbf{MeasureSpace} \rightarrow \mathbf{Complete}$

**completion**( $\Omega, \mathcal{F}, \mu$ ) =  $(\widehat{\Omega, \mathcal{F}, \mu}) := (\Omega, C(\mu), \hat{\mu} := \Lambda A \cup N \in_{\mathfrak{D}} C(\mu) . \mu(A)$

**ComplementaryCompletion** ::  $\forall \mu : \mathbf{FExtendible}(\Omega, \mathcal{F}) . \widehat{\mu^\circ} = (\Omega, H(P_\mu), P_\mu)$

**Proof** =

**Assume**  $\mu : \mathbf{FExtendible}(\Omega, \mathcal{F})$ ,

$P := P_\mu$ ,

**Assume**  $A \in H(P)$ ,

$\partial H(P)(A) \rightsquigarrow A^\complement \in H(P)$ ,

$a := \partial P(A) : \mathbb{N} \rightarrow U(\mathcal{F}) : P(a) \downarrow P(A)$ ,

$b := \partial P(A^\complement) : \mathbb{N} \rightarrow U(\mathcal{F}) : P(b) \downarrow P(A^\complement)$ ,

$a' := b^\complement : \mathbb{N} \rightarrow \sigma(\mathcal{F})$

$\partial H(P)(A) \rightsquigarrow (1) : P(\Omega) = P(A) + P(A^\complement) = \lim_{n \rightarrow \infty} P(a_n) + \lim_{n \rightarrow \infty} P(b_n)$ ;

$(2) : P(\Omega) = \lim_{n \rightarrow \infty} P(b_n) + P(a'_n)$ ;

$(1, 2) \rightsquigarrow \lim_{n \rightarrow \infty} P(a'_n) = \lim_{n \rightarrow \infty} P(a_n) = P(A)$ ;

$\alpha := \bigcap_{n=1}^{\infty} a_n \in \sigma(\mathcal{F})$ ,

$\alpha' := \bigcup_{n=1}^{\infty} a'_n \in \sigma(\mathcal{F})$ ,

$\beta := \alpha \cap \alpha^\complement \in \sigma(\mathcal{F})$ ,

$P(A) = P(\alpha) = P(\alpha') + P(\beta) = P(A) + P(\beta) \rightsquigarrow P(\beta) = 0$ ,

$\alpha' \subset A \subset \alpha \rightsquigarrow A \cap \alpha^\complement \subset \beta \rightsquigarrow A = \alpha' \cup (A \cap \alpha^\complement) \in C(P, \sigma(\mathcal{F}))$ ;

$(1) : H(P) \subset C(P, \sigma(\mathcal{F}))$ ,

**Assume**  $A \in C(P, \sigma(\mathcal{F}))$ ,

$(B, N) := \partial C(P, \sigma(\mathcal{F})) : \sigma(\mathcal{F}) \times \{A \in 2^\Omega : \exists N \in \sigma(\mathcal{F}) : P(N) = 0 : A \subset P\} : A = B \cup N$ ,

$B \in \sigma(\mathcal{F}) \rightsquigarrow B \in H(P)$ ,

$M = \partial(N) \in \sigma(\mathcal{F}) : P(M) = 0 : N \subset M$ ,

$M \in \sigma(\mathcal{F}) \rightsquigarrow M \in H(P)$ ,

$\partial \mathbf{Complete}(\Omega, H(P), P)(N.M) \rightsquigarrow N \in H(P)$ ,

$B, N \in H(P) \rightsquigarrow A = B \cup N \in H(P)$ ;

$(2) : C(P, \sigma(\mathcal{F})) \subset H(P)$ ,

$(1, 2) \rightsquigarrow C(P, \sigma(\mathcal{F})) = H(P); \square$

**MonotoneClass** :: ??? $\Omega$

$M : \text{MonotoneClass} \iff \forall A : \mathbb{N} \rightarrow M : A \downarrow \alpha . \alpha \in M,$   
 $\forall A : \mathbb{N} \rightarrow M : A \uparrow \alpha . \alpha \in M,$

**MonotoneClassTHM** ::  $\forall M : \text{MonotoneClass}(\Omega) . \forall \mathcal{F} : \text{Algebra}(\Omega) : \mathcal{F} \subset M . \sigma(\mathcal{F}) \subset M$

**Proof** =

**Assume**  $M : \text{MonotoneClass}(\Omega)$

**Assume**  $\mathcal{F} : \text{Algebra}(\Omega) : \mathcal{F} \subset M,$

$N := \min\{N : \text{MonotoneClass}(\Omega) : \mathcal{F} \subset N\},$

$\delta N(M) \rightsquigarrow N \subset M$

**Assume**  $A \in \mathcal{F},$

$N' = \{B \in N : A \cap B \in N \wedge A^c \cap B \in N \wedge A \cap B^c \in N\} \subset N,$

$\delta \text{Algebra}(\mathcal{F}) \rightsquigarrow \mathcal{F} \subset N',$

$\delta(N'), \delta \bigcap, \delta \bigcup \rightsquigarrow N' : \text{MonotoneClass}(\Omega) \rightsquigarrow N = N';$

(1) :  $\forall A \in \mathcal{F} . \forall B \in N . A \cap B \in N \wedge A^c \cap B \in N \wedge A \cap B^c \in N,$

**Assume**  $A \in N,$

$N' = \{B \in N : A \cap B \in N \wedge A^c \cap B \in N \wedge A \cap B^c \in N\} \subset N,$

(1)  $\rightsquigarrow \mathcal{F} \subset N',$

$\delta(N'), \delta \bigcap, \delta \bigcup \rightsquigarrow N' : \text{MonotoneClass}(\Omega) \rightsquigarrow N = N';$

$N : \text{Algebra}(\Omega),$

$N : \text{MonotoneClass}(\Omega) \& \text{Algebra}(\Omega) \rightsquigarrow N : \sigma\text{-Algebra}(\Omega) \rightsquigarrow \sigma(\mathcal{F}) \subset N \rightsquigarrow \sigma(\mathcal{F}) \subset M; ; \square$

**$\sigma$ -Finite** :: ?  $\sum \mathcal{F} : \text{Algebra}(\Omega) . \mathcal{F} \rightarrow \mathbb{R}_+^\infty$

$(\mathcal{F}, \mu) : \sigma\text{-Finite}(\Omega) \iff \exists A : \mathbb{N} \rightarrow \mathcal{F} : \forall n \in \mathbb{N} . \mu(A_n) < \infty : A \uparrow \Omega$

**CaratheodoryExtension** ::  $\forall(\mathcal{F}, \mu) : \sigma\text{-Finite}(\Omega) : \mu : \text{Extendible}(\mathcal{F}, \Omega) .$   
 $. \exists! \lambda : \text{Measure}(\Omega, \sigma(\mathcal{F})) : \lambda|_{\mathcal{F}} = \mu$

**Proof** =

**Assume**  $(\mathcal{F}, \mu) : \sigma\text{-Finite}(\Omega) : \mu : \text{Extendible}(\mathcal{F}, \Omega),$

$\omega := \text{toDisjoint } \partial\sigma\text{-Finite}(\Omega)(\mu, \mathcal{F}),$

**Assume**  $n \in \mathbb{N},$

$p := \Lambda A \in \mathcal{F} . \mu(\omega_n \cap A) : \text{Extendible}(\Omega, \mathcal{F}),$

$\mu(\omega_n) < \infty \rightsquigarrow p : \text{FExtendible}(\Omega, \mathcal{F}),$

$P_n := \text{extension}(p) : \text{FiniteMeasure}(\Omega, \sigma(\mathcal{F}));$

$\lambda := \Lambda A \in \sigma(\mathcal{F}) . \sum_{n=1}^{\infty} P_n(A) : \sigma(\mathcal{F}) \rightarrow \mathbb{R}_+^{\infty};$

**Assume**  $\lambda' : \text{Measure}(\Omega, \sigma(\mathcal{F})) : \lambda'|_{\mathcal{F}} = \mu,$

$\partial\lambda \rightsquigarrow (1) : \lambda'|_{\mathcal{F}} = \lambda|_{\mathcal{F}},$

**Assume**  $m \in \mathbb{N},$

$P'_n := \Lambda A \in \sigma(\mathcal{F}) . \lambda'(A \cap \omega_n) : \text{FiniteMeasure}(\Omega, \sigma(\mathcal{F})),$

$M := \{A \in \sigma(\mathcal{F}) : P(A) = P'(A')\},$

$P, P' : \text{Measure}(\Omega, \mathcal{F}) \rightsquigarrow M : \text{MonotomeClass}(\Omega),$

$(1) \rightsquigarrow \mathcal{F} \subset M$

, **MonotoneClassTHM**  $\rightsquigarrow \sigma(\mathcal{F}) \subset M \rightsquigarrow P_n = P'_n;$

$(2) : \forall n \in \mathbb{N} . P_n = P'_n,$

$(\partial(\lambda', P', \omega), 2) \rightsquigarrow \lambda' = \sum_{n=1}^{\infty} P'_n = \sum_{n=1}^{\infty} P_n = \lambda; \square$

**ApproximationI** ::  $\forall \mathcal{F} : \text{Algebra} . \forall P : \text{FiniteMeasure}(\Omega, \sigma(\mathcal{F})) .$

$. \forall A \in \sigma(\mathcal{F}) . \forall \epsilon \in \mathbb{R}_{++} . \exists B \in \mathcal{F} : P(A \triangle B) \leq \epsilon$

**Assume**  $F : \text{Algebra},$

**Assume**  $P : \text{FiniteMeasure}(\Omega, \sigma(\mathcal{F})),$

**Assume**  $A \in \sigma(\mathcal{F}),$

**Assume**  $\epsilon \in \mathbb{R}_{++},$

$B := \text{ExtensionII}(A) : \mathbb{N} \rightarrow U(\mathcal{F}) : \forall n \in \mathbb{N} . A \subset B_n . P(B) \downarrow P(A),$

$P(B) \downarrow P(A) \rightsquigarrow \exists n \in \mathbb{N} : P(B_n) \leq P(A) + \epsilon/2, \text{ Extract}$

$C = \partial U(\mathcal{F})(B_n) : \mathbb{N} \rightarrow \mathcal{F} : C \uparrow B_n,$

$P : \text{UpperContinuous}(\sigma(\mathcal{F})) \rightsquigarrow \lim_{m \rightarrow \infty} P(C_m) = P(B_n) \rightsquigarrow$

$\rightsquigarrow \exists n \in \mathbb{N} : P(B_n) \leq P(C_m) + \epsilon/2,$

$P(A \triangle C_m) \leq P(A \triangle B_n \cup B_n \triangle C_m) =$

$= P(A \triangle B_n) + P(B_n \triangle C_m) - P(A \triangle B_n \cap B_n \triangle C_m) \leq$



$$\leq P(A \triangle B_n) + P(B_n \triangle C_m) = P(B_n) - P(A) + P(B_n) - p(C_m) = \epsilon; ; ; \square$$

**ApproximationII** ::  $\forall \mathcal{F} : \text{Algebra}(\Omega) . \forall \mu : \text{Measure}(\Omega, \sigma(\mathcal{F})) : (\mu, \mathcal{F}) : \sigma\text{-Finite}(\Omega) .$   
 $. \forall A \in \sigma(\mathcal{F}) . \forall \epsilon \in \mathbb{R}_{++} . \exists B \in \mathcal{F} : \mu(A \triangle B) \leq \epsilon$

**Assume**  $F : \text{Algebra}$ ,

**Assume**  $\mu : \text{Measure}(\Omega, \sigma(\mathcal{F})) : (\mu, \mathcal{F}) : \sigma\text{-Finite}(\Omega) ,$

**Assume**  $A \in \sigma(\mathcal{F}) ,$

**Assume**  $e \in \mathbb{R}_{++} ,$

**Assume**  $\epsilon \in \mathbb{R}_{++} ,$

$\omega := \text{toDisjoint } \text{d}\sigma\text{-Finite}(\Omega)(\mathcal{F}, \mu) : \text{DisjointElems}(\mathcal{F}, \mathbb{N}) : \forall n \in \mathbb{N} . \mu(\omega_n) < \infty : \bigcup_{n=1}^{\infty} \omega_n = \Omega ,$

**Assume**  $n \in \mathbb{N} ,$

$P_n := \Lambda A \in \sigma(\mathcal{F}) . \mu(A \cup \omega_n) : \text{FiniteMeasure}(\Omega, \mathcal{F}) ,$

$B_n := \text{ApproximationI}(\mathcal{F})(P_n)(A)(\epsilon/e(n-1)!) \in F : P(B_n \triangle A) \leq \epsilon/e(n-1)!,$

$P_n(A \triangle B_n) = \mu((A \triangle B_n) \cap \omega_n) = \mu((A \triangle (B_n \cap \omega_n)) \cap \omega_n) = P_n(A \triangle (B_n \cap \omega_n));$

$C := \bigcup_{n=1}^{\infty} B_n ,$

**Assume**  $n \in \mathbb{N};$

$P_n(A \triangle C) = P_n(A \triangle B_n);$

$\mu(A \triangle C) = \sum_{n=1}^{\infty} P_n(A \triangle C) = \sum_{n=1}^{\infty} (A \triangle B_n) \leq \epsilon;$

$0 = \sum_{n=1}^{\infty} P_n(A \triangle B_n) = \sum_{n=1}^{\infty} P_n(A \setminus B_n \cup B_n \setminus A) = \sum_{n=1}^{\infty} P_n(A \setminus B_n) + P_n(B_n \setminus A) =$   
 $= \sum_{n=1}^{\infty} P_n(A \setminus B_n) + \sum_n P_n(B_n \setminus A) = \lim_{n \rightarrow \infty} \mu \left( A \setminus \bigcup_{i=1}^n B_i \right) + \lim_{n \rightarrow \infty} \mu \left( \bigcup_{i=1}^n B_i \setminus A \right) \rightsquigarrow$   
 $\rightsquigarrow \exists n \in \mathbb{N} : \mu \left( \bigcup_{i=1}^n B_i \setminus A \right) \leq e/2 \wedge \mu \left( A \setminus \bigcup_{i=1}^n B_i \right) \leq e/2 \text{Extract},$

$\beta := \bigcup_{i=1}^n B_i \in \mathcal{F},$

$\mu(A \triangle \beta) = \mu(A \setminus \beta) + \mu(\beta \setminus A) \leq e; ; ; \square$

## 1.4 Lebesgue-Stieltjes Measures and Distributions on the Real Line

$\text{borelSets} :: \prod X : \text{TopologicalSpace} . \sigma\text{-Algebra}(X)$   
 $\text{borelSets} = \mathcal{B}(X) := \sigma(\mathcal{T}_X)$

$\text{Lebesgue-Stieltjes}(\mathbb{R}) :: ?\text{Measure}(\mathbb{R}, \mathcal{B}(\mathbb{R}))$ ,  
 $\mu : \text{Lebesgue-Stieltjes} \iff \forall I \in \mathcal{B}(\mathbb{R}) : \text{Bounded} . \mu(I) < \infty$

$\text{DistributionFunction}(\mathbb{R}) :: ?\text{RightContinuous\&Increasing} \left( \overset{\infty}{\mathbb{R}}, \overset{\infty}{\mathbb{R}} \right)$   
 $F : \text{DistributionFunction}(\mathbb{R}) \iff \lim_{x \rightarrow \infty} F(x) = F(\infty)$

$\text{MeasureAsDistribution} :: \forall \mu : \text{Lebesgue-Stieltjes}(\mathbb{R}) . \forall x, c \in \mathbb{R} .$   
 $. \exists F : \text{DistributionFunction}(\mathbb{R}) : F(x) = c : \forall (a, b] \in \text{SemiClosed}(\mathbb{R}) .$   
 $. \mu(a, b] = F(b) - F(a)$

**Proof** =

**Assume**  $\mu : \text{Lebesgue-Stieltjes}(\mathbb{R})$ ,

**Assume**  $x, c \in \mathbb{R}$ ,

$F :: \mathbb{R} \rightarrow \mathbb{R}$

$F(x) = c$

$F(a) =$

$|a < x = -\mu(a, x] + F(x)$

$|a > x = \mu(x, a] - F(x),$

**Assume**  $a, b \in \mathbb{R} : b > a,$

$\partial F \rightsquigarrow F(b) - F(a) = \mu(a, b] \geq 0 \rightsquigarrow F(b) \geq F(a);$

$F : \text{Increasing}(\mathbb{R}, \mathbb{R}),$

**Assume**  $a : \mathbb{N} \rightarrow \mathbb{R} : a \downarrow A,$

**Assume**  $n \in \mathbb{N},$

$(A, a] \downarrow \emptyset \rightsquigarrow \lim_{n \rightarrow \infty} F(a_n) - F(A) = \lim_{n \rightarrow \infty} \mu(A, a_n] = 0 \rightsquigarrow \lim_{n \rightarrow \infty} F(a_n) = F(A);$

$F : \text{RightContinuous}(\mathbb{R}, \mathbb{R}),$

$F : \text{DistributionFunction}(\mathbb{R});; \square$

$\text{toDistribution} :: \text{Lebesgue-Stieltjes}(\overset{\infty}{\mathbb{R}}) \rightarrow \text{DistributionFunction}(\overset{\infty}{\mathbb{R}})$   
 $\text{toDistribution}(\mu) = F_\mu := \text{MeasureAsDistribution}(\mu, 0, 0)$

**DistributionAsMeasure** ::  $\forall F : \text{DistributionFunction}(\mathbb{R}) . \exists ! \mu : \text{Lebesgue-Stieltjes}(\mathbb{R}) :$   
 $: \text{MeasureAsDistribution}(\mu, 0, F(0)) = F$

**Proof** =

**Assume**  $F : \text{DistributionFunction}(\mathbb{R})$

$\mu :: \text{Preborel} \rightarrow \mathbb{R}_+$

$\mu(\emptyset) = 0$

$\mu(\mathbb{R}) = \lim_{x \rightarrow \infty} F(x) - \lim_{x \rightarrow -\infty} F(x)$

$\mu(a, \infty] = \lim_{x \rightarrow \infty} F(x) - F(a)$

$\mu(\infty, a] = F(a) - \lim_{x \rightarrow -\infty} F(x)$

$\mu(a, b] = F(b) - F(a)$

$\mu\left(\bigsqcup_{i=1}^n I_n\right) = \sum_{i=1}^n \mu(I_n)$

**Assume**  $n \in \mathbb{N}$ ,

**Assume**  $I : \text{DisjointElems}(\text{Preborel}, A)$ ,

$\delta\mu \rightsquigarrow \mu\left(\bigsqcup_{i=1}^n I_n\right) = \sum_{i=1}^n \mu(I_n);$

$\mu : \text{CountablyAdditive}(\mathbb{R}, \text{Preborel})$ ,

**Assume**  $A : \mathbb{N} \rightarrow \text{Preborel} : A \uparrow \alpha : \alpha \in \text{Preborel}$

$\bigsqcup_{i=1}^n I_n := \alpha$

$\mu\left(\bigsqcup_{i=1}^n I_n\right) = \sum_{k=1}^n \mu(I_k);$

**Assume**  $k \in n$ ,

$B^k := A \cap I_k$ ,

$a^k := \inf B^k$ ,

$b^k := \sup B^k$ ,

$C^k := (a^k, b^k) \setminus B_k$ ,

$\lim_{m \rightarrow \infty} \mu(B_m^k) = \lim_{m \rightarrow \infty} \mu(a_m^k, b_m^k] - \mu(C_m^k) = F(\lim_{m \rightarrow \infty} b_m^k) - \lim_{m \rightarrow \infty} F(a_m^k),$

$F : \text{RightContinuos} \rightsquigarrow \lim_{m \rightarrow \infty} \mu(B_m^k) = F(\lim_{m \rightarrow \infty} b_m^k) - F(\lim_{m \rightarrow \infty} a_m^k) = \mu(I_k);$

$\lim_{m \rightarrow \infty} \mu(A_m) = \sum_{k=1}^n \lim_{m \rightarrow \infty} \mu(B_m^k) = \sum_{k=1}^n \mu(I_n) = \mu(\alpha);$

$\mu : \text{Extendible}(\mathbb{R}, \text{Preborel})$ ,

$\lim_{n \rightarrow \infty} (-n, n] = \mathbb{R} \rightsquigarrow (\mu, \text{Preborel}) : \sigma\text{-Finite}(\mathbb{R}),$

$\lambda := \text{CarathedoryExtension}(\mu),$   
 $\text{Assume } I : \mathcal{B}(\mathbb{R}) : I : \text{Bounded}(\mathbb{R}),$   
 $\text{Bounded}(\mathbb{R}) \leadsto \exists a, b \in \mathbb{R} : a < b : I \subset (a, b] \text{ Extract},$   
 $\lambda(I) \leq \lambda(a, b] = F(b) - F(a) < \infty;$   
 $\lambda : \text{Lebesgue-Stieltjes}(\mathbb{R}),$   
 $j\lambda \leadsto \text{MeasureAsDistribution}(\lambda, 0, F(0)) = F; \square$

$\text{toMeasure} :: \text{DistributionFunction}(\mathbb{R}) \rightarrow \text{Lebesgue-Stieltjes}(\mathbb{R})$   
 $\text{toMeasure}(F) = \mu_F := \text{DistributionAsMeasure}(F)$

$\text{LebesgueMeasure} :: \text{Lebesgue-Stieltjes}(\mathbb{R})$   
 $\text{LebesgueMeasure} = \lambda := \text{ToMeasure}(\text{id})$

$\text{LebesgueMesurable} :: ??\mathbb{R}$   
 $\text{LebesgueMesurable} = \overline{\mathcal{B}}(\mathbb{R}) := \mathcal{B}(\mathbb{R}) \cup \{A \subset \mathbb{R} : \exists B \in \mathcal{B}(\mathbb{R}) : A \subset B : \lambda(B) = 0\}$

$\text{Lebesgue-Stieltjes}(\mathbb{R}^n) :: ?\text{Measure}(\mathbb{R}^n, \mathcal{B}(\mathbb{R}^n))$   
 $\mu : \text{Lebesgue-Stieltjes}(\mathbb{R}^n) \iff \forall a, b \in \mathbb{R}^n : a \prec b . \mu(a, b] < \infty$

$\text{DistributionFunction}(\mathbb{R}^n) :: \mathbb{R}^n \rightarrow \mathbb{R}$   
 $F : \text{DistributionFunction}(\mathbb{R}^n) :: \forall m \in \mathbb{N} . \forall a \in \mathbb{R}^{n-1} .$   
 $\quad . \Lambda x \in \mathbb{R} . F(a_{1 \dots (m-1)} \oplus x \oplus a_{m \dots n+1}) : \text{DistributionFunction}(\mathbb{R})$

$\text{Difference} :: (\mathbb{R}^n \rightarrow \mathbb{R}) \rightarrow \mathbb{R} \rightarrow \mathbb{R} \rightarrow n \rightarrow \mathbb{R}^{n-1} \rightarrow \mathbb{R}$   
 $\text{Difference}(F, a, b, m) = \Delta_{b,a}^m F :=$   
 $\quad := \Lambda x \in \mathbb{R} . F(x_{1 \dots (m-1)} \oplus b \oplus a_{m \dots n+1}) - F(x_{1 \dots (m-1)} \oplus a \oplus a_{m \dots n+1})$

$\text{toMeasure} :: \text{DistributionFunction}(\mathbb{R}^n) \rightarrow \text{Lebesgue-Stieltjes}(\mathbb{R}^n)$   
 $\text{toMeasure}(F) = \mu_F := \text{CaratheodoryExtension}(\mu)$   
 $\text{where}$   
 $\mu(a, b] = \left( \bigcirc_{i=1}^n \Delta_{a_i, b_i}^i \right) F$   
 $\dots$

$\text{CompactApproximationI} :: \forall \mu : \text{FiniteMeasure}(\mathbb{R}^n, \mathcal{B}(\mathbb{R}^n)) . \forall A \in \mathcal{B}(\mathbb{R}^n) .$   
 $\quad . \mu(A) = \sup \{ \mu(K) \mid K : \text{Compact}(\mathbb{R}^n) : K \subset A \}$   
 $\text{Proof} =$   
 $\text{Assume } (\mu, \mathcal{B}(\mathbb{R}^n)) : \sigma\text{-Finite}(\mathbb{R}^n) : \mu : \text{FiniteMeasure},$   
 $G := \left\{ A \in \mathcal{B}(\mathbb{R}^n) : \mu(A) = \sup \{ \mu(K) \mid K : \text{Compact}(\mathbb{R}^n) : K \subset A \} \right\},$   
 $\text{Assume } A \uparrow_G \alpha,$   
 $\text{Assume } \epsilon \in \mathbb{R}_{++},$   
 $\text{Assume } n \in \mathbb{N},$   
 $\partial A(n) \rightsquigarrow A_n \in G,$   
 $A_n \in G, \partial \sup(A_n, \epsilon) \rightsquigarrow \exists K \in \text{Compact} : K \subset B : \mu(A_n) \leq \mu(K) + \epsilon \text{ Extract as } K_n,$   
 $A_n \subset \alpha, K_n \subset \alpha \rightsquigarrow K_n \subset \alpha,$   
 $K_n \subset \alpha \rightsquigarrow \mu(K_n) \leq \mu(\alpha);$   
 $\lim_{n \rightarrow \infty} \mu(K_n) \leq \mu(\alpha) = \lim_{n \rightarrow \infty} \mu(A_n) \leq \lim_{n \rightarrow \infty} \mu(K_n) + \epsilon;$   
 $\mu(K) \uparrow \mu(\alpha) \rightsquigarrow \alpha \in G;$   
 $\text{Assume } A \downarrow_G \alpha,$   
 $\text{Assume } \epsilon \in \mathbb{R}_{++},$   
 $\text{Assume } n \in \mathbb{N},$   
 $\partial A(n) \rightsquigarrow A_n \in G,$   
 $A_n \in G, \partial \sup(A_n, \epsilon 2^{-n}) \rightsquigarrow \exists K \in \text{Compact} : K \subset B : \mu(A_n) \leq \mu(K) + \epsilon 2^{-n} \text{ Extract as } K_n;$   
 $C := \bigcap_{n=1}^{\infty} K_n,$   
 $\partial C \rightsquigarrow \forall n \in \mathbb{N} . C \subset A_n \rightsquigarrow C \subset A,$   
 $\mu(\alpha) - \mu(C) = \mu(\alpha \setminus C) \leq \mu \left( \bigcup_{n=1}^{\infty} (B_n \setminus K_n) \right) \leq \sum_{n=1}^{\infty} \mu(B_n \setminus K_n) \leq \epsilon;$   
 $\mu(\alpha) = \sup \{ \mu(K) \mid K : \text{Compact}(\mathbb{R}^n) : K \subset A \} \rightsquigarrow \alpha \in G;$   
 $G : \text{MonotoneClass},$   
 $\text{Assume } A \in \text{Preborel}(\mathbb{R}^n),$   
 $\partial \text{Preborel}(A) \rightsquigarrow \exists n \in \mathbb{N} : \exists (a, b] : n \rightarrow \text{Halfinterval}(\mathbb{R}^n) : A = \bigsqcup_{k=1}^n (a_k, b_k] \text{ Extract},$   
 $\text{Assume } k \in n,$   
 $x := \Lambda m \in \mathbb{N} . a_k + (b_k - a_k)/(2n),$   
 $\forall m \in \mathbb{N} . [x_m, b_k] : \text{Compact}(\mathbb{R}^n),$   
 $[x, b_k] \uparrow (a_k, b_k] \rightsquigarrow (a_k, b_k] \in G;$   
 $G : \text{MonotoneClass}(\mathbb{R}^n) \rightsquigarrow A \in G;$   
 $\text{Preborel} \subset G,$   
 $G : \text{MonotoneClass}(\mathbb{R}^n) \rightsquigarrow \mathcal{B}(\mathbb{R}^n) \subset G; \square$

**CompactApproximationII** ::  $\forall (\mathcal{B}(\mathbb{R}^n), \mu) : \sigma\text{-Finite}(\mathbb{R}^n) . \forall A \in \mathcal{B}(\mathbb{R}^n) .$   
 $\mu(A) = \sup \{ \mu(K) \mid K : \text{Compact}(\mathbb{R}^n) : K \subset A \}$

**Proof** =

**Assume**  $(\mathcal{B}(\mathbb{R}^n), \mu) : \sigma\text{-Finite}(\mathbb{R}^n),$

$A := \exists \sigma\text{-Finite}(\mathbb{R}^n) (\mathcal{B}(\mathbb{R}^n), \mu) : \mathbb{N} \rightarrow \mathcal{B}(\mathbb{R}^n) : \mathbb{R}^n = \bigcup_{n=1}^{\infty} A_n : \forall n \in \mathbb{N} . \mu(A_n) < \infty,$

**Assume**  $B \in \mathcal{B}(\mathbb{R}^n),$

**Assume**  $n \in \mathbb{N},$

$\beta_n := B \cap \bigcup_{k=1}^n A_k$

$M_n := \Lambda X \in \mathcal{B}(\mathbb{R}^n) . \mu(X \cap \beta_n)$

$K_n := \text{CompactApproximationI}(M_n, B, 1/n);$

$\lim_{n \rightarrow \infty} \mu(K_n) \leq \mu(B) = \lim_{n \rightarrow \infty} M_n(B) \leq \lim_{n \rightarrow \infty} M_n(K_n) + 1/n = \lim_{n \rightarrow \infty} M_n(K_n) \leq \lim_{n \rightarrow \infty} \mu(K_n) \rightsquigarrow$   
 $\rightsquigarrow \mu(K) \uparrow \mu(B); ; \square$

**OpenApproximationI** ::  $\forall \mu : \text{FiniteMeasure}(\mathbb{R}^n, \mathcal{B}(\mathbb{R}^n)) . \forall A \in \mathcal{B}(\mathbb{R}^n) .$

$\mu(A) = \sup \{ \mu(K) \mid K : \text{Compact}(\mathbb{R}^n) : K \subset A \}$

**Proof** =

**Assume**  $\mu : \text{FiniteMeasure}(\mathbb{R}^n, \mathcal{B}(\mathbb{R}^n)),$

**Assume**  $A \in \mathcal{B}(\mathbb{R}^n),$

$\exists \sigma\text{-Algebra}(\mathbb{R}^n, \mathcal{B}(\mathbb{R}^n)) (A) \rightarrow A^c \in \mathcal{B}(\mathbb{R}^n),$

$K := \text{CompactApproximationI}(\mu) (A^c) : \mathbb{N} \rightarrow \text{Compact}(\mathbb{R}^n) : \mu(K) \uparrow \mu(A^c),$

$\lim_{n \rightarrow \infty} \mu(K_n^c) = \lim_{n \rightarrow \infty} \mu(\Omega) - \mu(K_n) = \mu(\Omega) - \lim_{n \rightarrow \infty} \mu(K_n) = \mu(\Omega) - \mu(A^c) = \mu(A) \rightsquigarrow$   
 $\rightsquigarrow \mu(K^c) \downarrow \mu(A),$

$\exists \text{Closed}(\mathbb{R}^n)(\exists \text{Compact}(\mathbb{R}^n)(K)) \rightsquigarrow K^c : \mathbb{N} \rightarrow \text{Open}(\mathbb{R}^n)$

**OpenApproximationII** ::  $\forall (\mathcal{B}(\mathbb{R}^n), \mu) : \sigma\text{-Finite}(\mathbb{R}^n) : \mu : \text{Lebesgue-Stieltjes}(\mathbb{R}^n) . \forall A \in \mathcal{B}(\mathbb{R}^n) .$   
 $\mu(A) = \inf \{ \mu(U) \mid U : \text{Open}(\mathbb{R}^n) : A \subset U \}$

**Proof** =

**Assume**  $(\mathcal{B}(\mathbb{R}^n), \mu) : \sigma\text{-Finite}(\mathbb{R}^n) : \mu : \text{Lebesgue-Stieltjes}(\mathbb{R}^n)$

**Assume**  $A \in \sigma\text{-Algebra}(\mathbb{R}^n, \mathcal{B}(\mathbb{R}^n))$

$B := \exists \sigma\text{-Finite}(\mathbb{R}^n)(\mathcal{B}(\mathbb{R}^n), \mu) : \mathbb{N} \rightarrow \mathcal{B}(\mathbb{R}^n) : \mathbb{R}^n = \bigsqcup_{n=1}^{\infty} B_n :$

$: \forall n \in \mathbb{N} . \mu(B_n) < \infty : B_n : \text{Bounded}(\mathbb{R}^n),$

$C := \exists \text{Lebesgue-Stieltjes}(\mathbb{R}^n)(B) : \prod n \in \mathbb{N} . \mathcal{U}(B_n) : \forall n \in \mathbb{N} . \mu(C_n) < \infty,$

**Assume**  $\epsilon \in \mathbb{R}_{++},$

**Assume**  $n \in \mathbb{N},$

$M_n := \Lambda S \in \mathcal{B}(\mathbb{R}^n) . \mu(S \cap C_n) : \text{FiniteMeasure}(\mathbb{R}^n, \mathcal{B}(\mathbb{R}^n))$

$U_n := C_n \cap \text{OpenApproximationI}(M_n, A \cap B_n, \epsilon/2^n) : \text{Open}(\mathbb{R}^n) : M_n(U_n) - M_n(A \cap B_n) \leq \epsilon/2^n : A \subset U_n;$

$O := \bigcup_{n=1}^{\infty} U_n,$

$$\begin{aligned} \mu(A) &\leq \mu(O) \leq \sum_{n=1}^{\infty} \mu(U_n) = \sum_{n=1}^{\infty} M_n(U_n) \leq \\ &\leq \sum_{n=1}^{\infty} M_n(A \cap B_n) + \epsilon = \sum_{n=1}^{\infty} \mu(A \cap B_n) + \epsilon = \mu(A) + \epsilon; \end{aligned}$$

$\mu(A) = \inf \{ \mu(U) \mid U : \text{Open}(\mathbb{R}^n) : A \subset U \} ; ; \square$

**Example 1.3.1**

$$F(x) := \begin{cases} 0 & x < -1 \\ 1+x & -1 \leq x < 0 \\ 2+x^2 & 0 \leq x < 2 \\ 9 & x \geq 2 \end{cases} : \text{DistributionFunction}(\mathbb{R}, \mathcal{B}\mathbb{R})$$

$\mu := \text{ToMeasure} : \text{Measure}(\mathbb{R}, \mathcal{B}\mathbb{R})$

$\mu\{2\} = 3,$

$\mu[-1/2, 3) = 9 - 1/2 = 8.5,$

$\mu(-1, 0] \cup (1, 2) = 2 + (6 - 3) = 5$

$\mu[0, 1/2) \cup (1, 2] = (2.25 - 2) + (9 - 3) = 6.25$

$\mu(-\infty, -1/2) \cup (1/2, \infty) = 1/2 + (9 - 2.25) = 7.25$

**Example 1.3.2**

**Assume**  $\mu : \text{Lebesgue-Stieltjes}(\mathbb{R}) : F_\mu \in \mathcal{M}_{\text{TOP}}(\mathbb{R}, \mathbb{R}),$

**Assume**  $N : \text{Countable}(\mathbb{R}),$

$$\mu(N) = \sum_{n \in N} \mu\{n\} = \sum_{n \in N} F(n) - \lim_{x \rightarrow n-0} F(x) = 0;$$

**Assume**  $\mu = U[0, 1],$

$$\mu([0, 1] \cap \mathbb{Q}) = 1,$$

$$\mu\left([0, 1]^{\mathbb{C}}\right) = 0,$$

**Assume**  $A : \mathcal{BR} ! \text{Dense}(\mathbb{R}),$

$$I := \partial A : \text{Open}(\mathbb{R}) : I \subset A^{\mathbb{C}},$$

$$\lambda\left(A^{\mathbb{C}}\right) \geq \lambda(I) \geq 0;;$$

**BorelTranslationInvariance** ::  $\forall A \in \mathcal{BR}^n . \forall a \in \mathbb{R}^n . a + A \in \mathcal{BR}^n \wedge -A \in \mathcal{BR}^n$

**Proof** =

$$G := \{A \in \mathcal{BR}^n : \forall a \in \mathbb{R}^n : a + A \in \mathcal{BR}^n \wedge -A \in \mathcal{BR}^n\} : ?\mathcal{BR}^n,$$

**Assume**  $(a, b] \in \text{Halfinterval}(\mathbb{R}^n)$

**Assume**  $r \in \mathbb{R}^n,$

$$r + (a, b] = (a + r, b + r] : \text{Halfinterval}(\mathbb{R}^n) \rightsquigarrow r + (a, b] : \mathcal{BR}^n \text{ as } (1),$$

$$-(a, b] = [b, a) \in \mathcal{BR}^n \text{ as } (2),$$

$$(1, 2) \rightsquigarrow (a, b] \in G,$$

$$\text{HalfInterval}(\mathbb{R}^n) \subset G,$$

**Assume**  $X : \mathbb{N} \rightarrow G,$

**Assume**  $a \in \mathbb{R}^n,$

$$f := \Lambda v \in \mathbb{R}^n . v + a : \text{ISO}_{\text{SET}}(\mathbb{R}^n, \mathbb{R}^n),$$

$$g := \Lambda v \in \mathbb{R}^n . -v : \text{ISO}_{\text{SET}}(\mathbb{R}^n, \mathbb{R}^n),$$

$$-A^{\mathbb{C}} = g\left(A^{\mathbb{C}}\right) = g(A^{\mathbb{C}}) = (-A)^{\mathbb{C}} \text{ as } (1),$$

$$\partial(G)(X) \rightsquigarrow -A \in \mathcal{BR}^n \rightsquigarrow_{(1)} -A^{\mathbb{C}} \in \mathcal{BR}^n,$$

$$a + A^{\mathbb{C}} = f\left(A^{\mathbb{C}}\right) = f(A)^{\mathbb{C}} = (a + A)^{\mathbb{C}} \text{ as } (2),$$

$$\partial(G)(A) \rightsquigarrow a + A \in \mathcal{BR}^n \rightsquigarrow_{(2)} a + A^{\mathbb{C}} \in \mathcal{BR}^n,$$

$$-\bigcup_{n=1}^{\infty} X_n = g\left(\bigcup_{n=1}^{\infty} X_n\right) = \bigcup_{n=1}^{\infty} g(X_n) = \bigcup_{n=1}^{\infty} -X_n \text{ as } (3)$$

$$\partial(G)(X) \rightsquigarrow \forall n \in \mathbb{N} . -X_n \in \mathcal{BR}^n \rightsquigarrow_{(3)} -\bigcup_{n=1}^{\infty} X_n \in \mathcal{BR}^n,$$



$$a + \bigcup_{n=1}^{\infty} X_n = f \left( \bigcup_{n=1}^{\infty} X_n \right) = \bigcup_{n=1}^{\infty} f(X_n) = \bigcup_{n=1}^{\infty} a + X_n \text{ as (4)}$$

$$\mathfrak{d}(G)(X) \rightsquigarrow \forall n \in \mathbb{N} . a + X_n \in \mathcal{B}\mathbb{R}^n \rightsquigarrow_{(4)} a + \bigcup_{n=1}^{\infty} X_n \in \mathcal{B}\mathbb{R}^n;$$

$$A^{\mathfrak{c}} \in G,$$

$$\bigcup_{n=1}^{\infty} X_n \in G;$$

$$G : \sigma\text{-Algebra}(\mathbb{R}^n) \rightsquigarrow \mathcal{B}\mathbb{R}^n = \sigma\{\text{Halfinterval}\} \subset G,$$

$$\mathfrak{d}G \rightsquigarrow G \subset \mathcal{B}\mathbb{R} \rightsquigarrow G = \mathcal{B}\mathbb{R}^d \square$$

$$\text{LebesgueTranslationInvariance} :: \forall A \in \mathcal{B}\mathbb{R}^n . \forall a \in \mathbb{R}^n . \lambda(a + A) = \lambda(A)$$

**Proof** =

$$G := \{A \in \mathcal{B}\mathbb{R}^{\infty^n} : \forall a \in \mathbb{R}^n . \lambda(A) + \lambda(a + A)\}$$

$$\text{Assume } (a, b] : \text{Halfinterval},$$

$$\text{Assume } r \in \mathbb{R}^n,$$

$$\lambda(r + (a, b]) = \lambda(a + r, b + r] = b + r - a - r = b - a = \lambda(a, b];;$$

$$\text{Halfinterval} \subset G \text{ as (1),}$$

$$\text{Assume } A \uparrow_G \alpha,$$

$$\text{Assume } B \downarrow_G \beta,$$

$$\text{Assume } a \in \mathbb{R}^n,$$

$$\lambda(a + \alpha) = \lim_{n \rightarrow \infty} \lambda(a + A_n) = \lim_{n \rightarrow \infty} \lambda(A_n) = \lambda(\alpha),$$

$$\lambda(a + \beta) = \lim_{n \rightarrow \infty} \lambda(a + B_n) = \lim_{n \rightarrow \infty} \lambda(B_n) = \lambda(\beta);$$

$$G : \text{MonotoneClass}(\mathbb{R}^n) \rightsquigarrow_{(1)} G = \mathcal{B}\mathbb{R}^n \square$$

$$\text{InvariantIsLebesgue} :: \forall \mu : \text{Lebesgue-Stieltjes}(\mathbb{R}^n) .$$

$$\text{if } \forall A \in \mathcal{B}\mathbb{R}^n . \forall a \in \mathbb{R}^n . \mu(A + a) = \mu(A) \text{ then } \exists c \in \mathbb{R}_+ : \mu = c\lambda,$$

$$\text{Assume } \mu : \text{Lebesgue-Stieltjes}(\mathbb{R}^n),$$

$$\text{Assume } (*) : A \in \mathcal{B}\mathbb{R}^n . \forall a \in \mathbb{R}^n . \mu(A + a) = \mu(A),$$

$$\text{Assume } I, J : \text{Halfcube} : \lambda(I) = \lambda(J) : \mu(I) \neq \mu(J),$$

$$\mathfrak{d}I, J \rightsquigarrow \exists v \in \mathbb{R}^n . I = J + a \text{ Extract,}$$

$$\mu(I) = \mu(J + a) =_* \mu(J) \rightsquigarrow \perp;$$

$$(1) : \forall I, J : \text{Halfcube} . \lambda(I) = \lambda(J) \Rightarrow \mu(I) = \mu(J),$$

$$c := \mu(0, 1] : \mathbb{R}_+,$$

$$\mathfrak{d}(\text{Measure})(\mu) \rightsquigarrow (1) : \mu|_{\text{Halfcube}} = c\lambda|_{\text{Halfcube}},$$

$$\mathcal{B}(\mathbb{R}^n) = \sigma(\text{Halfcube}(\mathbb{R}^n)) \rightsquigarrow_{(1)} \mu = c\lambda; ; \square$$

**Example 1.3.3**

$R := \{(x, y) \in \mathbb{R}^2 : x - y \in \mathbb{Q}\} : \mathbf{Eq}(\mathbb{R}),$

$B := \mathbf{eqclasses}(R),$

$A := \mathbf{choice}(B, [0, 1]),$

**Assume**  $r, q \in \mathbb{Q} : r \neq q,$

**Assume**  $a \in A + r \cap A + q,$

$\bar{\partial}(a) \rightsquigarrow \exists x, y \in A : x + r = a = y + q,$

$x - y = q - r \in \mathbb{Q} \rightsquigarrow x = y \rightsquigarrow q = r \rightsquigarrow \perp,$

$\forall r, q \in \mathbb{Q} . A + r \cap A + q = \emptyset,$

**Assume**  $A : \mathbf{Measurable}(\lambda),$

$$\bigcup_{q \in \mathbb{Q} \cap [0, 1]} q + A \subset [0, 2] \rightsquigarrow 2 \geq \lambda \left( \bigcup_{q \in \mathbb{Q} \cap [0, 1]} q + A \right) = \sum_{q \in \mathbb{Q} \cap [0, 1]} \lambda(q + A) = \sum_{q \in \mathbb{Q} \cap [0, 1]} \lambda(A) \rightsquigarrow \lambda(A) = 0,$$

$$\lambda(\mathbb{R}) = \bigcup_{q \in \mathbb{Q}} \lambda(A + q) = \bigcup_{q \in \mathbb{Q}} \lambda(A) = 0 \rightsquigarrow \perp,$$

$A ! \mathbf{LebesgueMeasurable}(\mathbb{R}^n) \square$

**Example 1.3.4**

$F := \Lambda(x, y) \in \mathbb{R}^2 . \mathbf{if } x + y > 1 \mathbf{ then } \ln(x + y) \mathbf{ else } 0,$

$$\begin{aligned} \mu_F(0.5, 2.5] &= F(0.5, 0.5) - F(0.5, 2.5) - F(2.5, 0.5) + F(2.5, 2.5) = \ln(5) - 2\ln(3) = \\ &= \ln(5) - \ln(9) < 0 \rightsquigarrow F ! \mathbf{DistributionFunction}(\mathbb{R}^2), \end{aligned}$$

**Example 1.3.5**

$F := \Lambda(x, y) \in \mathbb{R}^2 . \mathbf{if } x < 0 | y < 0 \mathbf{ then } 0 \mathbf{ else } xy + 1 : \mathbb{R}^2 \rightarrow \mathbb{R},$

$\mathbf{discont}(F) = [(0, 0), (\infty, 0)] \cup [(0, 0), (\infty)] ! \mathbf{Countable}$

## 1.5 Lebesgue-Stieltjes Measures in Multivariable Context [!]

## 1.6 Categorical Viewpoint: Boolean Algebras [!]

$\text{Boolean} :: ?\text{Commutative}$

$$B : \text{Boolean} \iff \forall b \in B . b^2 = b$$

$$\mathcal{F} : \sigma\text{-Algebra} (()) \Rightarrow (\mathcal{F}, \cap, \triangle) : \text{Boolean}$$

$$\sigma\text{-Ideal} :: \prod \mathcal{F} : \sigma\text{-Algebra} (\Omega) . ?\text{Ideal}(\mathcal{F})$$

$$N : \sigma\text{-Ideal} \iff \forall A : \mathbb{N} \rightarrow N . \bigcup_{n=1}^{\infty} A_n \in N$$

$$\text{ZeroSpace} := \sum \Omega : \text{Set} . \mathcal{F} : \sum \sigma\text{-Algebra} (\Omega) . \sigma\text{-Ideal} (\mathcal{F})$$

$\text{Localizable} :: ?\text{ZeroSpace}$

$$(\Omega, \mathcal{F}, N) : \text{Localizable} \iff \forall A : ?\frac{\mathcal{F}}{N} . \sup A \in \frac{\mathcal{F}}{N}$$

$$\text{IdealMeasure} :: \prod (\Omega, \mathcal{F}, N) : \text{ZeroSpace} . ?\text{Measure}(\Omega, \mathcal{F})$$

$$\mu : \text{IdealMeasure} \iff \forall A \in \mathcal{F} . \mu(A) = 0 \iff A \in N$$

## 2 Lebesgue Integration

### 2.1 Measurable Functions

**Measurable** ::  $\prod (\Omega, \mathcal{F}), (\Omega', \mathcal{F}') . ?\Omega \rightarrow \Omega'$

$f : \text{Measurable} \iff \forall A \in \mathcal{F}' . f^{-1}(A) \in \mathcal{F}$

**BorelMeasurableCriterion** ::  $\forall f : (\Omega, \mathcal{F}) \rightarrow \mathbb{R} : \forall c \in \mathbb{R} . f^{-1}(-\infty, c) \in \mathcal{F} . f : \text{Measurable}(\Omega, \mathcal{F})$

**Proof** =

**Assume**  $f : (\Omega, \mathcal{F}) \rightarrow \mathbb{R} : \forall c \in \mathbb{R} . f^{-1}(-\infty, c) \in \mathcal{F}$ ,

**Assume**  $A \in \mathcal{B}\mathbb{R}$ ,

$\mathcal{B}\mathbb{R} = \sigma(\{(-\infty, c] \mid c \in \mathbb{R}\})$ ,

$(c, T) := \mathcal{J}\sigma(\{(-\infty, c] \mid c \in \mathbb{R}\})(A) : \sum \mathbb{N} \rightarrow \mathbb{R} . \text{SetAlgTransform} : T((-\infty, c]) = A$ ,

**Assume**  $n \in \mathbb{N}$ ,

$(1) := \mathfrak{D}(f)((-\infty, c_n]) : f^{-1}(-\infty, c_n] \in \mathcal{F}$ ,

$(1) : \forall n \in \mathbb{N} . f^{-1}(-\infty, c_n] \in \mathcal{F}$ ,

$f^{-1}(A) = f^{-1}(T((-\infty, c]) = T(f^{-1}[(-\infty, c_n)]_{n=1}^{\infty})) \in_{(1)} \mathcal{F}$ ;

$f : \text{Measurable}(\Omega, \mathcal{F}); \square$

**MeasurableMax** ::  $\forall f, g : \text{Measurable}(\Omega, \mathcal{F}) . \max(g, f) : \text{Measurable}(\Omega, \mathcal{F})$

**Proof** =

**Assume**  $f, g : \text{Measurable}(\Omega, \mathcal{F})$ ,

**Assume**  $c \in \mathbb{R}$ ,

$\max(f, g)^{-1}(-\infty, c] = f^{-1}(-\infty, c] \cap g^{-1}(-\infty, c] \in \mathcal{F}$ ;

**BorelMeasurableCriterion**  $\leadsto \max(g, f) : \text{Measurable}(\Omega, \mathcal{F}); \square$

**MeasurableMin** ::  $\forall f, g : \text{Measurable}(\Omega, \mathcal{F}) . \min(g, f) : \text{Measurable}(\Omega, \mathcal{F})$

**Proof** =

**Assume**  $f, g : \text{Measurable}(\Omega, \mathcal{F})$ ,

**Assume**  $c \in \mathbb{R}$ ,

$\min(f, g)^{-1}(-\infty, c] = f^{-1}(-\infty, c] \cup g^{-1}(-\infty, c] \in \mathcal{F}$ ;

**BorelMeasurableCriterion**  $\leadsto \min(g, f) : \text{Measurable}(\Omega, \mathcal{F}); \square$

**IndicatorMeasurable** ::  $\forall f : \text{Measurable}(\Omega, \mathcal{F}) \forall A \in \mathcal{F} . I_A f : \text{Measurable}(\Omega, \mathcal{F})$ ,

**Proof** =

**Assume**  $f : \text{Measurable}(\Omega, \mathcal{F})$

**Assume**  $A \in \mathcal{F}$

**Assume**  $B \in \mathcal{B}\mathbb{R}$ ,

(1) := **LawOfExcludeMiddle**( $B, 0$ ) :  $0 \in B \mid 0 \notin B$

**Assume** (2) :  $0 \in B$ ,

$I_A f^{-1}(B) = (A \cap f^{-1}(B)) \cup A^c \in \mathcal{F}$

**Assume** (2) :  $0 \notin B$

$I_A f^{-1}(B) = (A \cap f^{-1}(B)) \in \mathcal{F}$ ;

(1)  $\leadsto f^{-1}(B) \in \mathcal{F}$ ;

$I_A f : \text{Measurable}(\Omega, \mathcal{F}) \square$

**MeasurableConvergence** ::  $\forall f : \mathbb{N} \rightarrow \text{Measurable}(\Omega, \mathcal{F}) . \forall \phi : f_n \xrightarrow{p} \phi . \phi : \text{Measurable}(\Omega, \mathcal{F})$ ,

**Proof** =

**Assume**  $f : \mathbb{N} \rightarrow \text{Measurable}(\Omega, \mathcal{F})$ ,

**Assume**  $\phi : f_n \xrightarrow{p} \phi$ ,

**Assume**  $c \in \mathbb{R}$ ,

$\phi^{-1}(-\infty, c) = \{\omega \in \Omega : \phi(\omega) \leq c\} = \{\omega \in \Omega : \lim_{n \rightarrow \infty} f_n(\omega) \leq c\} =$

$= \bigcup_{n \in \mathbb{N}} \in \mathbb{N} \{\omega \in \Omega : \exists K : \text{Infinite}(\mathbb{N}) : \forall k \in K . f_k(\omega) \leq c - 1/n\} =$

$= \bigcup_{n \in \mathbb{N}} \liminf_{n \in \mathbb{N}} \{\omega \in \Omega : f_n(\omega) \leq c - 1/n\} = \bigcup_{n=1}^{\infty} \bigcup_{k=1}^{\infty} \bigcap_{m=k}^{\infty} f_m^{-1}(-\infty, c + 1/n] \in \mathcal{F}$ ;

**BorelMeasurableCriterion**  $\leadsto \phi : \text{Measurable}(\Omega, \mathcal{F})$ ; ;  $\square$

**Simple** ::  $? \text{Measurable}(\Omega, \mathcal{F}) \left( \mathbb{R}^{\infty}, \mathcal{B} \left( \mathbb{R}^{\infty} \right) \right)$

$f : \text{Simple} \iff \exists r \in \mathbb{N} : \exists A : r \rightarrow \mathcal{F} : \exists a : r \rightarrow \mathcal{F} : f = \sum_{k=1}^r a_k I_{A_k}$

**SimpleApproximationI** ::  $\forall f : \text{Measurable}(\Omega, \mathcal{F}) : f > 0 .$

$\exists S : \mathbb{N} \rightarrow \text{Simple}(\Omega, \mathcal{F}) : \forall n \in \mathbb{N} . S_n > 0 : S \xrightarrow{p} f$

**SimpleApproximationII** ::  $\forall f : \text{Measurable}(\Omega, \mathcal{F}) .$

$\exists S : \mathbb{N} \rightarrow \text{Simple}(\Omega, \mathcal{F}) : \forall n \in \mathbb{N} . |S_n| \leq |f| : S \xrightarrow{p} f$

**MeasurableAlgebra** ::  $\forall f, g : \text{Measurable}(\Omega, \mathcal{F}) . f + g, fg, f - g, f/g : \text{Measurable}(\Omega, \mathcal{F}) . f + g, fg, f - g, f/g$

**MeasurableComposition** ::  $\forall f : \text{Measurable } A \ B \forall g : \text{Measurable } B \ C . f \circ g : \text{Measurable } A \ B$

**LebesgueMeasurableMap** ::  $?( \mathbb{R}^m, \mathcal{B}\mathbb{R}^m ) \rightarrow ( \mathbb{R}^n, \mathcal{B}\mathbb{R}^n )$

$f : \text{LebesgueMeasurableMap} \iff \forall A \in \mathcal{B}\mathbb{R}^m . f^{-1}A : \text{LebesgueMeasurable}(\mathbb{R}^n)$

**MultivariateMeasurability** ::  $\forall f : \Omega \rightarrow \mathbb{R}^n .$

$f : \text{Measurable}(\Omega, \mathcal{F})(\mathbb{R}^n, \mathcal{B}\mathbb{R}^n) \iff \forall i \in n . f^i : \text{Measurable}(\Omega, \mathcal{F})$

**Example 2.2.1**

**Assume**  $f : \text{Measurable}(\Omega, \mathcal{F}) : f > 0,$

$S : \mathbb{N} \rightarrow \text{Simple}(\Omega, \mathcal{F})$

$S_n := \sum_{k=1}^{n2^n} \frac{k-1}{2^n} \left[ f^{-1} \left[ \frac{k-1}{2^n}, \frac{k}{2^n} \right) \right]_{\in} + n f^{-1}[n, \infty)$

**Assume**  $f : \text{Bounded} \rightsquigarrow \exists b : \forall \omega \in \Omega . f(\omega) \leq \Omega$  **Extract**,

**Assume**  $\epsilon \in \mathbb{R}_{++},$

$N = \lceil \max(b, \log_2 \epsilon^{-1}) \rceil,$

**Assume**  $n \in \mathbb{N} : n \geq N,$

**Assume**  $\omega \in \Omega,$

$f(\omega) - S_n(\omega) \leq \frac{1}{2^n} \leq \frac{1}{2^N} \leq \epsilon; ; ;$

$f : \text{Bounded} \Rightarrow S \Rightarrow f$

**Assume**  $\epsilon \in \mathbb{R}_{++},$

**Assume**  $\omega \in \Omega,$

$N = \lceil \max(f(\omega), \log_2 \epsilon^{-1}) \rceil,$

**Assume**  $n \in \mathbb{N} : n \geq N,$

$f(\omega) - S_n(\omega) \leq \frac{1}{2^n} \leq \frac{1}{2^N} \leq \epsilon; ; ;$

$S \xrightarrow{p} f; \square$

**ConditionalMeasurable** ::  $\forall f, g : \text{Measurable}(\Omega, \mathcal{F})(R, \mathcal{B}) . \forall A \in \mathcal{F} .$

$\Lambda \omega \in \Omega . \text{if } \omega \in A \text{ then } f(\omega) \text{ else } g(\omega) : \text{Measurable}(\Omega, \mathcal{F})(R, \mathcal{B})$

**Proof** =

**Assume**  $f, g : \text{Measurable}(\Omega, \mathcal{F})(R, \mathcal{B}),$

**Assume**  $A \in \mathcal{F},$

**Assume**  $h := \Lambda \omega \in \Omega . \text{if } \omega \in A \text{ then } f(\omega) \text{ else } g(\omega) : \Omega \rightarrow R$

**Assume**  $B \in \mathcal{B},$

$h^{-1}(B) = (f^{-1}(B) \cap A) \cup (g^{-1}(B) \cap A^c) \in \mathcal{F};$

**MeasurableInf** ::  $\forall f : \mathbb{N} \rightarrow \text{Measurable}(\Omega, \mathcal{F}) . \inf_{n \in \mathbb{N}} f_n, \sup_{n \in \mathbb{N}} f_n : \text{Measurable}(\Omega, \mathcal{F})$

**Proof** =

**Assume**  $f : \mathbb{N} \rightarrow \text{Measurable}(\Omega, \mathcal{F})$

**Assume**  $c \in \mathbb{R}$ ,

$$\left( \inf_{n \in \mathbb{N}} f_n \right)^{-1} (-\infty, c] = \bigcup_{n=1}^{\infty} f_n^{-1} (-\infty, c] \in \mathcal{F},$$

$$\left( \sup_{n \in \mathbb{N}} f_n \right)^{-1} (-\infty, c] = \bigcap_{n=1}^{\infty} f_n^{-1} (-\infty, c] \in \mathcal{F};$$

**BorelMeasurableCriterion**  $\leadsto \inf_{n \in \mathbb{N}} f_n, \sup_{n \in \mathbb{N}} f_n : \text{Measurable}(\Omega, \mathcal{F}); \square$

**MeasurableAlmost** ::  $\forall (\Omega, \mathcal{F}, \mu) : \text{CompleteMeasure} . \forall f : \text{Measurable}(\Omega, \mathcal{F})(R, \mathcal{B}) .$   
 $. \forall A \in \mathcal{F} : \mu(A) = 0 . \forall g : \Omega \rightarrow R : f|_{A^c} = g|_{A^c} . g : \text{Measurable}(\Omega, \mathcal{F})(R, \mathcal{B})$

**Proof** =

**Assume**  $(\Omega, \mathcal{F}, \mu) : \text{CompleteMS}$ ,

**Assume**  $f : \text{Measurable}(\Omega, \mathcal{F})(R, \mathcal{B})$ ,

**Assume**  $A \in \mathcal{F} : \mu(A) = 0$ ,

**Assume**  $g : \Omega \rightarrow R : f|_{A^c} = g|_{A^c}$ ,

**Assume**  $B \in \mathcal{B}$ ,

$$g^{-1}(B) \cap A \subset A \leadsto \text{as } (\Omega, \mathcal{F}, \mu) : \text{CompleteMeasure} \leadsto g^{-1}(B) \cap A \in \mathcal{F}$$

$$g^{-1}(B) = (g^{-1}(B) \cap A) \cup (g^{-1}(B) \cap A^c) = (g^{-1}(B) \cap A) \cup (f^{-1}(B) \cap A^c) \in \mathcal{F};$$

$g : \text{Measurable}(\Omega, \mathcal{F})(R, \mathcal{B}); ; ; ; \square$

**ClosedUnion** ::  $\prod X : \text{TopologicalSpace} . ??X$

$$A : \text{ClosedUnion} \iff A : F_\sigma \iff \exists K : \mathbb{N} \rightarrow \text{Closed}(X) . A = \bigcup_{n=1}^{\infty} K_n$$

**DiscontTHM**  $\forall f : \mathbb{R}^n \rightarrow \mathbb{R}^m : \text{discont}(f) : F_\sigma$

**Proof** =

**Assume**  $f : \mathbb{R}^n \rightarrow \mathbb{R}^m$

$s :: \mathbb{R}^n \rightarrow \mathbb{R}$

$$s(p) = \sup \{ \lim_{n, m \rightarrow \infty} d(f(x_n), f(y_m)) \mid x, y : \mathbb{N} \rightarrow \mathbb{R}^n : \lim_{n \rightarrow \infty} x_n = a = \lim_{m \rightarrow \infty} y_m : f(x), f(y) : \text{Convergent} \}$$

**Assume**  $p \in \{p \in \mathbb{R}^n : s(p) < 1/n\}$ ,

**Assume** (1) :  $\forall U \in \mathcal{U}(p) . \exists u \in U : s(u) \geq 1/n$ ,

(1)  $\leadsto \exists x \in \mathbb{N} \rightarrow \mathbb{R}^n : \forall n \in \mathbb{N} . s(x_n) \geq 1/n$

**Assume**  $\epsilon \in \mathbb{R}_{++}$ ,

$\mathfrak{d}(x) \rightsquigarrow \exists a, b \in \mathbb{R}^n : d(a, p) < \epsilon : d(b, p) \leq \epsilon : d(f(a), f(b)) > 1/n + \epsilon;$

$s(p) \geq 1/n \rightsquigarrow \perp,$

$\exists U \in \mathcal{U}(p) : \forall u \in U . s(u) < 1/n$  **Extract as**  $U_p$ ;

$\{p \in \mathbb{R}^n : s(p) < 1/n\} = \bigcup_{p \in \mathbb{R}^n : s(p) < 1/n} U_p \rightsquigarrow \{p \in \mathbb{R}^n : s(p) < 1/n\} : \text{Open}(\mathbb{R}^n) \rightsquigarrow$

$\{p \in \mathbb{R}^n : s(p) \geq 1/n\} : \text{Closed}(\mathbb{R}^n);$

**discont**( $f$ ) :  $F_\sigma$   $\square$

**NoIrrationalDisconts** :  $\forall f : \mathbb{R} \rightarrow \mathbb{R} . \text{discont}(f) \neq \mathbb{Q}^\mathbb{C}$

**Assume**  $f : \mathbb{R} \rightarrow \mathbb{R} : \text{discont}(f) = \mathbb{Q}^\mathbb{C},$

**DiscontTHM**  $\rightsquigarrow \mathbb{Q}^\mathbb{C} : F_\sigma \rightsquigarrow \exists K : \mathbb{N} \rightarrow \text{Closed}(\mathbb{R}) : \mathbb{Q}^\mathbb{C} = \bigcup_{n=1}^{\infty} K_n$  **Extract**,

$\mathbb{Q} = \bigcap_{n=1}^{\infty} K_n^\mathbb{C} \rightsquigarrow \forall n \in \mathbb{N} . \mathbb{Q} \subset K_n^\mathbb{C},$

**Assume**  $n \in \mathbb{N},$

$\mathbb{Q} \subset K_n^\mathbb{C},$

$K_n : \text{Closed}(\mathbb{R}) \rightsquigarrow K_n^\mathbb{C} : \text{Open}(\mathbb{R}),$

$\mathbb{Q} : \text{Dense}(\mathbb{R}) \rightsquigarrow K_n^\mathbb{C} = \mathbb{R};$

$\mathbb{Q} = \mathbb{R} \rightsquigarrow \perp; \square$

**SteinhausLemma** ::  $\forall a : \mathbb{N} \rightarrow \mathbb{N} \rightarrow \mathbb{R} : \exists c \in \mathbb{R}_{++} : \forall n \in \mathbb{N} . \sum_{k=1}^{\infty} a_{n,k} = 1 \wedge$

$\wedge \sum_{k=1}^{\infty} |a_{n,k}| < c \wedge : \lim_{k \rightarrow \infty} x_{k,n} = 0 . \exists x : \mathbb{N} \rightarrow \{0, 1\} :$

$: \wedge n \in \mathbb{N} : \sum_{k=1}^{\infty} x_k a_{n,k} ! \text{Convergent} \left( \mathbb{R} \right)$

**Proof** =

**Assume**  $a : \mathbb{N} \rightarrow \mathbb{N} \rightarrow \mathbb{R} : \exists c \in \mathbb{R}_{++} : \forall n \in \mathbb{N} . \sum_{k=1}^{\infty} a_{n,k} = 1 \wedge \sum_{k=1}^{\infty} |a_{n,k}| < c \wedge : \lim_{k \rightarrow \infty} x_{k,n} = 0,$

**Iterate**  $n, k \in \mathbb{N}$  **Over**  $i \in \mathbb{N}$  **With**  $n_1 = 1, k_1 = 1,$

**Assume**  $j \in k_i,$

$\mathfrak{d}(a)(3) \rightsquigarrow \exists N \in \mathbb{N} : \forall m \in \mathbb{N} . |a_{m,k_i}| \leq \frac{1}{8k_i}$  **Extract as**  $N_j;$

$n_{i+1} := \max\{n_i + 1\} \cup N[k_i],$



$$\mathfrak{O}(n_{i+1}) \rightsquigarrow \sum_{j=1}^{k_i} |a_{n_{i+1},j}| \leq \sum_{j=1}^{k_i} \frac{1}{8k_i} = 1/8,$$

$$\mathfrak{O}(a)(2) \rightsquigarrow \exists K \in \mathbb{N} . \sum_{j=K+1}^{\infty} |a_{n_{i+1},j}| < 1/8 \text{ \textcolor{blue}{Extract}},$$

$$k_{i+1} := \max k_i + 1, K;$$

$$n : \text{\textcolor{blue}{Subsequer}} : \forall i \in \mathbb{N} . \sum_{j=1}^{k_i} |a_{n_{i+1},j}| \leq 1/8,$$

$$k : \text{\textcolor{blue}{Subsequer}} : \forall i \in \mathbb{N} . \sum_{j=k_{i+1}+1}^{\infty} |a_{n_{i+1},j}| < 1/8,$$

$$x : \mathbb{N} \rightarrow \{0, 1\}$$

$$x(m)$$

$$|\exists s \in \mathbb{N} . k_{2s-1} < m < k_{2s} = 0$$

$$\text{\textcolor{blue}{otherwise}} = 1$$

$$\tau := \Lambda n \in \mathbb{N} : \sum_{k=1}^{\infty} x_k a_{n,k},$$

$$\text{\textcolor{blue}{Assume}} \ m \in \mathbb{N},$$

$$\text{\textcolor{blue}{Assume}} \ m : \text{\textcolor{blue}{Odd}},$$

$$\mathfrak{O}(t, x) \rightsquigarrow |\tau_{n_{m+1}}| \leq \sum_{i=1}^{\infty} x_i |a_{n_{m+1},i}| \leq \sum_{i=1}^{k_n} |a_{n_{m+1},i}| + \sum_{i=k_{n+1}+1}^{\infty} |a_{n_{m+1},i}| \leq 1/4,$$

$$\tau_{n_{m+1}} \leq 1/4;$$

$$|\lim_{m \rightarrow \infty} \tau_m| \neq \infty,$$

$$z \text{\textcolor{blue}{Assume}} \ m : \text{\textcolor{blue}{Even}},$$

$$\sum_{i=k_n+1}^{k_{n+1}} a_{n_{m+1},i} \geq 3/4$$

$$\tau_{n_{m+1}} \geq 3/4 - \left| \sum_{i=1}^{k_n} |a_{n_{m+1},i}| + \sum_{i=k_{n+1}+1}^{\infty} |a_{n_{m+1},i}| \right| \geq 1/2;;$$

$$\tau ! \text{\textcolor{blue}{Convergent}} \left( \mathbb{R}^{\infty} \right); \square$$

**VitaliHahnSaksI** ::  $\forall P : \mathbb{N} \rightarrow \text{Probability}(\Omega, \mathcal{F}) . \forall \mathbb{P} : \mathcal{F} \rightarrow \mathbb{R} :$   
 $: \forall A \in \mathcal{F} . \lim_{n \rightarrow \infty} P_n(A) = \mathbb{P}(A) . P : \text{Probability}(\Omega, \mathcal{F}),$

**Proof** =

**Assume**  $P : \mathbb{N} \rightarrow \text{Probability}(\Omega, \mathcal{F}),$

**Assume**  $\mathbb{P} : \mathcal{F} \rightarrow \mathbb{R} : \forall A \in \mathcal{F} . \lim_{n \rightarrow \infty} P_n(A) = \mathbb{P}(A),$

**Assume**  $n \in \mathbb{N},$

**Assume**  $A : \text{DisjointElem}(\mathcal{F}, n),$

$$\alpha := \bigcup_{k=1}^n A_k \in \mathcal{F},$$

$$1 = \lim_{n \rightarrow \infty} P_n(\Omega) = \lim_{n \rightarrow \infty} P_n(\alpha) + P_n(\alpha^c) = \lim_{n \rightarrow \infty} P_n(\alpha) + \lim_{n \rightarrow \infty} P_n(\alpha^c) = \mathbb{P}(\alpha) + \mathbb{P}(\alpha^c) \rightsquigarrow \\ \rightsquigarrow (1) : 1 - \mathbb{P}(\alpha^c) = \mathbb{P}(\alpha)$$

$$1 = \lim_{n \rightarrow \infty} P_n(\alpha) + \lim_{n \rightarrow \infty} P_n(\alpha^c) = \lim_{m \rightarrow \infty} \sum_{k=1}^n P_m(A_k) + \mathbb{P}(\alpha^c) = \sum_{k=1}^n \lim_{m \rightarrow \infty} P_m(A_k) + \mathbb{P}(\alpha^c) = \\ = \sum_{k=1}^n \mathbb{P}(A_k) + \mathbb{P}(\alpha^c) \rightsquigarrow (2) : 1 - \mathbb{P}(\alpha^c) = \sum_{k=1}^n \mathbb{P}(A_k),$$

$$(1, 2) \rightsquigarrow \sum_{k=1}^n \mathbb{P}(A_k) = \mathbb{P}(\alpha);$$

$\mathbb{P} : \text{Charge}(\Omega, \mathcal{F}),$

**Assume**  $A : \text{DisjointElems}(\mathcal{F}, \mathbb{N}),$

$$c := \sum_{n=1}^{\infty} \mathbb{P}(A_n) \in \mathbb{R}_+ : c \leq 1,$$

$$\alpha := \bigcup_{k=1}^{\infty} A_k \in \mathcal{F}$$

**Assume**  $c = 1$

$$\mathbb{P}(\Omega) = \sum_{n=1}^{\infty} \mathbb{P}(A_n \sum_{n=1}^{\infty} \mathbb{P}(A_n)),$$

**Assume**  $\mathbb{P}(\alpha) < 1,$

**Assume**  $m \in \mathbb{N},$

$$\mathbb{P}(\alpha) = \lim_{n \rightarrow \infty} P_n(\alpha) \geq \lim_{n \rightarrow \infty} P_n\left(\bigcup_{k=1}^m A_k\right) = \sum_k^m = 1 \lim_{n \rightarrow \infty} P_n(A_k) = \sum_k^m = 1 \mathbb{P}(A_k);$$

$$\sum_{k=1}^m \mathbb{P}(A_k) < 1 \rightsquigarrow \perp;$$

$$\mathbb{P}(\alpha) = 1 \rightsquigarrow \mathbb{P}(\alpha) = \sum_{n=1}^{\infty} \mathbb{P}(A_n);$$

Assume  $c < 1$ ,

$\alpha = \Omega$ , Use  $\mathbb{P} : \text{Charge}(\Omega, \mathcal{F})$ ,

$a :: \mathbb{N} \rightarrow \mathbb{N} \rightarrow \mathbb{R}$

$a(n, k) := (1 - c)^{-1}(P_n(A_k) - \mathbb{P}(A_k))$ ,

Assume  $n \in \mathbb{N}$ ,

$$\begin{aligned} \sum_{k=1}^{\infty} a_{n,k} &= (1 - c)^{-1} \left( \sum_{k=1}^{\infty} P_n(A_k) - \sum_{k=1}^{\infty} \mathbb{P}(A_k) \right) = (1 - c)^{-1} \left( P_n(\alpha) - \sum_{k=1}^{\infty} \mathbb{P}(A_k) \right) = 1; \\ \sum_{k=1}^{\infty} |a_{n,k}| &\leq (1 - c)^{-1} \left( \sum_{k=1}^{\infty} P_n(A_k) + \sum_{k=1}^{\infty} \mathbb{P}(A_k) \right) = (1 - c)^{-1} \left( P_n(\alpha) + \sum_{k=1}^{\infty} \mathbb{P}(A_k) \right) = \frac{1 + c}{1 - c}, \\ \lim_{k \rightarrow \infty} a_{k,n} &= (1 - c)^{-1}(P_k(A_n) - \mathbb{P}(A_n)) = (1 - c)^{-1}(\mathbb{P}(A_n) - \mathbb{P}(A_n)) = 0; \end{aligned}$$

**SteinhausLemma**  $\rightsquigarrow \exists x : \mathbb{N} \rightarrow \{0, 1\} : \Lambda n \in \mathbb{N} : \sum_{k=1}^{\infty} x_k a_{n,k} ! \text{Convergent} \left( \mathbb{R} \right) \text{Extract},$

$$\begin{aligned} \lim_{n \rightarrow \infty} \sum_{k=1}^{\infty} x_k a_{n,k} &= (1 - c)^{-1} \left( \lim_{n \rightarrow \infty} P_n \left( \bigcup_{k \in \mathbb{N} : x_k = 1} A_k \right) \right) - (1 - c)^{-1} \sum_{k=1}^{\infty} x_k \mathbb{P}(A_k) = \\ &= (1 - c)^{-1} \mathbb{P} \left( \bigcup_{k \in \mathbb{N} : x_k = 1} A_k \right) - (1 - c)^{-1} \sum_{k=1}^{\infty} x_k \mathbb{P}(A_k) \rightsquigarrow \\ &\rightsquigarrow \sum_{k=1}^{\infty} x_k a_{n,k} : \text{Convergent}(\mathbb{R}) \rightsquigarrow \perp; \end{aligned}$$

$$\mathbb{P}(\alpha) = \sum_{n=1}^{\infty} \mathbb{P}(A_n);$$

$\mathbb{P} : \text{Probability}(\Omega, \mathcal{F}); ; \square$

## 2.2 Integration: Definition and Basic Results

$$\text{integrate} :: \text{Simple}(\Omega, \mathcal{F}) \rightarrow \text{Measure}(\Omega, \mathcal{F}) \rightarrow \overset{\infty}{\mathbb{R}}$$

$$\text{integrate} \left( \sum_{n=1}^r a_n I_{A_n} \right) (\mu) = \int_{\Omega} \sum_{n=1}^r a_n I_{A_n} d\mu = \sum_{n=1}^r a_n \mu(A_n)$$

$$\text{Extend integrate} :: \text{Measurable}(\Omega, \mathcal{F}) \& \text{Positive} \rightarrow \text{Measure}(\Omega, \mathcal{F}) \rightarrow \overset{\infty}{\mathbb{R}}$$

$$\int_{\Omega} f d\mu = \sup \left\{ \int_{\Omega} S d\mu \mid S : \text{Simple}(\Omega, \mathcal{F}) : 0 < S \leq f \right\}$$

$$\text{IntegralExist} :: \prod \mu : \text{Measure}(\Omega, \mathcal{F}) . ?\text{Measurable}(\Omega, \mathcal{F})$$

$$f : \text{IntegralExist} \iff \int_O f^+ d\mu < \infty \mid \int_O f^- d\mu < \infty$$

$$\text{Extend integrate} :: \prod \mu : \text{Measure}(\Omega, \mathcal{F}) . \text{IntegralExist} \rightarrow \overset{\infty}{\mathbb{R}}$$

$$\int_{\Omega} f d\mu = \int_{\Omega} f^+ d\mu - \int_{\Omega} f^- d\mu$$

$$\text{integrateOver} :: \mathcal{F} \rightarrow \text{Measurable}(\Omega, \mathcal{F}) \rightarrow \text{Measure}(\Omega, \mathcal{F}) \rightarrow \overset{\infty}{\mathbb{R}}$$

$$\text{integrateOver}(\text{Assume}, f, \mu) = \int_A f d\mu = \int_{\Omega} f I_A d\mu$$

$$\text{IntegralHomogeneity} :: \forall A \in \mathcal{F} .$$

$$\forall f : \text{Measurable}(\Omega, \mathcal{F}) . \forall \mu : \text{Measure}(\Omega, \mathcal{F}) . \forall c \in \mathbb{R} . \int_A c f d\mu = c \int_A f d\mu$$

$$\text{IntegralInequality} : \forall A \in \mathcal{F} .$$

$$\forall f, g : \text{Measurable}(\Omega, \mathcal{F}) : f \geq g . \forall \mu : \text{Measure}(\Omega, \mathcal{F}) . \int_A f d\mu \geq \int_A g d\mu$$

$$\text{IntegralModuleInequality} : \forall A \in \mathcal{F} .$$

$$\forall f : \text{Measurable}(\Omega, \mathcal{F}) . \forall \mu : \text{Measure}(\Omega, \mathcal{F}) . \int_A |f| d\mu \geq \left| \int_A f d\mu \right|$$

$$\text{Integrable} :: \prod \mu : \text{Measure}(\Omega, \mathcal{F}) . ?\text{IntegralExists}(\Omega, \mathcal{F})$$

$$f : \text{Integrable}(\Omega, \mathcal{F}) \iff \int_{\Omega} f d\mu < \infty$$

**FunToMeasureI** ::  $\forall f : \text{Simple}(\Omega, \mathcal{F}) . \forall \mu : \text{Measure}(\Omega, \mathcal{F}) .$

$$\Lambda A \in \mathcal{F} . \int_A f d\mu : \text{CountablyAdditive}(\Omega, \mathcal{F})$$

**Proof** =

**Assume**  $f : \text{Simple}(\Omega, \mathcal{F})$ ,

**Assume**  $\mu : \text{Measure}(\Omega, \mathcal{F})$ ,

$$f : \text{Simple}(\Omega, \mathcal{F}) \rightsquigarrow \exists n \in \mathbb{N} . \exists x : n \rightarrow \mathbb{R} . \exists A : n \rightarrow \mathcal{F} : f = \sum_{k=1}^n x_k I_{A_k} \text{ Extract},$$

**Assume**  $B : \text{DisjointElems}(\mathcal{F}, \mathbb{N})$ ,

$$\beta := \bigcup_{m=1}^{\infty} B_m \in \mathcal{F},$$

$$\int_{\beta} f d\mu = \sum_{k=1}^n x_k \mu(A_k \cap \beta) = \sum_{k=1}^n x_k \sum_{m=1}^{\infty} \mu(A_k \cap B_m) = \sum_{m=1}^{\infty} \sum_{k=1}^n x_k \mu(A_k \cap B_m) = \sum_{m=1}^{\infty} \int_{B_m} f d\mu;$$

$$\Lambda A \in \mathcal{F} . \int_A f d\mu : \text{CountablyAdditive}(\Omega, \mathcal{F}); ; \square$$

**FunToMeasureII** ::  $\forall f : \text{Measurable}(\Omega, \mathcal{F}) : f > 0 . \forall \mu : \text{Measure}(\Omega, \mathcal{F}) .$

$$\Lambda A \in \mathcal{F} . \int_A f d\mu : \text{CountablyAdditive}(\Omega, \mathcal{F})$$

**Proof** =

**Assume**  $f : \text{Simple}(\Omega, \mathcal{F})$ ,

**Assume**  $\mu : \text{Measure}(\Omega, \mathcal{F})$ ,

**Assume**  $B : \text{DisjointElems}(\mathcal{F}, \mathbb{N})$ ,

$$\beta := \bigcup_{m=1}^{\infty} B_m \in \mathcal{F},$$

$$\mathfrak{d}(\text{integrate}) \rightsquigarrow \int_{\beta} f d\mu \leq \sum_{n=1}^{\infty} \int_{B_n} f d\mu$$

**Assume**  $n \in \mathbb{N}$ ,

$$\mathfrak{d}(\text{integrate}) \exists S : \mathbb{N} \rightarrow \text{Simple}(\Omega, \mathcal{F}) : \int_{B_n} S d\mu \uparrow \int_{B_n} d\mu \text{ Extract as } S^n;$$

$$R : \mathbb{N} \rightarrow \text{Simple}(\Omega, \mathcal{F})$$

$$R_n(x)$$

$$| \exists k \in n : x \in B^k = S_n^k(x)$$

$$| \text{otherwise} = 0$$

$$\mathfrak{d}(R) \rightsquigarrow \lim_{n \rightarrow \infty} \int R_n d\mu = \sum_{n=1}^{\infty} \int_{B_n} f d\mu,$$

$$\forall_{n \in \mathbb{N}} R_n \in \{S : \text{Simple}(\Omega, \mathcal{F}) : 0 < S \leq f\} \rightsquigarrow \int_{\beta} f d\mu \geq \sum_{n=1}^{\infty} \int_{B_n} f d\mu \rightsquigarrow \int_{\beta} f d\mu = \sum_{n=1}^{\infty} \int_{B_n} f d\mu$$

$$\Lambda A \in \mathcal{F} . \int_A f d\mu : \text{CountablyAdditive}(\Omega, \mathcal{F}); ; \square$$

$$\text{FunToMeasureIII} :: \forall f : \text{IntegralExists}(\Omega, \mathcal{F}, \mu) . \Lambda A \in \mathcal{F} . \int_A f d\mu : \text{CountablyAdditive}(\Omega, \mathcal{F})$$

**Proof** =

**Assume**  $f : \text{IntegralExists}(\Omega, \mathcal{F}, \mu)$

**Assume**  $B : \text{DisjointElems}(\mathcal{F}, \mathbb{N})$

$$\beta := \bigcup_{m=1}^{\infty} B_m \in \mathcal{F},$$

$$\begin{aligned} \int_{\beta} f d\mu &= \int_{\beta} f^+ d\mu - \int_{\beta} f^- d\mu = \sum_{n=1}^{\infty} \int_{B_n} f^+ d\mu - \sum_{n=1}^{\infty} \int_{B_n} f^- d\mu = \sum_{n=1}^{\infty} \int_{B_n} f^+ d\mu - \int_{B_n} f^- d\mu = \\ &= \sum_{n=1}^{\infty} \int_{B_n} f d\mu; \end{aligned}$$

$$\Lambda A \in \mathcal{F} . \int_A f d\mu : \text{CountablyAdditive}(\Omega, \mathcal{F}); ; \square$$

$$\text{IntegrableCriterionI} :: \forall f : \text{Measurable}(\Omega, \mathcal{F}) .$$

$$f : \text{Integrable}(\Omega, \mathcal{F}, \mu) \iff |f| : \text{Integrable}(\Omega, \mathcal{F}, \mu)$$

$$\text{IntegrableCriterionII} :: \forall f : \text{Measurable}(\Omega, \mathcal{F}) . \forall g : \text{Integrable}(\Omega, \mathcal{F}) .$$

$$. |f| < g \Rightarrow f : \text{Integrable}$$

$$\text{almostEverywhere} :: \prod (\Omega, \mathcal{F}, \mu) : \text{MeasureSpace} . (? \Omega \rightarrow \text{Type}) \rightarrow \text{Type}$$

$$A : \text{almostEverywhere}(\Omega, \mathcal{F}, \mu)(T) \iff A : \text{a.e.} . [\mu]T \iff \exists Z \in \mathcal{F} : \mu(Z) = 0 : A : T(\Omega \setminus Z)$$

$$\text{ZeroAlmostEverywhere} :: \forall f : \text{Measurable}(\Omega, \mathcal{F}) . \forall P : \text{a.e.} . [\mu] f = 0 . \int_{\Omega} f d\mu = 0$$

$$\text{EqualAlmostEverywhere} :: \forall f : \text{Measurable}(\Omega, \mathcal{F}) . \forall g : \text{IntegralExists}(\Omega, \mathcal{F}, \mu) .$$

$$. \forall P : \text{a.e.} . [\mu] f = g \Rightarrow f : \text{IntegralExists}(\Omega, \mathcal{F}, \mu) \wedge \int_{\Omega} f d\mu = \int_{\Omega} g d\mu$$

$$\text{IntegrableAEFinite} :: \forall f : \text{Integrable}(\Omega, \mathcal{F}, \mu) . \text{a.e.} . [\mu] f < \infty$$

$$\text{ZeroAEZero} :: \forall f : \text{Integrable}(\Omega, \mathcal{F}, \mu) : f \geq 0 . \forall P : \int_{\Omega} f d\mu = 0 . \text{a.e.} . [\mu] f = 0$$

$$\text{IntegralAdditive} :: \forall f, g : \text{Measurable}(\Omega, \mathcal{F}) : f + g : \text{IntegralExists}(\Omega, \mathcal{F}, \mu) : \\ : \int_{\Omega} f d\mu + \int_{\Omega} g d\mu \in \widetilde{\mathbb{R}} \ . \int_{\Omega} (f + g) d\mu = \int_{\Omega} f d\mu + \int_{\Omega} g d\mu$$

$$\text{SumIntegral} :: \forall f : \mathbb{N} \rightarrow \text{Measurable}(\Omega, \mathcal{F}) : \forall n \in \mathbb{N} f_n \geq 0 \ . \sum_{n=1}^{\infty} \int_{\Omega} f_n d\mu = \int_{\Omega} \sum_{n=1}^{\infty} f_n d\mu$$

$$\text{AEInequality} :: \forall \mu : \text{FiniteMeasure}(\Omega, \mathcal{F}) \ . \forall f, g : \text{IntegralExists}(\Omega, \mathcal{F}, \mu) :$$

$$\forall A \in \mathcal{F} \ . \int_A f d\mu \leq \int_A g d\mu \ . \text{a.e.} \ . [\mu] f \leq g$$

**Proof** =

$$\text{Assume } f, g : \text{IntegralExists}(\Omega, \mathcal{F}, \mu) : \forall A \in \mathcal{F} \ . \int_A f d\mu \leq \int_A g d\mu$$

$$A :: \mathbb{N} \rightarrow \mathcal{F}$$

$$A_n = \{\omega \in \Omega : f(\omega) \geq g(\omega) + 1/n : |g(\omega)| \leq n\},$$

$$\bar{\partial}(A) \rightsquigarrow A \uparrow \{\omega \in \Omega : f(\omega) > g(\omega) : g(\omega) > -\infty\}$$

$$\text{Assume } n \in \mathbb{N},$$

$$\int_{A_n} g_n d\mu \geq \int_{A_n} f_n d\mu \geq \int_{A_n} g_n d\mu + \frac{1}{n} \mu(A_n),$$

$$\left| \int_{A_n} g d\mu \right| \leq \int_{A_n} |g| d\mu \leq n \mu(A_n) < \infty \rightsquigarrow \mu(A_n) = 0;$$

$$\mu\{\omega \in \Omega : f(\omega) > g(\omega) : g(\omega) > -\infty\} = \sum_{n=1}^{\infty} \mu(A_n) = 0,$$

$$C :: \mathbb{N} \rightarrow \mathcal{F}$$

$$C_n = \{\omega \in \Omega : g(\omega) = -\infty : f(\omega) > -n\},$$

$$C \uparrow \{\omega \in \Omega : f(\omega) > g(\omega) : g(\omega) = -\infty\},$$

$$\text{Assume } n \in \mathbb{N},$$

$$-\infty \mu(C_n) = \int_{C_n} g d\mu \geq \int_{C_n} f d\mu \geq -n \mu(C_n) \rightsquigarrow \mu(C_n) = 0;$$

$$\mu\{\omega \in \Omega : f(\omega) > g(\omega) : g(\omega) = -\infty\} = \sum_{n=1}^{\infty} \mu(C_n) = 0 \rightsquigarrow \text{a.e.} \ . [\mu] f \leq g; ; \square$$

## 2.3 Convergence of Integrals

**MonotoneConvergence** ::  $\forall h : \mathbb{N} \rightarrow \text{Measurable}(\Omega, \mathcal{F}) : \forall n \in \mathbb{N} . h_n \geq 0 . \forall h \uparrow_p H . \int h d\mu \uparrow \int H d\mu$

**Proof** =

**Assume**  $h : \mathbb{N} \rightarrow \text{Measurable}(\Omega, \mathcal{F}) : \forall n \in \mathbb{N} . h_n > 0$ ,

**Assume**  $h \uparrow_p H \rightsquigarrow \lim_{n \rightarrow \infty} \int_{\Omega} h_n d\mu \leq \int_{\Omega} H d\mu$

**Assume**  $S : \text{Simple}(\Omega, \mathcal{F}) : 0 \leq S \leq H$ ,

**Assume**  $c \in (0, 1)$ ,

$h \uparrow_p H \rightsquigarrow \lim_{n \rightarrow \infty} \int_{\Omega} f_n d\mu > c \int_{\Omega} S d\mu$ ;

$\lim_{n \rightarrow \infty} \int_{\Omega} f_n d\mu \geq \sup_c \sup_S \lim_{n \rightarrow \infty} c \int_{\Omega} S d\mu = \int_{\Omega} H d\mu \rightsquigarrow \int h d\mu \uparrow \int H d\mu ; \square$

**ExtendedMonotoneConvergenceUp** ::  $\forall h : \text{IntegralExists}(\Omega, \mathcal{F}, \mu) : \int_{\Omega} h d\mu > -\infty .$   
 $\forall f : \mathbb{N} \rightarrow \text{IntegralExists}(\Omega, \mathcal{F}, \mu) : \forall n \in \mathbb{N} . f_n \geq h . \forall f \uparrow \phi . \int_{\Omega} f d\mu \uparrow \int_{\Omega} \phi d\mu$

**Proof** =

**Assume**  $h : \text{IntegralExists}(\Omega, \mathcal{F}, \mu) : \int_{\Omega} h d\mu > -\infty$ ,

**Assume**  $f : \mathbb{N} \rightarrow \text{IntegralExists}(\Omega, \mathcal{F}, \mu) : \forall n \in \mathbb{N} . f_n \geq h$ ,

**Assume**  $f \uparrow \phi$ ,

**Alternative**  $\int_{\Omega} h d\mu = \infty \mid \int_{\Omega} h d\mu < \infty$ ,

**Assume Alternative**  $\int_{\Omega} h d\mu = \infty$ ,

**Assume**  $n \in \mathbb{N}$ ,

$f_n \geq h, \int_{\Omega} h d\mu = \infty \rightsquigarrow \int_{\Omega} f_n d\mu \geq \int_{\Omega} h d\mu = \infty \rightsquigarrow \int_{\Omega} f_n d\mu = \infty$ ;

$\int_{\Omega} \phi d\mu = \infty \rightsquigarrow \int_{\Omega} f d\mu \uparrow \int_{\Omega} \phi d\mu$ ;



Close Alternative  $\int_{\Omega} h d\mu < \infty$ ,

IntegrableAEFinite( $h$ )  $\rightsquigarrow$  (1) : a . e .  $[\mu] h < \infty$ ,

$h' := \Lambda\omega \in \Omega$  . if  $h(\omega) < \infty$  then  $h(\omega)$  else 0,

EquallAlmostEverywhere( $h, h', (1)$ )  $\rightsquigarrow$  (2) :  $\int_{\Omega} h' d\mu = \int_{\Omega} h d\mu$ ,

Assume  $n \in \mathbb{N}$ ,

$f_n \geq h \geq h' \rightsquigarrow$  (3) :  $f_n \geq h'$ ,

(3)  $\rightsquigarrow f_n - h' \geq 0$ ;

(3) :  $f - h' \geq 0$ ,

MonotoneConvergence( $f - h', (3), f - h' \uparrow \phi - h'$ )  $\rightsquigarrow$  (4) :  $\int_{\Omega} (f - h') d\mu \uparrow \int_{\Omega} (\phi - h') d\mu$ ,

(4)  $\rightsquigarrow \int_{\Omega} f d\mu + \int_{\Omega} h' d\mu = \int_{\Omega} (f + h') d\mu \uparrow \int_{\Omega} (\phi + h') d\mu = \int_{\Omega} \phi d\mu + \int_{\Omega} h' d\mu \rightsquigarrow \int_{\Omega} f d\mu \uparrow \int_{\Omega} \phi d\mu; ; ; \square$

ExtendedMonotoneConvergenceDown ::  $\forall h : \text{IntegralExists}(\Omega, \mathcal{F}, \mu) : \int_{\Omega} h d\mu < \infty$  .

.  $\forall f : \mathbb{N} \rightarrow \text{IntegralExists}(\Omega, \mathcal{F}, \mu) : \forall n \in \mathbb{N} . f_n \leq h . \forall f \downarrow \phi . \int_{\Omega} f d\mu \downarrow \int_{\Omega} \phi d\mu$

Proof =

Assume  $h : \text{IntegralExists}(\Omega, \mathcal{F}, \mu) : \int_{\Omega} h d\mu < \infty$ ,

Assume  $f : \mathbb{N} \rightarrow \text{IntegralExists}(\Omega, \mathcal{F}, \mu) : \forall n \in \mathbb{N} . f_n \leq h$ ,

Assume  $\int_{\Omega} f d\mu \downarrow \int_{\Omega} \phi d\mu$ ,

ExtendedMonotoneConvergenceUp( $-h, -f, -f \uparrow -\phi$ )  $\rightsquigarrow$

$\rightsquigarrow \int_{\Omega} -f d\mu \uparrow \int_{\Omega} -\phi d\mu \rightsquigarrow \int_{\Omega} f d\mu \downarrow \int_{\Omega} \phi d\mu; ; ; \square$

limInf ::  $(\mathbb{N} \rightarrow \Omega \rightarrow \mathbb{R}) \rightarrow \Omega \rightarrow \mathbb{R}$

limInf( $f, \omega$ ) =  $\left( \liminf_{n \rightarrow \infty} f_n \right) (\omega) = \sup_{n \in \mathbb{N}} \inf_{k \geq n} f_k(\omega)$

limSup ::  $(\mathbb{N} \rightarrow \Omega \rightarrow \mathbb{R}) \rightarrow \Omega \rightarrow \mathbb{R}$

limSup( $f, \omega$ ) =  $\left( \liminf_{n \rightarrow \infty} f_n \right) (\omega) = \inf_{n \in \mathbb{N}} \sup_{k \geq n} f_k(\omega)$

**FatouLemma** ::  $\forall f : \mathbb{N} \rightarrow \text{IntegralExists}(\Omega, \mathcal{F}, \mu) .$

$$. \forall g : \text{IntegralExists}(\Omega, \mathcal{F}, \mu) : f \geq g : \int_{\Omega} g d\mu > -\infty . \liminf_{n \rightarrow \infty} \int_{\Omega} f_n d\mu \geq \int_{\Omega} \liminf_{n \rightarrow \infty} f_n d\mu$$

**Proof** =

**Assume**  $f : \mathbb{N} \rightarrow \text{IntegralExists}(\Omega, \mathcal{F}, \mu),$

**Assume**  $g : \text{IntegralExists}(\Omega, \mathcal{F}, \mu) : f \geq g : \int_{\Omega} g d\mu > -\infty,$

$h :: \mathbb{N} \rightarrow \text{IntegralExists}(\Omega, \mathcal{F}, \mu)$

$$h_n = \inf_{k \geq n} f_k,$$

$$H := \liminf_{n \rightarrow \infty} f_n,$$

$$\mathfrak{D}(\liminf) \rightsquigarrow h \uparrow H$$

**Assume**  $n \in \mathbb{N},$

$$\mathfrak{D}(g) \rightsquigarrow h_n \geq g,$$

$$\mathfrak{D}(h) \rightsquigarrow f_n \geq h;$$

$$(1) : h \geq g,$$

$$(*) : f \geq h$$

$$\text{ExtendedMonotoneConvergenceUp}(h, g, (1), h \uparrow H) \rightsquigarrow (2) : \int_{\Omega} H d\mu = \lim_{n \rightarrow \infty} \int_{\Omega} h_n d\mu,$$

$$(*) \rightsquigarrow \lim_{n \rightarrow \infty} \int_{\Omega} h_n d\mu = \liminf_{n \rightarrow \infty} \int_{\Omega} h_n d\mu \leq \liminf_{n \rightarrow \infty} \int_{\Omega} f_n d\mu \rightsquigarrow \int_{\Omega} H d\mu \leq \liminf_{n \rightarrow \infty} \int_{\Omega} f_n d\mu;; \square$$

**DominatedConvergence** ::  $\forall D : \text{Integrable}(\Omega, \mathcal{F}, \mu) . \forall f : \mathbb{N} \rightarrow \text{Measurable}(\Omega, \mathcal{F}, \mu) :$

$$: \forall n \in \mathbb{N} . |f_n| \leq D . \forall P : \text{a.e.} . [\mu] f \xrightarrow{P} \phi . \phi : \text{Integrable}(\Omega, \mathcal{F}, \mu) \wedge \lim_{n \rightarrow \infty} \int_{\Omega} f_n d\mu = \int_{\Omega} \phi d\mu \quad \text{Proof} =$$

**Assume**  $D : \text{Integrable}(\Omega, \mathcal{F}, \mu),$

**Assume**  $f : \mathbb{N} \rightarrow \text{Measurable}(\Omega, \mathcal{F}, \mu) :$

$$: \forall n \in \mathbb{N} . |f_n| \leq D,$$

**Assume**  $P : \text{a.e.} . [\mu] f \xrightarrow{P} \phi,$

$$P, \mathfrak{D}(f) \rightsquigarrow (1) : |\phi| < D$$

**IntegrableCriterionII** $(\phi, D, 1) \rightsquigarrow \phi : \text{Integrable}(\Omega, \mathcal{F}, \mu),$

$$\begin{aligned} \int_{\Omega} \phi d\mu &= \int_{\Omega} \liminf_{n \rightarrow \infty} f_n d\mu \leq \liminf_{n \rightarrow \infty} \int_{\Omega} f_n d\mu \leq \lim_{n \rightarrow \infty} \int_{\Omega} f_n d\mu \leq \\ &\leq \limsup_{n \rightarrow \infty} \int_{\Omega} f_n d\mu \leq \int_{\Omega} \limsup_{n \rightarrow \infty} f_n d\mu = \int_{\Omega} \phi d\mu \rightsquigarrow \lim_{n \rightarrow \infty} \int_{\Omega} f_n d\mu = \int_{\Omega} \phi d\mu;; \square \end{aligned}$$

**PDominatedConvergence** ::  $\forall p \in \mathbb{R}_{++} .$

$$. \forall D : \text{Integrable}(\Omega, \mathcal{F}, \mu) : |D|^p : \text{Integrable}(\Omega, \mathcal{F}, \mu) . \forall f : \mathbb{N} \rightarrow \text{Measurable}(\Omega, \mathcal{F}, \mu) :$$

$$: \forall n \in \mathbb{N} . |f_n| \leq D . \forall P : \text{a.e.} . [\mu] f \xrightarrow{P} \phi . |\phi|^p : \text{Integrable}(\Omega, \mathcal{F}, \mu) \wedge \lim_{n \rightarrow \infty} \int_{\Omega} |f_n - \phi| d\mu = 0$$

$$\begin{aligned}
& \text{LimIntegral} :: \forall f : (a, b) \times (d, c) \rightarrow \mathbb{R} : (\exists g : \text{Measurable}((d, c), \mathcal{B}(d, c)) : g > |f| : \int_a^b g d\mu < \infty) \\
& : \forall x \in (a, b) . \Lambda y \in (d, c) . f(x, t) : \text{Measurable}((d, c), \mathcal{B}(d, c)) . \forall x' \in (a, b) . \\
& . \forall P : \forall y \in (c, d) . \lim_{x \rightarrow x'} f(x, t) \in \mathbb{R} . \int_a^b f(x, y) d\mu(y) = \int_a^b \lim_{x \rightarrow x'} f(x, y) d\mu(y)
\end{aligned}$$

Proof

$$\text{Assume } \forall f : (a, b) \times (d, c) \rightarrow \mathbb{R} : (\exists g : \text{Measurable}((d, c), \mathcal{B}(d, c)) : g > |f| : \int_a^b g d\mu < \infty)$$

$$: \forall x \in (a, b) . \Lambda y \in (d, c) . f(x, t) : \text{Measurable}((d, c), \mathcal{B}(d, c)),$$

$$\text{Assume } x' \in (a, b),$$

$$\text{Assume } P : \forall y \in (c, d) . \lim_{x \rightarrow x'} f(x, t) \in \mathbb{R},$$

$$P \leadsto \exists X : \mathbb{N} \rightarrow (a, b) : \lim_{n \rightarrow \infty} X_n = x' : \forall y \in (a, b] . \lim_{n \rightarrow \infty} f(X_n, y) = \lim_{n \rightarrow \infty} f(x', y) \text{ Extract,}$$

$$\text{DominatedConvergence}(f(X_n, \cdot), g, \lim_{n \rightarrow \infty} f(X_n, \cdot) = \lim_{x \rightarrow \infty x'} f(x, \cdot)) :$$

$$: \int_c^d \lim_{x \rightarrow x'} f(x, y) d\mu(y) = \lim_{n \rightarrow \infty} \int_c^d f(X_n, y) d\mu(y) = \lim_{x \rightarrow x'} \int_c^d f(x, y) d\mu(y); ; ; \square$$

## 2.4 Lebesgue VS Riemann [!]

## 2.5 Categorical Viewpoint: Integral As a Functor [!]

$\text{Transport} :: \text{Measure}(\Omega, \mathcal{F}) \rightarrow \text{Measure}(R, \mathcal{B}) \rightarrow ?\text{Measurable}(\Omega, \mathcal{F})(R, \mathcal{B})$

$T : \text{Transport}(\alpha, \beta) \iff \forall B \in \mathcal{B} . \beta(B) = \alpha(T^{-1}B)$

$\text{TransportIntegral} :: \forall T : \text{Transport}(\Omega, \mathcal{F})(\Omega', \mathcal{F}')(\alpha, \beta) . \forall f : \text{IntegralExists}(\Omega', \mathcal{F}', \beta) .$

$$. \forall A \in \mathcal{F}' . \int_A f d\beta = \int_{T^{-1}A} f \circ T d\alpha$$

**Proof** =

**Assume**  $T : \text{Transport}(\Omega, \mathcal{F})(\Omega', \mathcal{F}')(\alpha, \beta)$ ,

**Assume**  $f : \text{IntegralExists}(\Omega', \mathcal{F}', \beta)$ ,

**Assume**  $A \in \mathcal{F}'$ ,

**Assume**  $B \in \mathcal{F}'$ ,

$$\int_A I_B d\beta = \beta(A \cap B) = \alpha(T^{-1}(A \cap B)) = \alpha(T^{-1}A \cap T^{-1}B) = \int_{T^{-1}A} I_{T^{-1}B} d\alpha = \int_{T^{-1}A} I_B \circ T d\alpha,$$

$$(1) : \forall B \in \mathcal{F}' . \int_A I_B d\beta = \int_{T^{-1}A} I_B \circ T d\alpha,$$

**Assume**  $S : \text{Simple}(\Omega, \mathcal{F})$ ,

$$\text{Simple}(S) \rightsquigarrow : \exists n \in \mathbb{N} . \exists x : n \rightarrow \mathbb{R} . \exists B : \text{DisjointElems}(\mathcal{F}', n) : S = \sum_{k=1}^n x_k I_{B_k},$$

$$\int_A S d\beta = \int_A \sum_{k=1}^n x_k I_{B_k} d\beta = \sum_{k=1}^n x_k \int_A I_{B_k} d\beta = \sum_{k=1}^n x_k \int_{T^{-1}A} I_{B_k} \circ T d\alpha = \int_{T^{-1}A} S \circ T d\alpha;$$

$$(2) : \forall S : \text{Simple}(\Omega, \mathcal{F}) . \int_A S d\beta = \int_{T^{-1}A} S \circ T d\alpha,$$

**Assume**  $g : \text{IntegralExists}(\Omega, \mathcal{F}, \beta) : g > 0$  .

$S := \text{SimpleApproximationI}(g) : \mathbb{N} \rightarrow \text{Simple}(\Omega, \mathcal{F}) : 0 \leq S \leq g : S \uparrow_p g$ ,

$$\begin{aligned} \text{MonotoneConvergence}(S, g, 0) &\rightsquigarrow \int_A g d\beta = \lim_{n \rightarrow \infty} \int_A S_n d\beta = \lim_{n \rightarrow \infty} \int_{T^{-1}A} S_n \circ T d\alpha = \\ &= \int_{T^{-1}A} g \circ T d\alpha; \end{aligned}$$

$$(3) : \forall f : \text{IntegralExists}(\Omega, \mathcal{F}, \mu) . \int_A g d\beta = \int_{T^{-1}A} g \circ T d\alpha,$$

$$\int_A f d\beta = \int_A f^+ d\beta - \int_A f^- d\beta = \int_{T^{-1}A} f^+ \circ T d\alpha - \int_{T^{-1}A} f^- \circ T d\alpha = \int_{T^{-1}A} f \circ T d\alpha; \square$$

$\text{MEAS} :: \text{Category}$

$\mathcal{O}_{\text{MEAS}} := \text{MeasureSpace}$

$\mathcal{M}_{\text{MEAS}}(A, B) := \text{Transport}(A, B)$

$\cdot_{\text{MEAS}} := \circ$

$\mathbf{I} :: \text{Functor}(\text{MEAS}, \text{VS}(\mathbb{R}))$

$\mathbf{I}(A) = (\text{Integrable}(A))^*$

$\mathbf{I}(T) = \Lambda v \in \mathbf{I}(A) . \Lambda f \in \text{Integrable}(B) . \langle v, f \circ T \rangle$

$\text{BOR} :: \text{Category}$

$\mathcal{O}_{\text{BOR}} := \text{MeasurableSpace}$

$\mathcal{M}_{\text{BOR}}(A, B) := \text{Measurable}(A, B)$

$\cdot_{\text{BOR}} := \circ$

$\text{BOR}_0 :: \text{Category}$

$\mathcal{O}_{\text{BOR}_0} := \text{Localizeble}$

$\mathcal{M}_{\text{BOR}_0}((\Omega, \mathcal{F}, N), (\Omega', \mathcal{F}', M)) := \frac{\{f : \text{Measurable}(\Omega, \mathcal{F})(\Omega', \mathcal{F}') : \forall Z \in M . f^{-1}(Z) \in N\}}{\{(f, g) \in \text{Measurable}(\Omega, \mathcal{F})(\Omega', \mathcal{F}')^2 : \{\omega \in \Omega : f(\omega) \neq g(\omega)\} \in N\}}$

$\cdot_{\text{BOR}_0} := \circ$

## 3 Radon-Nikodym Theory

### 3.1 Jordan-Hahn Decomposition

**Diffusion** :: ?Lebesgue-Stieltjes( $\mathbb{R}^n$ )

$\mu : \text{Diffusion} \iff F_\mu \in \mathcal{M}_{\text{TOP}}(\mathbb{R}^n, \mathbb{R})$

**AbsolutelyContinuous** :: SignedMeasure( $\Omega, \mathcal{F}$ )  $\rightarrow$  ?SignedMeasure( $\Omega, \mathcal{F}$ )

$\alpha : \text{AbsolutelyContinuous}(\beta) \iff \alpha \ll \beta \iff \forall A \in \mathcal{F} : \beta(A) = 0 \rightarrow \alpha(A) = 0$

**SupIsAttainable** ::  $\forall H : \text{CountablyAdditive}(\Omega, \mathcal{F}) . \exists A \in \mathcal{F} : H(A) = \sup_{S \in \mathcal{F}} H(S)$

**Proof** =

**Assume**  $H : \text{CountablyAdditive}(\Omega, \mathcal{F})$ ,

$s := \sup_{S \in \mathcal{F}} H(S) \in \mathbb{R}$ ,

$A := \bar{\partial}(\sup)(\sup_{S \in \mathcal{F}} H(S)) : \mathbb{N} \rightarrow \mathcal{F} : \lim_{n \rightarrow \infty} H(A_n) = s$

$\alpha = \bigcup_{n=1}^{\infty} A_n$ ,

**Assume**  $n \in \mathbb{N}$ ,

**Assume**  $b \in \mathbb{B}^n$ ,

$a_{n,b} := \bigcap_{n=1}^{\infty} \text{bToC}(A_n, b_n, \alpha)$ ;

$B_n = \bigcup \{a_{n,b} \mid b \in \mathbb{B} : a_{n,b} > 0\}$

**Assume**  $r \in \mathbb{N} : r > n$ ,

$H(A_n) \leq H(B_n) \leq H\left(\bigcup_{k=n}^r B_k\right)$ ,

$H(A_n) \leq H\left(\bigcup_{k=n}^{\infty} B_k\right)$ ,

$c := \liminf_{B \in \mathcal{F}}$

$\bigcup_{k=n}^{\infty} B_k \downarrow_n C$ ,

$\bar{\partial}(B) \rightsquigarrow \lim_{n \rightarrow \infty} H\left(\bigcup_{k=n}^{\infty} B_k\right) = H(C)$ ,

$s = \lim_{n \rightarrow \infty} H(A_n) \leq \lim_{n \rightarrow \infty} H\left(\bigcup_{k=n}^{\infty} B_k\right) = H(C) \leq s \rightsquigarrow H(C) = s; \square$

$\text{upperVariation} :: \text{CountablyAdditive}(\Omega, \mathcal{F}) \rightarrow \text{Measure}(\Omega, \mathcal{F})$   
 $\text{upperVariation}(H) = H^+ := \Lambda A \in \mathcal{F} . \sup\{H(B) | B \in \mathcal{F} : B \subset A\}$

$\text{lowerVariation} :: \text{CountablyAdditive}(\Omega, \mathcal{F}) \rightarrow \text{Measure}(\Omega, \mathcal{F})$   
 $\text{lowerVariation}(H) = H^- := \Lambda A \in \mathcal{F} . -\inf\{H(B) | B \in \mathcal{F} : B \subset A\}$

$\text{absVariation} :: \text{CountablyAdditive}(\Omega, \mathcal{F}) \rightarrow \text{Measure}(\Omega, \mathcal{F})$   
 $\text{absVariation}(H) = |H| := H^+ + H^-$

$\text{JordanHahnDecomposition} :: \forall H : \text{CountablyAdditive}(\Omega, \mathcal{F}) . H = H^+ - H^-$

Proof =

Assume  $H : \text{CountablyAdditive}(\Omega, \mathcal{F})$ ,

Assume  $P : H > -\infty$ ,

$D := \text{SupIsAttainable}(H) \in \mathcal{F} : H(D) = \inf_{A \in \mathcal{F}} H(A)$ ,

Assume  $A \in \mathcal{F}$ ,

Assume  $R : -H^-(A) < H(A \cap D)$ ,

$R \rightsquigarrow \exists C \in \mathcal{F} : C \subset A : H(A \cap D) > H(C)$  **Extract**,

$H(A \cap D) > H(C) > H(D) \rightsquigarrow H(D) > H(D) - H(A \cap D) + H(C) = H((D \setminus (A \cap D)) \cup C)$ ,

$(D \setminus (A \cap D)) \cup C \in \mathcal{F} \rightsquigarrow \perp$ ;

$-H^-(A) \geq H(A \cap D) \rightsquigarrow_{\text{dH}} -H^-(A) = H(A \cap D)$ ,

$-H^-(A) \leq H(\emptyset) = 0$ ;

$H^- = \Lambda A \in \mathcal{F} . -H(A \cap D)$

$H^- \geq 0 \rightsquigarrow H^- : \text{Measure}(\Omega, \mathcal{F})$

Assume  $A \in \mathcal{F}$ ,

$H(A) = H(A \cap D) + H(A \cap D^c) = -H^- + H(A \cap D^c)$ ,

Assume  $R : H(A \cap D^c) < 0$ ,

$H^-(A) = H(A \cap D) < H(A \cap D) + H(A \cap D^c) \rightsquigarrow \perp$ ;

$H(A \cap D^c) \geq 0$ ;

(1) :  $\forall A \in \mathcal{F} . H(A \cap D^c) \geq 0$ ,

Assume  $B \in \mathcal{F} : B \subset A$ ,

$H(B) = H(B \cap D) + H(B \cap D^c) \leq H(B \cap D^c) \leq_1 H(A \cap D^c)$ ,

$H^+(A) \leq H(A \cap D^c) \rightsquigarrow H^+(A) = H(A \cap D^c)$ ;

$H = H^+ - H^-$ ;  $\square$



$$\begin{aligned}
|\lambda_1 + \lambda_2| &= (\lambda_1 + \lambda_2)^+ + (\lambda_1 + \lambda_2)^- = \\
&= (\lambda_1^+ + \lambda_2^+ - \lambda_1^- - \lambda_2^-)^+ + (\lambda_1^+ + \lambda_2^+ - \lambda_1^- - \lambda_2^-)^- \leq \lambda_1^+ + \lambda_2^+ + \lambda_1^- + \lambda_2^- = |\lambda_1| + |\lambda_2|
\end{aligned}$$

## 3.2 Radon-Nikodym Theorem

**RadonNykodymI** ::  $\forall P : \text{FiniteMeasure}(\Omega, \mathcal{F}) . \forall \mu : \text{FiniteMeasure} : \mu \ll P .$

$$\exists f : \text{Measurable}(\Omega, \mathcal{F}) : \forall A \in \mathcal{F} : \mu(A) = \int_A f dP$$

**Proof** =

**Assume**  $P : \text{FiniteMeasure}(\Omega, \mathcal{F})$ ,

**Assume**  $\mu : \text{FiniteMeasure} : \mu \ll P$ ,

$$F := \{f : \text{Integrable}(\Omega, \mathcal{F}, P) : \forall A \in \mathcal{F} . \int_A f dP \leq \mu(A)\}$$

**Assume**  $f : \mathbb{N} \rightarrow F$ ,

$$\phi := \sup_{n \in \mathbb{N}} f : \text{Integrable}(\Omega, \mathcal{F}, P),$$

**Assume**  $n \in \mathbb{N}$ ,

$$g_n := \Lambda \omega \in \Omega . \max\{f_k(\omega) : k\} \in F;$$

$$M = \partial g : g \uparrow \phi,$$

$$B = \partial(F, \Im g \subset F) : \forall n \in \mathbb{N} . \forall A \in \mathcal{F} . \int_A g_n dP \leq \mu(A),$$

$$MC = \text{MonotoneConvergence}(g, M) : \int_{\Omega} g dP \uparrow \int_{\Omega} \phi dP$$

**Assume**  $A \in \mathcal{F}$ ,

$$B \rightsquigarrow \mu(A) \geq \int_A g dP = \int_{\Omega} I_A g dP \uparrow \int_{\Omega} I_A \phi dP = \int_A \phi dP;$$

$$\phi \in F;$$

$$0 \in F \rightsquigarrow F \neq \emptyset \rightsquigarrow \sup F = \max F \neq \emptyset,$$

**Assume**  $f \in \max F$ ,

$$\lambda := \Lambda A \in \mathcal{F} . \mu(A) - \int_A \phi dP : \text{Measure}(\Omega, \mathcal{F}),$$

**Assume**  $A \in \mathcal{F} : P(A) = 0$

$$\mu \ll P \rightsquigarrow (1) : \mu(A) = 0,$$

$$(1) \rightsquigarrow \lambda(A) = \mu(A) - \int_A \phi dP = 0;$$

$$(1) : \lambda \ll P,$$

**Assume**  $A1 : \lambda \neq 0$

$$A1, f \in F \rightsquigarrow \lambda(\Omega) > 0 \rightsquigarrow \exists k \in \mathbb{R}_{++} : P(\Omega) - k\lambda(\Omega) < 0 \text{ **Extract**,$$

$$H := P - k\lambda : \text{SignedMeasure}(\Omega, \mathcal{F})$$

$$D := \text{SupIsAttainable}(H) \in \mathcal{F} : H(D) = \inf_{A \in \mathcal{F}} H(A)$$

$$D1 : \partial(h, H)D \neq \emptyset \wedge H(D) < 0$$

**Assume**  $Z : P(D) = 0$ ,

$$H(D) = P(D) - \lambda(D) =_{Z,1} 0 \rightsquigarrow_{D1} \perp$$

$$(2) : P(D) > 0,$$

$$h := (1/k)I_D : \text{Measurable}(\Omega, \mathcal{F}),$$

Assume  $A \in \mathcal{F}$ ,

$$\int_A h dP = 1/k P(A \cap D) \leq \lambda(A \cap D) \leq \lambda(A) = \mu(A) - \int_A f dP,$$

$$\int_A (h + g) dP \leq \mu(A);$$

$$h + f \in F,$$

$$h + f > f \rightsquigarrow f \notin \max F \rightsquigarrow \perp;$$

$$H = 0 \rightsquigarrow \forall A \in \mathcal{F} : \mu(A) = \int_A f dP; ; \square$$

**RadonNykodym** ::  $\forall \mu : \sigma\text{-Finite}(\Omega, \mathcal{F}) . \forall H : \text{SignedMeasure}(\Omega, \mathcal{F}) : H \ll \mu .$

$$. \exists f \in \text{Measurable}(\Omega, \mathcal{F}) : \forall A \in \mathcal{F} . \int_A f d\mu = H(A)$$

**Density** ::  $\text{Measure}(\Omega, \mathcal{F}) \rightarrow \text{SignedMeasure}(\Omega, \mathcal{F}) \rightarrow ?\text{Measurable}(\Omega, \mathcal{F})$

$$f : \text{Density}(\mu, \lambda) \iff \text{forall } A \in \mathcal{F} . \int_A f d\mu = \lambda(A)$$

**density** ::  $\prod \mu : \text{Measure}(\Omega, \mathcal{F}) . \prod \lambda : \text{SignedMeasure} : \lambda \ll \mu . \text{Density}(\mu, \lambda)$

$$\text{density} = f_\lambda = \text{RadonNykodym}(\mu, \lambda)$$

**Singular** ::  $\text{Measure}(\Omega, \mathcal{F}) \rightarrow ?\text{Measure}(\Omega, \mathcal{F})$

$$\lambda : \text{Singular}(\mu) \iff \lambda \perp \mu \iff \exists A \in \mathcal{F} : \lambda(A) = 0 \wedge \mu(A^c) = 0$$

**MutSingular** ::  $\text{SignedMeasure}(\Omega, \mathcal{F}) \rightarrow ?\text{SignedMeasure}(\Omega, \mathcal{F})$

$$\lambda : \text{Singular}(\mu) \iff \lambda \perp \mu \iff |\mu| \perp |\lambda|$$

**BorelCantelli** ::  $\forall \mu : \text{Measure}(\Omega, \mathcal{F}) . \forall A : \mathbb{N} \rightarrow \mathcal{F} . \forall S : \sum_{n=1}^{\infty} \mu(A_n) < \infty . \mu(\limsup A) = 0$

**Proof** =

Assume  $\mu : \text{Measure}(\Omega, \mathcal{F})$ ,

Assume  $A : \mathbb{N} \rightarrow \mathcal{F}$ ,

Assume  $S : \sum_{n=1}^{\infty} \mu(A_n) < \infty$ ,

Assume  $\epsilon \in \mathbb{R}_{++}$ ,

$$k = \text{SumConverge}(S, \epsilon) : \mathbb{N} : \sum_{n=1}^{\infty} \mu(A_n) < \epsilon,$$

$$D1 : \epsilon > \sum_{n=1}^{\infty} \mu(A_n) \geq \mu \left( \bigcup_{n=k}^{\infty} A_n \right) \geq \mu \left( \bigcap_{m=k}^{\infty} \bigcup_{n=m}^{\infty} A_n \right) \geq \mu \left( \bigcap_{m=1}^{\infty} \bigcup_{n=m}^{\infty} A_n \right) = \mu(\limsup A);$$

$$\mu(\limsup A) = 0; ; \square$$

**RNProperties** ::  $\forall \mu : \text{Measure}(\Omega, \mathcal{F}) . \forall \lambda_1, \lambda_2 : \text{SignedMeasure}(\Omega, \mathcal{F})$

$\forall P1 : \lambda_1 \perp \mu \& \lambda_2 \perp \mu . \mu = \lambda_1 + \lambda_2,$

$\forall P2 : \lambda_1 \ll \mu . |\lambda_1| \ll \mu,$

$\forall P3 : \lambda_1 \ll \mu \& \lambda_2 \perp \mu . \lambda_1 \perp \lambda_2,$

$\forall P4 : \lambda_1 \ll \mu \& \lambda_1 \perp \mu . \lambda_1 = 0,$

$\forall P5 : |\lambda_1| < 0 . \lambda_1 \ll \mu \iff \lim_{\mu(A) \rightarrow 0} \lambda_1(A) = 0;$

**LebesgueDecomposition** ::  $\forall \mu : \text{Measure}(\Omega, \mathcal{F}) . \forall H : \text{SignedMeasure}(\Omega, \mathcal{F}) :$

$(|H| : \sigma\text{-Finite}(\Omega, \mathcal{F})) . \exists \alpha, \beta : \text{SignedMeasure}(\Omega, \mathcal{F}) : \alpha \ll \mu : \beta \perp \mu : H = \alpha + \beta$

**Proof** =

**Assume**  $\mu : \text{Measure}(\Omega, \mathcal{F}),$

**Assume**  $P : \text{FiniteMeasure}(\Omega, \mathcal{F}),$

$\mathcal{B} := \{A \in \mathcal{F} : \mu(A) = 0\} : ?\mathcal{F},$

$s := \sup\{P(b) | b \in \mathcal{B}\} \in \mathbb{R}_+,$

$b := \partial(\sup)(s) : \mathbb{N} \rightarrow B : P(b) \uparrow s,$

$B := \bigcup_{n=1}^{\infty} b_n \in \mathcal{F},$

$MUC1 := \text{MeasureUpperContinuity}(\mu, b) : \mu(B) = \lim_{n \rightarrow \infty} \mu(b_n) = 0 \rightsquigarrow B \in \mathcal{B},$

$MUC2 := \text{MeasureUpperContinuity}(P, b) : P(B) = \lim_{n \rightarrow \infty} P(b_n) = s,$

$D1 : s =_{MUC2} P(B) \leq P(\Omega) < \infty,$

$D1 \rightsquigarrow s \in \mathbb{R}_{++},$

$\alpha = \Lambda A \in \mathcal{F} := P(A \cap B^c) : \text{Measure}(\Omega, \mathcal{F}),$

$\beta = \Lambda A \in \mathcal{F} := P(A \cap B) : \text{Measure}(\Omega, \mathcal{F}),$

$S1 := \partial(\perp)(\beta, (B, MUC1)) : \beta \perp \mu,$

**Assume**  $A \in \mathcal{F} : \mu(A) = 0,$

**Assume**  $C1 : \alpha(A) > 0,$

$\partial(\mathcal{B})(\partial A) \rightsquigarrow A \in \mathcal{B} \rightsquigarrow B \cup A \in \mathcal{B},$

$P(A \cup B) = P(B) + P(A \cap B^c) > P(B) \rightsquigarrow B \neq s \rightsquigarrow \perp,$

$\alpha(A) = 0;$

$\alpha \ll \mu,$

$P = \alpha + \beta,$

$A1 : \forall P : \text{FiniteMeasure}(\Omega, \mathcal{F}) . \exists \alpha, \beta : \text{SignedMeasure}(\Omega, \mathcal{F}) : \alpha \ll \mu : \beta \perp \mu : P = \alpha + \beta,$

Assume  $\lambda : \sigma\text{-Finite}(\Omega, \mathcal{F})$ ,

$$a := \text{d}\sigma\text{-Finite}(\Omega, \mathcal{F}) : \mathbb{N} \rightarrow \mathcal{F} : \bigcup_{n=1}^{\infty} a_n = \Omega : \forall n \in \mathbb{N} . \lambda(a) < \infty,$$

Assume  $n \in \mathbb{N}$ ,

$$P := \Lambda A \in F . \lambda(a_n \cap A) : \text{FiniteMeasure}(\Omega, \mathcal{F}),$$

$$(\alpha_n, \beta_n) := A1(P) : \text{SignedMeasure}^2(\Omega, \mathcal{F}) : \alpha_n \ll \mu : \beta_n \perp \mu : P = \alpha_n + \beta_n;$$

$$\alpha := \sum_{n=1}^{\infty} \alpha_n,$$

$$\beta := \sum_{n=1}^{\infty} \beta_n,$$

$$\lambda = \alpha + \beta;$$

$$A2 : \forall \lambda : \sigma\text{-Finite}(\Omega, \mathcal{F}) . \exists \alpha, \beta : \text{SignedMeasure}(\Omega, \mathcal{F}) : \alpha \ll \mu : \beta \perp \mu : \lambda = \alpha + \beta,$$

Assume  $H : \text{SignedMeasure}(\Omega, \mathcal{F}) : (|H| : \sigma\text{-Finite}(\Omega, \mathcal{F}))$ ,

$$|H| : \sigma\text{-Finite}(\Omega, \mathcal{F}) \rightsquigarrow H^+, H^- : \sigma\text{-Finite}(\Omega, \mathcal{F}),$$

$$(\alpha^+, \beta^+) := A2(H^+),$$

$$(\alpha^-, \beta^-) := A2(H^-),$$

$$\alpha := \alpha^+ - \alpha^-,$$

$$\beta := \beta^+ - \beta^-,$$

$$H = H^+ - H^- := \alpha^+ + \beta^+ - \alpha^- - \beta^- := \alpha - \beta; ; \square$$

ChainRule ::  $\forall \mu, \lambda : \text{Measure}(\Omega, \mathcal{F}) . \forall f : \text{Density}(\mu, \lambda) . \forall g : \text{Measurable}(\Omega, \mathcal{F}) .$

$$. \int_{\Omega} g d\lambda = \int_{\Omega} g f d\mu$$

Proof =

$$g \geq 0,$$

$$\begin{aligned} \int_{\Omega} g d\lambda &= \lim_{n \rightarrow \infty} \sum_{k=1}^{m_n} a_k^n \lambda(A_k^n) = \lim_{n \rightarrow \infty} \sum_{k=1}^{m_n} a_k^n \int_{A_k^n} f d\mu = \lim_{n \rightarrow \infty} \int_{\Omega} \sum_{k=1}^{m_n} a_k^n I_{A_k^n} f d\mu = \\ &= \lim_{n \rightarrow \infty} \int_{\Omega} S_n f d\mu = \int_{\Omega} g f d\mu \end{aligned}$$

—

$$\int_{\Omega} g d\lambda = \int_{\Omega} g^+ d\lambda - \int_{\Omega} g^- d\lambda = \int_{\Omega} g^+ f d\mu - \int_{\Omega} g^- f d\mu = \int_{\Omega} g f d\mu$$

### 3.3 Absolutely Continuous Functions

$\text{AbsCont} :: \prod [a, b] : \text{CInterval} . ?([a, b] \rightarrow \mathbb{R})$

$f : \text{AbsCont} \iff \forall \epsilon \in \mathbb{R}_{++} . \exists \delta \in \mathbb{R} . \forall n \in \mathbb{N} .$

$. \forall [c, d] : \mathbb{N} \rightarrow \text{CInterval} : \sum_{k=1}^n d_k - c_k \leq \delta . \sum_{k=1}^n |f(c_k) - f(d_k)| \leq \epsilon$

$\text{AbsContDistr} :: \forall \alpha, \beta : \text{Measure}([a, b], \mathcal{B}[a, b]) . F_\alpha - F_\beta : \text{AbsCont}[a, b] \iff \alpha - \beta \ll \lambda$

**Proof** =

**Assume**  $\alpha, \beta : \text{Measure}([a, b], \mathcal{B}[a, b]),$

$H := \alpha - \beta : \text{SignedMeasure}([a, b], \mathcal{B}[a, b]),$

**Assume**  $[c, d] : \text{SubInterval}[a, b],$

$H[c, d] = F_\alpha(d) - F_\alpha(c) - F_\beta(d) + F_\beta(c) = G(d) - G(c);$

$R1 : G = F_H,$

**Assume**  $P1 : F_\alpha - F_\beta : \text{AbsCont}[a, b],$

$G := F_\alpha - F_\beta : \text{AbsCont}[a, b],$

**Assume**  $A \in \mathcal{F} : \lambda(A) = 0,$

$V := \text{OuterRegularity}(A) : \mathbb{N} \rightarrow \mathcal{T}_{[a, b]} . \lambda(V) \downarrow 0 : \alpha(V) \downarrow \alpha(A) : \beta(V) \downarrow \beta(A)$

$W := \text{toDecreasing}(V),$

**Assume**  $\epsilon \in \mathbb{R},$

$\delta := \text{AbsCont}[a, b](G) \in \mathbb{R}_{++} : \forall n \in \mathbb{N} . \forall (c, d) : \mathbb{N} \rightarrow \text{CInterval} :$

$: \forall k \in n . d_k - c_k \leq \delta . \sum_{k=1}^n |G(c_k) - G(d_k)| \leq \epsilon,$

$N := \mathfrak{O}(w, \lambda(W) \downarrow 0) \in \mathbb{N} : \forall n \geq N . \exists m \in \mathbb{N} : \exists (c, d) : m \rightarrow \text{Subinterval}[a, b] : W_n = \bigsqcup_{k=1}^m (c_k, d_k) :$

$: \sum_{k=1}^m \lambda(c_k, d_k) \leq \delta,$

$(m, (c, d)) := \mathfrak{O}(N) : \sum m \in \mathbb{N} \rightarrow \mathbb{N} . m \rightarrow \text{Subinterval}[a, b] : W_n \subset \bigsqcup_{k=1}^m (c_k, d_k) : \sum_{k=1}^m \lambda(c, d) \leq \delta,$

$\mathfrak{O}(\delta, N) \rightsquigarrow \forall n \geq N . |H(W_n)| \leq \left| \sum_{k=1}^{m_n} H(c_k, d_k) \right| \leq \sum_{k=1}^{m_n} |H(c_k, d_k)| = \sum_{k=1}^{m_n} |G(d_k) - G(c_k)| \leq \epsilon \rightsquigarrow$

$\lim_n \rightarrow \infty |H(W_n)| \leq \epsilon,$

$H(A) = \lim_{n \rightarrow \infty} H(A_n) = 0;$

$R2 : \forall P : F_\alpha - F_\beta : \text{AbsCont}[a, b] . H \ll \lambda,$

**Assume**  $P : H \ll \lambda,$

$R3 := \mathfrak{O}(\ll)(P) \rightsquigarrow |H| \ll \lambda,$

**Assume**  $\epsilon \in \mathbb{R}_{++}$ ,

$\delta := \text{RNProperties}(|H|, R3)(\epsilon) : \forall A \in \mathcal{B}[a, b] : \lambda(A) \leq \delta \cdot |H|(A) \leq \epsilon$ ,

$n \in \mathbb{N}$

**Assume**  $[c, d] : n \rightarrow \text{SubInterval}[c, d] : \sum_{k=1}^n d_k - c_k \leq \delta$ ,

$R4 \leq \text{Subadditivity}(\lambda) : \lambda \left( \bigcup_{k=1}^n [c_k, d_k] \right) \leq \sum_{k=1}^n d_k - c_k \leq \delta$

$R5 : \mathfrak{O}(\delta) \left( \bigcup_{k=1}^n [c_k, d_k], R4 \right) : |H| \left( \bigcup_{k=1}^n [c_k, d_k] \right) \leq \epsilon$ ,

$R5 \rightsquigarrow \epsilon \geq |H| \left( \bigcup_{k=1}^n [c_k, d_k] \right) = \sum_{k=1}^n |H|[c_k, d_k] \geq \sum_{k=1}^n |H[c_k, d_k]| = \sum_{k=1}^n |G(d_k) - G(c_k)| ; ;$

$G : \text{AbsCont}[a, b]$ ;

$R3 : \forall P : H \ll \lambda \cdot F_\alpha - F_\beta : \text{AbsCont}[a, b]$ ,

$\text{IFFI}(R2, R3) : F_\alpha - F_\beta : \text{AbsCont}[a, b] \iff \alpha - \beta \ll \lambda ; \square$

**variation** :  $([a, b] \rightarrow \mathbb{R}) \rightarrow \mathbb{R}_+^\infty$

**variation**( $f$ ) =  $V_f := \sup \left\{ \sum_{k=1}^{\text{size}(P)-1} |f(P_k) - f(P_{k+1})| \mid P : \text{Partition}[a, b] \right\}$

**BoundedVariation** ::?  $([a, b] \rightarrow \mathbb{R})$

$f : \text{BoundedVariation} \iff V_f < \infty$

**ACIsBV** ::  $\forall f : \text{AbsCont}[a, b] \cdot f : \text{BoundedVariation}$

**Proof** =

**Assume**  $f : \text{AbsCont}[a, b]$ ,

**Assume**  $[a, b] : \text{Compact}$ ,

$(y, y') := \text{ExtremeValue}(f, [a, b]) : \mathbb{R}^2 : y = \max_{x \in [a, b]} f(x) : y' = \min_{x \in [a, b]} f(x)$ ,

$c := y - y' \in \mathbb{R}_+$ ,

$\delta := \mathfrak{O}\text{AbsCont}(f)(c) \in \mathbb{R}_{++} : \forall [c, d] : \forall n \in \mathbb{N} \cdot \forall [c, d] : n \rightarrow \text{Subinterval}[a, b]$

$:: \sum_{k=1}^n d_k - c_k \geq \delta \cdot \sum_{k=1}^n |f(c_k) - d(d_k)| \leq c$ ,

$n := \left\lceil \frac{b-a}{\delta} \right\rceil \in \mathbb{N}$ ,

Assume  $P : \text{Partition}[a, b]$ ,

$m := \text{size}(P) \in \mathbb{N}$ ,

$R1 := \text{Partition}(P) : \sum_{k=1}^{m-1} P_{k+1} - P_k = b - a$ ,

$K := \{i \in m - 1 : P_{i+1} - P_i > \delta\} : ?(m - 1)$ ,

$k := \#K \in \mathbb{Z}_+$ ,

$R1 \rightsquigarrow k \leq n$ ,

$L := K^{\mathbb{L}} : ?(m - 1)$ ,

$(l, A) := \text{JollyPartitioningLemma}(\delta, L) : \sum l \in \mathbb{N} : l + k \leq n . l \rightarrow \mathcal{B}[a, b] : \forall i \in l . \lambda(A_i) \leq \delta : \\ : \forall i \in L . \exists j \in l : [P_i, P_{i+1}] \subset A_j$ ,

Assume  $j \in l$

$I_j := \{i \in m - 1 : i \in A_j\} : ?(m - 1)$

$\lambda(A_j) \leq \delta \rightsquigarrow \sum_{i \in I_j} P_{i+1} - P_i \leq \delta \rightsquigarrow \sum_{i \in I_j} |f(P_{i+1}) - f(P_i)| \leq c$ ;

$\sum_{i=1}^{m-1} |f(P_i) - f(P_{i+1})| = \sum_{i \in K} |f(P_i) - f(P_{i+1})| + \sum_{i \in L} |f(P_i) - f(P_{i+1})| \leq \\ \leq kc + \sum_{j=1}^l \sum_{i \in I_j} |f(P_i) - f(P_{i+1})| \leq kc + lc \leq nc < \infty$ ;

$f : \text{BoundedVariation}$ ;  $\square$

**BVDecomposition** ::  $\forall f : \text{BoundedVariation}[a, b] . \exists F, G : \text{Increasing}[a, b] : f = F - G$

**Proof** =

Assume  $f : \text{BoundedVariation}[a, b]$ ,

$F := \Lambda x \in [a, b] . V_{f|_{[a, x]}} : \text{Increasing}[a, b]$ ,

$G := F - f : [a, b] \rightarrow \mathbb{R}$ ,

Assume  $x, y \in [a, b] : y > x$ ,

$G(y) - G(x) = V_{f|_{[a, y]}} - V_{f|_{[a, x]}} + f(x) - f(y) \leq V_{f|_{[x, y]}} - |f(x) - f(y)| \geq 0$ ;

$G : \text{Increasing}[a, b]$ ,

$f = F - G$ ;  $\square$

**ACDecomposition** ::  $\forall f : \text{AbsCont}[a, b] . \exists F, G : \text{Increasing} \ \& \ \text{AbsCont}[a, b] : f = F - G$



**ACIsIntegral** ::  $\forall F : \text{AbsCont}[a, b] . \exists f : \text{Integrable}([a, b], \mathcal{B}[a, b], \lambda) : \forall x \in [a, b] .$

$$. F(x) - F(a) = \int_a^x f d\lambda$$

**Proof** =

**Assume**  $F : \text{AbsCont}[a, b],$

$V, W := \text{ACDecomposition}(F) :: \text{Increasing} \ \& \ \text{AbsCont}[a, b] : F = V - W$

$V, W : \text{AbsCont}[a, b] \rightsquigarrow V, W : \mathcal{M}_{\text{TOP}}([a, b], \mathbb{R}),$

$V, W : \text{Increasing} \ \& \ \mathcal{M}_{\text{TOP}}([a, b], \mathbb{R}) \rightsquigarrow V, W : \text{Distribution}([a, b], \mathcal{B}[a, b]),$

$V, W : \text{AbsCont}[a, b] \rightsquigarrow V - W : \text{AbsCont}[a, b],$

$R1 := \text{AbsContDistr}(\mu_V, \mu_W) : \mu_V - \mu_W \ll \lambda,$

$f := \text{RadonNikodym}(\mu_V - \mu_W, \lambda, R1) : \text{Integrable}([a, b], \mathcal{B}[a, b], \lambda) : \forall A \in \mathcal{B}[a, b] .$

$$. (\mu_V - \mu_W)(A) = \int_A f d\lambda$$

**,Assume**  $x \in [a, b],$

$[a, x] \in \mathcal{B}[a, b],$

$$\int_a^x f d\lambda = \mu_V[a, x] - \mu_W[a, x] = V(x) - V(a) - W(x) + W(a) = F(x) - F(a); ; \square$$

### 3.4 Differentiation of Measures

$\text{RadonCharge} :: ?\text{Charge}(\mathbb{R}^d, \mathcal{B}\mathbb{R}^d)$

$H : \text{RadonCharge} \iff \forall A : \text{Bounded } \mathbb{R}^d . |H(A)| < \infty$

$\text{UpperRadonDifferential} :: \text{RadonCharge}(d) \rightarrow \mathbb{R}^d \rightarrow \mathbb{R}$

$\text{UpperRadonDifferential}(H)(x) = (\overline{D}\mu)(x) = \lim_{r \rightarrow 0} \sup_{C : \text{Cube}(d,r)} \frac{H(C)}{\lambda(C)}$

$\text{UpperRadonDifferential} :: \text{RadonCharge}(d) \rightarrow \mathbb{R}^d \rightarrow \mathbb{R}$

$\text{UpperRadonDifferential}(H)(x) = (\overline{D}\mu)(x) = \lim_{r \rightarrow 0} \sup_{C : \text{Cube}(d,r)} \frac{H(C)}{\lambda(C)}$

$\text{LowerRadonDifferential} :: \text{RadonCharge}(d) \rightarrow \mathbb{R}^d \rightarrow \mathbb{R}$

$\text{LowerRadonDifferential}(H)(x) = (\underline{D}\mu)(x) = \lim_{r \rightarrow 0} \inf_{C : \text{Cube}(d,r)} \frac{H(C)}{\lambda(C)}$

$\text{DifferentiableChargeAt} :: \mathbb{R}^d \rightarrow ?\text{RadonCharge}(d)$

$H : \text{DifferentiableChargeAt}(x) \iff (\underline{D}H)(x) = (\overline{D}H)(x)$

$\text{DifferentiableCharge} :: ?\text{RadonCharge}(d)$

$H : \text{DifferentiableCharge} \iff \forall x \in \mathbb{R}^d . H : \text{DifferentiableChargeAt}(x)$

$\text{RadonDifferentiate} :: \text{DifferentiableCharge}(d) \rightarrow \mathbb{R}^d \rightarrow \mathbb{R}$

$\text{RadonDifferentiate}(H)(x) = D H x := \overline{D} H x$

**DisjointCubesLemma** ::  $\forall n \in \mathbb{N} . \forall C : n \rightarrow \text{Cube}(d) .$

$$. \exists s \in n . \exists i : s \hookrightarrow n : \lambda \left( \bigcup_{j=1}^n C_j \right) \leq 3^d \sum_{k=1}^s \lambda(C_{i_k})$$

**Proof** =

**Assume**  $\in \mathbb{N}$

**Assume**  $C : n \rightarrow \text{Cube}(d),$

$C' := \text{sort}(C, <, \text{diam}),$

**Iterate over**  $k \in n$  **with**  $\mathbf{C}^1 = C', A_1 = \emptyset$

$i_k := \text{index}(\mathbf{C}_1^k)(C),$

$$A_k := \bigcup_{j=1}^k C_{i_j},$$

$$C^{k+1} := [\mathbf{C}_j^k : \mathbf{C}_j^k \cap A_k = \emptyset];$$

**until**  $\mathbf{C}^k = [];$

$s := \text{Dom } i,$

**Assume**  $j \in N,$

**Assume**  $c : \forall k \in s . C_j \cap C_{i_k} = \emptyset,$

$c \rightsquigarrow C^{s+1} \neq [] \rightsquigarrow \perp;;$

(1) :  $\forall j \in n . \exists k \in s . C_j \cap C_{i_k} \neq \emptyset,$

**Assume**  $k \in s,$

$B_k := \text{cube}(\text{center } C_{i_k} \text{ } 3\text{diam } C_{i_k}),$

**Assume**  $\ell \in n : \lambda(C_\ell) \leq \lambda(C_{i_k}) : C_\ell \cap C_{i_k} \neq \emptyset,$

$C_\ell \subset B_k;$

$$\bar{\mathfrak{O}}(B_k), \bar{\mathfrak{O}}(C), (1) \rightsquigarrow \lambda \left( \bigcup_{j=1}^n C_j \right) \leq \lambda \left( \bigcup_{j=1}^s B_j \right) \leq \sum_{j=1}^s \lambda(B_j) = 3^d \sum_{j=1}^s \lambda(C_{i_j});; \square$$

**ZeroDifferentialLemma** ::  $\forall \mu : \text{Lebesgue-Stieltjes}(\mathbb{R}^d, \mathcal{B}\mathbb{R}^d) . \forall A \in \mathcal{B}\mathbb{R}^d .$   
 $. \forall a : \mu(A) = 0 . D\mu|_A = 0 \text{ a . e . } [\lambda]$

**Proof** =

**Assume**  $\mu : \text{Lebesgue-Stieltjes}(\mathbb{R}^d, \mathcal{B}\mathbb{R}^d),$

**Assume**  $\forall A \in \mathcal{B}\mathbb{R}^d,$

**Assume**  $a : \mu(A) = 0$

**Assume**  $t \in \mathbb{R}_{++},$

$B := \{x \in A : D\mu x > t\},$

**Assume**  $K : \text{Compact}(B),$

**Assume**  $r \in \mathbb{R}_{++},$

**Assume**  $x \in K,$

$x \in K \rightsquigarrow x \in B \rightsquigarrow D\mu x > t \rightsquigarrow \exists q : \mathbb{N} \rightarrow \mathbb{R}_{++} : q \downarrow 0 : \forall n \in \mathbb{N} .$

$\frac{\mu \text{cube}(x, q_n)}{\lambda \text{cube}(x, q_n)} > t \text{ Extract},$

$a_n \downarrow 0 \rightsquigarrow \exists n \in \mathbb{N} . q_n < r \text{ Extract},$

$C_x := \text{cube}(x, q_n);$

$C : K \rightarrow \text{Cube}(\mathbb{R}^d) : \forall x \in K . x \in C_x,$

$C : \text{OCover}(K), K : \text{Compact}(B) \rightsquigarrow \exists O : \text{FSubOCover}(K, C) \text{ Extract},$

$n := \#O \in \mathbb{N},$

$k := \text{DisjointCubesLemma}(O),$

$\lambda(K) \leq \lambda\left(\bigcup_{i=1}^n O_i\right) \leq 3^d \sum_{i=1}^n \lambda(O_{k_i}) \leq \frac{3^k}{t} \sum_{i=1}^n \mu(O_{k_i}) = \frac{3^k}{t} \mu\left(\bigcup_{i=1}^n C_{k_i}\right) \leq \frac{3^k}{t} \mu(\mathbb{B}(K, r));$

$\lambda(K) \leq \frac{3^k}{t} \mu(K) \leq \frac{3^k}{t} \mu(A) = 0;$

$\lambda(B) = 0;$

$D\mu|_A = 0 \text{ a . e . } [\lambda];$

$\text{LebesgueDecompositionDerivatives} :: \forall H : \text{RadonCharge}(d) .$   
 $. \forall \alpha, \beta : \text{RadonCharge}(d) : (\alpha, \beta) = \text{LebesgueDecomposition } H .$   
 $. H : \text{RadonDifferentiableAt} \ \& \ DH = Df_\alpha \quad \text{a . e . } [\lambda]$

Proof =

Assume  $H : \text{RadonCharge}(d)$ ,

Assume  $\alpha, \beta : \text{Measure}(\mathbb{R}^d, \mathcal{B}\mathbb{R}^d) : (\alpha, \beta) = \text{LebesgueDecomposition } H$ ,

Assume  $a \in \mathbb{R}_{++}$ ,

$A := \{x \in \mathbb{R}^d : f_\alpha(x) < a\} : \mathcal{B}\mathbb{R}^d$ ,

$B := A^c : \mathcal{B}\mathbb{R}^d$

$\mu := \Lambda E \in \mathcal{B}\mathbb{R}^d . \int_{E \cap B} (f_\alpha - a) d\lambda : \text{Measure}(\mathbb{R}^d, \mathcal{B}\mathbb{R}^d),$

Assume  $r \in \mathbb{R}_{++}$ ,

Assume  $C : \text{Cube}(D) : \text{diam } C < r$ ,

$R1 :: \alpha(C) - a\lambda(C) = \int_C (f_\alpha - a) d\lambda \leq \int_{C \cap B} (f_\alpha - a) d\lambda,$

$\partial\mu \rightsquigarrow R_2 : \mu(A) = 0,$

$R_3 := \text{ZeroDifferentialLemma}(R_2) : D\mu|_A = 0 \quad \text{a . e . } [\lambda],$

$R_1 \rightsquigarrow R_4 : \frac{\alpha(C)}{\lambda(C)} \leq a + \frac{\mu(C)}{\lambda(C)};$

$R_5 : \overline{D}\alpha|_A \leq a \quad \text{a . e . } [\lambda|_{\mathcal{B}A}],$

$E_a := \{x \in \mathbb{R}^d : f_\alpha(x) < a < \overline{D}\alpha(x)\} \in \mathcal{B}\mathbb{R}^d,$

$R_5 \rightsquigarrow \lambda(E_a) = 0,$

$R_6 :: \{\overline{D}\alpha > f_\alpha\} \subset \bigcup_{q \in \mathbb{Q}_{++}} E_q,$

$R_6 \rightsquigarrow R_7 : \lambda\{\overline{D}\alpha > f_\alpha\} \leq \lambda\left(\bigcup_{q \in \mathbb{Q}_{++}} E_q\right) \leq \sum_{q \in \mathbb{Q}_{++}} \lambda(E_q) = 0,$

...

$D\alpha = f_\alpha \quad \text{a . e . } [\lambda],$

...

$D\beta = 0 \quad \text{a . e . } [\lambda],$

$DH = D\alpha + D\beta = f_\alpha \quad \text{a . e . } [\lambda] \square$

**MonotoneIsAlmostDiffrentiable** ::  $\forall f : \text{Increasing}[a, b] . f : \text{DifferentiableAt} \quad \text{a.e.} \quad [\lambda_{|\mathcal{B}[a, b]}]$

**Proof** =

**Assume**  $f : \text{Increasing}[a, b]$ ,

$R_1 := \text{MonotoneDisconts}(f) : \# \text{Discont}(f) \leq \aleph_0$ ,

$R_1 \leadsto \exists G : \text{DistributionFunction}[a, b] : f = G \quad \text{a.e.} \quad [\lambda_{|\mathcal{B}[a, b]}]$  **Extract**,

$R_2 := \text{LebesgueDecompositionDerivatives}(\mu_G) : (\mu_G : \text{RadonDifferentiableAt} \quad \text{a.e.} \quad [\lambda_{|\mathcal{B}[a, b]}])$

**Assume**  $x \in [a, b] : (\mu_G : \text{RadonDifferentiableAt}(x))$ ,

$\mu_G : \text{RadonDifferentiableAt}(x) \leadsto G : \text{ContAt}(x)$ ,

$$\lim_{h \rightarrow 0} \frac{G(x+h) - G(x)}{h} = \lim_{h \rightarrow 0} \lim_{k \rightarrow 0} \frac{G(x+h) - G(x-k)}{h+k} = D\mu_G(x);$$

$G : \text{DifferentiableAt} \quad \text{a.e.} \quad [\lambda_{|\mathcal{B}[a, b]}] \leadsto f : \text{DifferentiableAt} \quad \text{a.e.} \quad [\lambda_{|\mathcal{B}[a, b]}] \quad \square$

**ACHasNewtonProperty** ::  $\forall F : [a, b] \rightarrow \mathbb{R} .$

$$F : \text{AbsCont}[a, b] \iff \forall x \in [a, b] . F(x) - F(a) = \int_a^x F'(t) dt$$

**Proof** =

**Assume**  $F : \text{AbsCont}[a, b]$ ,

$(A, B) := \text{ACDecomposition} : \text{Increasing} \ \& \ \text{AbsCont}[a, b] : F = A - B$ ,

$dA := \text{MonotoneIsAlmostDiffrentiable}(A) : (A : \text{DifferentiableAt} \quad \text{a.e.} \quad [\lambda_{|\mathcal{B}[a, b]}])$ ,

$dB := \text{MonotoneIsAlmostDiffrentiable}(B) : (B : \text{DifferentiableAt} \quad \text{a.e.} \quad [\lambda_{|\mathcal{B}[a, b]}])$ ,

$dA, dB, \mathfrak{D}(A, B) \leadsto dF : (F : \text{DifferentiableAt} \quad \text{a.e.} \quad [\lambda_{|\mathcal{B}[a, b]}])$ ,

$$f := \text{ACIsIntegral}(F) : \text{Integrable}([a, b], \mathcal{B}[a, b], ) : F(x) - F(a) = \int_a^x f(t) dt,$$

$$H := \Lambda A \in \mathcal{B}[a, b] . \int_A f d\lambda : \text{RadonCharge}(1),$$

$E_1 := \text{LebesgueDecompositionDerivatives}(H) :: f = DH \quad \text{a.e.} \quad [\lambda_{|\mathcal{B}[a, b]}]$ ,

**Assume**  $x \in [a, b] : (F : \text{DifferentiableAt}(x))$ ,

$$F'(x) = \lim_{h \rightarrow 0} \frac{F(x+h) - F(x)}{h} = \lim_{h \rightarrow 0} \lim_{k \rightarrow 0} \frac{F(x+h) - F(x-k)}{h+k} = DH;$$

$E_1 \leadsto E_2 : f = F' \quad \text{a.e.} \quad [\lambda_{|\mathcal{B}[a, b]}]$ ,

$$E_2 \leadsto \forall x \in [a, b] . F(x) - F(a) = \int_a^x F'(t) dt;$$

$$I : F : \text{AbsCont}[a, b] \Rightarrow \forall x \in [a, b] . F(x) - F(a) = \int_a^x F'(t) dt$$

$$\text{Assume } a : \text{forall } x \in [a, b] . F(x) - F(a) = \int_a^x F'(t) dt,$$

**ACIsIntegral**(a)  $\leadsto F : \text{AbsCont}[a, b]$ ;

$$F : \text{AbsCont}[a, b] \iff \forall x \in [a, b] . F(x) - F(a) = \int_a^x F'(t) dt \square$$

**Assume**

**VectorChangeOfVariable** ::  $\forall V, W : \text{Open}(\mathbb{R}^d) . \forall f : \text{Integrable}(W, \mathcal{B}W, \lambda) . \forall T : \text{Iso}_{\text{DIFF}}(V, W) .$

$$\int_W f d\lambda = \int_V |\det \nabla T|(f \circ T) d\lambda$$

### 3.5 Categorical Viewpoint:Developing Borel-Null [!]



## 4 Convergence in Measure

### 4.1 Measure Topology

`measurePseudoMetrics` ::  $\prod (\Omega, \mathcal{F}, \mu) \in \text{MEAS} . \text{PseudoDistance}(\text{Measurable}(\Omega, \mathcal{F}))$

`measurePseudoMetrics`( $f, g$ ) =  $d_\mu(f, g) := \inf_{\delta > 0} \mu\{\omega \in \Omega : |f(\omega) - g(\omega)| > \delta\} + \delta$

`implicit` ::  $\text{Measure}(\Omega, \mathcal{F}) \rightarrow \text{Topology}(\text{Measurable}(\Omega, \mathcal{F}))$

$\mu := \text{pseudometricTopology}(d_\mu)$

`ConvergenceInMeasure` ::  $\forall f : \mathbb{N} \rightarrow \text{Measurable}(\Omega, \mathcal{F}) . \forall \phi : \text{Measurable}(\Omega, \mathcal{F}) .$

$f \rightarrow_\mu \phi \iff \forall \epsilon \in \mathbb{R}_{++} . \lim_{n \rightarrow \infty} \mu\{\omega \in \Omega : |f_n(\omega) - \phi| > \epsilon\} = 0$

`Proof` =

`Assume`  $f : \mathbb{N} \rightarrow \text{Measurable}(\Omega, \mathcal{F})$ ,

`Assume`  $\phi : \text{Measurable}(\Omega, \mathcal{F})$ ,

`Assume`  $C : f \rightarrow_\mu \phi$ ,

`Assume`  $\epsilon \in \mathbb{R}_{++}$ ;

`Assume`  $a \in \mathbb{R}_{++} : a < \epsilon$ ,

`Assume`  $L : \lim_{n \rightarrow \infty} \mu\{\omega \in \Omega : |f_n(\omega) - \phi| > \epsilon\} > a$ ,

`Assume`  $\delta \in \mathbb{R}_{++}$ ,

`Assume Alternative`  $\delta \geq a$ ,

$\forall n \in \mathbb{N} . \mu\{\omega \in \Omega : |f_n(\omega) - \phi(\omega)| > \delta\} + \delta \geq \delta \geq a$ ;

`Close Alternative`  $\delta < a$ ,

$\exists N : \text{Infinite}(\mathbb{N}) . \forall n \in N . \mu\{\omega \in \Omega : |f_n(\omega) - \phi(\omega)| > \delta\} + \delta \geq \mu\{\omega \in \Omega : |f_n(\omega) - \phi(\omega)| > \delta\} \geq a$ ;

$\exists N : \text{Infinite}(\mathbb{N}) . \forall n \in N . \mu\{\omega \in \Omega : |f_n(\omega) - \phi(\omega)| > \delta\} + \delta \geq a$ ;

$\exists N : \text{Infinite}(\mathbb{N}) . \forall n \in N . d_\mu(f_n, \phi) \geq a \rightsquigarrow \lim_{n \rightarrow \infty} f_n \neq \phi \rightsquigarrow \perp ; ; ;$

$\forall \epsilon \in \mathbb{R}_{++} . \lim_{n \rightarrow \infty} \mu\{\omega \in \Omega : |f_n(\omega) - \phi| > \epsilon\} = 0$ ;

...

`Assume`  $A : \forall \epsilon \in \mathbb{R}_{++} . \lim_{n \rightarrow \infty} \mu\{\omega \in \Omega : |f_n(\omega) - \phi| > \epsilon\} = 0$ ,

`Assume`  $\epsilon \in \mathbb{R}_n$ ,

$L := A(\epsilon/2) : \lim_{n \rightarrow \infty} \mu\{\omega \in \Omega : |f_n(\omega) - \phi| > \epsilon/2\} = 0$ ,

$N := L(\epsilon/2) \in \mathbb{N} : \forall n \in \mathbb{N} : n \geq N . \mu\{\omega \in \Omega : |f_n(\omega) - \phi| > \epsilon/2\} < \epsilon/2$ ,

`Assume`  $n \in \mathbb{N} : n \geq N$ ,

$d_\mu(f_n, \phi) = \inf_{\delta > 0} \mu\{\omega \in \Omega : |f_n(\omega) - \phi(\omega)| > \delta\} + \delta \leq \mu\{\omega \in \Omega : |f_n(\omega) - \phi(\omega)| > \epsilon/2\} + \epsilon/2 < \epsilon ; ; ;$

□

## 4.2 Comparisson with other modes of convergence

**LpImpleyMeasure** ::  $\forall f : \mathbb{N} \rightarrow L^p(\Omega, \mathcal{F}, \mu) . \forall \phi : L^p(\Omega, \mathcal{F}, \mu) . f \rightarrow_{L^p} \phi \Rightarrow f \rightarrow_{\mu} \phi$

**Proof** =

**Assume**  $f : \mathbb{N} \rightarrow L^p(\Omega, \mathcal{F}, \mu)$ ,

**Assume**  $\phi : L^p(\Omega, \mathcal{F}, \mu)$ ,

**Assume**  $C : f \rightarrow_{L^p} \phi$ ,

**Assume**  $\epsilon : \mathbb{R}_{++}$ ,

**Assume**  $n : \mathbb{N}$ ,

$I := \text{ChebishevIneq}(|f_n - \phi|, \epsilon) : \mu\{\omega \in \Omega : |f_n(\omega) - \phi(\omega)| < \epsilon\} \leq \frac{\|f_n(\omega) - \phi(\omega)\|_p}{\epsilon^p};$

$\leadsto R := \text{LimIneq}(\text{d}(\rightarrow_{L^p})(C), \text{IneqLim}(\cdot)) : \lim_{n \rightarrow \infty} \mu\{\omega \in \Omega : |f_n(\omega) - \phi(\omega)| < \epsilon\} \leq$

$\leq \lim_{n \rightarrow \infty} \frac{\|f_n - \phi\|_p}{\epsilon^p} = 0;$

$\leadsto R := \text{ConvergenceInMeasure}(f, \phi, \cdot) : f \rightarrow_{\mu} \phi;;;$

□

**UniformAEConvergence** ::  $\prod(\Omega, \mathcal{F}, \mu) : \text{MEAS} .$

$. ? \left( (\mathbb{N} \rightarrow \text{Measurable}(\Omega, \mathcal{F}, \mu)) \times \text{Measurable}(\Omega, \mathcal{F}, \mu) \right)$

$(f, \phi) : \text{UniformAEConvergence} \iff f \rightrightarrows_{\mu} \phi \iff \forall \epsilon : \mathbb{R}_{++} .$

$. \exists A \in \mathcal{F} : \mu(A) < \epsilon : f|_{A^c} \rightrightarrows \phi|_{A^c}$

**AEUnifImpleyMeasure** ::  $\forall f : \mathbb{N} \rightarrow \text{Measurable}(\Omega, \mathcal{F}, \mu) . \forall \phi : \text{Measurable}(\Omega, \mathcal{F}, \mu) .$

$. f \rightrightarrows_{\mu} \phi \Rightarrow f \rightarrow_{\mu} \phi$

**Proof** =

**Assume**  $f : \mathbb{N} \rightarrow \text{Measurable}(\Omega, \mathcal{F}, \mu)$ ,

**Assume**  $\phi : \text{Measurable}(\Omega, \mathcal{F}, \mu)$ ,

**Assume**  $C : f \rightrightarrows_{\mu} \phi$ ,

**Assume**  $\epsilon : \mathbb{R}_{++}$ ,

**Assume**  $a : \mathbb{R}_{++}$ ,

$A := C(a) : \mathcal{F} : \mu(A) < a : f|_{A^c} \rightrightarrows \phi|_{A^c},$

$N := \text{d}(A)(\epsilon) : \forall n \in \mathbb{N} : n \geq N . \{\omega \in \Omega : |f_n(\omega) - \phi(\omega)| > \epsilon\} \subset A,$

**Assume**  $n : \mathbb{N} : n \geq N,$

$S := \text{d}(N)(n) : \{\omega \in \Omega : |f_n(\omega) - \phi(\omega)| > \epsilon\} \subset A,$

$I := \text{MeasureMonotonicity}(\mu, S) \text{d}(A) : \mu\{\omega \in \Omega : |f_n(\omega) - \phi(\omega)| > \epsilon\} \leq \mu(A) < a;;$

$\leadsto L := \text{d}[\text{Limit}](\cdot) : \lim_{n \rightarrow \infty} \mu\{\omega \in \Omega : |f_n(\omega) - \phi(\omega)| > \epsilon\} = 0;$

$\leadsto R := \text{ConvergenceInMeasure}(f, \phi, \cdot) : f \rightarrow_{\mu} \phi;;;$

□

**AEUnifImPLYAE** ::  $\forall f : \mathbb{N} \rightarrow \text{Measurable}(\Omega, \mathcal{F}, \mu) . \forall \phi : \text{Measurable}(\Omega, \mathcal{F}, \mu) .$   
 $. f \Rightarrow_{\mu} \phi \Rightarrow f \rightarrow \phi \quad \text{a.e.} \quad [\mu]$

**Proof** =

**Assume**  $f : \mathbb{N} \rightarrow \text{Measurable}(\Omega, \mathcal{F}, \mu),$

**Assume**  $\phi : \text{Measurable}(\Omega, \mathcal{F}, \mu),$

**Assume**  $C : f \Rightarrow_{\mu} \phi,$

**Assume**  $n : \mathbb{N},$

$A_n := C(1/n) : \mathcal{F} : \mu(A_n) < \frac{1}{n} : f|_{A_n^c} \Rightarrow \phi|_{A_n^c},$

$L := \text{UnifImPLYPointwise}(\partial_2 A_n) : f|_{A_n^c} \rightarrow \phi|_{A_n^c};$

$\leadsto A := (\cdot) : \prod n : \mathbb{N} . \mathcal{F} : \mu(A_n) < \frac{1}{n} : f|_{A_n^c} \rightarrow \phi|_{A_n^c},$

$B := \bigcup_{n=1}^{\infty} A_n^c : \mathcal{F},$

$L := \partial(B.\partial(A)) : f|_B \rightarrow \phi|_B,$

$Z := \text{IneqLim}(\text{LimIneq} \lambda n \in \mathbb{N} . \text{MeasurMonotonicity} \left( A_n, \bigcap_{k=1}^{\infty} A_k \right) \text{UnionCompliment}(B),$

$, \text{LimEq}(\text{Lim}(\lambda n \in \mathbb{N} . 1/n), \mu(A), \partial_1 A)) : \mu(B^c) = \mu\left(\bigcap_{n=1}^{\infty} A_n\right) \leq \lim_{n \rightarrow \infty} \mu(A_n) = 0;$

$\leadsto R := \partial \text{a.e.} \quad [\mu] (\Lambda \omega \in \Omega . f(\omega) \rightarrow \phi(\omega))(\cdot) : f \rightarrow \phi \quad \text{a.e.} \quad [\mu]; ; ,$

□

**AEUnifSubseq** ::  $\forall f : \mathbb{N} \rightarrow \text{Measurable}(\Omega, \mathcal{F}, \mu) . \forall \phi : \text{Measurable}(\Omega, \mathcal{F}) : f \rightarrow_{\mu} \phi .$   
 $. \exists n : \text{Subsequer} : f_n \Rightarrow_{\mu} \phi$

**Proof** =

**Assume**  $f : \mathbb{N} \rightarrow \text{Measurable}(\Omega, \mathcal{F}),$

**Assume**  $\phi : \text{Measurable}(\Omega, \mathcal{F}, \mu) : f \rightarrow_{\mu} \phi,$

$(T) := \partial(\mu) \text{ as Topology} : (\mu : \text{Semimetrizable}),$

$(C) := \text{ConvergingIsCauchy}(T, f, \partial(\phi)) : (f : \text{Cauchy}),$

**Assume**  $k : \mathbb{N},$

$M := \partial \text{Cauchy}(f)(2^{-k}) : \mathbb{N} : \forall n, m \in \mathbb{N} : n \geq M : m \geq M . d_{\mu}(f_n, f_m) < 2^{-k},$

$N_k := \max(M, N_{k-1} + 1) : \mathbb{N};$

$\leadsto N := (\cdot) : \text{Subsequer} : \forall k : \mathbb{N} . \forall n, m : \mathbb{N} : n \geq N_k : m \geq N_k . d_{\mu}(f_n, f_m) < 2^{-k},$

$g := f_N : \mathbb{N} \rightarrow \text{Measurable} : \forall k : \mathbb{N} . d_{\mu}(g_k, g_{k+1}) < 2^{-k},$

**Assume**  $k : \mathbb{N},$

$A_k := \{\omega \in \Omega : |g_k(\omega) - g_{k+1}(\omega)| > 2^{-k}\} : \mathcal{F},$

$(I) := \mathfrak{D}(g, A) : 2^{-k} \geq d_\mu(g_k, g_{k+1}) + \epsilon =$   
 $= [x < 2^{-k}] = \{\omega \in \Omega : |g_k(\omega) - g_{k+1}(\omega)| > x\} + x \geq \mu(A_k);$   
 $\leadsto A := (\cdot) : \mathbb{N} \rightarrow \mathcal{F} : \forall k : \mathbb{N} . \mu(A_k) \leq 2^{-k},$   
 $\alpha := \limsup A : \mathcal{F},$   
 $Z := \text{BorellCanteli}(\alpha, \mathfrak{D}(A)) : \mu(\alpha) = 0,$   
**Assume**  $\omega : \alpha^{\mathbb{C}},$   
 $FF := \mathfrak{D}(\limsup)(\alpha, A, \omega) : \#\{k \in \mathbb{N} : \omega \in A_k\} < \aleph_0,$   
 $L := \text{CauchyCriterion}(f(\omega), A, FF) : (f(\omega) : \text{Cauchy}),$   
 $R := \mathfrak{D}\text{Complete}(\mathbb{R}, f(\omega)) : (f(\omega) : \text{Converge}(\mathbb{R}));$   
 $\leadsto AC := \mathfrak{D}(\text{a.e.} \ [\mu])(\cdot, Z) : (f : \text{Converge}(\mathbb{R})) \quad \text{a.e.} \ [\mu],$   
 $\gamma := [AC] \lim_{n \rightarrow \infty} g_n : \text{Measurable}(\Omega, \mathcal{F}),$   
**Assume**  $k : \mathbb{N},$   
 $B_k := \bigcup_{n=k}^{\infty} A_n : \mathcal{F};$   
 $\leadsto B := (\cdot) : \mathbb{N} \rightarrow \mathcal{F},$   
 $L := \mathfrak{D}(B, A) : \lim_{n \rightarrow \infty} \mu(B_n) = 0,$   
 $W := \text{WeierstrassMTest}(\mathfrak{D}(B), L) : g \rightrightarrows_\mu \gamma,$   
 $LM := \text{AEUnifImplyMeasure}(g, \gamma, W) : g \rightarrow_\mu \gamma,$   
 $LL := \text{ConvergingSubseqAgrees}(f, g, \mathfrak{D}f, LM) : g \rightarrow_\mu \phi,$   
 $E := \text{TopoEquelInMeasure}(LM, LL) : \phi = \gamma \quad \text{a.e.} \ [\mu],$   
 $R := \text{TopoEquelAEUniform}(W, E) : g \rightrightarrows_\mu \phi,$   
 $\square$

**AEProbabilityLemma** ::  $\forall P : \text{Probability}(\Omega, \mathcal{F}) . \forall f : \mathbb{N} \rightarrow \text{Measurable}(\Omega, \mathcal{F}) .$   
 $\quad . \forall \phi : \text{Measurable}(\Omega, \mathcal{F}) . f \rightarrow \phi \quad \text{a.e.} \ [P] \iff$   
 $\iff \forall \delta : \mathbb{R}_{++} . \lim_{n \rightarrow \infty} P \left( \bigcup_{k=n}^{\infty} \{\omega \in \Omega : |f_n(\omega) - \phi(\omega)| \geq \delta\} = 0 \right)$

**Proof** =

**Assume**  $P : \text{Probability}(\Omega, \mathcal{F}),$   
**Assume**  $f : \mathbb{N} \rightarrow \text{Measurable}(\Omega, \mathcal{F}),$   
**Assume**  $\phi : \text{Measurable}(\Omega, \mathcal{F}),$   
**Assume**  $\delta : \mathbb{R}_{++},$   
**Assume**  $n : \mathbb{N},$   
 $B_n^\delta := \{\omega \in \Omega : |f_n(\omega) - \phi(\omega)| \geq \delta\} : \mathcal{F};$   
 $\leadsto B^\delta := (\cdot) : \mathbb{N} \rightarrow \mathcal{F},$   
 $\beta^\delta := \limsup B : \mathcal{F},$   
 $CD := \mathfrak{D}(\limsup)(B^\delta)(\beta^\delta) : \bigcup_{k=n}^{\infty} B_k^\delta \downarrow_n \beta^\delta,$

$$MC := \text{MeasureLowerContinuity}(P, CD) : \lim_{n \rightarrow \infty} P \left( \bigcup_{k=n}^{\infty} B_k^{\delta} \right) = P(\beta^{\delta});$$

$$\leadsto E := \dots : \{\omega \in \Omega : f(\omega) \not\rightarrow \phi(\omega)\} = \bigcup_{\delta \in \mathbb{R}_{++}} \beta_{\delta},$$

$$\text{Assume } C : f \rightarrow \phi \text{ a.e. } [P],$$

$$\text{Assume } \delta : \mathbb{R},$$

$$R := \dots : 0 = P(\{\omega \in \Omega : f(\omega) \not\rightarrow \phi(\omega)\}) = P \left( \bigcup_{x \in \mathbb{R}_{++}} \beta_x \right) \geq P(\beta_{\delta}) \leadsto$$

$$\leadsto P(\beta_{\delta}) = 0 \leadsto \lim_{n \rightarrow \infty} \left( \bigcup_{k=n}^{\infty} B_k^{\delta} \right) = 0;;$$

$$\text{Assume } A : (*),$$

$$\begin{aligned} R &:= \dots : P(\{\omega \in \Omega : f(\omega) \not\rightarrow \phi(\omega)\}) = P \left( \bigcup_{x \in \mathbb{R}_{++}} \beta^{\delta} \right) = \\ &= P \left( \bigcup_{x \in \mathbb{Q}_{++}} \beta^{\delta} \right) \leq \sum_{x \in \mathbb{Q}_{++}} P(\beta^{\delta}) = \sum_{x \in \mathbb{Q}_{++}} \lim_{n \rightarrow \infty} \left( \bigcup_{k=n}^{\infty} B_k^x \right) = 0 \leadsto \\ &\leadsto f \rightarrow \phi \text{ a.e. } [P];; \end{aligned}$$

□

$$\text{Egoroff} :: \forall P : \text{Probability}(\Omega, \mathcal{F}) . \forall f : \mathbb{N} \rightarrow \text{Measurable}(\Omega, \mathcal{F}) .$$

$$\forall \phi : \text{Measurable} \Omega, \mathcal{F} : f \rightarrow \phi \text{ a.e. } [P] . f \rightrightarrows_P \phi$$

$$\text{Proof} =$$

$$\text{Assume } P : \text{Probability}(\Omega, \mathcal{F}),$$

$$\text{Assume } f : \mathbb{N} \rightarrow \text{Measurable}(\Omega, \mathcal{F}),$$

$$\text{Assume } \phi : \text{Measurable} \Omega, \mathcal{F} : f \rightarrow \phi \text{ a.e. } [P],$$

$$\text{Assume } \epsilon : \mathbb{R}_{++},$$

$$\text{Assume } i : \mathbb{N},$$

$$A_j := \bigcup_{k=n}^{\infty} \{\omega \in \Omega : |f_k(\omega) - \phi(\omega)| \geq i^{-1}\} : \mathcal{F},$$

$$N_j := \text{AEProbabilityLemma}(P, f, \phi)(\epsilon 2^{-i}) : \mathbb{N} : \forall n \in \mathbb{N} : n \geq N_j . P(A_j) < \epsilon 2^{-i};$$

$$\leadsto N := (\cdot) : \mathbb{N} \rightarrow \mathbb{N},$$

$$\alpha := \bigcup A : \mathcal{F},$$

$$I := \text{Measure}(P)(\text{Measure}(A, \alpha)) : P(\alpha) \leq \sum_{i=1}^{\infty} P(A_i) < \epsilon,$$

$$U := \text{Measure} A : \forall \omega \in A^{\mathbb{C}} . f(\omega) \rightrightarrows \phi(\omega);$$

$$\leadsto R := \text{Measure}[\rightrightarrows_P](\cdot) : f \rightrightarrows_P \phi;$$

□

## 5 Products over Measure spaces

## 5.1 Product Measure Theorem

$$\text{SigmaAlgebraProduct} :: \sigma\text{-Algebra}(A) \rightarrow \sigma\text{-Algebra}(B) \rightarrow \text{Set}(A \times B)$$
$$\text{SigmaAlgebraProduct}(\mathcal{A}, \mathcal{B}) = \mathcal{A} \times \mathcal{B} := \{a \times b \mid a \in \mathcal{A}, b \in \mathcal{B}\}$$

**BorProduct** :: BOR  $\rightarrow$  BOR  $\rightarrow$  BOR

$$\text{BorProduct}((A, \mathcal{A}), (B, \mathcal{B})) = (A, \mathcal{A}) \times (B, \mathcal{B}) := (A \times B, \sigma(\mathcal{A} \times \mathcal{B}))$$

Uniformly  $\sigma$ -Finite :: ?( $X \rightarrow \text{Measure}(Y)$ )

$$\mu : \text{Uniformly } \sigma\text{-Finite} \iff \exists b : \mathbb{N} \rightarrow \mathcal{F}_Y : \exists k : \mathbb{N} \rightarrow \mathbb{R}_+ : \bigcup_{n=1}^{\infty} b_n = Y : \\ : \forall x \in X . \forall n \in \mathbb{N} . \mu(x, b_n) \leq k_n$$
$$\text{SlicingMeasure} :: \prod X \in \text{MEAS} . \prod Y \in \text{BOR} . X \rightarrow \text{Measure}(Y)$$
$$\mu : \text{SlicingMeasure} \iff \forall b \in \mathcal{F}_Y . \Lambda x \in X . \mu(x, b) : \text{Measurable}(F_{\text{BOR}} X)$$
$$\text{RectangularAlgrebraTHM} :: \forall X, Y : \text{BOR} . \forall G : \text{MonotoneClass}(X \times Y) : \mathcal{F}_X \times \mathcal{F}_Y \subset G . \sigma(\mathcal{F}_X \times \mathcal{F}_Y) \subset G$$

Proof =

Assume  $x \times y : \mathcal{F}_X \times \mathcal{F}_Y$ ,

$$(1) := \text{ProductComplement}(x \times y) : (x \times y) = x^{\complement} \times y \cap x \times y^{\complement} \cap x^{\complement} \times y^{\complement},$$
$$(2) := \mathfrak{d}\text{MonotoneClass}(1, \mathfrak{d}(G)) : (x \times y)^{\mathfrak{C}};$$
$$\sim (\mathcal{F}_X \times \mathcal{F}_Y, 1) := (\mathcal{F}_X \times \mathcal{F}_Y, \text{ComplementClosed}(\cdot) : \text{ComplementClosed}(G),$$
$$(2) := \text{MonotoneClassTHM}(1) : \sigma(\mathcal{F}_X \times F_Y) \subset G;$$
☐
$$\text{MeasurableSection} :: \forall X, Y : \text{BOR} . \forall A : \mathcal{F}_{X \times Y} . \forall x : X . \text{section}(A, x) \in \mathcal{F}_Y$$

Proof =

$$B := \{A \in \mathcal{F}_{X \times Y} : \text{section}(A, x) \in \mathcal{F}_Y\} : \sigma\text{-Algebra}(X \times Y),$$
$$(I) := \mathfrak{F}B : \{a \times b | a \in F_X, b \in F_Y\} \subset B,$$
$$(II) := \breve{\partial}(\mathcal{F}_X \times F_Y)(\breve{\partial}(\sigma)(I)) : \mathcal{F}_{X \times Y} \subset B \rightsquigarrow \mathcal{F}_{X \times Y} = B; ; ;$$
☐

**MeasurableSlicing** ::  $\forall S : \text{SlicingMeasure}(X, U) . \forall A : \mathcal{F}_{X \times Y} .$

$\Lambda x \in X . S(x, \text{section}(A, x)) : \text{Measurable}(F_{\text{BOR}}X)$

**Proof** =

$B := \{A \in \mathcal{F}_{X \times Y} : \Lambda x \in X . S(x, \text{section}(A, x)) : \text{Measurable}(F_{\text{BOR}}X)\} :$

$: \text{Set}(F_{\text{BOR}}X \times Y),$

**Assume**  $b : \mathbb{N} \rightarrow B,$

**Assume**  $\beta : \mathcal{F}_{X \times Y} : b \uparrow \beta,$

(1) := **SectionIsMonotonic**( $b, \beta$ ) :  $\forall x : X . \text{section}(x, b_n) \uparrow \text{section}(x, \beta),$

(2) := **MeasureUpperContinuity**( $\Lambda x \in X . S(x, b), (1)$ ) :  $\Lambda x \in X . S(x, b_n) \uparrow \Lambda x \in X . S(x, \beta),$

(3) := **MonotoneConvergenceTHM**(2) :  $(x \in X . S(x, \beta) : \text{Measurable}(F_{\text{BOR}}X)),$

(4) :=  $\exists(B)(3) : \beta \in B;;$

$\leadsto (1\star) := \text{UniversalIntroduction}(\cdot) : \forall b : \mathbb{N} \rightarrow B . \forall \beta : \mathcal{F}_{X \times Y} : b \uparrow \beta . \beta \in B,$

**Assume**  $b : \mathbb{N} \rightarrow B,$

**Assume**  $\beta : \mathcal{F}_{X \times Y} : b \downarrow \beta,$

(1) := **SectionIsMonotonic**( $b, \beta$ ) :  $\forall x : X . \text{section}(x, b_n) \downarrow \text{section}(x, \beta),$

(2) := **MeasureLowerContinuity**( $\Lambda x \in X . S(x, b), (1)$ ) :  $\Lambda x \in X . S(x, b_n) \downarrow \Lambda x \in X . S(x, \beta),$

(3) := **MonotoneConvergenceTHM**(2) :  $(x \in X . S(x, \beta) : \text{Measurable}(F_{\text{BOR}}X)),$

(4) :=  $\exists(B)(3) : \beta \in B;;$

$\leadsto (2\star) := \text{UniversalIntroduction}(\cdot) : \forall b : \mathbb{N} \rightarrow B . \forall \beta : \mathcal{F}_{X \times Y} : b \downarrow \beta . \beta \in B,$

(1) :=  $\exists \text{MonotoneClass}(1\star, 2\star) : B : \text{MonotoneClass}(X \times Y),$

**Assume**  $a : \mathcal{F}_X,$

**Assume**  $b : \mathcal{F}_Y,$

(2) :=  $\exists \text{section}(a \times b) : \Lambda x \in X . S(x, \text{section}(x, a \times b)) = \Lambda x \in X . S(x, b),$

(3) :=  $\exists \text{SlicingMeasure}(S)(b)(2) : (\Lambda x \in X . S(x, \text{section}(x, a \times b)) : \text{Measurable}(F_{\text{BOR}}X)),$

(4) :=  $\exists B(3) : a \times b \in B;$

$\leadsto (2) := \exists \mathcal{F}_X \times \mathcal{F}_Y(\cdot) : \mathcal{F}_X \times \mathcal{F}_Y \subset B,$

(3) := **RectangularAlgebraTHM**( $X, Y, B$ )(2) :  $\text{Alg}(F_X \times F_Y) \subset B,$

(4) := **MonotoneClassTHM**(1, 3) :  $\sigma(\mathcal{F}_X \times \mathcal{F}_Y) \subset B,$

(5) := **SetEqIntroduction**(4,  $\exists B$ ) :  $\mathcal{F}_{X \times Y} = B;;$

□

**ProductMeasureTheorem** ::  $\forall X : \text{MEAS} . \forall Y : \text{BOR} . \forall S : \text{SlicingMeasure}(X, Y) .$

$$. \exists ! \gamma : \text{Measure}(F_{\text{BOR}} X \times Y) : \forall A : \mathcal{F}_{F_{\text{BOR}} X \times Y} . \gamma(A) = \int_X S(x, \text{section}(A, x)) d\mu_X$$

**Proof** =

$$\gamma := \Lambda A \in \mathcal{F}_{F_{\text{BOR}} X \times Y} . \int_X S(x, \text{section}(A, x)) d\mu_X(x) : \mathcal{F}_{F_{\text{BOR}} X \times Y} \rightarrow \mathbb{R}_+^\infty,$$

**Assume**  $A : \text{Disjoint}(\mathbb{N}, \mathcal{F}_{F_{\text{BOR}} X \times Y}),$

$$(1) := \text{Measure}(S(x, \cdot)) : \int_X S(x, \text{section}\left(\bigcap_{n=1}^\infty A_n, x\right)) d\mu_X(x) = \int_X \sum_{i=1}^n S(x, \text{section}(A_n, x)) d\mu_X(x),$$

$$(2) := \text{IntegralSum}(2) : \int_X S(x, \text{section}\left(\bigcap_{n=1}^\infty A_n, x\right)) d\mu_X(x) = \sum_{n=1}^\infty \int_X S(x, \text{section}(A_n, x)) d\mu_X(x),$$

$$(3) := \text{Measure}(\gamma) : \gamma\left(\bigcup_{n=1}^\infty A_n\right) = \sum_{n=1}^\infty \gamma(A_n);$$

$$\leadsto (1) := \text{Measure}(\gamma) : (\gamma : \text{Measure}(\mathcal{F}_{F_{\text{BOR}} X \times Y}));$$

□

**productMeasure** ::  $\text{SlicingMeasure}(X, Y) \rightarrow \text{Measure}(\mathcal{F}_{X \times Y})$

**productMeasure** ( $S$ ) := **ProductMeasureTHM**( $S$ )

**ProductProbabilityTheorem** ::  $\forall X : \text{ProbabilitySpace} . \forall Y : \text{BOR} . \forall P : \text{SlicingMeasure} :$

$$\forall x : X . S(x, Y) = 1 . \text{productMeasure}(P) : \text{Probability}(X \times Y)$$

**Proof** =

$$\mathbb{P} := \text{productMeasure}(P) : \text{Measure}(\mathcal{F}_{X \times Y}),$$

$$(1) := \text{EqE}(\text{section}(\text{SlicingMeasure}(P), X \times Y)) :$$

$$: \int_X P(x | \text{section}(X \times Y, x)) d\mu_X(x) = \int_X P(x | Y) d\mu_X(x),$$

$$(2) := (1) \text{EqE}(\text{section}(P)) : \int_X P(x | \text{section}(X \times Y, x)) d\mu_X(x) = \int_X d\mu_X(x),$$

$$(3) := (2) \text{Probability}(\mu_X) : \int_X P(x | \text{section}(X \times Y, x)) d\mu_X(x) = 1,$$

$$(4) := \text{P}(X \times Y(3)) : \mathbb{P}(X \times Y) = 1,$$

$$(*) := \text{Probability} : (\mathbb{P} : \text{Probability}(X \times Y));$$

□



**ProductSFTHM** ::  $\forall S : \text{SlicingMeasure} \ \& \ \text{Uniformly } \sigma\text{-Finite} (X, Y) : (\mu_X : \sigma\text{-Finite} (X)) .$   
 $. \text{productMeasure}(S) : \sigma\text{-Finite} (X \times Y)$

**Proof** =

$$(B, b) := \mathfrak{d}(\text{Uniformly } \sigma\text{-Finite} (X \times Y))(S) : \sum B : \mathbb{N} \rightarrow \mathcal{F}_Y : \bigcup_{n=1}^{\infty} B_n = Y .$$

$$. \mathbb{N} \rightarrow \mathbb{R}_+ : \forall x : X . \forall n : \mathbb{N} . S(x, B_n) \leq b_n,$$

$$A := \mathfrak{d}\sigma\text{-Finite} (X) (\mu_X) : \mathbb{N} \rightarrow \mathcal{F}_X : \bigcup_{n=1}^{\infty} A_n = X : \mu_X(A) < \infty,$$

$$(1) := \text{ProductPartition}(A, B) : \bigcup_{n=1}^{\infty} \bigcup_{m=1}^{\infty} A_n \times B_m = X \times Y,$$

$$\gamma := \text{productMeasure}(S) : \text{Measure}(X \times Y),$$

**Assume**  $n, m : \mathbb{N}$ ,

$$(2) := \mathfrak{d}\gamma(\text{Assume } n \times B_n) \text{IntIneq}(\mathfrak{d}b_m) \text{MeasureAsIntegral}(\mu_X, A_n) \mathfrak{d}(A_n) :$$

$$: \gamma(A_n \times B_m) = \int_{A_n} S(x, B_m) \mathrm{d}\mu_X(x) \leq \int_{A_n} b_m \mathrm{d}\mu_X = b_m \mu_X(A) < \infty;$$

$$\leadsto (2) := \text{UI} : \forall n, m : \mathbb{N} . \gamma(A_n \times B_m) < \infty,$$

$$(*) := \mathfrak{d}^{-1}\sigma\text{-Finite} (X \times Y) (\gamma, A \times B, 1, 2) : (\gamma : \sigma\text{-Finite} (X \times U));$$

□

**productOfMeasures** ::  $\text{MEAS} \rightarrow \text{MEAS} \rightarrow \text{MEAS}$

$$\text{productOfMeasures} ((X, \mathcal{F}, \mu), (Y, \mathcal{G}, \nu)) = \mu \times \nu := \left( X \times Y, \sigma(\mathcal{F} \times \mathcal{G}), A \mapsto \int_X \nu(\text{section}(A, x)) \mathrm{d}\mu(x) \right)$$

**ClassicalPMTHM** ::  $\forall X, Y : \text{MEAS} . \forall A \times B : F_X \times F_Y . \mu_X \times \mu_Y(A \times B) = \mu_X(A) \mu_Y(B)$

**Proof** =

$$(*) := \mathfrak{d}\text{productOfMeasure}(X, Y) \text{ProductSection}(A, B) \text{IntegralHomogeneity}(\mu_Y(B)$$

$$\text{MeasureAsIntegral}(\mu_X, A) : \mu_X \times \mu_Y(A \times B) = \int_X \mu_Y(\text{section}(A \times B), x) \mathrm{d}\mu_X(x) =$$

$$= \int_X \mu_Y(B) I_A \mathrm{d}\mu_X = \mu_Y(B) \int_X I_A \mathrm{d}\mu_X = \mu_Y(B) \mu_X(A);$$

□

**MeasureProductCommute** ::  $\forall X, Y : \text{MEAS} . \mu_X \times \mu_Y = \mu_Y \times \mu_X \circ \text{swap}$

**Proof** =

**Assume**  $A \times B : \mathcal{F}_X \times \mathcal{F}_Y$ ,

$$(1) := \text{ClassicalPMTHM}(X, Y)(A \times B) : \mu_X \times \mu_Y(A \times B) = \mu_X(A) \mu_Y(B),$$

$$(2) := \text{ClassicalPMTHM}(Y, X)(B \times A) : \mu_Y \times \mu_X(B \times A) = \mu_Y(B) \mu_X(A),$$

$$(3) := (1)(2) : \mu_X \times \mu_Y(A \times B) = \mu_Y \times \mu_X(B \times A);$$

$$\leadsto (*) := \text{SwapIntro}(\cdot) : \mu_X \times \mu_Y = \mu_Y \times \mu_X \circ \text{swap};$$

□

## 5.2 Fubini Theorem

**MeasrableOnProduct** ::  $\forall X, Y : \text{BOR} . \forall f : \text{Masurable}(X \times Y) . \forall x : X . \Lambda y : Y . f(x, y) : \text{Measurable}(Y)$

**Proof** =

**Assume**  $A : \mathcal{B} \mathbb{R}^\infty$ ,

(1) := **InversePointProduct**( $f, x, A$ ) :  $f^{-1}(x, \cdot)(A) = \text{section}(f^{-1}(A), x)$ ,

(2) :=  $\partial \text{Measurable}(X \times Y)(f)(A) : f^{-1}(A) : F_{X \times Y}$ ,

(3) := (1)**MeasurableSection**( $x, f^{-1}(A)$ ) :  $f^{-1}(x, \cdot)$ ;

$\leadsto (*)$  :=  $\partial^{-1} \text{Measurable}(X)(\cdot) : \Lambda y : Y . f(x, y) : \text{Measurable}(Y)$ ;

□

$Y : \text{BOR}$

$X : \text{MEAS}$

$S : \text{SlicingMeasure}(X, Y) \ \& \ \text{Uniformly}\sigma\text{-Finite}(X, Y)$

$\nu = \text{productMeasure}(S)$

**FubiniI** ::  $\forall f : \text{Measurable}(X \times Y) : f > 0 . \forall A : \mathcal{F}_{X \times Y} .$

$\Lambda x : X . \int_{\text{section}(A, x)} f(x, y) \, dS(x, y) : \text{Measurable}(X)$

**Proof** =

**Assume**  $B : \mathcal{F}_Y$ ,

**Assume**  $\phi : \text{Simple}(X \times Y)$ ,

$(n, b, c) := \partial \text{Simple}(X \times Y) : \mathbb{N} \times n \rightarrow \mathcal{F}_{X \times Y} \times n \rightarrow \mathbb{R}_{++} : \phi = \sum_{i=1}^n c_i I_{b_i}$ ,

(1) :=  $\partial(n, b, c) \rightarrow \int_B \phi \, dS : \int_B \phi \, dS = \sum_{i=1}^n c_i S(x, \text{section}(X \times B \cup b_i, x))$ ,

(2) := (1)**MeasrableSlicing**( $S, X \times B \cup b$ ) :  $\int \phi \, dS : \text{Measurable}(X)$ ;

$\leadsto (1) := UI(\cdot) : \forall \phi : \text{Simple}(X \times Y) . \int_B f \, dS : \text{Measurable}(X)$ ,

$\phi := \text{SimpleApprox}(f) : \mathbb{N} \rightarrow \text{Simple}(X \times Y) : \phi_n \uparrow f$ ,

(2) := **MonotoneConvergence**  $\left( \int_B \phi \, dS, \int_B f \, dS \right) : \int_B \phi \, dS : \text{Measurable}(X)$ ;

$\leadsto (1) := \partial^{-1} \text{SlicingMeasure}(\cdot) : fS : \text{SlicingMeasure}(X \times Y)$ ,

(2) := **MeasurableSlicing**( $fS$ ) :  $\Lambda x \in X . \int_{A_x} f(x, y) \, dS(x, y) : \text{Measurable}(X)$ ;

□

$$\text{FubiniII} :: \forall f : \text{Measurable}(X \times Y) : f \geq 0 . \forall A : \mathcal{F}_{X \times Y} . \int_X \int_{A_x} f(x, y) \, dS(x, y) \, d\mu(x) = \int_A f \, d\nu(S)$$

Proof =

Assume  $B : \mathcal{F}_Y$ ,

$$\begin{aligned} (1) &:= \text{Indicator}(B) \text{productMeasure} \text{Indicator}(B) = \\ &: \int_X \int_{A_x} I_B \, dS \, d\mu = \int_X \int_{A_x \cap B_x} 1 \, dS \, d\mu = \nu(A \cap B) = \int_A I_B \, d\nu; \\ \leadsto (1) &:= \text{UI}(\cdot) : \forall B : \mathcal{F}_{X \times Y} . \int_X \int_{A_x} I_B \, dS \, d\mu = \int_A I_B \, d\nu, \end{aligned}$$

Assume  $\phi : \text{Simple}(X \times Y)$ ,

$$\begin{aligned} (n, b, c) &:= \text{Simple}(X \times Y) : \mathbb{N} \times n \rightarrow \mathcal{F}_{X \times Y} \times n \rightarrow \mathbb{R}_{++} : \phi = \sum_{i=1}^n c_i I_{b_i}, \\ (2) &:= \dots : \int_X \int_{A_x} \phi \, dS \, d\mu = \int_X \int_{A_x} \sum_{i=1}^n c_i I_{b_i} \, dS \, d\mu = \sum_{i=1}^n c_i \int_X \int_{A_x} I_{b_i} \, dS \, d\mu = \\ &= \sum_{i=1}^n c_i \int_A I_{b_i} \, d\nu = \int_A \sum_{i=1}^n I_{b_i} \, d\nu = \int_A \phi \, d\nu; \\ \leadsto (2) &:= \text{UI}(\cdot : \forall \phi : \text{Simple}(X \times Y) . \int_X \int_{A_x} \phi \, dS \, d\mu = \int_A \phi \, d\nu, \end{aligned}$$

$\phi := \text{SimpleApproximation}(f) : \mathbb{N} \rightarrow \text{Simple}(X \times Y) : \phi \uparrow f$ ,

$$\begin{aligned} (3) &:= \dots : \int_X \int_{A_x} f \, dS \, d\mu = \int_X \int_{A_x} \lim_{n \rightarrow \infty} \phi_n \, dS \, d\mu = \lim_{n \rightarrow \infty} \int_X \int_{A_x} \phi_n \, dS \, d\mu = \lim_{n \rightarrow \infty} \int_A \phi_n \, d\nu = \\ &= \int_A \lim_{n \rightarrow \infty} \phi_n \, d\nu = \int_A f \, d\nu; \end{aligned}$$

□

**TonelliI** ::  $\forall f : \text{IntegralExists}(X \times Y, \nu) . \Lambda x \in X . \Lambda y \in Y . f(x, y) : \text{IntegralExists}(Y, S(x)) \text{a.e.} [\mu]$

Proof =

$$\begin{aligned} (1) &:= \text{FubiniII}(f_+, X \times Y) : \int_X \int_Y f_+ \, dS \, d\mu = \int_{X \times Y} f_+ \, d\nu, \\ (2) &:= \text{FubiniII}(f_-, X \times Y) : \int_X \int_Y f_- \, dS \, d\mu = \int_{X \times Y} f_- \, d\nu, \\ (3) &:= \text{IntegralExists}(\mu) \text{Integrate}(f, \nu)((1), (2)) \text{Integrate}(f, S(x)) : \\ &: \text{Error} \neq \int_{X \times Y} f \, d\nu = \int_{X \times Y} f_+ \, d\nu - \int_{X \times Y} f_- \, d\nu = \int_X \int_Y f_+ \, dS \, d\mu - \int_X \int_Y f_- \, dS \, d\mu = \int_X \int_Y f \, dS \, d\mu, \\ (4) &:= \text{IntegralEq}\left(\mu, \int_Y f \, dS, \text{Error}\right) : \int_Y f \, dS \neq \text{Error} \text{a.e.} [\mu], \\ (*) &:= \text{IntegralExists}(4) : (f : \text{IntegralExists}(Y, S) \text{a.e.} [\mu]); \end{aligned}$$

□

**ToneliIII** ::  $\forall f : \text{Integrable}(X \times Y, \nu) . \Lambda x \in X . \Lambda y \in Y . f(x, y) : \text{Integrable}(Y, S(x)) \text{ a . e . } [\mu]$

**Proof** =

$$\begin{aligned}
 (1) &:= \text{FubiniIII}(f_+, X \times Y) : \int_X \int_Y f_+ \, dS \, d\mu = \int_{X \times Y} f_+ \, d\nu, \\
 (2) &:= \text{FubiniIII}(f_-, X \times Y) : \int_X \int_Y f_- \, dS \, d\mu = \int_{X \times Y} f_- \, d\nu, \\
 (3) &:= \text{IntegralExists}(\mu) \text{IntegralExists}(f, \nu)((1), (2)) \text{IntegralExists}(f, S(x)) : \\
 &\quad : \infty > \int_{X \times Y} |f| \, d\nu = \int_{X \times Y} f_+ \, d\nu + \int_{X \times Y} f_- \, d\nu = \int_X \int_Y f_+ \, dS \, d\mu + \int_X \int_Y f_- \, dS \, d\mu = \int_X \int_Y f \, dS \, d\mu, \\
 (4) &:= \text{IntegralIneq} \left( \mu, \int_Y f \, dS, \infty \right) : \int_Y |f| \, dS < \infty \text{ a . e . } [\mu], \\
 (*) &:= \text{IntegralExists}^{-1}(4) : (f : \text{Integrable}(Y, S) \text{ a . e . } [\mu]); \\
 &\square
 \end{aligned}$$

**Toneli0** ::  $\forall f : \text{IntegralExists}(X \times Y, \nu) .$

$$. \exists \phi : \text{IntegralExists}(X \times Y, \nu) : \int_Y \phi \, dS : \text{Measurable}(X) : \phi =_\mu f$$

**Proof** =

$$\begin{aligned}
 (1) &:= \text{ToneliI}(f) : f : \text{Integrable}(Y, S) \text{ a . e . } [\mu], \\
 \phi &:= \Lambda(a, b) \in X \times Y . \text{if } \int_Y f(a, y) \, dS(x, y) = \text{Error then } 0 \text{ else } f(a, b) : \text{Integralexists}, \\
 (2) &:= \text{FubiniI}(\phi_+) : \int_Y f_+ \, dS : \text{Measurable}(X), \\
 (3) &:= \text{FubiniI}(\phi_-) : \int_Y f_- \, dS : \text{Measurable}(X), \\
 (4) &:= \text{AdditiveIntegral}(\phi_+, -\phi_-) : \int_Y \phi_+ \, dS - \int_Y \phi_- \, dS = \int_Y \phi \, dS, \\
 (*) &:= \text{ContinuousPreserveMeasureable}(2, 3, 4) : \int_Y \phi \, dS : \text{Measurable}(X), \\
 &\square
 \end{aligned}$$

**FubiniToneli** ::  $\forall f : \text{Measurable}(X \times Y) : \int_{X \times Y} |f| \, d\nu < \infty$

$$. \int_{X \times Y} f \, d\nu = \int_X \int_Y f \, dS \, d\mu$$

**Proof** =

**ClassicalFubini** ::  $\forall \nu : \text{Measure}(Y) . \forall f : \text{IntegralExists}(X \times Y, \mu \times \nu) .$

$$. \int_{X \times Y} f \, d\mu \times \nu = \int_X \int_Y f \, d\mu \, d\nu = \int_Y \int_X f \, d\nu \, d\mu$$

**Proof** =

### 5.3 Iterated Integrals

$$\text{MeasureSystem} :: \prod n \in \mathbb{N} . \prod X : n \rightarrow \text{BOR} . ?(\prod m : n . \prod_{i=1}^{m-1} X_i \rightarrow \text{Measure}(X_m))$$

$$P : \text{MeasureSystem} \iff \forall m \in n . \forall A \in \mathcal{F}_{X_m} . P(\cdot, A) : \text{Measurable}(X_m)$$

$$\text{iteratedMeasure} :: \prod n \in \mathbb{N} . \prod X : n \rightarrow \text{BOR} . \text{MeasureSystem}(X) \rightarrow \mathbb{R}_+^\infty$$

$$\text{iteratedMeasure}(P) = \int_X dP := \int_{X_1} \int_{X_{|\overline{2}, n}} dP_x dP_1(x)$$

$$n : \mathbb{N}$$

$$X : n \rightarrow \text{BOR}$$

$$P : \text{MeasureSystem}(X)$$

$$\text{IteratedMPTHM} :: \exists \mu : \text{Measure} \left( \prod_{i=1}^n X_i \right) : \forall A : \prod m : n . \mathcal{F}_{X_m} . \mu \left( \prod_{i=1}^n A_i \right) = \int_A dP$$

**Proof** =

Use MPTHM repeatedly

□

$$\text{iteratedProductMeasure} :: \text{MeasureSystem}(X) \rightarrow \text{MEAS}$$

$$\text{iteratedProductMeasure}(P) = \left( \prod_{i=1}^n X_i, P \right) := \left( \prod_{i=1}^n X_i, \text{IteratedMPTHM}(P) \right)$$

$$\text{Uniformly } \sigma\text{-Finite}(\cdot) \text{ System} :: ?\text{MeasureSystem}(X)$$

$$P : \text{Uniformly } \sigma\text{-Finite}(X) \text{ System} \iff \forall m : n . P_m : \text{Uniformly } \sigma\text{-Finite} \left( \prod_{i=1}^{m-1} X_i \right)$$

$$\text{SFSystem} :: P : \text{uniformly } \sigma\text{-Finite}(\cdot) \text{ System} \Rightarrow \left( \prod_{i=1}^n X_i, P \right) : \sigma\text{-Finite} \left( \prod_{i=1}^n X_i \right)$$

**Proof** =

$$\text{iteratedIntegral} :: \text{IntegralExists} \left( \prod_{i=1}^n X_i, P \right) \rightarrow \mathbb{R}^\infty$$

$$\text{iteratedIntegral}(f) = \int_X f dP := \int_{X_1} \int_{X_{|\overline{2}, n}} f_x dP_x dP_1(x)$$

$\text{ProbabilitySystem} :: ?\text{MeasureSystem}(X)$

$P : \text{ProbabilitySystem} \iff \forall m : n . \forall x \in \prod_{i=1}^{m-1} X_i . P(X, \cdot) : \text{Probability}(X_i)$

$P : \text{ProbabilitySystem}(X)$

$\text{IteratedPPTHM} :: (\prod_{i=1}^n X_i, P) : \text{Probability}\left(\prod_{i=1}^n X_i\right)$

$\text{Proof} =$

## 5.4 Infinite Products

$$\text{Cylinder} :: \prod X : \mathbb{N} \rightarrow \text{Set} . \prod n \in \mathbb{N} . ? \left( \prod_{i=1}^n X_i \right) \rightarrow ? \prod_{i=1}^{\infty} X_i$$

$$C : \text{Cylinder}(A) \iff \pi_{1,\dots,n} C = A$$

$$\text{MeasurableCylinder} :: \prod X : \mathbb{N} \rightarrow \text{BOR} . \prod n \in \mathbb{N} . \mathcal{F}_{\prod_{i=1}^n X_i} \rightarrow ? \prod_{i=1}^{\infty} X_i$$

$$C : \text{MeasurableCylinder}(A) : C : \text{Cylinder}(A) \iff$$

$$\text{infiniteBorProduct} :: (\mathbb{N} \rightarrow \text{BOR}) \rightarrow \text{BOR}$$

$$\text{InfiniteBorProduct}(X_i, \mathcal{F}_i) = \prod_{i=1}^n (X_i, \mathcal{F}_i) := \left( \prod_{i=1}^{\infty} X_i, \sigma(\text{MeasurableCylinder}(X)) \right)$$

$$\text{cylinder} :: \prod X : \mathbb{N} \rightarrow \text{Set} . \prod n \in \mathbb{N} . \prod A \subset \prod_{i=1}^n X_i \rightarrow \text{Cylinder}(X, n, A)$$

$$\text{cylinder}(A) := A \times \prod_{i=n+1}^{\infty} X_i$$

$$\text{DiscreteRandomProcess} :: \prod X : \mathbb{N} \rightarrow \text{BOR} . ? \left( \prod n : \mathbb{N} . \prod_{i=1}^n X_i \rightarrow \text{Probability} X_{n+1} \right)$$

$$P : \text{DiscreteRandomProcess} \iff$$

$$\iff \forall n \in \mathbb{N} . \forall A \in \mathcal{F}_{X_n} . \Lambda x \in \prod_{i=1}^{n-1} . P(x, A) : \text{Measureble} \prod_{i=1}^{n-1} X_i$$

$$\text{InfiniteProductTheoremI} :: \forall X : \mathbb{N} \rightarrow \text{BOR} . \forall P : \text{DiscreteRandomProcess}(X) . \forall n : \mathbb{N} .$$

$$. \Lambda B \in \mathcal{F}_{\prod_{i=1}^n X_i} . \int_X I_B dP_{|n} : \text{Probability} \left( \prod_{i=1}^n X_i \right)$$

**Proof** =

$$(1) := \mathfrak{D}^{-1}(\mathfrak{D} \text{DiscreteRandomProcess}(P)) : (P_{|n} : \text{ProbabilitySystem}(X_{|n})) ,$$

$$(*) := \text{IteretadPPTHM}(P_{|n}) : \left( \left( \prod_{i=1}^n X_i, P_{|n} \right) : \text{Probability} \left( \prod_{i=1}^n X_i \right) \right) ;$$

□

$$X : \mathbb{N} \rightarrow \text{BOR}$$

$$P : \text{DiscreteRandomProcess}(X)$$

$$\text{finiteTimeProbability} :: \prod n \in \mathbb{N} . \text{Probability} \left( \prod_{i=1}^n X \right)$$

$$\text{finiteTimeProbability}(t) = P_t := \text{InfiniteProductTheoremI}(X, P, t)$$

$$\text{InfiniteProductTheoremII} :: \exists \mathbb{P} : \text{Probability} \left( \prod_{i=1}^{\infty} X_i \right) : \forall t : \mathbb{N} . \forall B : \prod_{i=1}^n X_i . \mathbb{P}(\text{cylinder}(B)) = P_t(B)$$

Proof =

...

□

$$\text{ClassicalIPTHM} :: \forall (X, \mathcal{F}, P) : \mathbb{N} \rightarrow \text{ProbabilitySpace} . \prod_{i=1}^{\infty} P_i : \text{Probability} \left( \prod_{i=1}^{\infty} (X_i, \mathcal{F}_i) \right)$$

Proof =

...

□

$$\text{GeneralCylinder} :: \prod T : \text{Set} . \prod X : T \rightarrow \text{Set} . \prod \tau : \text{Finite}(T) . ? \prod_{t \in \tau} X_t \rightarrow ?? \prod_{t \in T} X_t$$

$$C : \text{GeneralCylinder}(A) \iff C = A \times \prod_{t \in \tau^c}$$

$T : \text{Set}$

$X : T \rightarrow \text{BOR}$

$$\text{GeneralMeasurableCylinder} :: \prod \tau : \text{Finite}(T) . \mathcal{F}_{\prod_{t \in \tau} X_t} \rightarrow ?? \prod_{t \in T} X_t$$

$$C : \text{GeneralMeasurableCylinder}(A) \iff \text{MeasurableCylinder}(A)$$

$\text{generalBorProduct} :: (T \rightarrow \text{BOR}) \rightarrow \text{BOR}$

$$\text{generalBorProduct}((X, \mathcal{F})) = \prod_{t \in T} (X_t, \mathcal{F}_t) := \left( \prod_{t \in T} X_t, \sigma(\text{GeneralMeasurableCylinder}(X, \mathcal{F})) \right)$$

$$\text{generalCylinder} :: \prod \tau : \text{Finite}(T) . \mathcal{F}_{\prod_{t \in \tau} X_t} \rightarrow \mathcal{F}_{\prod_{t \in T} X_t}$$

$$\text{generalCylinder}(B) := B \times \prod_{t \in \tau^c} X_t$$

$$\text{KolmogorovConsistent} :: ? \left( \prod \tau : \text{Finite}(T) . \text{ProbabilitySystem} \left( \prod_{t \in \tau} \right) \right)$$

$$P : \text{KolmogorovConsistent} \iff \forall \tau : \text{Finite}(T) . \forall \theta \subset \tau . \pi_{\theta}(P_{\tau}) = P_{\theta}$$



**KolmogorovExtension** ::  $\forall X : T \rightarrow \text{Polish} . \forall P : \text{KolmogorovConsistent}(X, \mathcal{B}X) .$

$. \exists \mathbb{P} : \text{Probability} \left( \prod_{t \in T} (X_t, \mathcal{B}X_t) \right) : \forall \tau : \text{Finite}(T) . \pi_\tau \mathbb{P} = P_\tau$

**Proof** =

$\mathcal{F}_0 := \text{GeneralMeasurableCylinder}(X, \mathcal{B}X) : \text{Set},$

$\mathbb{P} := \Lambda A \times \prod_{t \in \tau^0} X_t : \text{GeneralMeasurableCylinder}(X, \mathcal{B}X)(\tau) . P_\tau(A) :$

$: \text{GeneralMeasurableCylinder}(X, \mathcal{B}X) \rightarrow [0, 1],$

**Assume**  $A : \text{DisjointElems}(\mathcal{F}_0),$

$\tau := \bigcup_{i=1}^n \tau_A : \text{Finite}(T),$

$B := \partial \text{GeneralMeasurableCylinder}(A) : n \rightarrow \mathcal{F}_{\prod t \text{ in } \tau} : \forall i \in n . A_i = B_i \times \prod_{t \in \tau} X_i,$

$(1) := \partial B \partial \mathbb{P} : \mathbb{P} \left( \bigcup_{i=1}^n A_i \right) = P_\tau \left( \bigcup_{i=1}^n B_i \right),$

$(2) := (1) \partial \text{Measure}(P_\tau) : \mathbb{P} \left( \bigcup_{i=1}^n A_i \right) = \sum_{i=1}^n P_\tau(B_i),$

$(3) := (2) \partial \mathbb{P} : \mathbb{P} \left( \bigcup_{i=1}^n A_i \right) = \sum_{i=1}^n \mathbb{P}(A_i);$

$\leadsto (1) := (\cdot) : \mathbb{P} : \text{FinitelyAdditive},$

**Assume**  $A : \mathbb{N} \rightarrow \mathcal{F}_0 : A \downarrow \emptyset,$

**Assume**  $\epsilon : \mathbb{R}_{++} : \forall n : \mathbb{N} . \mathbb{P}(A_n) > \epsilon,$

$(\tau, B) := \partial \mathcal{F}_0(A) : \mathbb{N} \rightarrow \sum \tau : \text{Finite}(T) . \text{GeneralMeasurableCylinder}(X, \mathcal{B}X, ),$

$C := \text{PolishISTight}(X)(P)(B)(\Lambda k \in \mathbb{N} . \frac{\epsilon}{2^{k+1}}) : \prod n \in \mathbb{N} . \text{Compact} \left( \prod_{t \in \tau} X_i \right) : \forall n \in \mathbb{N} . P_{\tau_n},$

$\alpha := \text{generalMeasurableCylinder}(C) : \prod n \in \mathbb{N} . \text{GeneralMeasurableCylinder}(\tau_n),$

$(2) := \partial(D)(1) \partial(\alpha) \partial \alpha(\partial C) : \forall n \in \mathbb{N} . \mathbb{P}(A_n \setminus D_n) = \mathbb{P} \left( A_n \cap \bigcup_{i=1}^n \alpha_i^c \right) \leq \sum_{i=1}^n \mathbb{P}(A_i \cap \alpha_i) =$

$= \sum_{i=1}^n P_{\tau_i}(B_i \setminus C_i) < \sum_{i=1}^n \frac{\epsilon}{2^{n+i}} \leq \epsilon/2,$

$(3) := \text{IntersectionIsSubset}(\partial(D)) : \forall n \in \mathbb{N} . D_n \subset A_n,$

$(4) := \text{SubsetDifference}((3))(2) : \forall n \in \mathbb{N} . \mathbb{P}(D_n) > \mathbb{P}(P_n) - \frac{\epsilon}{2},$

$(5) := \partial \text{Probability}(4, \partial(\epsilon)) : \forall n : \mathbb{N} . D_n \neq \emptyset,$

$x := \partial \text{NonEmpty}(D, 5) : \prod n : \mathbb{N} . D_n,$

**Assume**  $n : \mathbb{N},$

$(6) := \partial D_n(\partial x) : \forall m : \mathbb{N} : m \geq n . \pi_{\tau_n} x_n \in C_n,$

$(7) := \text{PolishIsSeqCompact} \left( \prod_{t \in \tau_n} X_t, C_n \right) : (C_n : \text{SeqCompact}),$

$$(m, y) := \text{\textcolor{teal}{SeqCompact}}(C_n, \pi_{\tau_n} x) : \text{\textcolor{teal}{Subsequer}} \times C_n : \lim_{n \rightarrow \infty} x_{m_n} = y,$$

$$y_n := y : C_n;$$

$$\leadsto y := [\cdot] : \prod_{n \in \mathbb{N}} C_n,$$

$$(6) := \text{\textcolor{teal}{Seq}} y : \forall n : \mathbb{N} . \forall m : \mathbb{N} : m > n . \pi_{\tau_m}(y_n) = y_m,$$

$$Y := \text{\textcolor{teal}{restore}}(y, 6) : \bigcap_{n=1}^{\infty} D_n,$$

$$(7) := \text{\textcolor{teal}{NonEmpty}} \left( \bigcap_{n=1}^{\infty} D_n, Y \right) : \bigcap_{n=1}^{\infty} D_n \neq \emptyset,$$

$$(8) := \text{\textcolor{teal}{SubsetIntersection}}(D, A) : \bigcap_{n=1}^{\infty} D_n \subset \bigcap_{n=1}^{\infty} A_n,$$

$$(9) := \text{\textcolor{teal}{EmptySubset}}(8, \text{\textcolor{teal}{Seq}} A) : \bigcap_{n=1}^{\infty} D_n = \emptyset,$$

$$(10) := (7)(9) : \perp;$$

$$\leadsto (2) := \text{\textcolor{teal}{Convergent}}(\mathbb{R}_+)(\mathbb{P}(A_n) : \lim_{n \rightarrow \infty} \mathbb{P}(A_n) = 0;$$

$$\leadsto (2) := \text{\textcolor{teal}{CountablyAdditive}}(\mathbb{P}) : \left( \mathbb{P} : \text{\textcolor{teal}{CountablyAdditive}} \left( \prod_{t \in T} X_t, \mathcal{F}_0 \right) \right),$$

$$Q := \text{\textcolor{teal}{CaratheodoryExtension}}(\mathbb{P}) : \text{\textcolor{teal}{Probability}} \left( \prod_{t \in T} (X_t, \mathcal{B}X_t) \right) : \forall \tau : \text{\textcolor{teal}{Finite}}(T) . \pi_{\tau} Q = P_{\tau};$$

□

$$X : T \rightarrow \text{\textcolor{teal}{Polish}}$$

$$\text{\textcolor{teal}{RandomFieldLaw}} :: \text{\textcolor{teal}{KolmogorovConsistent}}(X, \mathcal{B}X) \rightarrow \text{\textcolor{teal}{Probability}} \prod_{t \in T} X_t$$

$$\text{\textcolor{teal}{RandomFieldLaw}}(P) = [P] := \text{\textcolor{teal}{KolmogorovExtension}}(P)$$



## 6 Convergence of Measures

### 6.1 Duals of Measures

$\text{UnboundedSpace} :: ?\text{MetricSpace}$

$X : \text{UnboundedSpace} \iff X \text{ ! Compact}$

$X : \text{UnboundedSpace}$

$Y : \text{MetricSpace}$

$\text{LimitAtInfinity} :: (X \rightarrow Y) \rightarrow ?Y$

$a : \text{LimitAtInfinity}(f) \iff \lim_{x \rightarrow \infty} f(x) = a \iff$   
 $\iff \forall U \in \mathcal{U}(a) . \exists K : \text{Compact}(X) . f^{-1}(U^c) : \text{Compact}(X)$

$\text{PuncturedSpace} :: ?\text{MetricSpace}$

$X : \text{PuncturedSpace} \iff \exists Y : \text{MetricSpace} \ \& \ \text{Compact} : \exists y \in Y : \mathcal{T}_X = \mathcal{T}_{Y \setminus \{y\}}$

$\text{IsPuncturedSpace} :: \forall X : \text{MetricSpace} \ \& \ \text{Separable} \ \& \ \text{LocallyComapact} . X : \text{PuncturedSpace}$

$\text{Proof} =$

$q := \text{Sepable}(X) : \mathbb{N} \rightarrow X : \text{Dense}(X),$

$K := \text{LocallyComapact}(X) : \mathbb{N} \rightarrow \text{Compact}(X) : K \uparrow X,$

$\delta := \Lambda n : \mathbb{N} . \Lambda x : X . \min(1, d(x, q_n)) : \mathbb{N} \rightarrow \mathcal{M}_{\text{TOP}}(X, \mathbb{R}_+),$

$g := \Lambda x : X . \sum_{n=1}^{\infty} \min(d(x, K_n^c), 2^{-n}) : X \rightarrow \mathbb{R}_+,$

$f := \Lambda n : \mathbb{N} . \Lambda x : X . \min(g(x), d(x, q_n), 2^{-n}) : \mathbb{N} \rightarrow X \rightarrow \mathbb{R}_+,$

$D := \Lambda x, y : X . \sum_{n=1}^{\infty} |f_n(x) - f_n(y)| : \text{Distance}(X),$

$(1) := \text{DistanceConstruction}(\delta(D)) : (X, d) \cong (X, D),$

$\text{Extend } D := \Lambda x \in X . D(\infty, x) = D(x, \infty) = g(x) : \text{Distance}(X \cup \{\infty\}),$

$(2) := \text{PuncturedSpace}(X, d)(X \cup \{\infty\}, D) : ((X, d) : \text{PuncturedSpace}),$

□

$\text{measures} :: \text{MetricSpace} \rightarrow \text{VectorSpace}(\mathbb{R})$

$\text{measures}(X) = \mathfrak{M}(X) := \{\mu : \text{Charge}(X, \mathcal{B}X) : |\mu|(X) < \infty\}$

$\text{localSupNorm} :: \prod X : \text{MetricSpace} . \prod A : \mathbb{N} \rightarrow \text{Compact}(X) : \bigcup_{n=1}^{\infty} A_n = X . \text{Norm}(\mathcal{M}_{\text{TOP}}(X, \mathbb{R}))$

$\text{localSupNorm}(A) = \|f\|_{\text{loc}(A)} := \sum_{n=1}^{\infty} \min(\sup_{x \in A_n} |f|, 2^{-n})$

$\text{localSupSpace} :: \prod X : \text{MetricSpace} . \prod A : \mathbb{N} \rightarrow \text{Compact}(X) : \bigcup_{n=1}^{\infty} A_n = X . \text{NormedSpace}(\mathbb{R})$

$\text{localSupSpace}(A) = C_{\text{loc}(A)} := (\{f \in \mathcal{M}_{\text{TOP}}(X, \mathbb{R})\}, \|\cdot\|_{\text{loc}(A)})$

$\text{SeparabilityOfSup} :: \forall X : \text{MetricSpace} \ \& \ \text{Compact} \ \& \ \text{Separable} . C_{\text{sup}}(X) : \text{Separable}$

**Proof** =

$q := \text{SepSeparable}(X) : \text{Dense}(X),$

$F := \mathbb{Q}[d(q_n, x) | n : \mathbb{N}] : \mathbb{R}\text{-Algebra}(\mathcal{M}_{\text{TOP}}(X, \mathbb{R})),$

$(1) := \text{polyfunctionalFunc}(F) : \text{Constants}(X, \mathbb{R}) \subset F,$

$(2) := \text{SepC}_{\text{sup}}(\text{CIsBoundedOnCompact}(X, F)) : F \subset C_{\text{sup}},$

**Assume**  $a, b : X : a \neq b,$

$(3) := \text{SepDistance}(X)(a, b)(\text{Sep}(a, b)) : d(a, b) \neq 0,$

$(\alpha) := \text{SepSeparable}(X)(a, b)(3)(a) : \text{Im } q : d(a, \alpha) < \frac{d(a, b)}{2},$

$f := \Lambda x : X . d(x, \alpha) : F,$

$(4) := \text{SepStictIneq}(\text{Sep } f)(\text{Sep } \alpha) : f(a) \neq f(b);$

$\leadsto (3) := \text{Sep}^{-1}\text{Separates}(F, \cdot) : (F : \text{Separates}(X)),$

$(4) := \text{StoneWeierstrass}(1, 2, 3) : (F : \text{Dense}(C_{\text{sup}}(X))),$

$(5) := \text{Sep}^{-1}\text{Separable}(C_{\text{sup}}(X))(F, 4) : (C_{\text{sup}}(X) : \text{Separable});$

□

$\text{compactSupp} :: \text{MetricSpace} \rightarrow \text{NormedSpace}$

$\text{compactSupp}(X) = C_{00}(X) := (\{f : \mathcal{M}_{\text{TOP}}(X, \mathbb{R}) : (\text{supp } f : \text{Compact}(X))\}, \|\cdot\|_{\text{sup}})$

$\text{zeroAtInf} :: \text{UnboundedSpace} \rightarrow \text{NormedSpace}$

$\text{zeroAtInf}(X) = C_0(X) := (\{f : \mathcal{M}_{\text{TOP}}(X, \mathbb{R}) : (\lim_{x \rightarrow \infty} f(x) = 0)\}, \|\cdot\|_{\text{sup}})$

$\text{ZeroAtInfSeparable} :: \forall X : \text{UnboundedSpace} \ \& \ \text{LocallyComapact} \ \& \ \text{Separable} . C_0(X) : \text{Separable}$

**Proof** =

$(X', \infty) := \text{IsPuncturedSpace}(X) : \sum X' : \text{MetricSpace} \ \& \ \text{Compact} . X' : X' \setminus \{\infty\} = X,$

**Assume**  $f : C_0(X),$

$f' := \text{Extend } f \text{ On } X' \text{ By } f'(\infty) = 0 : C_{\text{sup}}(X');$

$\leadsto F := \{f' | f \in C_0(X)\} : \text{Subset}(C_{\text{sup}}(X')),$

$(1) := \text{SeparabilityOfSup}(X') : (C_{\text{sup}}(X') : \text{Separable}),$

$(2) := \text{Sep}^{-1}\text{Separable}(C_0(X))((\text{Sep } F)^{-1}\text{SepSeparable}(C_{\text{sup}}(X')))) : C_0(X) : \text{Separable};$

□

**ZeroAtInfIsClosed** ::  $\forall X : \text{TopologicalSpace} . C_0(X) : \text{Closed}(X)$

**Proof** =

$(X', \infty) := \text{Alexandroff}(X) : \left( \sum X' : \text{TopologicalSpace} . X' \right) : X \cong_{\text{TOP}} X' \setminus \{\infty\},$

**Assume**  $f : C_0(X),$

$f^* := \text{Extend } f \text{ On } X \text{ By } f^*(\infty) = 0 : C_b(X');$

$\leadsto (\cdot)^* := \text{FI} : C_0(X) \hookrightarrow C_b(X),$

**Assume**  $f : \text{Convergent}(C_0(X)),$

$\phi := \lim_{n \rightarrow \infty} f_n : C_b(X),$

(1) :=  $\text{Convergent}(C_b(X'))(\text{FI}^*(\cdot)^* \text{Convergent}(C_0(X))) : f^* : \text{Convergent}(C_b(X')),$

$\phi' := \lim_{n \rightarrow \infty} f_n^* : C_b(X'),$

(2) :=  $\text{FI}^*(\phi')(0) : f^*(\infty) = 0,$

(3) :=  $\text{Limit}(f^*, \phi') : \phi'(\infty) = 0,$

(4) :=  $\text{M}_{\text{TOP}}(X', \mathbb{R}) : \lim_{x \rightarrow \infty} \phi'(x) = 0,$

(5) :=  $\text{FI}^{-1}(C_0(X))(\phi'_X)(\text{Limit}(4)) : \phi'_X \in C_0(X),$

(6) :=  $\text{UniformIsPointwise}(f, f', \phi, \phi', \text{FI}^*(\cdot)^*) : \phi'_X \in C_0(X),$

(7) :=  $= \text{E}(5, 6) : \phi \in C_0(X);$

$\leadsto (1) := \text{SeqClosed} : C_0(X) : \text{Closed}(C_b(X));$

□

**CompactSuppIsDense** ::  $\forall X : T_4 . C_{00}(X) : \text{Dense}(C_0(X))$

**Proof** =

**Assume**  $\varphi : C_0(X),$

**Assume**  $n : \mathbb{N},$

$K_n := \varphi^{-1}(-\infty, -2^{-n}] \cup [2^{-n}, \infty) : \text{Compact}(X),$

$U_n := \varphi^{-1}(-\infty, -2^{-n-1}) \cup (2^{-n}, \infty) : \text{Open}(X),$

$f_n := \text{Uryshon}(\phi, K_n, U_n) : \mathcal{M}_{\text{TOP}}(X, \mathbb{R}) : f_n|_{K_n} = \varphi : f_n|_{U_n^c} = 0,$

(1) :=  $\text{d}_X(\text{FI} f_n, \phi) : d(\varphi, f_n) \leq 2^{-n};$

$\leadsto (1) := \text{FI}^{-1} \text{Limit}(f, \phi) : \exists f \in \mathbb{N} \rightarrow C_{00}(X) . \lim_{n \rightarrow \infty} f_n = \varphi;$

$\leadsto (*) := \text{FI}^{-1} \text{Dense}(C_0(X)) : C_{00}(X) : \text{Dense}(C_0(X));$

□

**VagueLimit** ::  $\prod X : \text{MetricSpace} . ?((\mathbb{N} \rightarrow \mathfrak{M}(X)) \times \mathfrak{M}(X))$

$(\mu, \nu) : \text{VagueLimit} \iff \mu \rightarrow_v \nu \iff \forall f \in C_{00}(X) . \lim_{n \rightarrow \infty} \int_X f \, d\mu_n = \int_X f \, d\nu$

**VagueConvergent** ::  $\prod X : \text{MetricSpace} . ?(\mathbb{N} \rightarrow \mathfrak{M}(X))$

$\mu : \text{VagueConvergent} \iff \exists \nu \in \mathfrak{M}(X) : \mu \rightarrow_v \nu$

## 6.2 Extending Linear Functionals

**ExtensionFromOpenSubset** ::  $\forall X : \text{TopologicalSpace} . \forall U : \text{Open}(X) : \text{cl}U : \text{Compact}(X) .$   
 $. \forall f \in C_0(U) . \text{exists } g \in C_{00}(X) : g|_U = f$

**Proof** =

$g := \text{Extend } f \text{ On } x \in X \text{ By } g(x) = 0 : C_{00}(X) : g|_U = f;$

□

**OpenSubsetFunctionalContraction** ::  $\forall X : \text{TopologicalSpace} . \forall U : \text{Open}(U) .$   
 $. \forall L : \text{PositiveLinearFunctional}(C_{00}(X)) . \exists L' :$

$\text{PositiveLinearFunctional}(C_0(U)) : L \cdot \text{ExtensionFromOpenSubset}(X)(U) = L'$

**Proof** =

...

□

**ExtendFromCompactSuppToZeroAtInf** ::  $\forall X : T_4 . \forall L : \text{PositiveLinearFunctional}(C_{00})(X) .$   
 $. L : \text{ExtendsTo}(C_0(X))$

**Proof** =

**Assume**  $\varphi : C_0(X),$

$f := \text{CompactSuppIsDense}(X)(\varphi) : \mathbb{N} \rightarrow C_{00}(X) : \lim_{n \rightarrow \infty} f_n = \varphi,$

$b := \text{PLFIIsBounded}(L) : \text{BoundConstant}(L),$

**Assume**  $\epsilon : \mathbb{R}_+,$

$N := \text{Convergent}(f) : \mathbb{N} : \forall n, m : \mathbb{N} : n, m \geq N \implies |f_n - f_m| \leq \epsilon/b,$

$(1) := \text{Mvs}(\mathbb{R})(C_{00}(X))(L) \text{BoundConstant}(b, L) N :$

$: \forall n, m : \mathbb{N} : n, m \geq N . |L(f_n) - L(f_m)| = |L(f_n - f_m)| \leq \epsilon;$

$\leadsto (1) := \text{Complete}(C_0(X)) : L(f) : \text{Convergent}(C_0(X)),$

$M(\varphi) := \lim_{n \rightarrow \infty} L(f_n) : C_0(X);$

$\leadsto (*) := \text{ExtendsTo}(C_0(X)) : (L : \text{Extends}(C_0(X)));$

□





## 6.3 Vague Convergence over Compacts

$X$  : **MetricSpace** & **Compact**

**AsFunctional** ::  $\mathfrak{M}(X) \rightarrow \mathcal{M}_{\text{VS}(\mathbb{R})}(C_b(X), \mathbb{R})$

**AsFunctional** ( $\mu$ ) =  $[\mu] := \Lambda f \in C_b(X) . \int_X f \, d\mu$

**CompactMeasureBoundedFunctional** ::  $\forall \mu \in \mathfrak{M}(X) . [\mu] : \text{BoundedFunctional}$

**Proof** =

**Assume**  $f : C_b(X)$ ,

(1) :=  $\mathfrak{D}\text{AsFunctional}(\mu) \text{IntegralIneq}(f, \|f\|) \text{IntegralLinear}(\mu, \|f\|)$

$\text{MeasureAsIntegra}(\mu) : [\mu](f) = \int_X f \, d\mu \leq \int_X \|f\| \, d\mu = \|f\| \mu(X);$

$\leadsto (*) := \mathfrak{D}^{-1} \text{BoundedFunctional}([\mu], \mu(X)) : [\mu] : \text{BoundedFunctional};$

□

**EveryPLFIsMeasure** ::  $\forall L \in \text{PositiveLinearFunctional}(C_b(X)) . \exists \mu : \text{Radon}(X) : L = [\mu]$

**Proof** =

**Assume**  $K : \text{Closed}(X)$ ,

$F := \{f \in C_b(X) : f|_K = 1\} : \text{Set}(C_b(X))$ ,

$\mu(K) := \inf_{f \in F} L(f) : \mathbb{R}_+$ ;

$\leadsto \mu := (\cdot) : \text{Closed}(X) \rightarrow \mathbb{R}_+$ ,

**Assume**  $K, G : \text{Closed}(X) : K \cap G = \emptyset$ ,

$F_K := \{f \in C_b(X) : f|_K = 1 : f|_{K^c} < 1\} : \text{Set}(C_b(X))$ ,

$F_G := \{f \in C_b(X) : f|_G = 1 : f|_{G^c} < 1\} : \text{Set}(C_b(X))$ ,

$F := \{f \in C_b(X) : f|_{K \cup G} = 1\} : \text{Set}(C_b(X))$ ,

(1) :=  $\dots : \mu(K \cap G) = \inf_{f \in F} L(f) =$

$\inf_{f \in F_K + F_G} L(f) = \inf_{f \in F_K} \inf_{g \in F_G} L(f + g) = \inf_{f \in F_K} L(f) + \inf_{f \in F_G} L(f) = \mu(K) + \mu(G);$

$\leadsto (1) := \mathfrak{D}^{-1} \text{FinitelyAdditive}(\mu) : \mu : \text{FinetlyAdditive},$

$\lambda := \text{RadonExtension}(\mu) : \text{Radon}(X)$ ,

**Assume**  $K : \text{Closed}(X)$ ,

$f := \Lambda n \in \mathbb{N} . \Lambda x \in X . \exp(-nd(x, K)) : \mathbb{N} \rightarrow C_b(X)$ ,

(2) :=  $\mathfrak{D}^{-1} \text{Limit}(\mathfrak{D}(f)) : f \downarrow I_K$ ,

(3) :=  $\text{MonotoneConvergence}(\lambda, I_K) : \lim_{n \rightarrow \infty} \int_X f \, d\lambda = \lambda(K)$ ,

(2) :=  $\dots : \dots;$

(3) :=  $\mathfrak{D}\lambda(2) : L(1) = \lambda(X)$ ,

(4) :=  $\mathfrak{D}\mathcal{M}_{\text{VS}(\mathbb{R})}(C_b(X), \mathbb{R}) : \forall c : \text{Constant}(X) . L(c) = c\lambda(X)$ ,

**Assume**  $f : C_b(X) : f > 0$ ,

$a := \mathfrak{D}(f > 0)(a) \text{CompactOpt}(f) : (0, \min_{x \in X} f(x))$ ,

**Assume**  $\epsilon : \mathbb{R}_+$ ,

$k := \text{CompactOpy}(f) \mathfrak{D}\text{Archimedean}(\mathbb{R}) : \mathbb{N} : a + (k + 1)\epsilon > \max_{x \in X} f(x)$ ,

Assume  $j : k$ ,

$$A_j := f^{-1}[a + j\epsilon, a + (j + 1)\epsilon] : \text{Closed}X,$$

$$I_j := \dots (2) : C_b(X) : \lambda(A_j) \leq \int_{A_j} I_j \, d\lambda \leq \lambda(A_j) + \epsilon : \int_{A_j^c} I_j \, d\lambda \leq \epsilon;$$

$$\leadsto g := \sum_{j=1}^k (a + (j + 1)\epsilon) I_j : C_b(X),$$

$$(5) := \dots : L(f) \leq L(g) = \sum_j^k (a + (j + 1)\epsilon) L(I_j),$$

...

□