





STATISTICAL PROCEDURE DECISION TEMPLATE



Measures of Central Tendency	Measures of Variability	Measures of Association	Measures of Linear Relationship
Mean: indicator of central tendency sensitive to the exact position of each score in the distribution	Range: difference between the lowest and highest scores	Spearman rank-order correlation coefficient: variant of Pearson statistic for two rank-ordered variables	Point-Biserial correlation coefficient: variant of Pearson statistic where one variable is dichotomous
Mode: most commonly occurring value	Standard Deviation: indicator of average deviation from mean	Pearson product-moment correlation coefficient: strength and direction of the relationship between two variables	Phi coefficient: a variant of the Pearson statistic for two dichotomous variables
Median: middle score in a distribution	Variance: average squared deviation from mean		Simple regression: indicated the best estimate of one variable from the values of another

What statistical techniques will we use?
 Have our hypotheses been proved or disproved?
 Is our information valid and reliable?
 What are the implications of our study?

Who will read the report?
 What is their level of familiarity with the subject?
 How should we construct the report?



*For one group tests
 Nominal: Chi-square goodness of fit (binomial test if two categories and category size = 5 or more)
 Ordinal: Kolmogorov-Smirnov one sample test
 Interval or ratio: t-test for one sample