

EXERCISES GALOIS THEORY AND SOME SOLUTIONS

CARSTEN DIETZEL, SILVIA PROPERZI

1. WEEK 1

Exercise 1.1. Let $\sigma : K \rightarrow L$ be a field homomorphism, prove that σ is injective.

Solution. [using ring theory] We know that $\ker \sigma$ is an ideal of K . But K is a field so the only possibilities are that $\ker \sigma = \{0\}$ or $\ker \sigma = K$. As $\sigma(1) = 1 \neq 0$, we know that $1 \notin \ker \sigma$, hence $\ker \sigma = \{0\}$ which means that σ is injective. ■

Solution. [by hands] Let $a, b \in K$ such that $\sigma(a) = \sigma(b)$. Then

$$\sigma(a - b) = \sigma(a) - \sigma(b) = 0.$$

If we assume that $a \neq b$, then $a - b$ is a non-zero element in a field, hence it is invertible. Then

$$1 = \sigma(1) = \sigma((a - b)(a - b)^{-1}) = \sigma(a - b)\sigma((a - b)^{-1}) = 0,$$

which is a contradiction. Therefore $a = b$ and so σ is injective. ■

Exercise 1.2. Let K be a field, K_0 be its prime field and $\sigma : K \rightarrow K$ be a field homomorphism. Prove that $\sigma \in \text{Hom}(K/K_0, K/K_0)$.

Exercise 1.3. Let $\mathbb{Q}[i] = \{a + ib \mid a, b \in \mathbb{Q}\}$, $\mathbb{Q}[\sqrt{2}] = \{a + \sqrt{2}b \mid a, b \in \mathbb{Q}\}$ and $\mathbb{Q}(i) = \{\frac{a+ib}{c+di} \mid a, b, c, d \in \mathbb{Q}\}$, $\mathbb{Q}(\sqrt{2}) = \{\frac{a+\sqrt{2}b}{c+\sqrt{2}d} \mid a, b, c, d \in \mathbb{Q}\}$.

- (i) Prove that $\mathbb{Q}(\sqrt{2}) = \mathbb{Q}[\sqrt{2}]$ and $\mathbb{Q}(i) = \mathbb{Q}[i]$.
- (ii) Prove that $\mathbb{Q}(i)$ and $\mathbb{Q}(\sqrt{2})$ are not isomorphic.

Exercise 1.4.

- (i) Let $a = \sqrt{2}$ and $b = \sqrt[3]{3}$. Prove that ab is algebraic over \mathbb{Q} .
- (ii) Show that $\sqrt{2} + i$ is algebraic over \mathbb{Q} by finding a nonzero polynomial $f \in \mathbb{Q}[X]$ with $\deg(f) = 4$ such that $f(\sqrt{2} + i) = 0$. What are the other roots of f ?

Solution.

- (i) Observe that $ab^6 = \sqrt{2}^6 \sqrt[3]{3}^6 = 8 \cdot 9 = 72$. Therefore ab is a root of the polynomial $X^6 - 72 \in \mathbb{Q}[X]$.
- (ii) let $\alpha = \sqrt{2} + i$. Then $\alpha^2 = 2 + 2\sqrt{2}i - 1 = 1 + 2\sqrt{2}i$ and $\alpha^2 - 1 = 2\sqrt{2}i$. Squaring both sides of the last equality we have that

$$\alpha^4 - 2\alpha^2 + 1 = (\alpha^2 - 1)^2 = (2\sqrt{2}i)^2 = -8.$$

Therefore the polynomial

$$f(X) = X^4 - 2X^2 + 1 + 8 = X^4 - 2X^2 + 9 \in \mathbb{Q}[X]$$

is such that $f(\alpha) = 0$.

Looking for other roots of f means to find all $x \in C$ such that $f(x) = 0$, i.e. $x^4 - 2x^2 + 9 = 0$. Going back to how we construct f , this equation can also be written as $(x^2 - 1)^2 = (2\sqrt{2}i)^2$. Therefore $x^2 - 1 = 2\sqrt{2}i$ or $x^2 - 1 = -2\sqrt{2}i$. So

$$x^2 = 2\sqrt{2}i + 1 = \alpha^2 \quad \text{or} \quad x^2 = -2\sqrt{2}i + 1 = \overline{\alpha^2} = \overline{\alpha}^2,$$

where $\bar{\alpha}$ indicates the complex conjugate of α . Thus $x = \pm\alpha$ or $x = \pm\bar{\alpha}$ and the 4 roots of f are

$$\begin{aligned}x_1 &= \alpha = \sqrt{2} + i, x_2 = -\alpha = -\sqrt{2} - i, \\x_3 &= \bar{\alpha} = \sqrt{2} - i, x_4 = -\bar{\alpha} = -\sqrt{2} + i.\end{aligned}$$

■

Exercise 1.5. Let p be a prime number. Denote by $\binom{n}{k}$ the binomial coefficient “ n over k ”.

- (i) Prove that p divides $\binom{p}{k}$ for $1 \leq k \leq p-1$.
- (ii) Let K be a field of characteristic p . Show that the map $\Phi : K \rightarrow K$; $x \mapsto x^p$ is indeed a field endomorphism. This map is called the *Frobenius endomorphism* of K .

Exercise 1.6. Let L/K be a field extension and let M be a subring of L that contains K . Suppose that $\dim_K M < \infty$.

- (i) Prove that for any $\alpha \in M$, there is a nonzero $f \in K[X]$ with $f(\alpha) = 0$.
Hint: The elements α^n ($n \geq 0$) are linearly dependent.
- (ii) Prove that M is a field.
Hint: For any $0 \neq \alpha \in K$, let $f \in \mathbb{Q}[X]$ be a nonzero polynomial with $f(\alpha) = 0$ whose degree is as small as possible. If $f = \sum_{i=0}^{\infty} a_i X^i$, prove that $a_0 \neq 0$ and use this to construct an inverse of α that lies in M .

Exercise 1.7. Let K be field.

- (i) Prove that $K[X]$ is a PID.
- (ii) Let $I \neq \{0\}$ be an ideal of $K[X]$, then there exists a unique monic polynomial that generates I as an ideal.

2. WEEK 2

Exercise 2.1. Let L/K be a field extension and let $\alpha, \beta \in L$ such that

$$[K(\alpha) : K] = [K(\beta) : K] = 2.$$

Assume that the characteristic of K is not 2.

- (i) Prove that there is an $\alpha' \in L$ such that $K(\alpha') = K(\alpha)$ and $\alpha'^2 \in K$.
- (ii) Assume that $\alpha, \beta \in L$ satisfy $\alpha^2, \beta^2 \in K$. Prove that $K(\alpha) = K(\beta)$ if and only if $\frac{\alpha^2}{\beta^2}$ is a square in K .
- (iii) Prove that there is a bijective map

$$K^\times / (K^\times)^2 \longrightarrow \{L \mid L/K \text{ is a field extension with } [L : K] \leq 2\}.$$

Solution.

- (i) Since $[K(\alpha) : K] = 2$, $\alpha \notin K$ and the 3 elements $1, \alpha, \alpha^2$ are linearly dependent over K . Thus there exist $a_0, a_1, a_2 \in K$ not all zero such that

$$a_0 + a_1\alpha + a_2\alpha^2 = 0.$$

If $a_2 = 0$, then $a_0 + a_1\alpha = 0$, hence either $a_1 = 0$ or $\alpha = -a_0/a_1$. If $a_1 = 0$, then also $a_0 = 0$. But this is not possible because $(a_0, a_1, a_2) \neq (0, 0, 0)$. On the other hand, if $\alpha = -a_0/a_1$, then $\alpha \in K$, a contradiction.

So we have that $a_2 \neq 0$ and we can divide by a_2 , obtaining

$$b + a\alpha + \alpha^2 = 0, \text{ i.e. } \alpha^2 + a\alpha = -b,$$

where $a = a_2^{-1}a_1 \in K$ and $b = a_2^{-1}a_0 \in K$. Since we assumed that K has not characteristic two, we can also complete the square:

$$(\alpha + a/2)^2 = \alpha^2 + a\alpha + a^2/4 = -b + a^2/4 \in K.$$

Therefore $\alpha' = \alpha + a/2$ is such that $\alpha'^2 = -b + a^2/4 \in K$. Moreover $\alpha' = \alpha + a/2 \in K(\alpha)$, so $K(\alpha') \subseteq K(\alpha)$ and $\alpha = \alpha' - a/2 \in K(\alpha')$ so $K(\alpha) \subseteq K(\alpha')$. Therefore $K(\alpha) = K(\alpha')$ and $\alpha'^2 \in K$.

- (ii) Assume that $\frac{\alpha^2}{\beta^2}$ is a square in K , i.e. there is a $k \in K$ such that $\frac{\alpha^2}{\beta^2} = k^2$. (Note that $k \neq 0$ otherwise $\alpha = 0 \in K$ and $[K(\alpha) : K] = [K : K] = 1$.)

Then $\alpha = k^2\beta^2$, hence $\alpha = \pm k\beta \in K(\beta)$. So $K(\alpha) \subseteq K(\beta)$. On the other hand, $\beta^2 = \frac{\alpha^2}{k^2}$, hence $\beta = \pm \frac{\alpha}{k} \in K(\alpha)$. So $K(\beta) \subseteq K(\alpha)$. Having proved both inclusions we deduce that $K(\alpha) = K(\beta)$.

Vice versa, assume that $K(\alpha) = K(\beta)$. Knowing that $[K(\alpha) : K] = 2$, we have that $\{1, \alpha\}$ is a generating set of the K -vector space $K(\alpha) = K(\beta)$. Therefore there exist $a, b \in K$ such that $\beta = a + b\alpha$. Squaring both sides of the equality, we get $\beta^2 = a^2 + 2ab\alpha + \alpha^2$. So $2ab\alpha = \beta^2 - a^2 - \alpha^2$ is a sum of elements in K . Hence $2ab\alpha \in K$, but $\alpha \notin K$. Therefore the only possibility is that $2ab = 0$, i.e. (since we are not in characteristic 2) $ab = 0$. But $b \neq 0$, otherwise $\beta = a \in K$, which is not possible (otherwise and $[K(\beta) : K] = [K : K] = 1$). Thus $a = 0$, i.e. $\beta = b\alpha$ and so $\frac{\beta^2}{\alpha^2} = a^2$ is a square in K .

- (iii) Let L/K be a field extension with $[L : K] = 2$. Take $\alpha \in L \setminus K$, then

$$1 < [K(\alpha) : K] \leq [L : K] = 2.$$

Hence $L = K(\alpha)$. Moreover, the first part of this exercise allows us to choose α such that $\alpha^2 \in K$. Now we can define the following map

$$\psi : \{L \mid L/K \text{ is a field extension with } [L : K] \leq 2\} \rightarrow K^\times / (K^\times)^2$$

as $\psi(K) = [1]$ and for $[L : K] = 2$ as $\psi(L) = [\alpha^2] \in K^\times / (K^\times)^2$, for $\alpha \in L \setminus K$ (so $L = K(\alpha)$) such that $\alpha^2 \in K$.

This map is well-defined: take L/K of degree 2 and $\alpha, \beta \in L \setminus K$ such that $\alpha^2 \in K$. Then, by the remark made at the beginning, $L = K(\alpha) = K(\beta)$. As shown in the second part of this exercise, this is equivalent to $\frac{\alpha^2}{\beta^2} \in (K^\times)^2$, so $[\alpha^2] = [\beta^2] \in K^\times / (K^\times)^2$. (Note also that $[\alpha^2] = [1]$ if and only if $\alpha^2 = k^2 \in K^2$, so $\alpha = \pm k \in K$ and $K(\alpha) = L$.)

Let now prove that ψ is injective. Assume that we have two extension L/K and L'/K of degree ≤ 2 , such that $[\alpha^2] = \psi(L) = \psi(L') = [\beta^2]$.

If $[\beta^2] = [\alpha^2] = [1]$, then $L = K(\alpha) = K = K(\beta) = L'$.

Otherwise, $\beta^2/\alpha^2 \in (K^\times)^2$, so, by the previous part of this exercise, $L = K(\alpha) = K(\beta) = L'$.

Finally, for the surjectivity, let $x \in K^\times$. If $x \in (K^\times)^2$, then $x = \alpha^2$ for some $\alpha \in K^\times$ and so $L = K(\alpha)$ is an extension of K of degree ≤ 2 and $\psi(L) = [x]$.

If $x \notin (K^\times)^2$, then we can find an extension $L = K(\alpha)$ such that $\alpha^2 = x \in K^\times$ and so $\psi(L) = [x] \in K^\times / (K^\times)^2$. To construct this extension consider the polynomial $f(X) = X^2 - x \in K[X]$. It has to be irreducible, otherwise $X^2 - x = (aX + b)(cX + d) = acX^2 + (ad + bc)X + bd$, for some $a, c \in K^\times$ and $b, d \in K$. So $1 = ac$, $0 = ad + bc$ and $-x = bd$, i.e. $c = a^{-1}$, $d = -a^{-1}bc = bc^2$ and $-x = bd = b^2c^2 \in (K^\times)^2$, a contradiction to the assumption $x \notin (K^\times)^2$. Then we can consider the field $L = K[X]/(X^2 - x)$ that contains K (it is a field because the ideal $(X^2 - x)$ is maximal, since it is generated by an irreducible polynomial). Defining α as the class of X in L we get that $L = K(\alpha)$ and $\alpha^2 = x \in K$.

■

Exercise 2.2. Let E/K be a field extension and a and b be algebraic over K .

- (1) Assume that $[K(a) : K] = m$, $[K(b) : K] = n$. Prove that $K[a, b] \subseteq E$ is generated, as a vector space over K , by the elements $a^i b^j$ ($1 \leq i \leq m$, $1 \leq j \leq n$).
- (2) Prove that $a + b$ and ab are algebraic over K . Can you estimate the quantities $[K(a + b) : K]$ and $[K(ab) : K]$?
- (3) Find a polynomial $f \in \mathbb{Q}[X]$ such that $\deg(f) \leq 6$ and $f(\sqrt[3]{3} + \sqrt{5})$.

The following lemma can be used, without proof, in the following exercise.

Lemma (Gauss' Lemma). *Let A be a unique factorization domain and K be its fraction field. A non-constant polynomial $f \in A[X]$ is irreducible if and only if it is primitive and is irreducible in $K[X]$.*

Exercise 2.3 (Eisenstein's irreducibility criterion). Let A be a unique factorization domain and K be its fraction field. Let $f = \sum_{i=0}^n a_i X^i \in K[X]$ be a polynomial of degree $n > 0$. Assume that there exists a prime element $p \in A$ such that $p \mid a_i$ for all $i \in \{0, 1, \dots, n-1\}$, $p \nmid a_n$ and $p^2 \nmid a_0$. Then f is irreducible in $K[X]$.

Exercise 2.4. Let $\zeta \in \mathbb{C}$ be a primitive cubic root of one. Set $E = \mathbb{Q}[\sqrt[3]{2}]$, $F = \mathbb{Q}(\zeta)$ and $L = \mathbb{Q}[i]$.

- (i) Prove that $[E : \mathbb{Q}] = 3$ and $[F : \mathbb{Q}] = 2$ and compute the minimal polynomial of ζ over \mathbb{Q} and over L .
- (ii) Prove that $EF = \mathbb{Q}(\sqrt[3]{2}, \zeta)$.
- (iii) Compute $[EF : \mathbb{Q}]$ and $[E \cap F : \mathbb{Q}]$.

Exercise 2.5. Let E/K be a field extension and let L/K and M/K be subextensions.

- (i) Prove that $[LM : K] \cdot [L \cap M : K] \leq [L : K] \cdot [M : K]$.
- (ii) Can you find examples where $[LM : K] \cdot [L \cap M : K] < [L : K] \cdot [M : K]$?
Hint: Use two different roots of the polynomial $X^3 - 2$.

Exercise 2.6. Let E/K be a field extension and let $f \in K[X]$ be a polynomial such that f factorizes in $E[X]$ as $f = \prod_{i=1}^n (x - \alpha_i)$. Prove by induction that $[K(\alpha_1, \alpha_2, \dots, \alpha_n) : K] \leq n!$.

3. WEEK 3

Exercise 3.1. For every polynomial $p(X) = \sum_{i=0}^n a_i X^i \in K[X]$ of degree n , define its *reciprocal polynomial* as

$$\widehat{p}(X) = \sum_{i=0}^n a_{n-i} X^i.$$

Let $p(X), q(X) \in K[X]$ be polynomials of degree n and m respectively such that $p(0) \neq 0$ and $q(0) \neq 0$. Prove that

- (i) $\widehat{\widehat{p}}(X) = X^n p(1/X)$ in $K(X)$,
- (ii) $\widehat{\widehat{p}}(X) = p(X)$,
- (iii) $\widehat{pq}(X) = \widehat{p}(X)\widehat{q}(X)$,
- (iv) $\widehat{p}(X)$ is irreducible if and only if $p(X)$ is irreducible.

Solution. Suppose that $p(X) = \sum_{i=0}^n a_i X^i$ has degree n and $q(X) = \sum_{i=0}^m b_i X^i$ has degree m .

- (i) $X^n p(1/X) = X^n \sum_{i=0}^n a_i (1/X)^i = \sum_{i=0}^n a_i X^{n-i} = \sum_{j=0}^n a_{n-j} X^j = \widehat{p}(X)$
- (ii) Since $p(0) \neq 0$, $a_0 \neq 0$, therefore $\deg(\widehat{p}) = \deg(p) = n$. Hence, using the previous part of this exercise,

$$\widehat{\widehat{p}}(X) = X^n \widehat{p}(1/X) = X^n (1/X)^n p(1/(1/X)) = p(X).$$

- (iii) Since $p(0) \neq 0$ and $q(0) \neq 0$ we deduce that also $pq(0) \neq 0$. Hence we can apply the previous results for p, q and pq . So, using also that $pq(1/X) = p(1/X)q(1/X)$ and that $\deg pq = n + m$, we have that

$$\widehat{pq}(X) = X^{n+m}pq(1/X) = X^n p(1/X) X^m q(1/X) = \widehat{p}(X)\widehat{q}(X).$$

- (iv) Suppose $\widehat{p}(X)$ is irreducible and let $p = q_1 q_2$ for $q_1(X), q_2(X) \in K[X]$. Note that $0 \neq p(0) = q_1(0)q_2(0)$ implies that both $q_1(0) \neq 0$ and $q_2(0) \neq 0$. Considering now the reciprocal polynomials and using the previous properties, $\widehat{p}(X) = \widehat{q_1}(X)\widehat{q_2}(X)$. Since $\widehat{p}(X)$ is irreducible, there is $i \in \{1, 2\}$ such that $\widehat{q_i}$ is a constant, i.e. $\deg q_i = \deg \widehat{q_i} = 0$. Thus, q_i is a constant too, and therefore $p(X)$ is irreducible.

Vice versa, assume that $p(X)$ is irreducible. Since we know that $p(X) = \widehat{\widehat{p}}$, by the property just proved, we obtain that \widehat{p} is also irreducible. ■

Exercise 3.2. Let E/K be a field extension and $x \in E$ be an algebraic element and let $f = f(x, K)$ is the minimal polynomial of x over K of degree $\deg(f) = n$.

- (i) Prove that $[K(x) : K] = n$.
(ii) Prove that $\frac{1}{f(0)}\widehat{f}$ is the minimal polynomial of $1/x$ over K .
(iii) Write $f(X) = \sum_{i=0}^n a_i X^i = p(X^2) + Xd(X^2)$, where

$$p(X) = \sum_{j=0}^{\lfloor n/2 \rfloor} a_{2j} X^j \text{ and } d(X) = \sum_{j=0}^{\lfloor n/2 \rfloor} a_{2j+1} X^j.$$

Let $g(X) = p(X)^2 - Xd(X)^2$ and prove that

- if $d(x^2) = 0$, then the minimal polynomial of x^2 over K is $p(X)$,
- if $d(x^2) \neq 0$, then the minimal polynomial of x^2 over K is $(-1)^n g(X)$.

Solution.

- (i) We will prove that $B = \{1, x, \dots, x^{n-1}\}$ is a basis of $K(x)$ as a K vector space which implies that $[K(x) : K] = \dim_K(K(x)) = |B| = n$.

- B is a generating set for $K(x)$ as a K vector space.

To prove this recall that $K(x) = K[x]$, since x is algebraic over K .

(Similar proof to 1) \Rightarrow 2) of Theorem 2.7 of the lecture notes.)

Let $z \in K(x) = K[x]$, say $z = h(x)$ for some $h \in K[X]$. Divide h by f to obtain polynomials $q, r \in K[X]$ such that $h = fq + r$, where $r = 0$ or $\deg r < \deg f = n$. This implies that

$$z = h(x) = f(x)q(x) + r(x) = r(x).$$

Moreover we can write $r = \sum_{i=0}^{n-1} c_i X^i$ for some $c_0, \dots, c_{n-1} \in K$. Thus $z = \sum_{i=0}^{n-1} c_i x^i \in \langle 1, x, \dots, x^{n-1} \rangle$ and hence $K[x]$ is generated by $\{1, x, \dots, x^{n-1}\}$ as a K -vector space.

- B is linearly independent over K .

If B is linearly dependent over K then there exists a linear combination

$0 = \sum_{i=0}^{n-1} c_i x^i$ over K , with not all c_i equal to 0. Then the polynomial $h(X) = \sum_{i=0}^{n-1} c_i X^i$ is in $K[X] \setminus \{0\}$ and has x as a root. So

$$n - 1 = \deg(h) \leq \deg(f) = n,$$

a contradiction.

- (ii) First of all we note that $f(0) \neq 0$. Otherwise we can write $f(X) = Xg(X)$ for some $g(X) \in K[X]$, but f is monic and irreducible in $K[X]$, hence $g(X) = 1$ and $f(X) = X$. Evaluating f in x we obtain $0 = f(x) = x$, a contradiction with the hypothesis $x \neq 0$.

Since $f(0) \neq 0$, we can use the previous exercise and obtain that \widehat{f} is also irreducible and $\widehat{f} = X^n f(1/X)$. Hence

$$\frac{1}{f(0)} \widehat{f}(1/x) = \frac{1}{f(0)} (1/x)^n f\left(\frac{1}{1/x}\right) = \frac{1}{f(0)x^n} f(x) = 0.$$

So we have that $\frac{1}{f(0)} \widehat{f}$ is an irreducible polynomial in $K[x]$ with $1/x$ as a root. To prove that $\frac{1}{f(0)} \widehat{f}$ is the minimal polynomial of $1/x$ over K it remains to prove that it is monic. Looking at the definition of \widehat{f} we see that its leading coefficient is the constant term of f , i.e. $f(0)$. Therefore the leading coefficient of $\frac{1}{f(0)} \widehat{f}$ is $\frac{1}{f(0)} f(0) = 1$, hence $\frac{1}{f(0)} \widehat{f}$ is monic.

(iii) First of all note that

$$0 = f(x) = p(x^2) + xd(x^2), \quad (1)$$

therefore

$$g(x^2) = p(x^2)^2 - x^2 d(x^2)^2 = (p(x^2) + xd(x^2))(p(x^2) - xd(x^2)) = 0.$$

So x^2 is a root of g and the degree of g is

$$\begin{cases} 2 \deg(p) & \text{if } n \text{ is even} \\ 1 + 2 \deg(d) & \text{if } n \text{ is odd} \end{cases} = \begin{cases} 2 \lfloor \frac{n}{2} \rfloor & \text{if } n \text{ is even} \\ 1 + 2 \lfloor \frac{n}{2} \rfloor & \text{if } n \text{ is odd} \end{cases} = n.$$

Moreover, by the first point of this exercise and the fact that the degree is multiplicative

$$n = \deg(f) = [K(x) : K] = [K(x) : K(x^2)][K(x^2) : K].$$

Hence, to compute $[K(x^2) : K]$ (which is also the degree of $f(x^2, K)$, the minimal polynomial of x^2 over K), we need to know $[K(x) : K(x^2)]$. Observe that $X^2 - x^2$ is a polynomial in $K(x^2)[X]$ which has x as a root. Thus

$$[K(x^2) : K] = \deg(x, K(x^2)) \leq \deg(X^2 - x^2) = 2$$

Therefore

$$[K(x^2) : K] = \frac{[K(x) : K]}{[K(x) : K(x^2)]} \in \left\{n, \frac{n}{2}\right\}. \quad (2)$$

- If $d(x^2) = 0$, by Equation (1), also $p(x^2) = 0$ and $\deg(p) = \lfloor \frac{n}{2} \rfloor < n$. So

$$[K(x^2) : K] = \deg(f(x^2, K)) \leq \deg(p) < n,$$

thus, by (2), $[K(x^2) : K] = \frac{n}{2}$, which implies that n has to be even, $p(X)$ monic and

$$[K(x^2) : K] = \frac{n}{2} = \left\lfloor \frac{n}{2} \right\rfloor = \deg(p).$$

Therefore $p(X)$ is a monic polynomial in $K[X]$ which has x^2 as a root and of degree $[K(x^2) : K] = \deg(f(x^2, K))$, hence it is $\deg(f(x^2, K))$, the minimal polynomial of x^2 over K .

- If $d(x^2) \neq 0$, then, by Equation (1),

$$x = -\frac{p(x^2)}{d(x^2)} \in K(x^2).$$

Therefore $K(x) \subseteq K(x^2) \subseteq K(x)$ and so $K(x^2) = K(x)$, which means $[K(x) : K(x^2)] = 1$ and, by (2),

$$[K(x^2) : K] = \frac{[K(x) : K]}{[K(x) : K(x^2)]} = [K(x) : K] = n = \deg(g).$$

Moreover the leading coefficient of $g(X)$ is

$$\begin{cases} a_n & \text{if } n \text{ is even} \\ -a_n & \text{if } n \text{ is odd} \end{cases} = (-1)^n a_n = (-1)^n.$$

Therefore we have the monic polynomial $(-1)^n g(X) \in K[X]$ that vanishes in x^2 , of degree $n = [K(x^2) : K] = \deg(f(x^2, K))$. Hence $(-1)^n g(X) = f(x^2, K)$. ■

The following lemma can be used, without proof, in the following exercise.

Lemma (Gauss' Lemma). *Let A be a unique factorization domain and K be its fraction field. A non-constant polynomial $f \in A[X]$ is irreducible if and only if it is primitive and irreducible in $K[X]$.*

Exercise 3.3 (Eisenstein's irreducibility criterion). Let $f = \sum_{i=0}^n a_i X^i \in \mathbb{Z}[X]$ be a polynomial of degree $n > 0$. Assume that there exists a prime p such that $p \mid a_i$ for all $i \in \{0, 1, \dots, n-1\}$, $p \nmid a_n$ and $p^2 \nmid a_0$. Then f is irreducible in $\mathbb{Q}[X]$.

More general version

Let A be a unique factorization domain and K be its fraction field. Let $f = \sum_{i=0}^n a_i X^i \in A[X]$ be a polynomial of degree $n > 0$. Assume that there exists a prime element $p \in A$ such that $p \mid a_i$ for all $i \in \{0, 1, \dots, n-1\}$, $p \nmid a_n$ and $p^2 \nmid a_0$. Then f is irreducible in $K[X]$.

Exercise 3.4. Let $\zeta \in \mathbb{C}$ be a primitive cubic root of one. Set $E = \mathbb{Q}[\sqrt[3]{2}]$, $F = \mathbb{Q}(\zeta)$ and $L = \mathbb{Q}[i]$.

- (i) Prove that $[E : \mathbb{Q}] = 3$ and $[F : \mathbb{Q}] = 2$ and compute the minimal polynomial of ζ over \mathbb{Q} and over L .
- (ii) Prove that $EF = \mathbb{Q}(\sqrt[3]{2}, \zeta)$.
- (iii) Compute $[EF : \mathbb{Q}]$ and $[E \cap F : \mathbb{Q}]$.

Exercise 3.5. Let E/K be a field extension and let L/K and M/K be subextensions.

- (i) Prove that $[LM : K] \cdot [L \cap M : K] \leq [L : K] \cdot [M : K]$.
- (ii) Can you find examples where $[LM : K] \cdot [L \cap M : K] < [L : K] \cdot [M : K]$?
Hint: Use two different roots of the polynomial $X^3 - 2$.

Exercise 3.6. Let E/K be a field extension and let $f \in K[X]$ be a polynomial such that f factorizes in $E[X]$ as $f = \prod_{i=1}^n (x - \alpha_i)$. Prove by induction that $[K(\alpha_1, \alpha_2, \dots, \alpha_n) : K] \leq n!$.

(Carsten Dietzel) DEPARTMENT OF MATHEMATICS AND DATA SCIENCE, VRIJE UNIVERSITEIT BRUSSEL, PLEINLAAN 2, 1050 BRUSSEL, BELGIUM
Email address: Carsten.Dietzel@vub.be

(Silvia Properzi) DEPARTMENT OF MATHEMATICS AND DATA SCIENCE, VRIJE UNIVERSITEIT BRUSSEL, PLEINLAAN 2, 1050 BRUSSEL, BELGIUM
Email address: Silvia.Properzi@vub.be