Each condition: variety of its cases and distance from the others

Condition	Split proposed	Distinct work-ups of 30	Required items per case	Silhouette required studies	1-NN required studies	1-NN narrative style	1-NN narrative meaning	Copy F1 same condition	Copy F1 other condition	No study shared cases of 30	Closest work-up condition, cases of 30
Cerebrovascular											
Ischaemic stroke	train	20	6.3	0.19	1.00	0.67	0.63	0.79	**0.02**	0	Subarachnoid haemorrhage 21
Subarachnoid haemorrhage	train	13	5.6	0.47	1.00	0.90	0.97	0.88	**0.07**	0	Ischaemic stroke 30
Infection and inflammation											
Bacterial meningitis	**test**	20	4.4	0.29	1.00	0.23	0.80	0.68	**0.01**	2	Early Alzheimer's disease 14
Anti-NMDAR encephalitis	train	11	9.2	0.50	1.00	0.53	0.60	0.87	**0.14**	0	Temporal lobe epilepsy 28
Multiple sclerosis	**validation**	13	4.6	0.43	1.00	0.63	0.53	0.77	**0.07**	0	Bacterial meningitis 30
Neurodegeneration											
Early Alzheimer's disease	train	9	4.4	0.53	1.00	0.67	0.97	0.92	**0.04**	0	Frontotemporal dementia 30
Frontotemporal dementia	train	7	4.2	0.71	1.00	0.60	0.70	0.93	**0.53**	0	Early Alzheimer's disease 30
Vascular dementia	train	18	4.6	0.27	1.00	0.27	0.73	0.80	**0.27**	0	Parkinson's disease 13
Normal-pressure hydrocephalus	train	3	4.1	0.91	1.00	0.43	0.90	0.97	**0.22**	0	Parkinson's disease 30
Parkinson's disease	train	7	1.5	0.57	0.97	0.80	0.93	0.84	**0.03**	0	Migraine with aura 24
ALS	train	11	5.2	0.53	1.00	0.73	0.67	0.85	**0.10**	0	Parkinson's disease 24
Seizures											
Temporal lobe epilepsy	train	13	2.7	0.33	0.93	0.63	0.60	0.73	**0.04**	0	Anti-NMDAR encephalitis 17
Status epilepticus	train	29	6.4	0.15	1.00	0.60	0.43	0.62	**0.14**	0	Temporal lobe epilepsy 12
Neuromuscular											
Guillain–Barré syndrome	train	9	5.4	0.61	1.00	0.47	0.47	0.89	**0.10**	0	Myasthenia gravis 12
Myasthenia gravis	**test**	19	5.9	0.30	1.00	0.43	0.67	0.76	**0.03**	19	Guillain–Barré syndrome 7
Oncology											
High-grade glioma	**test**	21	5.7	0.26	1.00	0.57	0.73	0.71	**0.01**	27	Multiple sclerosis 1
Headache											
Migraine with aura	train	20	5.2	0.17	0.97	0.67	0.63	0.53	**0.04**	18	Vascular dementia 5
Functional											
Functional neurological disorder	train	2	1.2	0.73	1.00	0.67	0.70	0.90	**0.00**	23	Parkinson's disease 7
Systemic mimics											
Cardiac syncope	**validation**	17	6.3	0.40	1.00	0.73	0.87	0.74	**0.02**	0	Ischaemic stroke 29
Hepatic encephalopathy	**test**	27	6.0	0.08	0.83	0.60	0.63	0.43	**0.09**	4	Functional neurological disorder 7

Groups as in Fig. 2b. Proposed split: test conditions in blue, validation in cyan. Distinct work-ups: different sets of required studies (tool + arguments) among the 30 cases. Silhouette and 1-NN per condition, computed over all 600 cases (Extended Data Table on views). Copy F1: token F1 of the required studies when a case copies the required work-up of its most similar narrative (TF-IDF), within its own condition or from any other condition. Closest work-up: for each case, the other condition holding the case with the most similar required studies (Jaccard); the most frequent such condition and how many of the 30 cases point