

※ 注意：請於答案卷上依序作答，並應註明作答之題號。

Problem 1 (4%) ____ The number of independent row vectors in a matrix is the same as the number of independent column vectors.

- T. true,
F. false.

Problem 2 (4%) ____ If H is a row-echelon form of a matrix A , then the nonzero column vectors in H form a basis for the column space of A .

- T. true,
F. false.

Problem 3 (4%) ____ $\text{rank}(AC) \leq \text{rank}(A)$, where A and C are matrices such that the product AC is defined.

- T. true,
F. false.

Problem 4 (4%) ____ Let V and W be vector spaces of dimensions n and m , respectively. A linear transformation $T: V \rightarrow W$ is invertible if and only if $m = n$.

- T. true,
F. false.

Problem 5 (4%) ____ Let v and w be independent vectors in V , and let $T: V \rightarrow W$ be a one-to-one linear transformation of V into W , then $T(v)$ and $T(w)$ are independent vectors in W .

- T. true,
F. false.

Problem 6 (10%) Let $V \subset \mathbb{R}^n$ be a subspace. Let $T: \mathbb{R}^n \rightarrow \mathbb{R}^n$ and $S: \mathbb{R}^n \rightarrow \mathbb{R}^n$ be the linear transformations defined respectively by projecting onto and reflecting across V . What are the eigenvalues and eigenvectors of T and S , respectively?

接背面

Problem 7 (20%) Suppose A is an $n \times n$ matrix with the property that $A^2 = A$. Let $a_1, \dots, a_n \in \mathbb{R}^n$ be the column vectors of A and $A_1, \dots, A_n \in \mathbb{R}^n$ be the row vectors of A . Let $C(A) = \text{span}(a_1, \dots, a_n)$ and $R(A) = \text{span}(A_1, \dots, A_n)$ be the column space and the row space of A , respectively. Define

$$E(A) = \{x \in \mathbb{R}^n | x = Ax\},$$

$$F(A) = \{x \in \mathbb{R}^n | x = u - Au \text{ for some } u \in \mathbb{R}^n\}.$$

Find the following four sets: $C(A) \cap E(A)$, $N(A) \cap F(A)$, $C(A) \cap N(A)$, $C(A) + N(A)$.

Problem 8 (5%) $7! =$ _____.

Problem 9 (5%) Consider n distinct objects. Two circular arrangements are equivalent if one can be obtained from the other by rotation. The number of circular arrangements is _____.

Problem 10 (5%) For $m, n \geq 0$, $\sum_{k=m}^n \binom{k}{m} =$ _____.

Problem 11 (5%) $\binom{n}{0} - \binom{n}{1} + \dots + (-1)^n \binom{n}{n} =$ _____.

Problem 12 (5%) If $|A| = m$ and $|B| = n$, then there are _____ relations from A to B .

Problem 13 (10%) There are _____ ways to distribute m distinct objects into $m - 1$ identical containers with no container left empty.

Problem 14 (5%) A relation is called an equivalence relation if it is reflexive, symmetric, and _____.

Problem 15 (5%) ____ For $2^n - 1$ to be a prime, the necessary condition is that n be a prime.

T. true,

F. false.

Problem 16 (5%) The inverse of 5 modulo 11 is _____.