Homework - 10 COEN 240-Machine Learning

Pujitha Kallu ID: W1653660 pkallu@scu.edu

Consider a dataset with 2 points in 1D: $(x_1 = 0, y_1 = -1)$ and $(x_2 = \sqrt{2}, y_2 = 1)$. Map each point to 3D using the feature vector $\phi(x) = \left[1, \sqrt{2}x, x^2\right]^T$. This is equivalent to using a second order polynomial kernel. The SVM classifier has the form

$$\min \|\boldsymbol{w}\|^2 \, s. \, t.$$

$$y_1(\mathbf{w}^T\phi(x_1) + w_0) \ge 1$$

$$y_2(\boldsymbol{w}^T\phi(x_2)+w_0)\geq 1$$

(a). Find the corresponding points in 3D. That is, $\phi(x_1)$ and $\phi(x_2)$.

Sol:

Solution:
a) Given, 21=0, 41=-1
22=12, 42=1
$\phi(x) = [1,\sqrt{2}x,x^2]^{\top}$
Ø (x1) = [1, V2 (0), 02]
$\emptyset(x_1) = [1,0,0]^T$
The second secon
$\phi(x_2) = \begin{bmatrix} 1, \sqrt{2}\sqrt{2}, \sqrt{2} \end{bmatrix}$ $\phi(x_2) = \begin{bmatrix} 1, 2, 2 \end{bmatrix}^T$
Ø (22) = [1,2,2]T

(b). What is the value of the margin? Notice since there are only 2 points in the dataset, those points are the support vectors. Hence, the margin is the distance between each of them in 3D to the decision boundary, which lies in the middle.

Sol:

b) The decision boundary is in the mid way between 2 vectors. $\phi(x_1)$ & $\phi(x_2)$
Midpoint = $(1,0,0) + (1,2,2) = (2,2,2)$
$= (1,1).$ distance = $\sqrt{(1-1)^2 + (0-2)^2 + (0-2)^2} = \sqrt{0+4+4}$
distance = 1 (1-1)2+(0-2)2+(0-2)2 = 10+4+4
distance = 2 V2
The margin is the distance of each vector to the decision boundary which lies in middle
decision boundary which lies in middle
$Margin = 2\sqrt{2} = \sqrt{2}$
2

(c). The margin obtained from part (b) is in fact equal to $1/\|\boldsymbol{w}\|$. Determine the vector \boldsymbol{w} . Recall this vector is the line through $\phi(x_1)$ and $\phi(x_2)$, which is perpendicular to the decision boundary.

from the given assumption,	
110011 = 1	
V2_	
V 0+402+402 =	1
, , , , , , , , , , , , , , , , , , , ,	1/2_
2√2 92 =	
V V	5_
α - J	
$\alpha_2 = 1$	
by substituting w=	6
	1/2
	41.5

(d). Solve for w_0 using your value for \boldsymbol{w} and the above equations. Since the points are on the decision boundary, the inequalities will become equalities.

Sol:

| Variable of the SVM conditions $y(w^T \phi(x_1) + w_0) \ge 1$ | $y_2(w^T \phi(x_2) + w_0) \ge 1$ as equalities:

| $-for y_1(w^T \phi(x_1) + w_0) = 1 \Rightarrow$ | $-1(0, \frac{1}{2}, \frac{1}{2})(0) + w_0 = 1$ | $w_0 = -1$ | Same for $y_2(w^T \phi(x_2) + w_0) = 1$ | $y_0 = -1$ | $y_0 = -1$ | $y_0 = -1$

(e). Write the function $f(x) = w_0 + \mathbf{w}^T \phi(x)$ as an explicit function of x.

	:		AND THE RESERVE OF THE PARTY OF	
e) +(2)=w ₀ +	m, QC	x)		
we have ØC:	x)=	1		
	(0,0)	γ <u>2</u> 2		
	TANK TANK			
wi	,T _	6	1/2_ 1/2	
		The same		J
4	90 =-	J		
By substituting we	get:	The same		
가게 하면 하는 것이 되었다. 이 중에 가장 하는 것이 되었다면 하는 것이 없는 것이 없는 것이 없는 것이 없는데 없다면 없다.				
+(x) =		+ 0	1/2 42	1
			184	1/2
2	4 +	0.+	1 (122)	+ 1 2
No. of the control of	- Flat (100 cm cape) - (%)	The state of the s		
		+ 2	+ 22	
H(n) =	The second second	advantation of paragraphic		
H(a) =		1/2	2	
-H(n) =		1/2	2-1/2 mg/mg/mg/mg/mg/mg/mg/mg/mg/mg/mg/mg/mg/m	

References:

- Class notes:Support vector machines (SVMs)
 https://scikit-learn.org/stable/modules/svm.html
 https://www.geeksforgeeks.org/support-vector-machine-algorithm/