0.1 Q1 – matrix representation for $|\phi_k\rangle \langle \phi_j|$, in the orthonormal $|\nu_i\rangle$ basis

Perform change of basis.

0.2 Q2 – positive operator is Hermitian

We first show that a positive operator is normal, and this automatically implies that it is Hermitian.

To show that a positive operator is normal, we consider $A^{\dagger}A$

Now that we know that it is normal, by spectral decomposition, it possesses an eigenbasis. We now show that all of its eigenvalues are real. This is now a matrix with real entries on the diagonal, which is hermitian.

To show that the eigenvalues are real, let $|\lambda\rangle$ be an eigenvector with magnitude 1 and eigenvalue λ .

$$\langle \lambda | A | \lambda \rangle \geqslant 0$$
 $\lambda \langle \lambda | \lambda \rangle = \lambda \geqslant 0$

Hence, the eigenvalues are real and positive, and therefore it is Hermitian.

0.3 Q3 – $A^{\dagger}A$ is positive

$$\forall v \in V, \ \langle v | A^{\dagger} A | v \rangle = \langle A v | | A v \rangle = ||A v||^2 \geqslant 0$$

Hence, $A^{\dagger}A$ is positive.

0.4 Q4. Eigenvalues of a projector P are either 0 or 1

Let $|\lambda\rangle$ be an eigenvector of P with associated eigenvalue λ .

$$P^2(|\lambda\rangle) = \lambda(P\,|\lambda\rangle) = \lambda^2\,|\lambda\rangle \qquad P(|\lambda\rangle) = \lambda\,|\lambda\rangle$$

However, since P is a projector, $P^2 = P$, and therefore, $\lambda^2 = \lambda$. The roots of this equation are 0,1. Hence, $\lambda \in \{0,1\}$.

0.5 Q5. Tensor product of two unitary operators is unitary

Let U, V be unitary operators.

$$\begin{split} \langle Uu\otimes V\nu|\,|Uu\otimes V\nu\rangle &= \\ \langle u\otimes \nu|\,(U^\dagger\otimes V^\dagger)(U\otimes V)\,|u\otimes \nu\rangle &= \\ \langle u\otimes \nu|\,(U^\dagger U\otimes V^\dagger V)\,|u\otimes \nu\rangle &= \\ \langle u\otimes \nu|\,I\otimes I\,|u\otimes \nu\rangle &= \\ \langle u\otimes \nu|\,|u\otimes \nu\rangle &= \\ \langle u\otimes \nu|\,|u\otimes \nu\rangle &= \end{split}$$

Hence, $U \otimes V$ is unitary since it preserves inner products.

- 0.6 Q6. Tensor product of projectors is a projector
- 0.7 Q7. Find log and square root of matrix
- o.8 Q8. Trace properties

o.8.1
$$Tr(AB) = Tr(BA)$$

$$Tr(AB) = \sum_{z} (AB)_{zz} = \sum_{z} \sum_{k} A_{zk} B_{kz} = \sum_{z} \sum_{k} B_{kz} A_{kz} = \sum_{z} (BA)_{zz} = Tr(BA)$$

$$\textbf{0.8.2} \quad \mathsf{Tr}(A+B) = \mathsf{Tr}(A) + \mathsf{Tr}(B)$$

$$Tr(A + B) = \sum_{z} (A + B)_{zz} = \sum_{z} A_{zz} + B_{zz} = Tr(A) + Tr(B)$$

0.8.3
$$Tr(2A) = 2Tr(A)$$

$$Tr(2A) = \sum_{z} (2A)_{zz} = \sum_{z} 2A_{zz} 2\sum_{z} A_{zz} = 2Tr(A)$$

3

0.9 Commutator properties

0.9.1
$$[A, B] = -[B, A]$$

$$[A, B] = AB - BA = -(BA - AB) = -[B, A]$$

0.9.2
$$\frac{[A,B]+\{A,B\}}{2} = AB$$

$$\frac{[A,B] + \{A,B\}}{2} = \frac{(AB - BA) + (AB + BA)}{2} = AB$$

- 0.10 Express polar decomposition as outer product
- 0.11 Find left and right polar decomposition