

Below is how I would restructure the roadmap into a **research-grade master documentation** that combines the original implementation plan with the additional requirements you requested.

Differentiable Stellar Evolution for Bayesian Astronomical Inference

Executive Summary

This project aims to build a fully differentiable stellar evolution framework using JAX that can reproduce the physical behavior of traditional stellar evolution codes while enabling modern gradient-based optimization and Bayesian inference.

The system will:

- Model stellar structure and evolution
- Incorporate nuclear burning physics
- Integrate observational astronomy datasets
- Support uncertainty quantification
- Enable automatic differentiation
- Expose capabilities through an MCP server
- Allow AI agents to perform scientific reasoning over astronomical data

Development will be performed primarily in **Google Colab** with reproducible notebooks.

1. Problem Statement

Why This Project Exists

Modern astronomy increasingly relies on inverse problems.

Astronomers observe:

- Brightness
- Spectra
- Oscillation frequencies
- Surface temperatures
- Metal abundances

but wish to infer:

- Stellar mass

- Age
- Radius
- Interior composition
- Mixing efficiency
- Evolutionary state

Traditional stellar evolution tools such as MESA are extremely accurate but were not designed for:

- Automatic differentiation
- AI integration
- Bayesian inference at scale
- End-to-end gradient optimization

As a result, astronomers often rely on:

- Grid searches
- MCMC sampling
- Surrogate models

which are computationally expensive.

The goal is to replace those workflows with a differentiable framework capable of directly optimizing stellar parameters against observations.

2. Astronomy Fundamentals

What Is A Star?

A star is a self-gravitating sphere of plasma.

Gravity attempts to collapse the star inward.

Pressure generated by nuclear fusion pushes outward.

The balance between these forces determines stellar stability.

Stellar Lifecycle

Molecular Cloud

Gas collapses under gravity.

Protostar

Temperature rises.

Fusion has not started.

Main Sequence

Hydrogen fusion begins.

Star spends ~90% of its life here.

Red Giant

Hydrogen depleted.

Core contracts.

Outer layers expand.

White Dwarf / Neutron Star / Black Hole

Final state depends on initial mass.

Stellar Composition

Astronomers describe composition using:

X = Hydrogen fraction

Y = Helium fraction

Z = Metals fraction

Example:

$X = 0.70$

$Y = 0.28$

$Z = 0.02$

These directly influence stellar evolution.

Effective Temperature

Surface temperature inferred from emitted radiation.

Solar value:

$T_{\text{eff}} \approx 5777 \text{ K}$

Surface Gravity

Defined as:

$$g = GM/R^2$$

Usually reported as:

$\log(g)$

Used extensively in spectroscopic analysis.

Metallicity

Amount of elements heavier than helium.

Typically reported as:

$[\text{Fe}/\text{H}]$

Important because metals alter opacity and energy transport.

Asteroseismology

Stars oscillate similarly to musical instruments.

Observed oscillation frequencies reveal:

- Core structure
- Radius
- Density
- Age

These oscillations form one of the strongest constraints available for stellar inference.

3. Physics Foundations

Hydrostatic Equilibrium

A stable star satisfies:

$$dP/dr = -GM(r)\rho(r)/r^2$$

Where:

P = pressure

ρ = density

M(r) = enclosed mass

Interpretation:

Pressure decreases outward.

Gravity attempts to compress the star.

Pressure resists collapse.

Mass Conservation

$$dM/dr = 4\pi r^2 \rho$$

Meaning:

Mass enclosed increases with radius.

Energy Generation

Nuclear fusion generates energy.

Primary mechanisms:

Proton-Proton Chain

Dominates Sun-like stars.

CNO Cycle

Dominates massive stars.

Energy generation:

$\epsilon_{\text{nuclear}}$

Units:

$\text{erg g}^{-1} \text{s}^{-1}$

Energy Transport

Radiative Transport

Energy moves via photons.

Convective Transport

Energy moves via bulk fluid motion.

Both mechanisms must be modeled.

Boundary Conditions

The stellar interior must connect smoothly to the atmosphere.

Incorrect boundary conditions can destabilize the entire solution.

4. Numerical Methods

Why Numerical Methods Are Needed

The stellar structure equations cannot generally be solved analytically.

We discretize the star into layers.

Example:

100 layers

500 layers

1000 layers

Each layer contains:

- Temperature
 - Density
 - Pressure
 - Composition
-

Thomas Algorithm

Used for tridiagonal systems.

Advantages:

- $O(N)$ complexity
- Stable
- Efficient

Critical for hydrostatic equilibrium solutions.

ODE Integration

Implemented using:

- Diffrax
- JAX

Benefits:

- GPU acceleration
 - Automatic differentiation
 - Checkpointing
-

5. Bayesian Inference Foundations

Goal

Estimate stellar parameters from observations.

Parameters:

$\theta = [\text{Mass}, \text{Radius}, \text{Age}, Z, \alpha_{\text{mix}}, Y]$

Prior

Existing knowledge.

Example:

$\text{Mass} \sim \text{Normal}(1.0, 0.2)$

Likelihood

Probability of observations given parameters.

$$P(\text{Data} \mid \theta)$$

Posterior

$$P(\theta \mid \text{Data})$$

Represents updated scientific belief.

Variational Inference

Approximates posterior distributions.

Benefits:

- Faster than MCMC
 - GPU friendly
 - Differentiable
-

ELBO

Evidence Lower Bound

Optimization target:

Maximize ELBO

Equivalent to minimizing divergence from true posterior.

Parameter Degeneracy

Different parameter combinations can produce similar observations.

Example:

Older metal-rich star

and

Younger metal-poor star

may appear similar.

The framework must quantify these uncertainties.

6. Data Sources and Research References

MIST

Purpose:

Validation stellar tracks.

Source:

MIST project

Used for:

Forward model validation.

MESA

Purpose:

Reference stellar evolution code.

Used as physical benchmark.

Kepler

Purpose:

Asteroseismic observations.

Provides:

- Light curves
 - Oscillation frequencies
-

TESS

Purpose:

Modern all-sky asteroseismology.

Gaia

Purpose:

Distances and stellar properties.

APOGEE

Purpose:

Spectroscopic abundances.

SDSS

Purpose:

Large astronomical surveys.

SIMBAD

Purpose:

Object metadata and literature references.

7. System Architecture

Observational Sources

|

v

Astronomy Data Layer

|

v


Data Normalization

|

v

Differentiable Stellar Model





8. Google Colab Development Environment

Why Colab

Advantages:

- Free GPU access
- Easy sharing
- Reproducibility
- Notebook-based experimentation

Recommended:

- High RAM runtime
- T4 or A100 GPU
- Google Drive integration

Project Structure

```
project/
|
├─ docs/
├─ notebooks/
├─ src/
├─ tests/
├─ data/
├─ outputs/
├─ references/
└─ papers/
```



9. Implementation Roadmap

Phase 1 (Weeks 1-4)

Foundation and MIST validation.

Goals:

- Environment setup
- Data ingestion
- Thomas algorithm
- Hydrostatic equilibrium
- Nuclear burning

Success criterion:

<1% deviation from MIST.

Phase 2 (Weeks 5-12)

Bayesian inference.

Goals:

- ELBO implementation
- Variational inference
- Synthetic recovery tests
- Real Kepler/TESS data

Success criterion:

Recover synthetic parameters within 3σ .

Phase 3 (Weeks 13-16)

Validation and benchmarking.

Goals:

- Speed benchmarks
- Memory profiling
- Gradient validation

Success criterion:

Adjoint gradients substantially outperform finite differences.

Phase 4 (Weeks 17-20)

Publication preparation.

Goals:

- Figures
- Tables
- Comparisons
- Manuscript writing

Target journals:

- The Astrophysical Journal
 - Monthly Notices of the Royal Astronomical Society
 - Astronomy & Astrophysics
-

10. Validation Philosophy

Every test must answer:

1. What scientific claim is being validated?
2. What failure modes exist?
3. What is the acceptable error threshold?
4. What does failure imply physically?

Example:

Hydrostatic Equilibrium Test

Claim:

Pressure profile matches accepted stellar models.

Threshold:

<1% error against MIST.

Failure Meaning:

Incorrect force balance implementation.

11. MCP Server Design

The MCP server should expose:

Observational Queries

- Star metadata
- Spectra
- Light curves

Inference Queries

- Mass estimation
- Radius estimation
- Age estimation

Research Queries

- Literature references
- Validation reports
- Confidence intervals

Agent Tools

- Stellar evolution simulation
 - Posterior generation
 - Asteroseismic analysis
-

12. Reproducibility Requirements

Every milestone must include:

- Notebook
- Source code
- Validation plots
- Configuration file
- Dataset references
- Reproduction instructions

No result should exist without a reproducible notebook.

13. Publication Requirements

Every figure must include:

- Physical interpretation

- Statistical significance
- Validation source
- Reproducibility notes

Every result must contain:

- Mean estimate
 - Uncertainty interval
 - Assumptions
 - Reference dataset
-

Final Expected Outcome

At the end of 20 weeks, the project should produce:

1. A differentiable stellar evolution engine.
2. Bayesian stellar parameter inference framework.
3. Astronomy MCP server for AI agents.
4. Reproducible Colab notebooks.
5. Validation against MIST/MESA.
6. Publication-ready research manuscript.
7. Open-source framework usable by astronomers, ML researchers, and autonomous scientific agents.

This version is much closer to what a PhD reviewer, research scientist, or astrophysicist would expect: it explains the science, the mathematics, the software architecture, the validation strategy, and the publication path in a single coherent document rather than only providing implementation tasks.