

Wide limits – nonstandard analysis meets computational complexity

Ondra Ježil

May 16, 2022

Context

- Imagine a “growing” sequence of finite structures, can we define a limit object (some kind of infinite structure)? (e.g. ‘a structure’ = graph)

Context

- Imagine a “growing” sequence of finite structures, can we define a limit object (some kind of infinite structure)? (e.g. ‘a structure’ = graph)
- Yes! In many (sometimes overlapping) ways: arguments through compactness, ultraproduct, limits of dense graphs, graphons, ...

Context

- Imagine a “growing” sequence of finite structures, can we define a limit object (some kind of infinite structure)? (e.g. ‘a structure’ = graph)
- Yes! In many (sometimes overlapping) ways: arguments through compactness, ultraproduct, limits of dense graphs, graphons, ...
- Studying such structural limits can help us understand asymptotical behaviour of the growing sequence.

Context

- Imagine a “growing” sequence of finite structures, can we define a limit object (some kind of infinite structure)? (e.g. ‘a structure’ = graph)
- Yes! In many (sometimes overlapping) ways: arguments through compactness, ultraproduct, limits of dense graphs, graphons, ...
- Studying such structural limits can help us understand asymptotical behaviour of the growing sequence.
- In this talk I will attempt to explain the central construction of my Master’s thesis and present a few of my results.

Context

- Imagine a “growing” sequence of finite structures, can we define a limit object (some kind of infinite structure)? (e.g. ‘a structure’ = graph)
- Yes! In many (sometimes overlapping) ways: arguments through compactness, ultraproduct, limits of dense graphs, graphons, ...
- Studying such structural limits can help us understand asymptotical behaviour of the growing sequence.
- In this talk I will attempt to explain the central construction of my Master’s thesis and present a few of my results.
- Key method – Forcing

An example

- Consider φ a first order \emptyset -sentence.

An example

- Consider φ a first order \emptyset -sentence.
- e.g. $(\exists x)(\forall y)(x = y)$, $(\exists x)(\exists y)(x \neq y)$,
 $(\forall x)(\exists y)(\exists z)(\exists w)((x = y \leftrightarrow x \neq z) \wedge (w = y))$, ...

An example

- Consider φ a first order \emptyset -sentence.
- e.g. $(\exists x)(\forall y)(x = y)$, $(\exists x)(\exists y)(x \neq y)$,
 $(\forall x)(\exists y)(\exists z)(\exists w)((x = y \leftrightarrow x \neq z) \wedge (w = y))$, ...
- For any set S we can ask whether $S \models \varphi$. 'Is the sentence φ valid in S ?'

An example

- Consider φ a first order \emptyset -sentence.
- e.g. $(\exists x)(\forall y)(x = y)$, $(\exists x)(\exists y)(x \neq y)$,
 $(\forall x)(\exists y)(\exists z)(\exists w)((x = y \leftrightarrow x \neq z) \wedge (w = y))$, ...
- For any set S we can ask whether $S \models \varphi$. 'Is the sentence φ valid in S ?'
- **The problem:** Is $\{|S|; S \models \varphi, S \text{ is finite}\}$ always finite or cofinite?

Solution

Theorem

Let $A := \{|S|; S \text{ is finite}, S \models \varphi\}$. Then either A is finite or $\mathbb{N} \setminus A$ is finite.

Proof.

Assume A is not finite, we will prove, $\mathbb{N} \setminus A$ is infinite.

For contradiction, assume $\mathbb{N} \setminus A$ is infinite. Compactness theorem from mathematical logic lets us then construct countable sets $S_1 \models \varphi$ and $S_2 \models \neg\varphi$. But such sets without no structure are isomorphic! A contradiction. □

Wide limits – motivation

- Our goal is similar. That is to understand asymptotical behaviour of a class of finite structures through one limiting object.

Wide limits – motivation

- Our goal is similar. That is to understand asymptotical behaviour of a class of finite structures through one limiting object.
- A key difference – we have another parameter F , a set of functions with some restricted computational complexity. (e.g. polynomial time functions, function computed by decision trees, constant functions, ...)

Wide limits – motivation

- Our goal is similar. That is to understand asymptotical behaviour of a class of finite structures through one limiting object.
- A key difference – we have another parameter F , a set of functions with some restricted computational complexity. (e.g. polynomial time functions, function computed by decision trees, constant functions, ...)
- The wide limit $\lim_F \mathcal{G}_n$ represents how ‘an average’ graph from \mathcal{G}_k looks like to F .

Wide limits – motivation

- Our goal is similar. That is to understand asymptotical behaviour of a class of finite structures through one limiting object.
- A key difference – we have another parameter F , a set of functions with some restricted computational complexity. (e.g. polynomial time functions, function computed by decision trees, constant functions, ...)
- The wide limit $\lim_F \mathcal{G}_n$ represents how ‘an average’ graph from \mathcal{G}_k looks like to F .

Wide limits – motivation

- Our goal is similar. That is to understand asymptotical behaviour of a class of finite structures through one limiting object.
- A key difference – we have another parameter F , a set of functions with some restricted computational complexity. (e.g. polynomial time functions, function computed by decision trees, constant functions, ...)
- The wide limit $\lim_F \mathcal{G}_n$ represents how ‘an average’ graph from \mathcal{G}_k looks like to F .

existential formulas in the limit \leftrightarrow hardness of search problems

Wide limits – Setup

- Let us now specify for what kind of classes of structures we will define a limit object for.

Wide limits – Setup

- Let us now specify for what kind of classes of structures we will define a limit object for.
- We restrict ourselves to graphs, but in theory any kind of relational (and even algebraic) structures can be used.

Wide limits – Setup

- Let us now specify for what kind of classes of structures we will define a limit object for.
- We restrict ourselves to graphs, but in theory any kind of relational (and even algebraic) structures can be used.

Wide limits – Setup

- Let us now specify for what kind of classes of structures we will define a limit object for.
- We restrict ourselves to graphs, but in theory any kind of relational (and even algebraic) structures can be used.

Definition (Wide sequence)

A sequence $\{\mathcal{G}_k\}_{k=1}^{\infty}$ of non-empty sets of finite graphs a **wide sequence** if

- *there is a strictly increasing sequence of positive integers $\{g_k\}_{k=1}^{\infty}$ such that the vertex set of all $G \in \mathcal{G}_k$ is $\{0, \dots, g_k - 1\}$,*
- $\lim_{k \rightarrow \infty} |\mathcal{G}_k| = \infty$. (Hence, wide!)

Examples of wide sequences

- $\text{EDGE}_k = \{(\{0, \dots, k-1\}, E); |E| = 1\}$

Examples of wide sequences

- $\text{EDGE}_k = \{(\{0, \dots, k-1\}, E); |E| = 1\}$
- $\text{nonEDGE}_k = \{(\{0, \dots, k-1\}, E); |E| = \binom{k}{2} - 1\}$

Examples of wide sequences

- $\text{EDGE}_k = \{(\{0, \dots, k-1\}, E); |E| = 1\}$
- $\text{nonEDGE}_k = \{(\{0, \dots, k-1\}, E); |E| = \binom{k}{2} - 1\}$
- $\text{SK}_k^{1/2} = \{(\{0, \dots, k-1\}, E); E \text{ consist of exactly one } \lfloor \frac{k}{2} \rfloor \text{ clique}\}$

Examples of wide sequences

- $\text{EDGE}_k = \{(\{0, \dots, k-1\}, E); |E| = 1\}$
- $\text{nonEDGE}_k = \{(\{0, \dots, k-1\}, E); |E| = \binom{k}{2} - 1\}$
- $\text{SK}_k^{1/2} = \{(\{0, \dots, k-1\}, E); E \text{ consist of exactly one } \lfloor \frac{k}{2} \rfloor \text{ clique}\}$
- And so on...

Nonstandard models of arithmetic – definition

- Let us now introduce an important concept from mathematical logic which we will need later on.

Nonstandard models of arithmetic – definition

- Let us now introduce an important concept from mathematical logic which we will need later on.
- By $\text{Th}(\mathbb{N})$ we mean the set of all true sentences of natural numbers. So axioms of ordered discrete semirings, all instances of induction, the four squares theorem, Fermat's last theorem and so on.

Nonstandard models of arithmetic – definition

- Let us now introduce an important concept from mathematical logic which we will need later on.
- By $\text{Th}(\mathbb{N})$ we mean the set of all true sentences of natural numbers. So axioms of ordered discrete semirings, all instances of induction, the four squares theorem, Fermat's last theorem and so on.
- In essence every theorem we will ever prove and more.

Nonstandard models of arithmetic – definition

- Let us now introduce an important concept from mathematical logic which we will need later on.
- By $\text{Th}(\mathbb{N})$ we mean the set of all true sentences of natural numbers. So axioms of ordered discrete semirings, all instances of induction, the four squares theorem, Fermat's last theorem and so on.
- In essence every theorem we will ever prove and more.
- By standard theorems of model theory, there exist a semiring $\mathcal{M} \models \text{Th}(\mathbb{N})$ which is **not isomorphic** to \mathbb{N} .

Nonstandard models of arithmetic – definition

- Let us now introduce an important concept from mathematical logic which we will need later on.
- By $\text{Th}(\mathbb{N})$ we mean the set of all true sentences of natural numbers. So axioms of ordered discrete semirings, all instances of induction, the four squares theorem, Fermat's last theorem and so on.
- In essence every theorem we will ever prove and more.
- By standard theorems of model theory, there exist a semiring $\mathcal{M} \models \text{Th}(\mathbb{N})$ which is **not isomorphic** to \mathbb{N} .
- \mathcal{M} contains an isomorphic copy of \mathbb{N} and then some 'infinite numbers', we call those elements **nonstandard numbers**.

Nonstandard models of arithmetic – definition

- Let us now introduce an important concept from mathematical logic which we will need later on.
- By $\text{Th}(\mathbb{N})$ we mean the set of all true sentences of natural numbers. So axioms of ordered discrete semirings, all instances of induction, the four squares theorem, Fermat's last theorem and so on.
- In essence every theorem we will ever prove and more.
- By standard theorems of model theory, there exist a semiring $\mathcal{M} \models \text{Th}(\mathbb{N})$ which is **not isomorphic** to \mathbb{N} .
- \mathcal{M} contains an isomorphic copy of \mathbb{N} and then some 'infinite numbers', we call those elements **nonstandard numbers**.
- From now on we fix one $\mathcal{M} \models \text{Th}(\mathbb{N})$ which satisfies a technical condition. (It is \aleph_1 -saturated.)

Pseudofinite structures

- Let $n \in \mathcal{M} \setminus \mathbb{N}$ be a nonstandard number. We can then use it as an index in \mathcal{G}_k to obtain the n -th level of $\{\mathcal{G}_k\}_{k=0}^\infty$.

Pseudofinite structures

- Let $n \in \mathcal{M} \setminus \mathbb{N}$ be a nonstandard number. We can then use it as an index in \mathcal{G}_k to obtain the n -th level of $\{\mathcal{G}_k\}_{k=0}^{\infty}$.
- Graphs $G \in \mathcal{G}_n$ are what we call **pseudofinite**, they satisfy the theory of all finite graphs.

Pseudofinite structures

- Let $n \in \mathcal{M} \setminus \mathbb{N}$ be a nonstandard number. We can then use it as an index in \mathcal{G}_k to obtain the n -th level of $\{\mathcal{G}_k\}_{k=0}^{\infty}$.
- Graphs $G \in \mathcal{G}_n$ are what we call **pseudofinite**, they satisfy the theory of all finite graphs.
- Example: In EDGE_n we get graphs on the infinite vertex set $\{0, \dots, n-1\}$ such that there is exactly one edge.

The (random) vertex family F

- The other parameter of the limit yet to be specified is the family of functions F .

The (random) vertex family F

- The other parameter of the limit yet to be specified is the family of functions F .
- The main idea is this, imagine we uniformly sample a graph $\omega \in \mathcal{G}_n$, this would be impossible in a purely infinite setting, but we are in pseudofinite setting, so such sampling can take place in \mathcal{M} .

The (random) vertex family F

- The other parameter of the limit yet to be specified is the family of functions F .
- The main idea is this, imagine we uniformly sample a graph $\omega \in \mathcal{G}_n$, this would be impossible in a purely infinite setting, but we are in pseudofinite setting, so such sampling can take place in \mathcal{M} .
- The sampled graph is then used as an input to the function $\alpha \in F$ and outputs an element in $\{0, \dots, g_k - 1\}$ which is a vertex in each ω .


The (random) vertex family F

- The other parameter of the limit yet to be specified is the family of functions F .
- The main idea is this, imagine we uniformly sample a graph $\omega \in \mathcal{G}_n$, this would be impossible in a purely infinite setting, but we are in pseudofinite setting, so such sampling can take place in \mathcal{M} .
- The sampled graph is then used as an input to the function $\alpha \in F$ and outputs an element in $\{0, \dots, g_k - 1\}$ which is a vertex in each ω .
- The main vertex family we consider is F_{rud} the family of functions computed by decision trees of depth $n^{1/t}$ for t nonstandard.

The vertex family F_{rud}

Definition

The vertex family F_{rud} consists of all functions α with input \mathcal{G}_n and output computed by a directed tree of depth at most $g_n^{1/t}$, for some nonstandard t , whose form we present by the following example.



\mathcal{M} -rational numbers

- Let us elaborate on the uniform distribution on $\Omega := \mathcal{G}_k$.

\mathcal{M} -rational numbers

- Let us elaborate on the uniform distribution on $\Omega := \mathcal{G}_k$.
- We assign probability to sample each $\omega \in \mathcal{M}$ to be $1/|\mathcal{G}_n|$ as an ' \mathcal{M} -rational number'.

\mathcal{M} -rational numbers

- Let us elaborate on the uniform distribution on $\Omega := \mathcal{G}_k$.
- We assign probability to sample each $\omega \in \mathcal{M}$ to be $1/|\mathcal{G}_n|$ as an ' \mathcal{M} -rational number'.
- We can construct the \mathcal{M} -rational numbers $\mathbb{Q}^{\mathcal{M}}$ in an analogous way to how we construct \mathbb{Q} from \mathbb{N} . (Add negative elements and take the fraction field.)

\mathcal{M} -rational numbers

- Let us elaborate on the uniform distribution on $\Omega := \mathcal{G}_k$.
- We assign probability to sample each $\omega \in \mathcal{M}$ to be $1/|\mathcal{G}_n|$ as an ' \mathcal{M} -rational number'.
- We can construct the \mathcal{M} -rational numbers $\mathbb{Q}^{\mathcal{M}}$ in an analogous way to how we construct \mathbb{Q} from \mathbb{N} . (Add negative elements and take the fraction field.)
- Those \mathcal{M} -rationals bounded by some $n \in \mathbb{N}$ we call **finite**, and those which are bounded by all $\frac{1}{n}$ we call **infinitesimal**.

\mathcal{M} -rational numbers

- Let us elaborate on the uniform distribution on $\Omega := \mathcal{G}_k$.
- We assign probability to sample each $\omega \in \mathcal{M}$ to be $1/|\mathcal{G}_n|$ as an ' \mathcal{M} -rational number'.
- We can construct the \mathcal{M} -rational numbers $\mathbb{Q}^{\mathcal{M}}$ in an analogous way to how we construct \mathbb{Q} from \mathbb{N} . (Add negative elements and take the fraction field.)
- Those \mathcal{M} -rationals bounded by some $n \in \mathbb{N}$ we call **finite**, and those which are bounded by all $\frac{1}{n}$ we call **infinitesimal**.

\mathcal{M} -rational numbers

- Let us elaborate on the uniform distribution on $\Omega := \mathcal{G}_k$.
- We assign probability to sample each $\omega \in \mathcal{M}$ to be $1/|\mathcal{G}_n|$ as an ' \mathcal{M} -rational number'.
- We can construct the \mathcal{M} -rational numbers $\mathbb{Q}^{\mathcal{M}}$ in an analogous way to how we construct \mathbb{Q} from \mathbb{N} . (Add negative elements and take the fraction field.)
- Those \mathcal{M} -rationals bounded by some $n \in \mathbb{N}$ we call **finite**, and those which are bounded by all $\frac{1}{n}$ we call **infinitesimal**.

Theorem

There exists a surjective, order preserving ring homomorphism $\varphi : \mathbb{Q}_{fin}^{\mathcal{M}} \rightarrow \mathbb{R}$ whose kernel is the ideal of infinitesimal numbers. We call the image $st(q)$ the standard part of the \mathcal{M} -rational q .

Nonstandard analysis

How does this correspond to classical analysis?

Nonstandard analysis

How does this correspond to classical analysis?

Theorem

Let $\{b_k\}_{k=1}^{\infty}$ be a sequence of rational numbers then

$$\lim_{k \rightarrow \infty} b_k = r \in \mathbb{R}$$

if and only if for all nonstandard $n \in \mathcal{M}$

$$st(b_n) = r \in \mathbb{R}.$$

The Boolean algebra \mathcal{B}

- Boolean algebra is a complemented distributive lattice. The maximum element is denoted **1** and the minimum **0**.

The Boolean algebra \mathcal{B}

- Boolean algebra is a complemented distributive lattice. The maximum element is denoted **1** and the minimum **0**.
- Examples $(\{0, 1\}, \wedge, \vee, \neg, \mathbf{0}, \mathbf{1}), (\mathcal{P}(X), \cap, \cup, X \setminus -, \emptyset, X), \dots$

The Boolean algebra \mathcal{B}

- Boolean algebra is a complemented distributive lattice. The maximum element is denoted $\mathbf{1}$ and the minimum $\mathbf{0}$.
- Examples $(\{0, 1\}, \wedge, \vee, \neg, \mathbf{0}, \mathbf{1}), (\mathcal{P}(X), \cap, \cup, X \setminus -, \emptyset, X), \dots$
- We start with the boolean algebra $\mathcal{A} = \{A \subseteq \mathcal{G}_n; A \in \mathcal{M}\}$ of subsets of the sample space \mathcal{G}_n represented by elements in \mathcal{M} .

The Boolean algebra \mathcal{B}

- Boolean algebra is a complemented distributive lattice. The maximum element is denoted $\mathbf{1}$ and the minimum $\mathbf{0}$.
- Examples $(\{0, 1\}, \wedge, \vee, \neg, \mathbf{0}, \mathbf{1}), (\mathcal{P}(X), \cap, \cup, X \setminus -, \emptyset, X), \dots$
- We start with the boolean algebra $\mathcal{A} = \{A \subseteq \mathcal{G}_n; A \in \mathcal{M}\}$ of subsets of the sample space \mathcal{G}_n represented by elements in \mathcal{M} .
- The counting measure is defined as $A \mapsto |A|/|\mathcal{G}_n| \in \mathbb{Q}^{\mathcal{M}}$.

The Boolean algebra \mathcal{B}

- Boolean algebra is a complemented distributive lattice. The maximum element is denoted $\mathbf{1}$ and the minimum $\mathbf{0}$.
- Examples $(\{0, 1\}, \wedge, \vee, \neg, \mathbf{0}, \mathbf{1}), (\mathcal{P}(X), \cap, \cup, X \setminus -, \emptyset, X), \dots$
- We start with the boolean algebra $\mathcal{A} = \{A \subseteq \mathcal{G}_n; A \in \mathcal{M}\}$ of subsets of the sample space \mathcal{G}_n represented by elements in \mathcal{M} .
- The counting measure is defined as $A \mapsto |A|/|\mathcal{G}_n| \in \mathbb{Q}^{\mathcal{M}}$.
- The boolean algebra \mathcal{B} is defined as the factor-algebra \mathcal{A}/\mathcal{I} where \mathcal{I} is the ideal of \mathcal{A} of all elements with infinitesimal counting measure.

The Boolean algebra \mathcal{B} – what we need

- **Key point:** The truth values of the first order sentences of the wide limit $\lim_F \mathcal{G}_n$ are the \mathcal{M} -definable subsets of \mathcal{G}_n modulo sets of infinitesimal measure. We then have that \mathcal{B} is a σ -algebra and that $\mu : A/\mathcal{I} \rightarrow \text{st}(|A|/|\mathcal{G}_n|)$ is measure in the classical sense.

The Boolean algebra \mathcal{B} – what we need

- **Key point:** The truth values of the first order sentences of the wide limit $\lim_F \mathcal{G}_n$ are the \mathcal{M} -definable subsets of \mathcal{G}_n modulo sets of infinitesimal measure. We then have that \mathcal{B} is a σ -algebra and that $\mu : A/\mathcal{I} \rightarrow \text{st}(|A|/|\mathcal{G}_n|)$ is measure in the classical sense.
- Moreover, we have that \mathcal{B} is a complete Boolean algebra, it contains all infinite disjunctions and conjunctions.

The definition of a wide limit

Definition (Wide limit)

For a wide sequence \mathcal{G}_k , vertex family F and a nonstandard number n we define the wide limit

$$\lim_F \mathcal{G}_n$$

as a \mathcal{B} -valued graph whose vertex set is F and we interpret equality as

$$\lim_F \mathcal{G}_n[\![\alpha = \beta]\!] = \{\omega \in \mathcal{G}_n; \alpha(\omega) = \beta(\omega)\} / \mathcal{I},$$

and the edge relational symbol as

$$\lim_F \mathcal{G}_n[\![E(\alpha, \beta)]\!] = \{\omega \in \mathcal{G}_n; E_\omega(\alpha(\omega), \beta(\omega))\} / \mathcal{I},$$

and let it commute with \neg , \wedge , \vee and interpret \forall and \exists respectively as infinite conjunctions and disjunctions.

$$\mathcal{G}_k = \text{EDGE}_k$$

- Recall the wide sequence EDGE_k which roughly consist of all finite undirected graphs with one edge.

$$\mathcal{G}_k = \text{EDGE}_k$$

- Recall the wide sequence EDGE_k which roughly consist of all finite undirected graphs with one edge.
- Every $\omega \in \text{EDGE}_k$ has an edge.

$$\mathcal{G}_k = \text{EDGE}_k$$

- Recall the wide sequence EDGE_k which roughly consist of all finite undirected graphs with one edge.
- Every $\omega \in \text{EDGE}_k$ has an edge.
- What happens in the wide limit (with $F = F_{rud}$)?

$$\mathcal{G}_k = \text{EDGE}_k$$

- Recall the wide sequence EDGE_k which roughly consist of all finite undirected graphs with one edge.
- Every $\omega \in \text{EDGE}_k$ has an edge.
- What happens in the wide limit (with $F = F_{rud}$)?
- What is

$$\lim_{F_{rud}} \text{EDGE}_n \llbracket (\exists x)(\exists y)E(x, y) \rrbracket = ?$$

$$\mathcal{G}_k = \text{EDGE}_k$$

- Recall the wide sequence EDGE_k which roughly consist of all finite undirected graphs with one edge.
- Every $\omega \in \text{EDGE}_k$ has an edge.
- What happens in the wide limit (with $F = F_{rud}$)?
- What is

$$\lim_{F_{rud}} \text{EDGE}_n \llbracket (\exists x)(\exists y)E(x, y) \rrbracket = ?$$

$$\mathcal{G}_k = \text{EDGE}_k$$

- Recall the wide sequence EDGE_k which roughly consist of all finite undirected graphs with one edge.
- Every $\omega \in \text{EDGE}_k$ has an edge.
- What happens in the wide limit (with $F = F_{rud}$)?
- What is

$$\lim_{F_{rud}} \text{EDGE}_n \llbracket (\exists x)(\exists y)E(x, y) \rrbracket = ?$$

- It is

$$\lim_{F_{rud}} \text{EDGE}_n \llbracket (\exists x)(\exists y)E(x, y) \rrbracket = \mathbf{0}$$

Theorem

$$\lim_{F_{rud}} \text{EDGE}_n[(\exists x)(\exists y)E(x, y)] = \mathbf{0}$$

Proof.

The universe of the wide limit is F_{rud} the (random) vertices computed by decision trees of depth $n^{1/t} = g_n^{1/t}$ for some $t > \mathbb{N}$.

So we need to prove that any two such trees find the edge in uniformly sampled $\omega \in \text{EDGE}_n$ with uniform probability. We can just compose the trees into one tree T which actually outputs the edge.

Theorem

$$\lim_{F_{rud}} EDGE_n \llbracket (\exists x)(\exists y)E(x, y) \rrbracket = \mathbf{0}$$

Proof cont.

So assume T is a possibly successful tree. We will prove it is definitely not successful.

Go down the tree and answer $\mathbf{0}$ to every query of the tree. Then the T cannot possibly succeed, what is the ratio of $\omega \in EDGE_k$ which result in such answers?

Theorem

$$\lim_{F_{rud}} EDGE_n[(\exists x)(\exists y)E(x, y)] = 0$$

Proof cont.

So we compute the ratio of those graphs in which the edge is not among $n^{1/t}$ many specified pairs of vertices.

$$\frac{\binom{n-n^{1/t}}{2}}{\binom{n}{2}} = \frac{(n - n^{1/t})(n - n^{1/t} - 1)}{n(n - 1)} \quad (1)$$

$$= \left(1 - \frac{n^{1/t}}{n}\right) \left(1 - \frac{n^{1/t} + 1}{n - 1}\right) \quad (2)$$

$$\geq 1 - \frac{2n^{1/t} + 2}{n - 1} \quad (3)$$

but the standard part of (3) is 1! So the probability any tree succeeds is infinitesimally close to 0. □

Too sparse wide sequences

- We say a wide sequence \mathcal{G}_k is too sparse, if its wide limit is the empty graph.

Too sparse wide sequences

- We say a wide sequence \mathcal{G}_k is too sparse, if its wide limit is the empty graph.
- We can analyze this situation more generally for F_{rud} .

Too sparse wide sequences

- We say a wide sequence \mathcal{G}_k is too sparse, if its wide limit is the empty graph.
- We can analyze this situation more generally for F_{rud} .

Too sparse wide sequences

- We say a wide sequence \mathcal{G}_k is too sparse, if its wide limit is the empty graph.
- We can analyze this situation more generally for F_{rud} .

Theorem

If the probability that a uniformly random pair of vertices on a uniformly random $G \in \mathcal{G}_k$ is $o(1/\sqrt[j]{k})$ for any $j \in \mathbb{N}$ then

$$\lim_{F_{rud}} \mathcal{G}_n \llbracket (\exists x)(\exists y)E(x, y) \rrbracket = \mathbf{0}.$$

$$\mathcal{G}_k = \text{ALL}_k$$

- No edges? Too boring!

$$\mathcal{G}_k = \text{ALL}_k$$

- No edges? Too boring!

$$\mathcal{G}_k = \text{ALL}_k$$

- No edges? Too boring!
- Let us see some stuff!

$$\mathcal{G}_k = \text{ALL}_k$$

- No edges? Too boring!
- Let us see some stuff!

$$\mathcal{G}_k = \text{ALL}_k$$

- No edges? Too boring!
- Let us see some stuff!
- Consider ALL_k the wide sequence of all finite graphs, there ought to be something, right?

$$\mathcal{G}_k = \text{ALL}_k$$

- No edges? Too boring!
- Let us see some stuff!
- Consider ALL_k the wide sequence of all finite graphs, there ought to be something, right?

$$\mathcal{G}_k = \text{ALL}_k$$

- No edges? Too boring!
- Let us see some stuff!
- Consider ALL_k the wide sequence of all finite graphs, there ought to be something, right?

Theorem (Everything exists)

Let $\varphi(\bar{x})$ be a formula in the language of graphs (possibly with parameters), then

$$\lim_{F_{rud}} \text{ALL}_k \llbracket (\exists \bar{x}) \varphi(x) \rrbracket = 1.$$

Proof. (sketch).

Iterate search on $n^{1/t}$ different tuples of ω and use mutual independence. □

Large clique

- Let $SK_k^{1/2}$ be the wide sequence of all graphs containing one copy of a $\lfloor k/2 \rfloor$ -clique.

Large clique

- Let $SK_k^{1/2}$ be the wide sequence of all graphs containing one copy of a $\lfloor k/2 \rfloor$ -clique.

Theorem

$$\lim_{F_{rud}}^{G_{rud}} \mathcal{G}_n[\textit{there is a } \lfloor n/(2 \ln n) \rfloor \textit{ clique}] = \mathbf{1}$$

Large clique

- Let $SK_k^{1/2}$ be the wide sequence of all graphs containing one copy of a $\lfloor k/2 \rfloor$ -clique.

Theorem

$$\lim_{F_{rud}}^{G_{rud}} \mathcal{G}_n[\textit{there is a } \lfloor n/(2 \ln n) \rfloor \textit{ clique}] = \mathbf{1}$$

- Open problem: Is this the biggest clique here? What about in the wide sequence which contains a large clique and possibly other edges?

What's next?

- Further analyze when we get the truth values strictly in the set $\{0, 1\}$.

What's next?

- Further analyze when we get the truth values strictly in the set $\{0, 1\}$.
- Analyze the F_{poly} , connection to **P** vs. **NP**.

What's next?

- Further analyze when we get the truth values strictly in the set $\{\mathbf{0}, \mathbf{1}\}$.
- Analyze the F_{poly} , connection to **P** vs. **NP**.
- Easily generalizable to any relational and algebraic structure.

What's next?

- Further analyze when we get the truth values strictly in the set $\{0, 1\}$.
- Analyze the F_{poly} , connection to **P** vs. **NP**.
- Easily generalizable to any relational and algebraic structure.
- One can then take a wide limit of **CSP**(\mathbb{A}), connection to tractability.

What's next?

- Further analyze when we get the truth values strictly in the set $\{\mathbf{0}, \mathbf{1}\}$.
- Analyze the F_{poly} , connection to **P** vs. **NP**.
- Easily generalizable to any relational and algebraic structure.
- One can then take a wide limit of **CSP**(\mathbb{A}), connection to tractability.
- The ultimate direction to pursue is to characterize some wide limit independently of the direct definition – this immediately results in (probabilistic) lower and upper bounds for F .

$$\lim_F \mathcal{G}_n[\textbf{Thank you!}] = \mathbf{1}$$