

---

**PROTOCOL TITLE**

**Evaluation of clinical outcome in Asian patients with localised rectal adenocarcinoma managed with watch-and-wait approach by developing a multinational collaborative database.**

**Short title: Asian Watch-and-Wait Database (AWWD)**

---

## TABLE OF CONTENTS

1. INTRODUCTION AND RATIONALE .....	42.
OBJECTIVES .....	53.
STUDY DESIGN .....	<b>Error! Bookmark not defined.</b> 4.
STUDY POPULATION.....	7
4.1 Population.....	74.2
Inclusion criteria.....	74.3
Exclusion criteria.....	75.
METHODS .....	75.1
Study objectives/endpoints .....	85.1.1
Main study objective/endpoint.....	85.1.2
Secondary objective/endpoints (if applicable) .....	85.2
Study procedures .....	8
5.3 Withdrawal of individual subjects.....	95.4
Follow-up of subjects withdrawn from treatment .....	96.
SAFETY REPORTING.....	96.1
AEs, SAEs and SUSARs .....	9
6.2 Data Safety Monitoring Board (DSMB) / Safety Committee.....	97.
STATISTICAL ANALYSIS.....	97.1
Primary study parameter(s).....	107.2
Secondary study parameter(s) .....	107.3
Statistical Analysis plan .....	108.
ETHICAL CONSIDERATIONS.....	108.1
Recruitment and consent .....	118.2
Considerations for incapacitated subjects or vulnerable population (if applicable).....	118.3
Benefits and risks assessment.....	118.4
Compensation for injury .....	118.5
Incentives (if applicable) .....	119.
ADMINISTRATIVE ASPECTS, MONITORING AND PUBLICATION.....	119.1
Handling and storage of data and documents .....	129.2
Monitoring and Quality Assurance .....	129.3
Publication policy .....	1210.
REFERENCES .....	13

---

## **LIST OF ABBREVIATIONS AND RELEVANT DEFINITIONS**

cCR: Clinical Complete Response

DFS: Disease free survival

DICOM: Digital Imaging and Communications in Medicine

DSMB: Data safety and monitoring board

ECOG: Eastern Cooperative Oncology Group

EORTC: European Organisation for Research and Treatment of Cancer

IQR: Inter Quartile Range

OS: Overall Survival

PI: Principal Investigator

QLQ: Quality of Life Questionnaire

QOL: Quality of Life

REDCap: Research Electronic Data Capture

SD: Standard Deviation

SPSS: Statistical Package for the Social Sciences

TME: Total Mesorectal Excision

TNT: Total Neoadjuvant Therapy

---

## **SUMMARY**

The standard treatment approach of localised rectal adenocarcinoma involves neoadjuvant radiotherapy and chemotherapy followed by assessment for surgery and optional adjuvant chemotherapy. However, such patients who achieve complete or near complete clinical response after neoadjuvant therapy are increasingly being managed with the 'Watch-and-Wait strategy' in the recent years. Although safety of such policy has been established, the patient population in the published studies are not represented well by Asian population. Moreover, the data about safety of such policy in the setting of modern neoadjuvant strategies is limited. We propose a multinational collaborative prospective database study in Asia to systematically collect data on patients being managed with such a strategy. The target population is patients with localised rectal adenocarcinoma achieving near complete or complete response being managed by the Watch-and-Wait strategy. The primary end point being local regrowth and other survival and quality of life end points will be evaluated as well. A secure multinational database will be generated over 5 years with an estimated expected minimal sample size of 337 patients over 5 years across 7-10 Asian institutes to precisely evaluate the required endpoints. Ethical and legal requirements will be met at each participating institute.

---

## 1. INTRODUCTION AND RATIONALE

Neoadjuvant radiation and chemotherapy followed by total mesorectal excision (TME) remains a standard approach in most of the centres globally. Although the oncological outcomes have improved significantly with this multimodality approach over the last few decades, this approach adversely affects quality of life and functional outcomes especially for those needing permanent colostomy. More recently, patients achieving complete or near complete response are increasingly being managed by the 'Watch-and-Wait' approach where in, patients are followed using an intense follow up protocol after radiation and chemotherapy and avoid or delay surgery(1)(2)(3)(4)(5). However, this approach has not been widely adopted worldwide and there is limited data in the published medical literature. While the randomized trials in this setting does not appear to be feasible for obvious ethical reasons, robust data collected both retrospectively and prospectively can provide insights into this approach and help us refine the Watch-and-Wait approach in our routine clinics and in optimizing patient selection and surveillance strategies. Such an attempt was done by developing a multinational database from the Netherlands, Brazil, Portugal, and the UK(6). This data base consisted of more than a thousand patients and several important clinical questions were answered using this data(7)(8). However, this database largely consisted of patients treated using long course chemoradiation which was considered a standard treatment previously at that time. Moreover, there was no representation from Asian countries in this database. Currently, with the recent publications from the large, randomised trials like RAPIDO, UNICANCER PRODIGE 23, OPRA, more and more patients with locally advanced nonmetastatic rectal adenocarcinoma are managed with a total neoadjuvant therapy approach(9)(10)(11). Radiotherapy dose escalation using brachytherapy has also gained interest with recent publication of results from the OPERA trial(12). Hence, with these developments, management practice of locally advanced rectal adenocarcinomas has become heterogeneous across institutions globally. These patients are being treated with more aggressive neoadjuvant therapies including total neoadjuvant therapy (TNT) approach. Many of these regimens involve multiagent neoadjuvant chemotherapy in addition to chemoradiation or short course hypofractionated radiotherapy leading increased response rates. The knowledge about the risks and benefits of watch-and-wait strategy after newer neoadjuvant strategies is somewhat limited. It is not entirely known whether the watch-and-wait approach can be applied safely to these patients who achieve clinical complete response after more aggressive neoadjuvant therapy regimens. Also, the data on Asian patients managed with such a strategy is scarce in the literature. Not only the inherent biological and genetic factors associated with rectal adenocarcinoma in Asian patients may be different but also the compliance and feasibility of intense surveillance protocol can be challenging in some of the Asian countries. Considering all these, we intend to develop a comprehensive and secure database exclusively from Asian countries and to collect meaningful data systematically to answer important clinical questions on watch-and-wait strategy in Asian patients. This effort is likely to strengthen available evidence on this subject and generate evidence based on data exclusively from Asian patients. Collective efforts from participating nations can facilitate mutual learning and help in safe adoption of the watch-

---

and-wait strategy in a greater number of centres across Asia. This collaborative study among Asian countries may potentially pave the way for future studies of similar kind.

## **2. OBJECTIVES**

### Primary Objective

- To determine the local regrowth rates at 2 years in Asian patients managed with the watch-and-wait strategy after neoadjuvant therapy in rectal cancer.

### Secondary Objectives

- To determine the rates of overall survival (OS) at 2 years and 5 years
- To determine disease free survival (DFS) at 2 years and 5 years
- To determine the rates of colostomy free survival at 2 years and 5 years
- To determine the clinical complete response rates (cCR) and regrowth rates with respect to various neoadjuvant treatment approaches
- To determine Quality of life (QOL) of patients managed with Watch-and-wait policy.
- To determine compliance of these patients to recommended surveillance protocol.

## **3. STUDY DESIGN**

Prospective observational study

---

## **4. STUDY POPULATION**

### **4.1 Population**

### **4.2 Inclusion criteria**

- Histologically confirmed diagnosis of adenocarcinoma of the rectum
- Age  $\geq 18$  years
- Clinical stage T1-T4, N0-N2, M0 at diagnosis
- ECOG Performance status 0-1
- Neoadjuvant treatment with either short course or long course radiation therapy and / or chemotherapy
- Complete or near complete clinical response after neoadjuvant therapy
- Patients willing to be on watch-and-wait strategy.
- The distance from anal verge of tumour up to 10 cm (Mid and low rectal tumours)

### **4.3 Exclusion criteria**

- Recurrent rectal cancer
- Patients with any other concurrent medical or psychiatric condition or disease which would make them inappropriate candidates for entry into this study.
- Not willing to consent for the study or to follow up routinely

---

## 5. METHODS

### 5.1 Study objectives/endpoints

#### 5.1.1 Main study objective/endpoint:

- To determine the local regrowth rates at 2 years in Asian patients managed with the watch-and-wait strategy after neoadjuvant therapy in rectal cancer.

#### 5.1.2 Secondary objective/endpoints:

- To determine the rates of overall survival (OS) at 2 years and 5 years
- To determine disease free survival (DFS) at 2 years and 5 years
- To determine the rates of colostomy free survival at 2 years and 5 years
- To determine the clinical complete response rates (cCR) and regrowth rates with respect to various neoadjuvant treatment approaches
- To determine Quality of life (QOL) of patients managed with Watch-and-wait policy.
- To determine compliance of these patients to recommended surveillance protocol.

### 5.2 Study procedures

- a. Participation of centres: Centres practicing watch-and-wait approach and willing to contribute to the database are invited to participate in this collaborative study. A central review committee will evaluate and approve the eligibility of the centre participating to ensure the quality of the data collected. Participating centres will be responsible for clearance and approval from local ethics boards and other regulatory bodies. A designated local Principal Investigator (PI) for individual site will be responsible for the regulatory formalities, participant accrual, data collection, storage and sharing of the data.
- b. Participants satisfying eligibility criteria are invited to participate in this prospective observational study and will be included in the study after informed consenting process. Data will be collected prospectively by each participating centre and electronically uploaded on a secure database after pseudonymisation using REDCap software. A separate password protected datasheet for each institute will be maintained by respective local PI, however the Study Principal Investigator will be able to access the data.
- c. Data variables to be collected include patient demographics, family history, personal history, symptomatology, clinical findings, stage and location of the tumour, histology, biomarkers, staging details, diagnostic imaging datasets (DICOM format), lab parameters, serial tumour marker levels, details of neoadjuvant radiation and chemotherapy, response assessment findings, surveillance imaging and endoscopic images, treatment details for regrowth or



---

relapse, need for colostomy etc. The data is collected up to 5 years after enrolment.

- d. No investigations or therapeutic interventions will be done as a part of this study and the nature of the study remains observational only.

### **5.3 Withdrawal of individual subjects**

Participants have the right to withdraw any time after consenting and the data will not be collected any further after withdrawal. However, the data collected up till the withdrawal can be used for analysis.

### **5.4 Follow-up of subjects withdrawn from treatment**

The follow up of the patients after withdrawal will continue as per standard of care as this is an observational study.

## **6. SAFETY REPORTING**

**6.1 AEs, SAEs and SUSARs:** Not applicable as this is an observational study.

### **6.2 Data Safety Monitoring Board (DSMB) / Safety Committee**

The study will be monitored by an independent data safety monitoring committee from the respective institute as per guidance of the institutional review board.

---

## 7. STATISTICAL ANALYSIS

Sample Size justification: The primary objective of the study is to evaluate the local regrowth at 2 years reduces in watch-and-wait patients as compared to surgery patients. Considering a local regrowth rate of 25% it is estimated that approximately 306 patients are required to be included in the study to achieve a precision of 10% at 5% level of significance and 80% power without attrition. Assuming 10% attrition rate a total of 337 patients will be required for the study. This appears feasible expecting a contribution of 35-70 patients each over 3 years from about 7-10 centres across Asia.

### 7.1 Primary study parameter (Definitions)

- Local regrowth rate: Rate of local regrowth at 2 years from the date of diagnosis to date of regrowth

### 7.2 Secondary study parameters (Definitions)

- Rates of overall survival at 2 years and 5 years is measured as a time to event end point from the date of diagnosis to date of death from any cause
- Disease free survival at 2 years and 5 years measured as a time to event end point from the date of diagnosis to date of relapse or death from any cause
- Colostomy free survival at 2 years and 5 years is measured as a time to event end point from the date of diagnosis to date of colostomy or date of death from any cause
- Clinical complete response rates and regrowth rates with respect to various neoadjuvant treatment approaches
- Quality of life using EORTC QLQC30 and QLQCR29 questionnaire at baseline, 3 months, 6months then 6 monthly for 2 years.
- Compliance of these patients to recommended surveillance protocol

### 7.3 Statistical Analysis Plan:

Continuous variables will be summarized with descriptive statistics Mean (SD) or Median (IQR). Categorical variables will summarize with frequency and percentage.

The local regrowth rate at 2yrs will be obtained using Kaplan Meier Estimator and the local regrowth will be compared in two arms using log rank test.

Survival endpoints like OS and DFS rates at 2years will be estimated using Kaplan Meier Method and comparison for factors will be done using log rank Test. Multivariate analysis will be carried out using cox regression based on the factors identified in the univariate analysis by maintaining the variable to event ratio. The Clinical Complete Response will be reported using proportions and will be compared between two arms using Chi Square test. P value of less than 0.05 will be considered significant. All statistical analysis will be conducted using SPSS ver.25.

---

## **8. ETHICAL CONSIDERATIONS**

### **8.1 Recruitment and consent**

Before commencing the study, the protocol is subjected to review and approval by the local Research Ethics Committee in accordance with national guidelines and legislation. Declaration of Helsinki (Tokyo, Venice, Hong Kong, Somerset West, and Edinburgh amendments) or the laws and regulations of the country, whichever provides the greatest protection to the patient will be followed. Patient information sheets will be produced in all the relevant languages to eligible participants after screening in routine clinics. All patients will be informed of the aims of the study, the possible adverse events, the procedures, and possible hazards to which he/she will be exposed. It will be emphasized that the participation is voluntary and that the patient is allowed to refuse further participation in the protocol whenever he/she wants. This will not prejudice the patient's subsequent care.

### **8.2 Considerations for incapacitated subjects or vulnerable population (if applicable)**

Not applicable

### **8.3 Benefits and risks assessment.**

This is an observational study with no interventions done as a part of study. There are no direct benefits and risks to the participant. However, the data could add to the current knowledge about outcome of patients in such clinical settings and further help in optimizing the management of these patients in future based on the results of the study.

### **8.4 Compensation for injury**

As this is an observational study, there is no compensation applicable.

### **8.5 Incentives (if applicable)**

No incentive will be provided to participants.

---

## **9. ADMINISTRATIVE ASPECTS, MONITORING AND PUBLICATION**

### **9.1 Handling and storage of data and documents**

Subject identification: The name of the patient will not be asked for nor recorded at the Study Office. A sequential identification number will be automatically attributed to each patient registered in the trial. There will be an additional identifier denoting the institute. This number will identify the patient and must be included on all case record forms.

They will be informed as to the strict confidentiality of their patient data, but that their medical records may be reviewed for study purposes by authorized individuals other than their treating physician. Documented informed consent must be obtained for all patients included in the study before they are registered at the Study Office. This must be done in accordance with the national and local regulatory requirements.

### **9.2 Monitoring and Quality Assurance**

The study will be monitored by an independent data and safety monitoring committee.

### **9.3 Publication policy**

Investigator from the Institution contributing data for >10 patients will be considered for the authorship provided the investigator is willing to review and approve the final manuscript and satisfy authorship criteria and per journal requirements.

---

## 10. REFERENCES

1. Habr-Gama A, Perez RO, Nadalin W, Sabbaga J, Ribeiro U, Silva E Sousa AH, et al. Operative versus nonoperative treatment for stage 0 distal rectal cancer following chemoradiation therapy: long-term results. *Ann Surg* [Internet]. 2004 Oct [cited 2023 Sep 20];240(4):711–8. Available from: <https://pubmed.ncbi.nlm.nih.gov/15383798/>
2. Maas M, Beets-Tan RGH, Lambregts DMJ, Lammering G, Nelemans PJ, Engelen SME, et al. Wait-and-see policy for clinical complete responders after chemoradiation for rectal cancer. *J Clin Oncol* [Internet]. 2011 Dec 10 [cited 2023 Sep 20];29(35):4633–40. Available from: <https://pubmed.ncbi.nlm.nih.gov/22067400/>
3. Smith JD, Ruby JA, Goodman KA, Saltz LB, Guillem JG, Weiser MR, et al. Nonoperative management of rectal cancer with complete clinical response after neoadjuvant therapy. *Ann Surg* [Internet]. 2012 Dec [cited 2023 Sep 20];256(6):965–72. Available from: <https://pubmed.ncbi.nlm.nih.gov/23154394/>
4. Appelt AL, Pløen J, Harling H, Jensen FS, Jensen LH, Jørgensen JCR, et al. High-dose chemoradiotherapy and watchful waiting for distal rectal cancer: a prospective observational study. *Lancet Oncol* [Internet]. 2015 Aug 1 [cited 2023 Sep 20];16(8):919–27. Available from: <https://pubmed.ncbi.nlm.nih.gov/26156652/>
5. Renehan AG, Malcomson L, Emsley R, Gollins S, Maw A, Myint AS, et al. Watch-and-wait approach versus surgical resection after chemoradiotherapy for patients with rectal cancer (the OnCoRe project): a propensity-score matched cohort analysis. *Lancet Oncol* [Internet]. 2016 Feb 1 [cited 2023 Sep 20];17(2):174–83. Available from: <https://pubmed.ncbi.nlm.nih.gov/26705854/>
6. van der Valk MJM, Hilling DE, Bastiaannet E, Meershoek-Klein Kranenbarg E, Beets GL, Figueiredo NL, et al. Long-term outcomes of clinical complete responders after neoadjuvant treatment for rectal cancer in the International Watch & Wait Database (IWWD): an international multicentre registry study. *Lancet* [Internet]. 2018 Jun 23 [cited 2023 Sep 20];391(10139):2537–45. Available from: <https://pubmed.ncbi.nlm.nih.gov/29976470/>
7. Fernandez LM, São Julião GP, Figueiredo NL, Beets GL, van der Valk MJM, Bahadoer RR, et al. Conditional recurrence-free survival of clinical complete responders managed by watch and wait after neoadjuvant chemoradiotherapy for rectal cancer in the International Watch & Wait Database: a retrospective, international, multicentre registry study. *Lancet Oncol* [Internet]. 2021 Jan 1 [cited 2023 Sep 20];22(1):43–50. Available from: <https://pubmed.ncbi.nlm.nih.gov/33316218/>
8. Temmink SJD, Peeters KCMJ, Bahadoer RR, Kranenbarg EMK, Roodvoets AGH, Melenhorst J, et al. Watch and wait after neoadjuvant treatment in rectal cancer: comparison of outcomes in patients with and without a complete response at first reassessment in the International Watch & Wait Database (IWWD). *Br J Surg* [Internet]. 2023 Jun 1 [cited 2023 Sep 20];110(6):676–84. Available from: <https://pubmed.ncbi.nlm.nih.gov/36972213/>
9. Bahadoer RR, Dijkstra EA, van Etten B, Marijnen CAM, Putter H, Kranenbarg EMK, et al. Short-course radiotherapy followed by chemotherapy before total mesorectal excision (TME) versus preoperative chemoradiotherapy, TME, and optional adjuvant

- 
- chemotherapy in locally advanced rectal cancer (RAPIDO): a randomised, open-label, phase 3 trial. *Lancet Oncol* [Internet]. 2021 Jan 1 [cited 2023 Sep 20];22(1):29–42. Available from: <https://pubmed.ncbi.nlm.nih.gov/33301740/>
10. Conroy T, Bosset JF, Etienne PL, Rio E, François É, Mesgouez-Nebout N, et al. Neoadjuvant chemotherapy with FOLFIRINOX and preoperative chemoradiotherapy for patients with locally advanced rectal cancer (UNICANCER-PRODIGE 23): a multicentre, randomised, open-label, phase 3 trial. *Lancet Oncol* [Internet]. 2021 May 1 [cited 2023 Sep 20];22(5):702–15. Available from: <https://pubmed.ncbi.nlm.nih.gov/33862000/>
  11. Garcia-Aguilar J, Patil S, Gollub MJ, Kim JK, Yuval JB, Thompson HM, et al. Organ Preservation in Patients With Rectal Adenocarcinoma Treated With Total Neoadjuvant Therapy. *J Clin Oncol* [Internet]. 2022 Apr 1 [cited 2023 Sep 20];40(23). Available from: <https://pubmed.ncbi.nlm.nih.gov/35483010/>
  12. Gerard JP, Barbet N, Schiappa R, Magné N, Martel I, Mineur L, et al. Neoadjuvant chemoradiotherapy with radiation dose escalation with contact x-ray brachytherapy boost or external beam radiotherapy boost for organ preservation in early cT2-cT3 rectal adenocarcinoma (OPERA): a phase 3, randomised controlled trial. *Lancet Gastroenterol Hepatol* [Internet]. 2023 Apr 1 [cited 2023 Sep 20];8(4):356–67. Available from: <https://pubmed.ncbi.nlm.nih.gov/36801007/>