

RESEARCH

Open Access



# Immunomodulatory effects of $\beta$ -defensin 2 on macrophages induced immuno-upregulation and their antitumor function in breast cancer

Sonam Agarwal, Anita Chauhan, Khushwant Singh, Kunal Kumar, Rupinder Kaur, Marilyn Masih and Pramod Kumar Gautam\*

## Abstract

**Background:** Macrophages are mononuclear CD34<sup>+</sup> antigen-presenting cells of defense mechanism and play dual roles in tumor burden. The immunomodulatory and their antitumor function of  $\beta$ -defensin 2 is still unclear, despite the accumulating evidence of the response in infection. So, the aim of present study is to elucidate the role of  $\beta$ -defensin 2 on the level of ROS, cytokines, chemokine expression in macrophages and antitumor function in breast cancer.

**Method:** Swiss albino mice were used to harvest PEC macrophages and C127i breast cancer cells line for tumor model was used in this study. Macrophages were harvested and characterized by flow-cytometry using F4/80 and CD11c antibodies. MTT was performed to estimate cytotoxicity and dose optimization of  $\beta$ -defensin 2. Oxidative stress was analyzed by H<sub>2</sub>O<sub>2</sub> and NO estimation followed by iNOS quantified by q-PCR. Cytokines and chemokines estimation was done using q-PCR. Co-culture experiment was performed to study anti-tumor function using PI for cell cycle, Annexin -V and CFSE analysis for cell proliferation.

**Results:** PEC harvested macrophages were characterized by flow-cytometry using F4/80 and CD11c antibodies with the purity of 8% pure population of macrophages. It was found that 99% of cells viable at the maximum dose of 100 ng/ml of  $\beta$ -defensin 2 in MTT. Levels of NO and H<sub>2</sub>O<sub>2</sub> were found to be decreased in  $\beta$ -defensin 2 as compared to control. Expression of cytokines of IFN- $\gamma$ , IL-1 $\alpha$ , TNF- $\alpha$ , TGF- $\beta$  was found to be increased while IL-3 was decreased in  $\beta$ -defensin 2 group as compared to control. Levels of chemokines CXCL-1, CXCL-5 and CCL5 increased in treated macrophages while CCL24 and CXCL-15 expression decreased. Adhesion receptor (CD32) and fusion receptor (CD204) were decreased in the  $\beta$ -defensin 2 group as compared to control. Anti-tumor experiment was performed using co-culture experiment apoptosis (Annexin-V) was induced, cell cycle arrest in phase and cell proliferation of C127i cells was decreased.

**Conclusion:** This is the first report of  $\beta$ -defensin 2 modulates macrophage immunomodulatory and their antitumor function in breast cancer.  $\beta$ -defensin 2 as a new therapeutic target for immunotherapy as an adjuvant in vaccines.

**Keywords:** Macrophages,  $\beta$ -defensin 2, Cytokine profiling, Anti-tumor function, Oxidative stress

\*Correspondence: pramodgautam\_13@yahoo.com; gautam@aiims.edu

Department of Biochemistry, All India Institute of Medical Sciences, New Delhi 110029, India



© The Author(s) 2022. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.

## Background

Macrophages are mononuclear CD34<sup>+</sup> phagocytic cell known as potent antigen presenting cells of defense mechanisms which can promote specific immunity by recruitment and activation of different immune cell towards infection and malignancies [1]. Macrophages are classified according to their phenotype and cytokine release such as M1, M2, M2a, M2b, M2c, M2d and tumor associated (TAMs) [2]. Upon phagocytosis, it displays antigen via MHC class I and II bound antigen to T lymphocytes for effective response towards infection and cancer cell specific cell-mediated immunity. In different infective milieu and upon antigen encounter and anchoring with the cell via receptors, releases of a vast range of cytokines include tumor necrosis factor (TNF), IL-1 $\beta$ , IL-6, IL-8, and IL-12. Additionally, the contact of the T cell receptor with an antigen-bound MHC is necessary for the first signal (recognition), and the CD80 or CD86 on the antigen-presenting cell, as well as CD28 and CTLA-4 on the responding T cell, supply the second signal. [3]. It has been demonstrated that the interaction between B7 and CD28 produces a crucial signal for T cell activation.

Immunologically, defensins are associated with the disturbance of the numbers and function of peri- and intrafollicular inflammatory cells, and this disturbance is more abundant during the active stage of the disease. Defensins are a family of small 18–55 amino acids residues, cysteine-rich, cationic proteins expressed predominantly in innate immune cells such as neutrophils and epithelial cells [4]. Their vast research has confirmed their significance in innate immunity as a key host-protective factor against bacterial, viral, and fungal infections, as well as their broad antibacterial properties and diverse immunomodulatory actions. [5].  $\beta$ -defensin regulates the chemotactic activity for numerous innate immune cells, which also stimulate the release of secrete cytokines from other cells [6]. However, they can also capable of reduce inflammation by specifically attaching to molecular patterns associated with microbes. These patterns have ability to stimulate the expression of  $\beta$ -defensins in gingival epithelial cells, while many bacterial species are found to differ significantly from one another. Simultaneously these finding suggest that a complex paradigm of a host-defense related function in maintenance of bacterial homeostasis and pathogen response [7].

$\beta$ -defensin 2 are plausible molecules for stimulating not only innate immune but also adaptive immune responses [8]. It activates the immune system by modulation of signaling pathways and inflammatory response [9]. Recent studies have demonstrated that  $\beta$ -defensin 2 can modulate immune responses in various diseases [10]. Apart from their antimicrobial activities,  $\beta$ -defensin 2

is also involved in a variety of other activities, including pathogens killing, cell activation, proliferation, regulation of cytokine/chemokine production, migration, differentiation, angiogenesis, and wound healing. Moreover, they also participate in tumorigenesis, due to their proliferative or suppressive properties relative to different cells [9]. Mei et al. reported that  $\beta$ -defensin 2 may act as a positive regulator, promoting anti-tumor responses of T and NK cell in vivo, therefore it may be useful in immunotherapy [11]. NK cells and CD8<sup>+</sup> T cells attack against malignancies, while other cells accumulate into the tumor. This imbalance in the inflammatory cells, leading to a collapse of the immune surveillance and inflammatory cells, is not only restricted to the site but promotes tumor. It is also noticed to occur in the spleen, lymph nodes, and the peripheral blood cells affected in tumor patients. A gap in the knowledge of immunomodulatory and tumoricidal function of macrophages with a cross-talk with  $\beta$ -defensin 2.

However, few reports of  $\beta$ -defensin 2 in immunomodulation have been extensively studied and their involvement in other processes such as antitumor activity against cancer remaining unknown. So, the aim of this study was to evaluate the immunomodulatory function of  $\beta$ -defensin 2 on macrophages and to elaborate the anti-tumor activity on breast cancer. We analyzed the impact of  $\beta$ -defensin 2 on the level of oxidative stress, expression of cytokines and chemokines gene of macrophages. Consequently, the antitumor effect of  $\beta$ -defensin 2 on breast cancer during the active state of the macrophages was also determined in the current study.



## Results

### Isolation and characterization of macrophages (Fig. 1)

Normal macrophages were harvested from Swiss albino mice as peritoneal exudates cell by peritoneal lavage with chilled PBS and incubated for 2 h in plastic petri dishes (Tarson, India). After 2 h, adherent cells were collected and stored in media, washed thrice in serum free media at 1500 rpm for 10 min by centrifugation. Flow cytometry was performed for quantification and characterization of harvested cells (Fig. 1E). It was found that more than 96% cells of F4/80, CD11c positive are labeled with PE and further utilized for ROS estimation and gene expression analysis.

### Immunocytochemistry (ICC) (Fig. 2)

ICC was done to localize the CD14 receptor, which is a marker of macrophages. The  $\beta$  defensins-2 (BD-2) internalization/ phagocytosis by macrophages ICC assay were done. Purified macrophage cells were incubated with a 100 ng/ml dose of BD-2 for 24 h. Slides were prepared as per protocol followed by monoclonal antibodies. Figure 2



shows that internalization of BD-2 in macrophages uniform punctate patch in the cytoplasmic extensions of the cells, while untreated cells there are no any punctate pattern found in the cytoplasmic extension of the cells.

### MTT assay (Fig. 3)

Figure 3 shows that the effect of BD 2 on the viability of macrophages, cells were treated with 10–100 ng/ml BD-2 for 24 and 48 h and subjected to MTT assay. Figure 3 shows that no change was found on cell viability of macrophages when treated with BD-2 at various concentrations after 24 and 48 h.


### ROS production on BD-2 treated macrophages (Fig. 4)

DCFHDA is a membrane-permeable dye that enters into the cells and is converted into DCF, a fluorescent form when oxidized by ROS present in the cells. The intensity of fluorescence is proportional to the level of ROS present in the cells. Result shows that normally macrophages release ROS to remove the foreign particles present in


their surroundings. Therefore, we determined the level of ROS generation with the exposure of BD 2 by DCFHDA assay. Our results revealed that BD 2 could not induce the macrophage to generate ROS at given concentrations in the indicated time points as shown in Fig. 4A–D. Cells were treated with 100 ng/ml of BD-2 for 24 h. Results show that treatment of BD-2 does not induce the release of expression of ROS on normal phenotype of macrophages as compared to LPS treated and medium-only while a positive stimulator H<sub>2</sub>O<sub>2</sub> stimulates the ROS releases at a significant level. NAC used a negative control for the release of ROS which shows decreased expression of ROS as compared to medium only (Control).

### Nitric oxide measurement (Fig. 5A–C)


The Reactive oxygen species contains NO<sup>−</sup> and H<sub>2</sub>O<sub>2</sub> which was estimated by Griess method and TISO<sub>4</sub> method. The amount of NO production in the supernatant of macrophages after 24 h treatment with  $\beta$ -defensin 2 and LPS was determined with the Griess reagent.



**Fig. 3** Effects of BD 2 on cell viability of mice isolated peritoneal macrophage cells after 24 and 48 h of treatment. Results are expressed as mean ± SE



**Fig. 4** Effects of BD 2 on ROS induction in mice isolated peritoneal macrophage cells after 1 and 2 h of treatment. Results are expressed as mean ± SE. BD-2 (100 ng). Statistical analysis: \*\*\**P* < 0.001



**Fig. 5 A–C** Effects of BD 2 on the level of H<sub>2</sub>O<sub>2</sub>, NO and iNOS in mice isolated peritoneal macrophage cells after 24 h treatment. Results are expressed as mean ± SE. Statistical analysis: \**P* < 0.05, \*\*\**P* < 0.001

During this period, the NO level decreased remarkably in the  $\beta$ -defensin 2 group as compared to the control group (Fig. 5A). It was observed that in other experiments were conducted to examining the effect of  $\beta$ -defensin 2 on the production of  $H_2O_2$ . Production of  $H_2O_2$  was found to be decreased in  $\beta$ -defensin 2 treated group as compared to control (Fig. 5B). mRNA level of iNOS was downregulated in  $\beta$ -defensin 2 treated group as compared to control group (Fig. 5C).


#### RT-PCR for selected gene expression on macrophages (Figs. 6 and 7)

The effect of  $\beta$ -defensin 2 was studied on cytokine and chemokine genes in M1 macrophage with the help of real time PCR. Cytokine and chemokine expression profiles were examined. Macrophages ( $1 \times 10^6$ ) were purified and characterized in 12 well plates with complete media in triplicate. Cells were treated with 100 ng/ml of BD-2 for 24 h. We have found that BD-2 treated macrophages significantly increased the expression of cytokines IFN- $\gamma$ , IL-1 $\alpha$ , IL-6, TNF- $\alpha$  and TGF- $\beta$  while diminished the level of IL-3, CD32 and CD204 as compared to control group (7A and B). In chemokines, we have found increased the


level of CXCL-1, CXCL-5 and CCL-5 while decreased the level of CXCL-15 and CCL-24 as compared to control mice (Fig. 6A–E).

#### Apoptotic assay (Fig. 8)

In comparison to macrophage co-cultured and unstimulated c127i, co-culture of c127i with macrophage decreased apoptosis in c127i cells from 5.37 to 4.76%, and  $\beta$ -defensin 2 activated macrophages further decreased apoptosis to 2.66% from 4.76 to 5.37%, respectively. In contrast to unstimulated c127i, there was an increase in late apoptotic cells percent 7.26% from 1.42% in macrophage co-cultured c127i. Macrophage co-cultured c127i when compared to  $\beta$ -defensin 2 activated macrophage co-cultured c127i and to unstimulated c127i, increased percentage of late apoptotic cells 7.26% was observed. At the very same time in c127i cells co-cultured with  $\beta$ -defensin 2, we noticed an increase in the number of dead cells. The dead cell population was enhanced by  $\beta$ -defensin 2 activated macrophages to 10%, up from 5.4% in c127i co-cultured with macrophages and 2.8% in untreated cells. Macrophages increase late apoptotic cell counts but not direct apoptosis in c127i




**Fig. 6** A–E Effects of BD 2 on the expression of chemokines in mice isolated peritoneal macrophage cells after 24 h treatment. Results are expressed as mean  $\pm$  SE. Statistical analysis: \* $P < 0.05$ , \*\* $P < 0.01$



**Fig. 7 A, B** Effects of BD 2 on the expression of cytokines in mice isolated peritoneal macrophage cells after 24 h treatment. Results are expressed as mean  $\pm$  SE. \* $P$  < 0.05, \*\*\* $P$  < 0.001, \*\* $P$  < 0.01






cells, according to the apoptosis analyses. Furthermore,  $\beta$ -defensin 2 stimulated macrophages triggers necrosis or cell death in late apoptotic c127i cells.

#### Cell division analysis (Fig. 9A-C)

The number of undivided cells in  $\beta$ -defensin-activated-macrophage co-culture c127i cells was 512, compared to 5906 and 1367 cells in unstimulated and macrophage co-culture c127i cells, respectively. In comparison to the unstimulated and macrophage co-culture c127i cells,  $\beta$ -defensin-activated-macrophage co-culture c127i cells undergo more divisions and cells of the 4th and 5th generations were missing. Furthermore, it was discovered that  $\beta$ -defensin-activated-macrophage co-culture c127i cells had a higher percentage of dead cells (50%) than unstimulated (1.9%) and macrophage co-culture c127i cells (4.4%), respectively. The present findings of CFSE cell proliferation and apoptosis indicate that  $\beta$ -defensin-activated-macrophage cause cell death in c127i cells via necrosis or other alternative mechanism other than direct apoptosis.

#### Discussion

The immune system is an interactive network of immune cells, lymphoid organs, humoral factors, fibroblasts, tumor vasculature and extracellular matrix make up the stromal environment. In many ways, it is like an army equipped with different weapons to provide protection against infection and malignancies. It has been found that systemic regulation of immune responses is highly impaired when infection greatly increases. During tumor progression, infected cells release various immunosuppressive molecules that suppress immune response directly or activate the host suppressor mechanism in many ways. Tumor actively recruits cells into the tumor microenvironment, including bone marrow-derived mesenchymal stem cells (MSCs), macrophages, NK cell, T cells and GD cells whose function is suppressed in tumor milieu and promotes tumor growth [12, 13]. Several defense proteins expressed by cells and tissue play a cytoprotective role and control infections by recruiting immune cells for pathogen clearance.  $\beta$ -defensins are a family of epithelial cell



derived antimicrobial peptides (AMPs) that protect mucosal membranes from microbial challenges [14]. In addition to their antimicrobial activities, they possess other functions; e.g., cell activation, proliferation, regulation of cytokine/chemokine production, migration, differentiation, angiogenesis, and wound healing processes. It has also become apparent that defending levels change with the development of neoplasia. However, inconsistent observations published by various laboratories make it difficult to reach a consensus as to the direction of the dysregulation and role the hBDs may play in various cancers.

We discovered that consensus-driven findings indicate  $\beta$ -defensin 2 not only actively participates in infection, it activates macrophages to perform immunomodulatory and anti-tumor functions too. The effect of  $\beta$ -defensin 2 on macrophages at the dose at maximum 1 ng/ml shows 99% cell viability. It shows that cytoprotective molecules expressed by tissue and immune cell are non-toxic for self cells. Several other published reports support this hypothesis that hsp increases immunomodulatory function and increasing anti-tumor function of tumor associated macrophages by enhancing ROS, cell surface receptors expression such as B7 family receptors,



CD54, Cd47, SIRP  $\alpha$ , CTLA-4 etc. It also increases the antitumor by increasing il-2, il-4, il-6, tnf-alpha. In some amount ROS production facilities signaling pathway such as NF-kappa b which actively involved in cytokine production. Some data shows that during tumor progression ROS release an increase many fold which facilitates tumor growth.

The most commonly encountered free radicals in cell are hydroxyl radical, superoxide radical, nitric oxide radical, and lipid peroxy radical. NO and H<sub>2</sub>O<sub>2</sub> are produced in high quantities from macrophages activated by pro-inflammatory cytokines which leads to cell lesion [15]. NO is synthesized by iNOS, involved in suppression of tumor growth [16]. In accordance with these reports, in the current study, NO and H<sub>2</sub>O<sub>2</sub>, which is expressed by macrophages was found to be reduced when treated with  $\beta$ -defensin 2 as compared to control (Fig. 5), indicating the cytoprotective effects of BD-2.

TNF- $\alpha$  is a pro-inflammatory cytokine, plays an essential role in inflammation, apoptosis, lipid metabolism and immune system development [17]. TGF- $\beta$  is an anti-inflammatory cytokine, higher levels of anti inflammatory cytokine released by activated macrophages which promote tumor suppression leads to the antitumor function. Our results revealed that enhanced the expression of pro-inflammatory cytokines IFN- $\gamma$ , IL-1 $\alpha$ , IL-6, TNF- $\alpha$ , and TGF- $\beta$  while reduced the expression of IL-3 and IL-10 in BD-2 treated group as compared to control (Fig. 7). A high level of TGF- $\beta$  has been associated with the regression of growth and survival of tumor which leads to the stimulate apoptosis.

There are several studies reported that higher level of CCL-5 was found in ischemia–reperfusion injuries of brain, lung, heart, kidney, intestine, and skeletal muscle [18–23]. We have shown here that the level of chemokines CXCL-1, CXCL-5 and CCL5 was found higher while CCL24 and CXCL-15 decreased in BD-2 treated group as compared to control (Fig. 6). Higher expression of chemokines was correlated with inflammatory response which leads to the tumor suppression suggesting that BD-2 may act as antitumor function.

### Concluding remarks

Our present study provides the first evidence that BD-2 plays essential role as an antitumor function. M1 macrophages secrete pro-inflammatory cytokines (including TNF- $\alpha$ , IL-6, TGF- $\beta$ , and IFN- $\gamma$ ) and chemokines to induce apoptosis and reduce cell proliferation which leads to the antitumor function. The main findings of the present study are that BD-2 abridged the level of ROS and NO which is associated with tumor regression. In future research, we hope that we can study the mechanism and learn more about the immunomodulatory function of BD-2.

## Method

### Reagents

RPMI-1640 media and FBS were purchased from Gibco, Waltham, Massachusetts, USA. Antibiotics, LPS, DCF-HDA, TISO<sub>4</sub> and H<sub>2</sub>O<sub>2</sub> were procured from Sigma-Aldrich, St. Louis, Missouri, USA. MTT, DAPI and DABCO were purchased from SRL, India. Anti-mouse CD14-PE (Thermo-Scientific, Waltham, Massachusetts, USA),  $\beta$ -defensin 2 (Prospec, Hamada, Israel) and RNAiso plus (Takara, Japan) were purchased. cDNA synthesis kit and SYBR green were obtained from BIO-RAD, USA. AnnexinV-FITC/PI and CFSE were purchased from Biolegend, San Diego, USA. Anti-mouse CD14 conjugated with PE was purchased from Thermofisher, USA.

### Animals and ethical clearance

Inbred populations of Swiss albino strain of mice of either sex and at 8–12 weeks of age were used. Mice obtained from the Animal House, AIIMS, New Delhi, and housed in a pathogen-free specialized small animal facility with a 12 h dark–light cycle. All the animals were treated with utmost human care and had free access to food and water. Mice were euthanized by cervical dislocation, a method authorized by Institutional Animal Ethical Committee (file no- 100/IAEC-1/2018), AIIMS, New Delhi for sacrificing experimental animals, and observed until all muscle activity and breathing has ceased for at least 120 s. No mice died before meeting the endpoints described.

### Isolation and characterization of macrophages from mice

Macrophages were harvested from Swiss albino mice as previously described [24]. Briefly, mice were killed by cervical dislocation and macrophages were harvested by peritoneal lavage as peritoneal exudate cells (PECs) using chilled serum free culture medium (RPMI 1640). PECs were harvested by adherent purification in plastic petri dishes (Tarson, India) at room temperature for 2 h. Adherent cells were collected and seeded in a flat-bottom culture flask (Tarson, India) at a cell density of  $1 \times 10^6$  in the culture medium with or without LPS and BD-2 and incubated for time periods of 24 h in a CO<sub>2</sub> incubator for treatment.

### Flow cytometry assay

Macrophages were harvested from Swiss albino mice as described above [24] and at a cell density of  $1 \times 10^6$  in the culture medium with or without LPS and BD-2 for 24 h in RPMI 1640 containing 10% FBS at 37 °C in 5% CO<sub>2</sub> in humidified CO<sub>2</sub> incubator. Macrophages were suspended in RPMI-1640 with 10% FBS, 0.1% NaN<sub>3</sub> and incubated with anti-mouse CD14, an antibody conjugated with PE. Isotype is used as a control conjugated with FITC as per manufacturer's instructions. After washing, cells were

suspended in 0.1% PBS containing 0.1% NaN<sub>3</sub> and then analyzed with a flow cytometry (BD Biosciences, Mountain View, CA, USA).

### ICC

Macrophages were harvested from Swiss albino mice as previously described [25]. The macrophages were characterized by their cell surface marker CD14. After 24 h, the harvested cells were spread over the pre-coated air-dried poly-lysine coated glass slide, and the slide with cells was kept at room temperature to air dry and fix with 4% PFA for 2 h. The slide was washed 3 times with PBS and incubated with 0.1% Triton X-100 for 15 min, followed by washing with PBS thrice followed by staining with DAPI for 5 min and cells were mounted in anti-fade dye Dabco mounting media. Cell morphology was photographed under a fluorescent microscope and Olympus BX51 microscope in grayscale at 100X magnification.

### Peptide cellular uptake assay

To visualize the localization of BD 2 in macrophage cells, an ICC assay was done [25]. For this assay, peritoneal macrophage cells were isolated from Swiss albino mice. A pre-sterilized cover slip was placed in the culture plate and macrophage cells were cultured on it. Then cells were treated with BD 2 with specified concentrations. After 24 h of incubation, the cover slip that has adhered macrophage cells were isolated from the culture dish and washed with (1X PBS) and treated with 4% paraformaldehyde for 2 h. After three consecutive wash with (1X PBS) cells containing macrophage cells were air-dried for 30 min and treated with 0.1% Triton X for 10 min. After three wash with (1X PBS) cells were treated with BD 2 antibody and incubated overnight. After three wash with (1X PBS) treated with specific secondary antibody and localization of BD 2 was detected under microscopy.

### MTT assay for dose optimization

MTT assay was performed to determine relative cell viability at 24 and 48 h after treatment of BD-2. Macrophage cells were seeded in 96-well plates at  $1 \times 10^5$  cells/well for 24 h. After that, cells were treated with BD-2 (10–100 ng) and incubated for 24 and 48 h. The cells were washed with 1X PBS and treated with MTT dye (5 mg/ml) for 3 h. After aspirating MTT dye from the plates, cells were treated with 100  $\mu$ l/well DMSO and optical density (OD) at 570 nm was taken with a multi-plate reader within 15 min and required concentration for a 50% inhibition of viability (IC<sub>50</sub>) was calculated. Each experiment was performed in triplicates [25].

### ROS measurement

For this assay, macrophages (20000 cells/well) were seeded in a 96-well plate and incubated for 24 h. A solution of 20  $\mu$ M DCFHDA in 100  $\mu$ l serum-free media was added in each well and incubated for 1 h. After the removal of the DCFHDA solution, each well was washed twice with serum-free media. However, for the negative control, a pre-treatment of 5 mM NAC was done to block the ROS in the specified wells and later treated with BD 2 and H<sub>2</sub>O<sub>2</sub> (2 mM). Finally, the fluorescence at excitation/emission (485/538 nm) was detected by multiple readers at 1 and 2 h time points [24].

### Nitric oxide (NO) measurement

The level of NO produced was quantified by determining the accumulation of nitrite using the Griess reagent. The optical density of the samples was measured at 540 nm with an ELISA reader. Briefly, cell supernatant was centrifuged to remove debris and clear supernatant was mixed with an equal volume of Griess reagent. The mixture was incubated in the dark for 40 min. The absorbance was measured at 540 nm using a microplate reader. The concentration of nitrite was determined from the sodium nitrite standard curve [24].

### H<sub>2</sub>O<sub>2</sub> measurement

The production of H<sub>2</sub>O<sub>2</sub> was quantified by TISO<sub>4</sub>. Add 50  $\mu$ l of titanium sulfate reagent per 100  $\mu$ l of sample. If the sample contains peroxide, it will react with the titanium ions in solution to form perititanic acid, giving the solution a yellow color. The mixture with the reagent should be stable for several hours. Measure the absorbance at 407 nm using the ELISA plate reader [24].

### Anti-tumor assay

AnnexinV-FITC/PI and CFSE labeled c127i breast cancer cells were co-cultured with macrophages in a transwell chamber with the medium containing 100 ng/ml BD-2. Subsequently, flow cytometry was employed to reveal the percentage of apoptosis, cell division and cell proliferation.

Carboxy-fluorescein succinimidyl ester (CFSE) is cell permeable and covalently couples via its succinimidyl group, to intracellular molecules, -Lysine residues and other amine sources. Due to this covalent coupling reaction, fluorescent CFSE can be retained within cells for extremely long periods. Also, due to this stable linkage, once incorporated within cells, the dye is not transferred to adjacent cells. 50,000 c127i cells were incubated in 300  $\mu$ l of 5  $\mu$ M CFSE-FITC working solution in PBS for 15 min

at 37 °C, protected from light. Excess CFSE was quenched by adding 2 ml complete DMEM culture media containing 10% FBS and incubating for 10 min at room temperature. The cell pellets were resuspended in pre-warmed PBS and washed twice with PBS and centrifugation at 1200 rpm for 5 min at room temperature. C127i cells labeled with CFSE-FITC were resuspended in pre-warmed complete DMEM culture media and seeded in culture dish.

### Gene expression analysis

Total RNA was extracted from the MΦ harvested from control and experimental mice with RNAiso Plus reagent as instructed by the manufacturer (Takara, Japan). High-quality RNA (as estimated by absorbance ratio A260/280P1.8) from different groups were resolved on 1% agarose gel and stained with ethidium bromide to check the integrity of 18S and 28S rRNA using a UV transilluminator.

The total RNAs were used for cDNA synthesis using a kit (BIO-RAD, US), where total RNA (1 µg/20 µl reaction) was implied and further processes were followed as directed by the manufacturer. For real-time PCR, iTaq Universal SYBR Green Supermix was used (BIO-RAD, US). Reaction mix preparation and thermal cycling protocol was followed as directed by manufacturer guidelines. The primers used for the reaction are listed in Tables 1 and 2.

**Table 1** Cytokine sequences used in this study

Genes	Sequences	Tm (°C)
IFN- gamma	Forward TTC TTC AGC AAC AGC AAG GC	56.2
	Reverse ACT CCT TTT CCG CTT CCT GA	56.2
IL-1α	Forward CCG TGT TGC TGA AGG AGT TG	56.4
	Reverse GTG CAC CCG ACT TTG TTC TT	55.9
IL- 6	Forward ACA AAG CCA GAG TCC TTC AGA	56.0
	Reverse TGG TCC TTA GCC ACT CCT TC	56.3
CD 32	Forward GAC ACA GCA CCA GTC CAA GA	57.2
	Reverse CAG TTT TGG CAG CTT CTT CC	54.5
TNF-α	Forward ACC CTC ACA CTC ACA AAC CA	56.4
	Reverse GGC AGA GAG GAG GTT GAC TT	56.6
CD 204	Forward GCC CCA GAG CAA AAA TAT GA	53.1
	Reverse TGG CTC AAG CTG TTG TCA TC	55.5
IL-3	Forward GCCTGCCTACATCTGCGAAT	54
	Reverse GGTTAGGAGAGACGGAGCCA	56
IL-10	Forward GCTGCCTGCTCTTACTGACT	54
	Reverse GGGGCATCACTTCTACCAGG	56
TGF-β	Forward GCTGAGCACCTTTTGTCTCC	54
	Reverse GATGGCATTTCGGAGGGGA	54
iNOS	Forward CTATGGCCGCTTTGATGTGC	59.97
	Reverse CAACCTTGGTGTGAAGGCG	59.97
GAPDH	Forward CGT CCC GTA GAC AAA ATG GT	54.9
	Reverse TTG ATG GCA ACA ATC TCC AC	53.3

**Table 2** Chemokine sequences used in this study

Genes	Sequences	Tm (°C)
CXCL-1	Forward GGTGAGGACATGTGTGGGAG	56
	Reverse CGAGACCAGGAGAAACAGGG	56
CXCL-5	Forward CTGCTGCTTTCCTACCTCT	54
	Reverse TCTTCTCTGGGTGGCACAC	54
CXCL-15	Forward TTGGAGCCAAGGCAAGAACA	52
	Reverse AATGGAGAGGCATCCGGTTC	54
CCL-24	Forward GGTCCCTGTCATGCTTCTGG	56
	Reverse GAGTAGCAGCAGGTGAGTGG	56
CCL-5	Forward TGCTCCAATCTTGCACTCGT	52
	Reverse GCAAGCAATGACAGGGAAGC	54
GAPDH	Forward CGT CCC GTA GAC AAA ATG GT	54.9
	Reverse TTG ATG GCA ACA ATC TCC AC	53.3

### Statistical analysis

Data were expressed as the mean ± SE. *P* value < 0.05 was considered significant. All graphs were generated using the Graph Pad Prism software. The multigroup comparisons of means were carried out by one-way analysis of variance (ANOVA) test. Student *t*-test was used for comparison between groups.

### Abbreviations

TAMs: Tumor associated; TNF: Tumor necrosis factor; PECs: Peritoneal exudate cells; NO: Nitric oxide; PECs: Peritoneal exudate cells; ROS: Reaction oxygen species; CFSE: Carboxy-fluorescein succinimidyl ester.

### Acknowledgements

Authors are thankful to the Department of Biochemistry, AIIMS, CIF facility for tissue culture, fluorescent microscopy, flow cytometry and real time PCR for experiments. Dr. Gautam expresses his appreciation to SERB DST, New Delhi and IMRG, AIIMS, New Delhi for funding.

### Author contributions

Dr. S.A., A.C., K.S. and Dr. K.K. performed wet lab experiments. Dr. P.K.G. wrote the manuscript. R.K. and M.M. arranged the references of the manuscript. Dr. P.K.G. was the supervisor and designed the experimental plan. He analyzed all the data and corrected the manuscript. All authors read and approved the final manuscript.

### Funding

This research was supported by the following grants: AIIMS, New Delhi, DST and CSIR.

### Availability of data and materials

All data generated or analyzed during the current study available from the corresponding author on reasonable request.

### Declarations

#### Ethics approval and consent to participate

All animal experimental protocols were approved by Institutional Animal Ethics Committee (IAEC) of AIIMS, New Delhi. All methods were carried out in accordance with relevant guidelines and regulations (file no- 100/IAEC-1/2018).

Human participants, human data or human tissue: Not applicable

**Consent for publication**

Not applicable

**Competing interests**

The authors declare that they have no competing interests.

Received: 16 April 2022 Accepted: 27 September 2022

Published online: 02 November 2022

**References**

- Shapouri-Moghaddam A, Mohammadian S, Vazini H, Taghadosi M, Esmaili SA, Mardani F, Seifi B, Mohammadi A, Afshari JT, Sahebkar A. Macrophage plasticity, polarization, and function in health and disease. *J Cell Physiol*. 2018;233(9):6425–40.
- Zhou D, Huang C, Lin Z, Zhan S, Kong L, Fang C, Li J. Macrophage polarization and function with emphasis on the evolving roles of coordinated regulation of cellular signaling pathways. *Cell Signal*. 2014;26(2):192–7.
- Mosser DM, Edwards JP. Exploring the full spectrum of macrophage activation. *Nat Rev Immunol*. 2008;8(12):958–69.
- Ganz T. Defensins: antimicrobial peptides of vertebrates. *C R Biol*. 2004;327:539–49.
- Raschig J, Mailänder-Sánchez D, Berscheid A, Berger J, Strömstedt AA, Courth LF, Malek NP, Brötz-Oesterhelt H, Wehkamp J. Ubiquitously expressed Human Beta Defensin 1 (hBD1) forms bacteria-entrapping nets in a redox-dependent mode of action. *PLoS Pathog*. 2017;13(3):e1006261.
- Swanson K, Gorodetsky S, Good L, Davis S, Musgrave D, Stelwagen K, et al. Expression of a beta-defensin mRNA, lingual antimicrobial peptide, in bovine mammary epithelial tissue is induced by mastitis. *Infect Immun*. 2004;72:7311–4.
- Agier J, Brzezińska-Błaszczak E, Różalska S, Wiktorska M, Wawrocki S, Żelechowska P.  $\beta$ -defensin strengthens antimicrobial peritoneal mast cell response. *J Immunol Res*. 2020;2020:5230172.
- Kim J, Yang YL, Jang SH, et al. Human  $\beta$ -defensin 2 plays a regulatory role in innate antiviral immunity and is capable of potentiating the induction of antigen-specific immunity. *Viral J*. 2018;15:124.
- Xu D, Lu W. Defensins: a double-edged sword in host immunity. *Front Immunol*. 2020;11:764.
- Deptuła J, Tokarz-Deptuła B, Deptuła W. Defensins in humans and animals. *Postepy Hig Med Dosw*. 2019;73:152–8.
- Mei H-f, Jin X-b, Zhu J-y, Zeng A-h, Wu Q, Lu X-m, et al.  $\beta$ -defensin 2 as an adjuvant promotes anti-melanoma immune responses and inhibits the growth of implanted murine melanoma *In Vivo*. *PLoS ONE*. 2012;7(2):e31328.
- Ahanban-Esfahlan R, Seidi K, Zarghami N. Tumor vascular infarction: prospects and challenges. *Int J Hematol*. 2017;105:244–56.
- Jahanban-Esfahlan R, Seidi K, Banimohamad-Shotorbani B, Jahanban-Esfahlan A, Yousefi B. Combination of nanotechnology with vascular targeting agents for effective cancer therapy. *J Cell Physiol*. 2017;233:2982–92.
- Sechet E, Telford E, Bonamy C, Sansonetti PJ, Sperandio B. Natural molecules induce and synergize to boost expression of the human antimicrobial peptide  $\beta$ -defensin-3. *PNAS*. 2018;115(42):E9869–78.
- Castaneda OA, Lee SC, Ho CT, Huang TC. Macrophages in oxidative stress and models to evaluate the antioxidant function of dietary natural compounds. *JFDA*. 2017;25:111–8.
- McNeill E, Crabtree MJ, Sahgal N, Patel J, Chuaiphichai S, Iqbal AJ, Hale AB, Greaves DR, Channon KM. Regulation of iNOS function and cellular redox state by macrophage Gch1 reveals specific requirements for tetrahydrobiopterin in NRF2 activation. *Free Radical Biol Med*. 2015;79:206–16.
- Sethi JK, Hotamisligil GS. Metabolic messengers: tumour necrosis factor. *Nat Metab*. 2021;3:1302–12.
- Braunersreuther V, et al. Chemokine CCL5/RANTES inhibition reduces myocardial reperfusion injury in atherosclerotic mice. *J Mol Cell Cardiol*. 2010;48:789–98.
- Burne-Taney MJ, Yokota N, Rabb H. Persistent renal and extrarenal immune changes after severe ischemic injury. *Kidney Int*. 2005;67:1002–9.
- Yilmaz G, Granger DN. Leukocyte recruitment and ischemic brain injury. *Neuromolecular Med*. 2010;12:193–204.
- Sharma AK, Linden J, Kron IL, Laubach VE. Protection from pulmonary ischemia-reperfusion injury by adenosine A2A receptor activation. *Respir Res*. 2009;10:58.
- Watson M, et al. Intestinal ischemia/reperfusion injury triggers activation of innate toll-like receptor 4 and adaptive chemokine programs. *Transpl Proc*. 2008;40:3339–41.
- Gute DC, Ishida T, Yarimizu K, Korthuis RJ. Inflammatory responses to ischemia and reperfusion in skeletal muscle. *Mol Cell Biochem*. 1998;179:169–87.
- Zhang X, Goncalves R, Mosser DM. The isolation and characterization of murine macrophages. *Curr Protoc Immunol*. 2008;83:14.1.1–14.1.14.
- Gautam PK, Kumar S, Tomar MS, Singh RK, Acharya A, Kumar S, Ram B. Selenium nanoparticles induce suppressed function of tumor associated macrophages and inhibit Dalton's lymphoma proliferation. *Biochem Biophys Rep*. 2017;12:172–84.

**Publisher's Note**

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.


**Ready to submit your research? Choose BMC and benefit from:**

- fast, convenient online submission
- thorough peer review by experienced researchers in your field
- rapid publication on acceptance
- support for research data, including large and complex data types
- gold Open Access which fosters wider collaboration and increased citations
- maximum visibility for your research: over 100M website views per year

**At BMC, research is always in progress.**

Learn more [biomedcentral.com/submissions](https://biomedcentral.com/submissions)





# Immunomodulatory effects of $\beta$ -defensin 2 on tumor associated macrophages induced antitumor function in breast cancer

Sonam Agarwal, Anita Chauhan, Pramod Kumar Gautam \*

Department of Biochemistry, AIIMS, New Delhi, 110029, India

## ARTICLE INFO

**Keywords:**  
 $\beta$ -defensin 2  
 TAMs  
 Cytokines  
 Chemokines  
 ROS  
 Breast cancer

## ABSTRACT

**Background:** Tumor-associated macrophages (TAMs) express cytokines and chemokines that can suppress anti-tumor immunity and promote tumor progression. The immunomodulatory and antitumor function of  $\beta$ -defensin 2 is still unclear, despite the evidence of infection response. We previously reported that  $\beta$ -defensin 2 modulates immunomodulatory and their antitumor function of macrophages in breast cancer. We investigate the association between  $\beta$ -defensin 2 and TAMs and determined the role in tumor-promoting attributes of TAMs reversal of phenotype in tumor regression.

**Methods:** Swiss albino mice and C127i breast cancer cell line was used in this study. C127i conditioned media was prepared and generated macrophage-derived TAMs to study antitumor function. Flow cytometry was performed for phenotype identification of macrophages and TAMs. MTT assay was performed to estimate cytotoxicity and dose optimization of  $\beta$ -defensin 2. Oxidative stress was analyzed by  $H_2O_2$  and NO estimation, and qPCR was performed for iNOS, cytokines and chemokines expression.

**Results:** PEC harvested macrophages were characterized by flow-cytometry using F4/80, CD11c antibodies with 98% pure population of macrophages and cultured in C127i conditioned media for 7 days. TAMs markers were estimated, and it was found that 98% expression of F4/80, CD-206, and CD-115 expression compared to macrophages. Purified 100 ng/ml of  $\beta$ -defensin 2 was used to stimulate the TAMs population was viable, which was confirmed by cell viability assay. ROS levels decreased in TAMs treated with  $\beta$ -defensin 2 compared to control group. Interleukins (ILs)-6, 10, and 3, tumor necrosis factor (TNF)- $\alpha$ , and transforming growth factor (TGF)- $\beta$  and chemokine ligand (CXCL)-1, 5 and 15, chemokine ligand (CCL)-24 and 5 decreased drastically compared to control.

**Conclusion:** This is the first report of  $\beta$ -defensin 2 on TAMs to elucidate the immunomodulatory and anti-tumor function. It was found that the cytokines, chemokines, and reactive oxygen species (ROS) expression plably changed which facilitates tumor regression.  $\beta$ -defensin 2 must be targets as adjuvant for future cancer immunotherapeutic agent.

## 1. Background

Tumor microenvironment (TME) consists of immunosuppressive factors and tumor-promoting cells. It continuously recruits immune cells such as macrophages, NK cells, B cells, T cells, and MSCs into the tumor microenvironment. These cells may aid cancer progression by down-regulating their normal anti-tumor function. Tumor-associated macrophages (TAMs) are largest population of infiltrating inflammatory cells in malignant tumors which promote the growth [1]. TAMs facilitate the immunosuppressive TME by releasing cytokines, chemokines, and growth factors and triggering the inhibitory immune checkpoint

proteins released in T cells. These functions directly or indirectly promote tumor growth by increasing tumor cell survival, proliferation, invasiveness, and motility [2]. TAMs have been demonstrated to have comparable roles in recent investigations, as TAMs tend to be pro-tumoral, reacting to Interleukins (IL)-4, 10, 13, transforming growth factor (TGF)- $\beta$ , and stimulating tissue regeneration. TAMs are often located in the invasive front of cancer cells, where most cancer cells undergo epithelial-mesenchymal transition (EMT) [3]. TAMs secrete TGF- $\beta$ , as described above, which plays an important role in promoting EMT [4,5]. M2-polarized TAMs facilitate EMT in PC via the TLR4/IL10 signaling pathway [6]. Many studies report the cross-talk of M2 and

\* Corresponding author. Dept. of Biochemistry, AIIMS, New Delhi, 110029, India.

E-mail addresses: [sonam.2mits@gmail.com](mailto:sonam.2mits@gmail.com) (S. Agarwal), [gautam@aiims.edu](mailto:gautam@aiims.edu) (P.K. Gautam).

<https://doi.org/10.1016/j.adcanc.2023.100102>

Received 21 January 2023; Received in revised form 26 March 2023; Accepted 4 May 2023

Available online 15 May 2023

2667-3940/© 2023 Published by Elsevier B.V. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).



MSCs in breast cancer, where M2 and secretory factors have a significant impact by promoting the EMT profiling of TA-MSCs, which helps in the breast cancer progression.

Immunologically, defensins are associated with the disturbance of numbers and function of peri-and intrafollicular inflammatory cells, which is more abundant during the active stage of the disease [7]. Defensins are cysteine-rich, cationic proteins of small amino acids (18–55), which are expressed predominantly in innate immune cells, for example, epithelial cells and neutrophils [8]. Their multifaceted immunomodulatory functions and antimicrobial activities have been extensively studied, establishing their role as a core host-protective component against bacterial, fungal, and viral infections [9].  $\beta$ -defensins are expressed in humans, while  $\alpha$  and  $\Theta$ -defensins have different functions and are not well studied.  $\beta$ -defensin exhibits potent chemotactic activity for various innate immune cells and stimulates other cells to secrete cytokines [10]. However, they can also inhibit the inflammatory response by the specific binding of microbe-associated molecular patterns. These patterns can also induce the expression of  $\beta$ -defensins in gingival epithelial cell breast cancer cell lines, although significant differences are observed between different species of bacteria. Together these results suggest a complex model of a host-defense-related function in maintaining bacterial homeostasis and response to pathogens [11].

Previously, we reported that  $\beta$ -defensin 2 showed immunomodulatory and anti-tumor function of macrophages increased in the breast cancer tumor model. There is a gap in the knowledge of the immunomodulatory and tumoricidal part of tumor-associated macrophages with crosstalk with defensin 2. So, in this study, we aimed to investigate the association between  $\beta$ -defensin 2 and TAMs and determine the role in tumor-promoting attributes of TAMs reversal of phenotype in tumor regression. We explored: (1) whether  $\beta$ -defensin 2 (BD2) inductions have any toxicity on TAMs, (2) effect of  $\beta$ -defensins 2 on cellular viability and reactive oxygen species (ROS) generation (3) we also studied the cytokines such as IL-6, IL-10, IL-3, TNF- $\alpha$ , and TGF- $\beta$ , and chemokines CXCL-1, CXCL-5, CXCL-15, and CCL-24 expression in TAMs.

## 2. Methods

Supplementary materials 1.

## 3. Results

### 3.1. Isolation and characterization of macrophages and TAMs (Fig. 1A and B)


TAMs generated using tumor conditioned media from mouse breast cancer cell line C127i. Flow cytometry was performed for quantification and characterization of harvested cells (Fig. 1). Purity greater than 90% for macrophages and TAMs was confirmed via flow cytometric analysis according to surface marker expression such as F4/80, CD206 and CD115 conjugated with alexafluor, PE-A and PE-CY-7-A respectively.

### 3.2. Nitric oxide and $H_2O_2$ measurement (Fig. 2A–C)

The ROS contains  $NO^-$  and  $H_2O_2$  which was estimated by Griess method and  $TiSO_4$  method. The amount of  $NO$  production in the supernatant of TAMs after 24 h treatment with  $\beta$ -defensin 2 was determined by Griess reagent. During this period, the  $NO$  level decreased remarkably in the  $\beta$ -defensin 2 group as compared to the TAMs group (Fig. 2A) It was observed that in other experiments were conducted to examine the effect of  $\beta$ -defensin 2 on the production of  $H_2O_2$ . Production of  $H_2O_2$  was found to be decreased in the  $\beta$ -defensin 2 treated group as compared to the TAMs group (Fig. 2B). mRNA level of iNOS was upregulated in the  $\beta$ -defensin 2 treated group as compared to the TAMs group (Fig. 2C).


### 3.3. qRT-PCR for selected gene expression on TAMs (Figs. 3 and 4)

We treated TAMs with mouse  $\beta$ -defensin 2 (BD-2) *in vitro*, followed by RT-PCR analysis to assess mRNA levels of respective cytokines and chemokines. Gene expression of cytokines (IL-6, IL-10, IL-3, TNF- $\alpha$  and TGF- $\beta$ ) and chemokines (CXCL-1, CXCL-5, CXCL-15, CCL-24 and CCL-5) were evaluated (Figs. 3 and 4). The results showed that BD-2 significantly increased the expression of cytokines TNF- $\alpha$ , TGF- $\beta$ , IL-6 and IL-10, while diminished the expression of IL-3 as compared to only TAMs group (Fig. 3A–E). The results found that BD-2 reduced the expression of chemokines such as CXCL-1, CXCL-5, CXCL-15, CCL-24 and CCL-5 as compared to only TAMs group (Fig. 4A–E) suggesting that BD-2 may stimulate their anti-tumor promoting capacity.




**Fig. 1.** Characterization of tumor associated macrophages isolated from mice peritoneal cavity based on F4/80, CD-206, and CD-115 conjugated with alexa fluor, PE-A and PE-CY-7A expression through Flow cytometry (E). The cells were gated based on F4/80, CD-206, and CD-115 expression. Percentage refers to the mean of cell subpopulations gated on the total population of macrophage cells. Representative FACS means fluorescence intensity (MFI) histograms and mean MFI  $\pm$  SEM of F4/80, CD-206, and CD-115.






**Fig. 2.** (A–C): Effects of BD 2 on the level of H<sub>2</sub>O<sub>2</sub>, NO and iNOS in tumor associated macrophage cells after 24 h treatment.

Macrophages were seeded in 96-well plate and treated with  $\beta$ -defensin 2 (100 ng). After 24 h of incubation, the supernatants were collected to measure NO and H<sub>2</sub>O<sub>2</sub> level using microplate reader. iNOS mRNA levels were measured by qPCR. Results are expressed as mean  $\pm$  SE. Statistical analysis: \* $p$  < 0.05, \*\* $p$  < 0.01, \*\*\* $p$  < 0.001.



**Fig. 3.** (A–E): Effects of BD 2 on the expression of cytokines in tumor associated macrophage cells after 24 h treatment.

Macrophages were isolated from mouse and treated with  $\beta$ -defensin 2 (100 ng). After 24 h of incubation, total RNA from cells were extracted and subjected to RT-qPCR to evaluate the levels of cytokines IL-6, TNF- $\alpha$ , IL-3, IL-10, TGF- $\beta$  and GAPDH. GAPDH used as a normalization control. Results are expressed as mean  $\pm$  SE. Statistical analysis: \* $p$  < 0.05, \*\* $p$  < 0.01.



**Fig. 4.** (A–E): Effects of BD-2 on the expression of chemokines in tumor associated macrophage cells after 24 h treatment.

Macrophages were isolated from mouse and treated with  $\beta$ -defensin 2 (100 ng). After 24 h of incubation, total RNA from cells were extracted and subjected to RT-qPCR to evaluate the levels of chemokines CXCL-1, CXCL-5, CXCL-15, CCL-24, CCL-5 and GAPDH. GAPDH was used as a normalization control. Results are expressed as mean  $\pm$  SE. Statistical analysis: \*\*\* $p$  < 0.001, \*\* $p$  < 0.01.

#### 4. Discussion

Previously, we reported that  $\beta$ -defensin 2 modulates immunomodulatory and anti-tumor functions of macrophages [12]. The present study

establishes a new paradigm whereby the alteration in oxidative stress, chemokines and cytokines pattern of TAMs by the  $\beta$ -defensin 2 is an effective strategy as a therapeutic agent. It plays an essential role in tumor regression and antitumor function.

Macrophages differentiate into TAMs via their interaction with tumor and tumor-derived factors. TAMs express F4/80, CD206 and CD163, which consist of transcriptionally, phenotypically, and functionally diverse populations of macrophages [13,14]. TAMs promote tumor growth through a variety of approach such as secretion of growth factors, matrix degradation enzymes, and proangiogenic factors [15,16]. TAMs represent an intriguing cell phenotype in different tumor compartments [17]. Tumor cells releasing cytokines and chemokines attract macrophages (tumor promoting TAMs phenotype) and other inflammatory cells to the tumor stroma [18]. Tumor-derived factors in the TME can stimulate macrophages to produce a wide array of tumor-promoting molecules, such as chemokines, cytokines, and growth factors, to stimulate tumor cell proliferation, tumor angiogenesis and metastasis [19, 20]. TAMs express chemokines and cytokines that can promote tumor progression and suppress antitumor immunity.

The level of  $\text{NO}^-$  and  $\text{H}_2\text{O}_2$  was found to be decreased in  $\beta$  defensin-2 treated group as compared to only TAMs group (Fig. 2).  $\beta$  defensin-2 may play role in controlled ROS release during tumor progression. Nitric oxide ( $\text{NO}^-$ ) is a highly reactive free radical that involves in many inflammatory processes. Depending on the source of the  $\text{NO}^-$ , tumor cells can either be stimulated to grow or be encouraged to die. Similarly, the function of iNOS in cancer varies depending on the kind of cell. Consistent with previous reports that TAMs promote tumor initiation, progression and metastasis, it is evident that BD-2 suppressed the tumorigenesis in TAMs.  $\text{NO}^-$  is a main mediator of angiogenesis and neovascularization. It also activates immune responses against pathogens and increased expression of iNOS on macrophages represents a marker for M1 polarization [21]. Although ROS are typically thought to be detrimental to cells, they also play a crucial role in controlling gene expression and signal transduction pathways. This is the first demonstration that  $\beta$ -defensin 2 is a key protein on immune defense which regulates ROS generation and decreased  $\text{NO}^-$  and  $\text{H}_2\text{O}_2$ , pro-inflammatory cytokines and chemokines.

TAMs with different polarization phenotypes can produce a variety of cytokines, chemokines, and growth factors that help to regulate inflammation and immunity. Chemokines are critical secretors derived from TAMs that mediate cancer progress and metastasis. In our study, we have found that lower expression of CXCL-1 in TAMs when treated with BD-2 as compared to only TAMs group (Fig. 3). CXCL-1 secreted by breast cancer cells, recruits  $\text{CD11b}^+\text{Gr1}^+$  myeloid cells into the tumor, therefore supporting the cancer metastasis and survival by activating calprotectin expression [22]. CXCL-1, a chemotaxis-stimulating factor, recruits a variety of stromal cells into tumor surroundings to involve in tumor growth, angiogenesis and metastasis [23].

Here we showed that the expression of  $\text{TNF-}\alpha$ ,  $\text{TGF-}\beta$ , IL-6 and IL-10 is found to be increased in BD-2 treated group as compared to only TAMs group (Fig. 4). TAMs stimulate the expression of IL-6 and IL-10 in cancer cells through TLR signaling, serve in maintaining the tumor's inflammatory microenvironment and promoting development and progression of cancer [24]. IL-6 is an inflammatory cytokine, considered as a tumor-promoting and antiapoptotic factor. It contributes to the induction of skin tumors, triggers malignancy in breast tumor mamospheres, and participates in suppression of antigen-specific anti-tumor immunity. IL-3 known as hematopoietic cytokine which is secreted by activated T cells and mast cells (MCs). IL-10 is a pleiotropic cytokine produced by a variety of immune cells (e.g.,  $\text{T}_\text{H}2$  cells, macrophages and  $\text{CD8}^+$  cells) that can inhibit a broad array of immune and inflammatory responses.  $\text{TGF-}\beta$  secreted by tumor cells can also induce M2-type polarization of macrophages.  $\text{TNF-}\alpha$  is a pleiotropic cytokine that regulates a broad range of biological activities, including cell differentiation, proliferation and death, as well as inflammation, innate and adaptive immune responses, and tissue development [25]. It is produced by monocytes and macrophages.  $\text{TNF-}\alpha$  secreted by TAM leads to the activation of NF- $\kappa\text{B}$  in tumor cells. Activation of NF- $\kappa\text{B}$  pathway in tumor cells prevents tumor cell death and enhances tumor cell invasion [25].

In conclusion, this is the first demonstration that  $\beta$ -defensin 2 is a key

protein on immune defense. Our findings indicated that  $\beta$ -defensin 2 reduced the oxidative stress and expression of chemokines (CXCL-1, 5, 15, CCL-5 and 24) while induced the expression of cytokines (IL-6, 10,  $\text{TNF-}\alpha$  and  $\text{TGF-}\beta$ ). Our data clearly demonstrated that treatment of  $\beta$ -defensin 2 for targeting TAMs may be a potentially effective method for cancer treatment.

### Ethics approval and consent to participate

All animal experimental protocols were approved by the Institutional Animal Ethics Committee (IAEC) (file no- 100/IAEC-1/2018) of AIIMS, New Delhi. All methods and animal monitoring were carried out in accordance with relevant guidelines and regulations provided by the Committee for the Purpose of Control and Supervision of Experimentation of Animals (CPCSEA) (Reg. No. 10/GO/ReBiT/99/CPCSEA). Animal studies are conducted in compliance with the ARRIVE guidelines. Human subject consent was not required in the study because we did not used human samples.

### Consent for publication

Not applicable.

### Availability of data and materials

The datasets used and/or analyzed during the current study are available by the corresponding author Dr. Pramod Kumar Gautam (gautam@aiims.edu) on reasonable request.

### Funding

This research did not receive any specific grant from funding agencies.

### Authors' contributions

Dr. Sonam Agarwal and Anita Chauhan performed wet lab experiments. Dr. Sonam Agarwal and Dr. Pramod Kumar Gautam wrote the manuscript. Dr. Pramod Kumar Gautam was the supervisor and designed the experimental plan. He analyzed all the data and corrected the manuscript. All authors read and approved the final manuscript.

### Declaration of competing interest

The authors declare that they have no competing interests.

### Data availability

Data will be made available on request.

### Acknowledgements

We would like to acknowledge the following core facilities at All India Institute of Medical Sciences: Department of Biochemistry, CIF facility for tissue culture, fluorescent microscopy, flow cytometry and real time PCR for experiments. Dr. Gautam expresses his appreciation to ICMR, New Delhi and IMRG, AIIMS, New Delhi, India for funding.

### Abbreviations

(TAMs)	Tumor associated macrophages
(TNF)	Tumor necrosis factor
(PECs)	peritoneal exudate cells
(NO)	Nitric oxide
(PECs)	Peritoneal exudate cells
(ROS)	Reaction oxygen species

## Appendix A. Supplementary data

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.adcanc.2023.100102>.

## References

- [1] C. Belli, D. Trapani, G. Viale, et al., Targeting the microenvironment in solid tumors, *Cancer Treat Rev.* 65 (2018) 22–32.
- [2] R. Baghban, L. Roshangar, R. Jahanban-Esfahlan, et al., Tumor microenvironment complexity and therapeutic implications at a glance, *Cell Commun. Signal.* 18 (2020) 59.
- [3] G. Dranoff, Cytokines in cancer pathogenesis and cancer therapy, *Nat. Rev. Cancer* 4 (2004) 11–22.
- [4] D. Zhang, X. Qiu, J. Li, et al., TGF- $\beta$  secreted by tumor-associated macrophages promotes proliferation and invasion of colorectal cancer via miR-34a-VEGF axis, *Cell Cycle* 17 (24) (2018) 2766–2778.
- [5] I. Kryczek, L. Zou, P. Rodriguez, G. Zhu, S. Wei, et al., B7-H4 expression identifies a novel suppressive macrophage population in human ovarian carcinoma, *J. Exp. Med.* 203 (2006) 871–881.
- [6] C.Y. Liu, J.Y. Xu, X.Y. Shi, et al., M2-polarized tumor-associated macrophages promoted epithelial-mesenchymal transition in pancreatic cancer cells, partially through TLR4/IL-10 signaling pathway, *Lab. Invest.* 93 (2013) 844–854.
- [7] T. Chanmee, P. Ontong, K. Konno, N. Itano, T. Chanmee, Tumor-associated macrophages as major players in the tumor microenvironment, *Cancers* 6 (2014) 1670–1690.
- [8] L.R. Machado, B. Ottolini, An evolutionary history of defensins: a role for copy number variation in maximizing host innate and adaptive immune responses, *Front. Immunol.* 6 (2015) 115.
- [9] M. Salvatore, A. Garcia-Sastre, P. Ruchala, R.I. Lehrer, T. Chang, M.E. Klotman, alpha-Defensin inhibits influenza virus replication by cell-mediated mechanism(s), *J. Infect. Dis.* 196 (2007) 835–843.
- [10] M. Hollmén, F. Roudnický, S. Karaman, et al., Characterization of macrophage - cancer cell crosstalk in estrogen receptor positive and triple-negative breast cancer, *Sci. Rep.* 5 (2015) 9188.
- [11] G. Diamond, L.K. Ryan, Beta-defensins: what are they REALLY doing in the oral cavity? *Oral Dis.* 17 (7) (2011) 628–635.
- [12] S. Agarwal, A. Chauhan, K. Singh, K. Kumar, R. Kaur, M. Masih, P.K. Gautam, Immunomodulatory effects of  $\beta$ -defensin 2 on macrophages induced immunoregulation and their antitumor function in breast cancer, *BMC Immunol.* 23 (2022) 53.
- [13] E. Van Overmeire, D. Laoui, J. Keirsse, J.A. Van Ginderachter, A. Sarukhan, Mechanisms driving macrophage diversity and specialization in distinct tumor microenvironments and parallels with other tissues, *Front. Immunol.* 5 (2014) 127.
- [14] S. Chevrier, et al., An immune atlas of clear cell renal cell carcinoma, *Cell* 169 (2017) 736–749.
- [15] A. Mantovani, F. Marchesi, A. Malesci, L. Laghi, P. Allavena, Tumour-associated macrophages as treatment targets in oncology, *Nat. Rev. Clin. Oncol.* 14 (2017) 399–416.
- [16] L. Cassetta, J.W. Pollard, Targeting macrophages: therapeutic approaches in cancer, *Nat. Rev. Drug Discov.* 17 (2018) 887–904.
- [17] T. Chanmee, P. Ontong, K. Konno, N. Itano, T. Chanmee, Tumor-associated macrophages as major players in the tumor microenvironment, *Cancers* 6 (2014) 1670–1690.
- [18] C.E. Lewis, J.W. Pollard, Distinct role of macrophages in different tumor microenvironments, *Cancer Res.* 66 (2006) 605–612.
- [19] S. Kim, H. Takahashi, W.W. Lin, P. Descargues, S. Grivennikov, et al., Carcinoma-produced factors activate myeloid cells through TLR2 to stimulate metastasis, *Nature* 457 (2009) 102–106.
- [20] A. Mantovani, P. Allavena, A. Sica, F. Balkwill, Cancer-related inflammation, *Nature* 454 (2008) 436–444.
- [21] F. Klug, et al., Low-dose irradiation programs macrophage differentiation to an iNOS1/M1 phenotype that orchestrates effective T cell immunotherapy, *Cancer Cell* 24 (2013) 589–602.
- [22] S. Acharyya, et al., A CXCL1 paracrine network links cancer chemoresistance and metastasis, *Cell* 150 (2012) 165–178.
- [23] N. Wang, W. Liu, Y. Zheng, et al., CXCL1 derived from tumor-associated macrophages promotes breast cancer metastasis via activating NF- $\kappa$ B/SOX4 signaling, *Cell Death Dis.* 9 (2018) 880.
- [24] X. Ke, M. Wu, J. Lou, S. Zhang, P. Huang, R. Sun, et al., Activation of Toll-like receptors signaling in non-small cell lung cancer cell line induced by tumor-associated macrophages, *Chin. J. Cancer Res.* 27 (2015) 181–189.
- [25] C.H. Lee, C.L. Wu, A.L. Shiau, Toll-like receptor 4 signaling promotes tumor growth, *J. Immunother.* 33 (2010) 73–82.