### Prof. Sarman Singh, Director, AIIMS, Bhopal

# Statement of Research Achievements for which Sun Pharma Award is Claimed

Tuberculosis is the main killer infectious diseases in India. On the top of that multi-drug resistant (MDR) and extensively drug resistant (XDR) forms have emerged. Diagnosis of TB still remains a major challenge. Prof. Singh has done outstanding work in the field of TB diagnostics.


In the last 20 years several serological test kits were dumped in the Indian market for the diagnosis of PTB and EPTB. But these kits gave highly inaccurate results, leading to unnecessary treatment to hundreds of thousands of patients, and leaving several thousands of TB patients untreated, leading ban on these kits.


However, the ban on serology has created a complete gap in the TB diagnostic field. Dr. Singh worked untiringly to find out novel molecular tools and biomarkers for the diagnosis and differentiation of drug susceptible and drug resistant Mycobacterium tuberculosis. His laboratory made a revolutionary discovery in this field by screening hundreds of proteins of MTB, his team identified 5 novel proteins/antigens which are over-expressed only during the in-vivo drug resistance development. The genes were cloned and proteins expressed, purified and used on various categories of patients showing the sensitivity and specificity of these proteins between 98.2% -100% and 89.1 - 98.2%, respectively (Singh et al, 2015, Singh et al, 2017). Using these novel proteins, development of a point-of-care (POC) rapid test is underway. This innovation is expected to be a game changer in the area of TB diagnostics not only for India but globally. The Foundation for Newer Innovative Diagnostics (FIND) evaluated these antigens and has shown interest in this innovation. The prototype ICT based RDTs prepared in the laboratory (images shown below) have shown excellent results and technology is ready for transfer to a commercial partner for marketing this innovation.

Dr. Singh has also done pioneering work in the field of non-tuberculous mycobacteria (NTM) often neglected by clinicians and medical microbiologists as contaminants. In AIDS era these NTM have gained much importance but the conventional methods of identification are neither reproducible nor very specific. His team has developed novel multiplex-PCR (<u>images shown below</u>) for the diagnosis and differentiation of Mycobacterium tuberculosis, M. avium, M. kansasii and other NTM, directly from the clinical samples in a single tube (PCT/IN2004/000396). These PCR primers and the process have been found highly sensitive and specific and are being routinely used at various laboratories throughout the country including AIIMS (Singh et al, 2006, Gopinath et al, 2009, Kumar et al 2014a). This has made a paradigm shift in the understanding of NTM disease in India. Using the same gene targets, a new technology known as loop mediated isothermal amplification (LAMP) assay (<u>images shown below</u>) have also been developed by him. The technology is being evaluated across the country through ICMR and DBT, Government of India.

Currently molecular tools have become essential to understand the pathophysiology and drug targets. Dr. Singh has moved with the time. His laboratory is equipped with all modern tools including 6 laser flow cytometer and New Generation Sequencer (NGS), beside all routine diagnostic tools and services. We have done WGS of more than 200 MTB clinical isolates and have submitted several Genome projects and several thousands of the gene sequences in the GenBank for the benefit of other researchers.

Therefore, my contribution of research has been highly significant and ground breaking. I could translate my research into application for the patient care, justifying for this recognition.






# SCIENTIFIC REPORTS


Received: 12 July 2016 Accepted: 06 February 2017 Published: 24 March 2017


# **OPEN** Evaluation of 5 Novel protein biomarkers for the rapid diagnosis of pulmonary and extra-pulmonary tuberculosis: preliminary results

Amit Singh<sup>1,\*</sup>, Anil Kumar Gupta<sup>1,\*</sup>, Krishnamoorthy Gopinath<sup>1,†</sup>, Pawan Sharma<sup>2,‡</sup> & Sarman Singh<sup>1</sup>


Improved methods are required for the early and accurate diagnosis of tuberculosis, especially in the patients with smear-negative disease. Several biomarkers have been tried but most have shown poor sensitivity or specificity. In present study we aimed to evaluate the diagnostic utility of five novel antigens identified earlier by us. This is an initial study conducted on 250 subjects. The five recombinant antigens, named as rSS1 (Rv2145c), rSS2 (Rv0164), rSS3 (Rv1437), rSS4 (Rv1827) and rSS5 (Rv2970c), were expressed in pQE-30 expression vector, purified and their sero-diagnostic efficacy was evaluated in an unblinded manner using dot-blot and ELISA methods. The sensitivity and specificity of these








Laboratory Scale preparation of ICT Based RDTs for the TB diagnosis





Indian Patent Application for Biomarkers and International Patent for Multiplex PCR

Novel set of PCR primers developed for detecting and differentiating the MTB from *M. avium*, M. *Kansasii* and other NTM in a single reaction tube.



LAMP (Loop Mediated Isothermal Amplification) Test Developed using the same primers and new probes . This test does not require any detection device to report the test results.





# Tuberculosis Patents by Inventor Dr. Sarman Singh

Sarman Singh has filed for patents to protect the following inventions. This listing includes patent applications that are pending as well as patents that have already been granted by the United States Patent and Trademark Office (USPTO).

# • Constructing a DNA chimera for vaccine development against leishmaniasis and tuberculosis

Patent number: 8299232

Abstract: A novel recombinant chimera of DNA construct having esat-6 region of Mycobacterium tuberculosis and kinesin region of Leishmania donovani cloned together on two sides of self cleaving peptide in a DNA vaccine vector pVAX-1 wherein the chimeric construct is operatively linked to a transcriptional promoter thus capable of self replication and expression within the mammalian cell, and the process of preparation thereof comprising: analysis of the predicted protein sequence of kinesin motor domain and esat-6 domain using Promiscuous MHC Class-1 Binding Peptide Prediction Servers; amplification of gene coding for kinesin motor domain and esat-6 domain; cloning of kinesin esat-6 gene region in pGEM-T™ vector for sequence analysis; generation of chimeric construct by directional cloning in pVAX-1 vector.

Type: Grant

Filed: February 10, 2009

Date of Patent: October 30, 2012

Assignees: Department of Biotechnology, All India Institute of Medical Science

**Inventors:** Sarman Singh, Ayan Dey

# Constructing a DNA Chimera for Vaccine Development Against Leishmaniasis and Tuberculosis

Publication number: 20110150932

Abstract: A novel recombinant chimera of DNA construct having esat-6 region of Mycobacterium tuberculosis and kinesin region of Leishmania donovani cloned together on two sides of self cleaving peptide in a DNA vaccine vector pVAX-1 wherein the chimeric construct is operatively linked to a transcriptional promoter thus capable of self replication and expression within the mammalian cell, and the process of preparation thereof comprising: analysis of the predicted protein sequence of kinesin motor domain and esat-6 domain using Promiscuous MHC Class-1 Binding Peptide Prediction Servers; amplification of gene coding for kinesin motor domain and esat-6 domain; cloning of kinesin esat-6 gene region in pGEM-T™ vector for sequence analysis; generation of chimeric construct by directional cloning in pVAX-1 vector.

**Type:** Application

Filed: February 10, 2009

Publication date: June 23, 2011

Applicants: DEPARTMENT OF BIOTECHNOLOGY, ALL INDIA INSTITUTE OF MEDICAL

SCIENCE

**Inventors:** Sarman Singh, Ayan Dey

# Publications ONLY on Tuberculosis (88 out of 345)

- 1. <u>Profiles of Volatile Biomarkers Detect **Tuberculosis** from Skin. Vishinkin R, et al. Adv Sci (Weinh). 2021 Aug;8(15):e2100235. doi: 10.1002/advs.202100235. Epub 2021 Jun 2.PMID: 34075714</u>
- COVID-19 Blind Spots: A Consensus Statement on the Importance of Competent Political Leadership and the Need for Public Health Cognizance. Papadimos TJ, et al. J Glob Infect Dis. 2020 Nov 30;12(4):167-190. doi: 10.4103/jgid.jgid\_397\_20. eCollection 2020 Oct-Dec.PMID: 33888955
- 3. <u>Pulmonary infections after renal transplantation: a prospective study from a tropical country.</u>

  Mangalgi S, et al. Transpl Int. 2021 Mar;34(3):525-534. doi: 10.1111/tri.13817. Epub 2021 Feb 26.PMID: 33423313
- Mycobacterium tuberculosis bloodstream infection prevalence, diagnosis, and mortality risk in seriously ill adults with HIV: a systematic review and meta-analysis of individual patient data.
   Barr DA, et al. Lancet Infect Dis. 2020 Jun;20(6):742-752. doi: 10.1016/S1473-3099(19)30695-4.
- 5. The risk of **tuberculosis** in children after close exposure: a systematic review and individual-participant meta-analysis. Martinez L, et al. Lancet. 2020 Mar 21;395(10228):973-984. doi: 10.1016/S0140-6736(20)30166-5.PMID: 32199484
- 6. Performance of Genotype MTBDRsl V2.0 over the Genotype MTBDRsl V1 for detection of second line drug resistance: An Indian perspective. Rufai SB, Umay K, Singh PK, Singh S. PLoS One. 2020 Mar 4;15(3):e0229419. doi: 10.1371/journal.pone.0229419. eCollection 2020.PMID: 32130233
- 7. <u>Trehalose limits opportunistic mycobacterial survival during HIV co-infection by reversing HIV-mediated autophagy block.</u> Sharma V, et al. Autophagy. 2021 Feb;17(2):476-495. doi: 10.1080/15548627.2020.1725374. Epub 2020 Feb 20.PMID: 32079455
- 8. Raman spectroscopy reveals distinct differences between two closely related bacterial strains,

  Mycobacterium indicus pranii and Mycobacterium intracellulare. Verma T, et al. Anal Bioanal Chem.

  2019 Dec;411(30):7997-8009. doi: 10.1007/s00216-019-02197-z. Epub 2019 Nov 16.PMID: 31732785
- Understanding lipidomic basis of iron limitation induced chemosensitization of drug-resistant <u>Mycobacterium tuberculosis.</u> Pal R, Hameed S, Kumar P, Singh S, Fatima Z. 3 Biotech. 2019 Apr;9(4):122. doi: 10.1007/s13205-019-1645-4. Epub 2019 Mar 5.PMID: 30863701
- Whole-Genome Sequencing of Two Extensively Drug-Resistant Mycobacterium tuberculosis Isolates from India. Rufai SB, Singh S. Microbiol Resour Announc. 2019 Feb 14;8(7):e00007-19. doi: 10.1128/MRA.00007-19. eCollection 2019 Feb.PMID: 30801057
- 11. <u>Investigations into Isoniazid Treated Mycobacterium tuberculosis</u> by Electrospray Mass Spectrometry <u>Reveals New Insights into Its Lipid Composition.</u> Pal R, Hameed S, Sabareesh V, Kumar P, **Singh S**, Fatima Z.J Pathog. 2018 Jun 19;2018:1454316. doi: 10.1155/2018/1454316.
- Association of gyrA and rrs gene mutations detected by MTBDRsl V1 on Mycobacterium tuberculosis strains of diverse genetic background from India. Rufai SB, et al. Sci Rep. 2018 Jun 18;8(1):9295. doi: 10.1038/s41598-018-27299-z.
- 13. Diagnostic algorithm for low-volume CSF samples in tuberculous meningitis.

- **Singh S,** Sankar MM. Lancet Infect Dis. 2017 Dec;17(12):1236-1237. doi: 10.1016/S1473-3099(17)30639-4.PMID: 29173879
- 14. Comparative lipidomics of drug sensitive and resistant Mycobacterium **tuberculosis** reveals altered <u>lipid imprints.</u> Pal R, Hameed S, Kumar P, **Singh S**, Fatima Z.3 Biotech. 2017 Oct;7(5):325. doi: 10.1007/s13205-017-0972-6. Epub 2017 Sep 16.PMID:
- 15. Evaluation of antimycobacterial, leishmanicidal and antibacterial activity of three medicinal orchids of Arunachal Pradesh, India. Bhatnagar M, et al. BMC Complement Altern Med. 2017 Aug 1;17(1):379. doi: 10.1186/s12906-017-1884-z.PMID: 28764749
- Diagnostic usefulness of Xpert MTB/RIF assay for detection of tuberculous meningitis using cerebrospinal fluid. Rufai SB, et al. J Infect. 2017 Aug;75(2):125-131. doi: 10.1016/j.jinf.2017.04.010. Epub 2017 May 10.PMID: 28501491
- 17. Evaluation of 5 Novel protein biomarkers for the rapid diagnosis of pulmonary and extrapulmonary tuberculosis: preliminary results. Singh A, et al. Sci Rep. 2017 Mar 24;7:44121. doi: 10.1038/srep44121.PMID: 28337993
- 18. <u>Predominance of Beijing lineage among pre-extensively drug-resistant and extensively drug-resistant strains of Mycobacterium tuberculosis: A tertiary care center experience.</u> Rufai SB, et al. Int J Mycobacteriol. 2016 Dec;5 Suppl 1:S197-S198. doi: 10.1016/j.ijmyco.2016.07.005.
- Comparative proteomic analysis of sequential isolates of Mycobacterium tuberculosis sensitive and resistant Beijing type from a patient with pulmonary tuberculosis. Singh A, et al. Int J Mycobacteriol. 2016 Dec;5 Suppl 1:S123-S124. doi: 10.1016/j.ijmyco.2016.10.028. Epub 2016 Nov 16.
- Development of rapid immuno-diagnostic test for the early detection of tuberculosis.
   Gupta AK, Srivastava S, Singh A, Singh S. Int J Mycobacteriol. 2016 Dec;5 Suppl 1:S114-S115. doi: 10.1016/j.ijmyco.2016.11.008. Epub 2016 Dec 2.PMID: 28043495
- 21. <u>Performance of Xpert MTB/RIF on Ascitic Fluid Samples for Detection of Abdominal **Tuberculosis**.</u> Rufai SB, **Singh S**, Singh A, Kumar P, Singh J, Vishal A.J Lab Physicians. 2017 Jan-Mar;9(1):47-52. doi: 10.4103/0974-2727.187927.PMID: 28042217
- 22. BLR1 and FCGR1A transcripts in peripheral blood associate with the extent of intrathoracic tuberculosis in children and predict treatment outcome.
  Jenum S, et al. Sci Rep. 2016 Dec 12;6:38841. doi: 10.1038/srep38841.PMID: 27941850
- 23. Approaching a diagnostic point-of-care test for pediatric **tuberculosis** through evaluation of immune biomarkers across the clinical disease spectrum. Jenum S, et al. Sci Rep. 2016 Jan 4;6:18520. doi: 10.1038/srep18520.PMID: 26725873
- 24. Performance of Four Transport and Storage Systems for Molecular Detection of Multidrug-Resistant Tuberculosis. Rabodoarivelo MS, et al. PLoS One. 2015 Oct 2;10(10):e0139382. doi: 10.1371/journal.pone.0139382. eCollection 2015.PMID: 26431352
- Xpert MTB/RIF assay can be used on archived gastric aspirate and induced sputum samples for sensitive diagnosis of paediatric tuberculosis. Singh S, et al. BMC Microbiol. 2015 Sep 29;15:191. doi: 10.1186/s12866-015-0528-z.PMID: 26420261

- 26. Performance of Xpert MTB/RIF Assay in Diagnosis of Pleural Tuberculosis by Use of Pleural Fluid Samples. Rufai SB, Singh A, Kumar P, Singh J, Singh S.J Clin Microbiol. 2015 Nov;53(11):3636-8. doi: 10.1128/JCM.02182-15. Epub 2015 Aug 26.PMID: 26311855
- 27. <u>High prevalence of primary drug resistance in children with intrathoracic **tuberculosis** in India. Prajapati S, et al. Paediatr Int Child Health. 2016 Aug;36(3):214-8. doi: 10.1179/2046905515Y.0000000041. PMID: 26052730</u>
- 28. Challenges in tuberculosis diagnosis and management: recommendations of the expert panel. Singh S, Dey B, Sachdeva KS, Kabra SK, Chopra KK, Chaudhary VK, Sharma P, Katoch VM.J Lab Physicians. 2015 Jan-Jun;7(1):1-3. doi: 10.4103/0974-2727.154778.PMID: 25949051
- 29. <u>Genetic diversity and drug susceptibility profile of Mycobacterium **tuberculosis** isolated from <u>different regions of India.</u> Singh J, Sankar MM, Kumar P, Couvin D, Rastogi N, **Singh S**; Indian TB Diagnostics Network.J Infect. 2015 Aug;71(2):207-19. doi: 10.1016/j.jinf.2015.04.028.</u>
- 30. Comparative proteomic analysis of sequential isolates of Mycobacterium **tuberculosis** from a patient with pulmonary **tuberculosis** turning from drug sensitive to multidrug resistant.

  Singh A, et al. Indian J Med Res. 2015 Jan;141(1):27-45. doi: 10.4103/0971-5916.154492.
- 31. Evolution of M. bovis BCG Vaccine: Is Niacin Production Still a Valid Biomarker?

  Singh S, Kumar M, Singh P. Tuberc Res Treat. 2015;2015:957519. doi: 10.1155/2015/957519. Epub 2015 Jan 28.PMID:
- 32. <u>Is Xpert MTB/RIF assay in gastric lavage aspirate useful for diagnosis of smear-negative childhood pulmonary tuberculosis?</u> Mathew JL, Vijayasekharan D, Singh S. Indian Pediatr. 2014 Dec;51(12):1007-11. doi: 10.1007/s13312-014-0548-z.PMID: 25560161
- 33. <u>Vitamin D levels in Indian children with intrathoracic **tuberculosis**.

  Khandelwal D, et al. Indian J Med Res. 2014 Oct;140(4):531-7.PMID: 25488448</u>
- 34. <u>Early detection of multi-drug resistant **tuberculosis** in India using GenoType MTBDRplus assay & profile of resistance mutations in Mycobacterium **tuberculosis**. **Singh S.** Indian J Med Res. 2014 Oct;140(4):477-9.PMID: 25488439</u>
- 35. Effect of micronutrient supplementation on treatment outcomes in children with intrathoracic tuberculosis: a randomized controlled trial. Lodha R, et al. Am J Clin Nutr. 2014 Nov;100(5):1287-97. doi: 10.3945/ajcn.113.082255. Epub 2014 Sep 10.PMID: 25332327 Clinical Trial.
- 36. <u>Loop-mediated isothermal amplification assay for rapid and sensitive diagnosis of **tuberculosis**. Kumar P, Pandya D, Singh N, Behera D, Aggarwal P, **Singh S.** J Infect. 2014 Dec;69(6):607-15. doi: 10.1016/j.jinf.2014.08.017. Epub 2014 Sep 9.PMID: 25218428</u>
- 37. Reply to "molecular diagnosis of rifampin-monoresistant **tuberculosis** in Indian patients: problems with a discordance analysis". **Singh S**, Rufai SB, Kumar P, Singh J.J Clin Microbiol. 2014 Sep;52(9):3504-5. doi: 10.1128/JCM.01731-14.PMID: 25143419
- 38. A toolbox for **tuberculosis** (TB) diagnosis: an Indian multi-centric study (2006-2008); evaluation of serological assays based on PGL-Tb1 and ESAT-6/CFP10 antigens for TB diagnosis. Lagrange PH, et al. PLoS One. 2014 May 5;9(5):e96367. doi: 10.1371/journal.pone.0096367. eCollection 2014.

- Comparison of Xpert MTB/RIF with line probe assay for detection of rifampin-monoresistant <u>Mycobacterium tuberculosis.</u> Rufai SB, et al. J Clin Microbiol. 2014 Jun;52(6):1846-52. doi: 10.1128/JCM.03005-13. Epub 2014 Mar 19.PMID: 24648554
- 40. Comparison of an in-house multiplex PCR with two commercial immuno-chromatographic tests for rapid identification and differentiation of MTB from NTM isolates. Kumar P, Benny P, Jain M, Singh S. Int J Mycobacteriol. 2014 Mar;3(1):50-6. doi: 10.1016/j.ijmyco.2013.12.001.
- 41. <u>Deciphering the sequential events during in vivo acquisition of drug resistance in Mycobacterium tuberculosis.</u> Singh A, Gopinath K, Singh N, Singh S. Int J Mycobacteriol. 2014 Mar;3(1):36-40. doi: 10.1016/j.ijmyco.2013.10.006. Epub 2013 Nov 15.PMID: 26786220
- 42. <u>High degree of multi-drug resistance and hetero-resistance in pulmonary TB patients from Punjab state of India.</u> Kumar P, Balooni V, Sharma BK, Kapil V, Sachdeva KS, **Singh S.** Tuberculosis (Edinb). 2014 Jan;94(1):73-80. doi: 10.1016/j.tube.2013.10.001. Epub 2013 Oct 22.PMID: 24184256
- 43. A toolbox for **tuberculosis** (TB) diagnosis: an Indian multicentric study (2006-2008). Evaluation of QuantiFERON-TB gold in tube for TB diagnosis. Lagrange PH, et al. PLoS One. 2013 Sep 6;8(9):e73579. doi: 10.1371/journal.pone.0073579. eCollection 2013.PMID: 24039990
- 44. <u>Diagnostic performance of commercially available enzyme-linked immunosorbent assay kit in the diagnosis of extrapulmonary tuberculosis.</u> Sankar MM, Balooni V, Singh J, Singh S. J Lab Physicians. 2013 Jan;5(1):11-6. doi: 10.4103/0974-2727.115902.PMID: 24014961
- 45. <u>Heat fixed but unstained slide smears are infectious to laboratory staff.</u>
  Kumar P, Rufai SB, **Singh S.**Indian J Tuberc. 2013 Jul;60(3):142-6.PMID: 24000491
- 46. <u>Bio-safety precautions in **tuberculosis** laboratory.</u> **Singh S.**Indian J Tuberc. 2013 Jul;60(3):135-7.PMID: 24000489
- 47. Ambulatory gastric lavages provide better yields of Mycobacterium **tuberculosis** than induced sputum in children with intrathoracic **tuberculosis**. Mukherjee A, et al. Pediatr Infect Dis J. 2013 Dec;32(12):1313-7. doi: 10.1097/INF.0b013e31829f5c58.PMID: 23958816
- Isolated tuberculous liver abscess in a 3-year-old immunocompetent child. Nandan D, Bhatt GC, Dewan V, Yadav TP, Singh S. Paediatr Int Child Health. 2013 Aug;33(3):187-9. doi: 10.1179/2046905512Y.0000000033.PMID: 23930733
- 49. <u>Incidence and prevalence of **tuberculosis** among household contacts of pulmonary **tuberculosis** patients in a peri-urban population of South Delhi, India. Singh J, et al. PLoS One. 2013 Jul 26;8(7):e69730. doi: 10.1371/journal.pone.0069730. Print 2013.PMID: 23922784</u>
- 50. Genetic heterogeneity revealed by sequence analysis of Mycobacterium tuberculosis isolates from extra-pulmonary tuberculosis patients. Das S, et al. BMC Genomics. 2013 Jun 17;14:404. doi: 10.1186/1471-2164-14-404.PMID: 23773324
- 51. <u>Pre-antiretroviral therapy plasma levels of CCL2 may aid in the prediction of **tuberculosis**-associated immune reconstitution inflammatory syndrome in HIV patients after they commence antiretroviral therapy. Oliver BG, et al.J Acquir Immune Defic Syndr. 2013. doi: 10.1097/QAI.0b013e31828e6182.</u>

- 52. The search for a genetic factor associating with immune restoration disease in HIV patients coinfected with Mycobacterium tuberculosis. Affandi JS, et al. Dis Markers. 2013;34(6):445-9. doi: 10.3233/DMA-130991.PMID: 23594560
- 53. <u>Does neutralization of gastric aspirates from children with suspected intrathoracic **tuberculosis** affect mycobacterial yields on MGIT culture? Parashar D, et al. J Clin Microbiol. 2013 Jun;51(6):1753-6. doi: 10.1128/JCM.00202-13. Epub 2013 Mar 27.PMID: 23536406</u>
- 54. Rapid Identification and Drug Susceptibility Testing of Mycobacterium tuberculosis: Standard Operating Procedure for Non-Commercial Assays: Part 3: Colorimetric Redox Indicator Assay v1.3.12.
  Singh S, et al. J Lab Physicians. 2012 Jul;4(2):120-6. doi: 10.4103/0974-2727.105594.PMID: 23440615
- 55. Rapid Identification and Drug Susceptibility Testing of Mycobacterium tuberculosis: Standard Operating Procedure for Non-Commercial Assays: Part 2: Nitrate Reductase Assay v1.3.12.
  Singh S, et al. J Lab Physicians. 2012 Jul;4(2):112-9. doi: 10.4103/0974-2727.105593.PMID: 23440455
- 56. Rapid Identification and Drug Susceptibility Testing of Mycobacterium tuberculosis: Standard Operating Procedure for Non-Commercial Assays: Part 1: Microscopic Observation Drug Susceptibility Assay v2.4.12. Singh S, et al. J Lab Physicians. 2012 Jul;4(2):101-11. doi: 10.4103/0974-2727.105592.
- 57. <u>Isolated tuberculous liver abscess in immunocompetent children report of two cases.</u> Bhatt GC, Nandan D, Singh S. Pathog Glob Health. 2013; 107(1):35-7. doi: 10.1179/2047773212Y.0000000071. PMID: 23432862
- Anti-mycobacterial activity of plumericin and isoplumericin against MDR
   Mycobacterium tuberculosis.
   Kumar P, Singh A, Sharma U, Singh D, Dobhal MP, Singh S. Pulm Pharmacol Ther. 2013 Jun;26(3):332-5. doi: 10.1016/j.pupt.2013.01.003. Epub 2013 Jan 16.PMID: 23333815
- 59. Molecular characterization of Mycobacterium tuberculosis isolates from North Indian patients with extrapulmonary tuberculosis. Sankar MM, Singh J, Diana SC, Singh S. Tuberculosis (Edinb). 2013 Jan;93(1):75-83. doi: 10.1016/j.tube.2012.10.005. Epub 2012 Nov 7.PMID: 23140853
- 60. Prevalence of drug resistant **tuberculosis** among patients at high-risk for Hiv attending outpatient clinics in Delhi, India. Magee MJ, Blumberg HM, Broz D, Furner SE, Samson L, **Singh S**, Hershow RC. Southeast Asian J Trop Med Public Health. 2012 Mar;43(2):354-63.PMID: 23082587
- 61. <u>A toolbox for **tuberculosis** diagnosis: an Indian multicentric study (2006-2008): microbiological results.</u> Lagrange PH, et al. PLoS One. 2012;7(8):e43739. doi: 10.1371/journal.pone.0043739. Epub 2012 Aug 24.PMID: 22937088
- 62. Poor performance of serological tests in the diagnosis of pulmonary **tuberculosis**: evidence from a contact tracing field study. **Singh S**, Singh J, Kumar S, Gopinath K, Balooni V, Singh N, Mani K.PLoS One. 2012;7(7):e40213. doi: 10.1371/journal.pone.0040213. Epub 2012 Jul 10.PMID: 22808119
- 63. AIDS associated **tuberculosis**: a catastrophic collision to evade the host immune system.

  Munawwar A, **Singh S.** Tuberculosis (Edinb). 2012 Sep;92(5):384-7. doi: 10.1016/j.tube.2012.05.006.

  Epub 2012 Jul 12.PMID: 22795795.

- 64. <u>Short communication: Plasma levels of vitamin D in HIV patients initiating antiretroviral therapy do not predict immune restoration disease associated with Mycobacterium **tuberculosis**.</u>
- 65. Price P, et al. AIDS Res Hum Retroviruses. 2012 Oct;28(10):1216-9. doi: 10.1089/AID.2011.0272.
- Commercial serological tests for the diagnosis of active tuberculosis in India: time for introspection.
   Singh S, Katoch VM.Indian J Med Res. 2011 Nov;134(5):583-7. doi: 10.4103/0971-5916.90980.PMID: 22199094
- 67. Mycobacterial antibody levels and immune restoration disease in HIV patients treated in South East Asia. Sumatoh HR, et al. Biomark Med. 2011 Dec;5(6):847-53. doi: 10.2217/bmm.11.79.PMID: 22103621
- 68. Mycobacterium tuberculosis evades host immunity by recruiting mesenchymal stem cells. Raghuvanshi S, Sharma P, Singh S, Van Kaer L, Das G.Proc Natl Acad Sci U S A. 2010 Dec 14;107(50):21653-8. doi: 10.1073/pnas.1007967107. Epub 2010 Dec 6.PMID: 21135221
- 69. <u>Evaluation of risk factors for antituberculosis treatment induced hepatotoxicity.</u>
  Singla R, Sharma SK, Mohan A, Makharia G, Sreenivas V, Jha B, Kumar S, Sarda P, **Singh S.**Indian J Med Res. 2010 Jul;132:81-6.PMID: 20693595
- 70. Non-tuberculous mycobacteria in TB-endemic countries: are we neglecting the danger?

  Gopinath K, Singh S. PLoS Negl Trop Dis. 2010 Apr 27;4(4):e615. doi: 10.1371/journal.pntd.0000615.

  PMID: 20436962
- 71. Assessment of the N-PCR assay in diagnosis of pleural tuberculosis: detection of M. tuberculosis in pleural fluid and sputum collected in tandem. Kumar P, et al. PLoS One. 2010 Apr 19;5(4):e10220. doi: 10.1371/journal.pone.0010220.PMID: 20419090
- 72. <u>Genome-wide analysis of the host intracellular network that regulates survival of Mycobacterium tuberculosis.</u> Kumar D, Nath L, Kamal MA, Varshney A, Jain A, **Singh S**, Rao KV. Cell. 2010 Mar 5;140(5):731-43. doi: 10.1016/j.cell.2010.02.012.PMID: 20211141
- 73. Safety of 3 different reintroduction regimens of antituberculosis drugs after development of antituberculosis treatment-induced hepatotoxicity. Sharma SK, et al. Clin Infect Dis. 2010 Mar 15;50(6):833-9. doi: 10.1086/650576.PMID: 20156055 Clinical Trial.
- 74. <u>Acute viral hepatitis as a confounding factor in patients with antituberculosis treatment induced hepatotoxicity.</u> Sharma SK, Singla R, Sreenivas V, Kumar S, Jha B, Rathored J, **Singh S.**Indian J Med Res. 2009 Aug;130(2):200-1.PMID: 19797819
- 75. Recovery of Mycobacterium tuberculosis from sputum treated with cetyl pyridinium chloride. Sankar MM, Kumar P, Munawwar A, Singh J, Parashar D, Singh S. J Clin Microbiol. 2009 Dec;47(12):4189-90. doi: 10.1128/JCM.01295-09. Epub 2009 Sep 30.PMID: 19794041
- 76. Visual format for detection of Mycobacterium tuberculosis and M. bovis in clinical samples using molecular beacons. Kumar P, et al. J Mol Diagn. 2009 Sep;11(5):430-8. doi: 10.2353/jmoldx.2009.080135. Epub 2009 Aug 6.PMID: 19661384
- 77. Immunogenicity of candidate chimeric DNA vaccine against **tuberculosis** and leishmaniasis. Dey A, Kumar U, Sharma P, **Singh S.** Vaccine. 2009 Aug 13;27(37):5152-60. doi: 10.1016/j.vaccine.2009.05.100. Epub 2009 Jun 24.PMID: 19559111

- 78. <u>Variable-number tandem repeat 3690 polymorphism in Indian clinical isolates of Mycobacterium tuberculosis and its influence on transcription.</u> Akhtar P, et al . J Med Microbiol. 2009 Jun;58(Pt 6):798-805. doi: 10.1099/jmm.0.002550-0. PMID: 19429757
- 79. Role of acute viral hepatitis as a confounding factor in antituberculosis treatment induced hepatotoxicity. Sarda P, Sharma SK, Mohan A, Makharia G, Jayaswal A, Pandey RM, Singh S.Indian J Med Res. 2009 Jan;129(1):64-7.PMID: 19287059
- Urine as an adjunct specimen for the diagnosis of active pulmonary tuberculosis. Gopinath K, Singh
   Int J Infect Dis. 2009 May;13(3):374-9. doi: 10.1016/j.ijid.2008.07.022. Epub 2008 Nov 5.
   PMID: 18980851
- 81. Comparative evaluation of FASTPlaque assay with PCR and other conventional in vitro diagnostic methods for the early detection of pulmonary tuberculosis. Singh S, Saluja TP, Kaur M, Khilnani GC.J Clin Lab Anal. 2008;22(5):367-74. doi: 10.1002/jcla.20264.PMID: 18803271
- 82. Opportunistic pathogens in AIDS: trends, diagnosis and priorities.

  Grewal HM, Gupta S, Singh S. Expert Rev Anti Infect Ther. 2008 Apr;6(2):163-6. doi: 10.1586/14787210.6.2.163. PMID: 18380598
- 83. <u>High rate of extensively drug-resistant **tuberculosis** in Indian AIDS patients. **Singh S**, Sankar MM, Gopinath K.AIDS. 2007 Nov 12;21(17):2345-7. doi: 10.1097/QAD.0b013e3282f125c9.PMID: 18090285</u>
- 84. Novel method for clearing red blood cell debris from BacT/ALERT blood culture medium for improved microscopic and antimycobacterial drug susceptibility test results. Gopinath K, Kumar S, Mani Sankar M, Singh S.J Clin Lab Anal. 2007;21(4):220-6. doi: 10.1002/jcla.20169.PMID: 17621357
- 85. <u>Controlling multidrug-resistant **tuberculosis** in India.</u> Gopinath K, Manisankar M, Kumar S, **Singh S.**Lancet. 2007 Mar 3;369(9563):741-742. doi: 10.1016/S0140-6736(07)60358-4.PMID: 17336647
- 86. Nontuberculous mycobacterial infections in Indian AIDS patients detected by a novel set of ESAT-6 polymerase chain reaction primers. Singh S, Gopinath K, Shahdad S, Kaur M, Singh B, Sharma P.Jpn J Infect Dis. 2007 Feb;60(1):14-8.PMID: 17314419
- 87. <u>Down-regulation of T helper 1 responses to mycobacterial antigens due to maturation of dendritic cells by 10-kDa mycobacterium **tuberculosis** secretory antigen. Natarajan K, Latchumanan VK, Singh B, **Singh S**, Sharma P. J Infect Dis. 2003 Mar 15;187(6):914-28. doi: 10.1086/368173.</u>
- 88. Effect of Mycobacterium **tuberculosis**-specific 10-kilodalton antigen on macrophage release of tumor necrosis factor alpha and nitric oxide. Trajkovic V, Singh G, Singh B, **Singh S**, Sharma P.Infect Immun. 2002 Dec;70(12):6558-66. doi: 10.1128/IAI.70.12.6558-6566.2002.PMID: 12438325

24<sup>th</sup> September 2021 Bhopal **Prof. Sarman Singh**