

Lacrimal Gland Insufficiency in Aqueous Deficiency Dry Eye Disease: Recent Advances in Pathogenesis, Diagnosis, and Treatment

Mariya B Doctor & Sayan Basu

To cite this article: Mariya B Doctor & Sayan Basu (2022): Lacrimal Gland Insufficiency in Aqueous Deficiency Dry Eye Disease: Recent Advances in Pathogenesis, Diagnosis, and Treatment, Seminars in Ophthalmology, DOI: [10.1080/08820538.2022.2075706](https://doi.org/10.1080/08820538.2022.2075706)

To link to this article: <https://doi.org/10.1080/08820538.2022.2075706>



Published online: 19 May 2022.



Submit your article to this journal [↗](#)





View related articles [↗](#)



View Crossmark data [↗](#)



Lacrimal Gland Insufficiency in Aqueous Deficiency Dry Eye Disease: Recent Advances in Pathogenesis, Diagnosis, and Treatment

Mariya B Doctor ^{a,b} and Sayan Basu ^{b,c}

^aAcademy of Eye Care Education, L V Prasad Eye Institute, Hyderabad, India; ^bThe Cornea Institute, L V Prasad Eye Institute, Hyderabad, India; ^cBrien Holden Eye Research Centre, L V Prasad Eye Institute, Hyderabad, India

ABSTRACT

Background: Aqueous deficiency dry eye disease is a chronic and potentially sight-threatening condition, that occurs due to the dysfunction of the lacrimal glands. The aim of this review was to describe the various recent developments in the understanding, diagnosis and treatment of lacrimal gland insufficiency in aqueous deficiency dry eye disease.

Methods: A MEDLINE database search using PubMed was performed using the keywords: “dry eye disease/syndrome”, “aqueous deficient/deficiency dry eye disease”, “lacrimal gland” and “Sjogren’s syndrome”. After scanning through 750 relevant abstracts, 73 eligible articles published in the English language from 2016 to 2021 were included in the review.

Results: Histopathological and ultrastructural studies have revealed new insights into the pathogenesis of cicatrizing conjunctivitis-induced aqueous deficiency, where the lacrimal gland acini remain uninvolved and retain their secretory property, while significant ultrastructural changes in the gland have been observed. Recent advances in diagnosis include the techniques of direct clinical assessment of the lacrimal gland morphology and secretion, tear film osmolality, tear film lysozyme and lactoferrin levels, tear film interferometry and lacrimal gland confocal microscopy. Developments in the treatment of aqueous deficiency dry eye disease, apart from the nanoparticle-based tear substitutes, include secretagogues like diquafosol tetrasodium and rebamipide, anti-inflammatory topical agents like nanomicellar form of cyclosporine and lifitegrast, scleral contact lenses, neurostimulation, and acupuncture for increasing the amount of tear production, minor salivary gland transplantation, faecal microbial transplantation, lacrimal gland regeneration and mesenchymal stem cell therapy.

Conclusions: Significant advances in the understanding, diagnosis and management of lacrimal gland insufficiency and its role in aqueous deficiency dry eye disease have taken place within the second half of the last decade. Of which, translational breakthroughs in terms of newer drug formulations and regenerative medicine are most promising.

ARTICLE HISTORY

Received 9 February 2022
Revised 10 February 2022
Accepted 15 February 2022

KEYWORDS

Aqueous deficiency dry eye disease; Dry eye disease; Lacrimal gland; Lacrimal gland regeneration; Minor salivary gland transplantation

Introduction

Dry eye disease (DED) is a major cause of chronic ocular morbidity, affecting millions of individuals worldwide.^{1,2} It is estimated that about 40% of the Indian urban population or close to 270 million people would be affected by DED in the next decade.¹ The second Tear Film and Ocular Surface Society dry eye workshop (TFOS DEWS II) has defined DED as “a multifactorial disease of the ocular surface characterized by a loss of homeostasis of the tear film, accompanied by ocular symptoms, in which tear film instability and hyperosmolality, ocular surface inflammation and damage and neurosensory abnormalities play an etiological role”.² Old age, female sex, medication use, dry environment or extended screen time on electronic devices are the known risk factors which lead to the development of DED.^{1,3} DED has been classified into evaporative dry eye (EDE), and aqueous-deficient dry eye (ADDE) disease, based on Meibomian or lacrimal gland involvement, respectively, although the DEWS II guidelines suggest that these variants may lie along the continuum of the disease.² Irrespective of the underlying cause, instability of the tear

film leads to ocular surface inflammation which is responsible for the clinical features characteristic of the disease.⁴ Among the two broad categories, ADDE due to lacrimal gland insufficiency accounts for about a third of all cases of DED¹ but is the more serious and potentially blinding condition, since it can cause severe ocular surface inflammation and chronic keratopathy. This review summarizes the significant advances in the understanding, diagnosis and management of lacrimal gland insufficiency and its role in ADDE that have taken place within the second half of the last decade.

Methodology

A MEDLINE search using PubMed was performed with the keywords: “dry eye”, “dry eye disease”, “dry eye syndrome”, “aqueous deficient dry eye”, “aqueous deficiency dry eye”, “Sjögren’s syndrome” and “lacrimal gland” for articles published between 2016 and 2021. The search revealed a total of 20,945 articles. Articles not having abstracts, not in the English

language, not relevant to the topic of interest and repetitive articles were excluded. A total of 750 abstracts were scanned, and 73 relevant articles were included in the review.

Etiology and pathogenesis of lacrimal gland insufficiency in ADDE

Aqueous deficiency dry eye disease can occur if the lacrimal gland is congenitally absent, developed incompletely or there is an acquired damage to the gland and its periductular tissue due to the ongoing inflammation. Depending on the underlying etiology, ADDE is classified into Sjögren's and non-Sjögren's type (Figure 1).²

Sjögren's Syndrome-Associated ADDE

Sjögren's syndrome (SS) is a systemic autoimmune condition wherein lymphocytic infiltration of the exocrine glands including the lacrimal gland occurs, leading to persisting inflammation and symptoms of dry eye and dry mouth, besides various other systemic manifestations such as arthritis. For its diagnosis, the American-European Consensus Group 2012-revised classification criteria are the most commonly used (Figure 1).⁵⁻⁷ The pathogenesis of SS is multifactorial. Genetic susceptibility, female sex, viral infections, overactivation of B cells, role of Th1 and Th17 cells and increase in the pro-inflammatory cytokines such as matrix metalloproteinases (MMPs) are known to play a role in its pathogenesis.⁸ Role of ocular and gut microbiome in the pathophysiology of SS has been recently reported. The gut microbiota is closely related to the immune system during development.⁹ It maintains

a balance between the immune responses and is said to play a crucial role of immune regulation. Any dysbiosis in the gut microbiome may lead to auto-immune diseases such as SS.¹⁰⁻¹²

Non-Sjögren's Type of ADDE

This can further be classified into congenital and acquired types (Figure 1). Congenital alacrima is a rare condition of absence of tear production from birth. There is abnormal or deficient development of the lacrimal gland or its ductal tissues. It is associated with a multitude of systemic disorders and is an important and rare cause of paediatric dry eye disease.¹³ Presentation can vary from congenital absence or hyposecretion of tears, developmental delay, to severe multi-systemic involvement.¹⁴⁻¹⁶

Acquired causes of non-Sjögren's-type ADDE include immune-mediated conditions such as Stevens-Johnson syndrome (SJS), mucous membrane pemphigoid (MMP) and graft versus host disease (GVHD).^{3,17,18} Stevens-Johnson syndrome is a chronic mucocutaneous blistering disease, which is triggered by drugs or viral infections.¹⁹⁻²² It is said to occur as a result of dysregulated immune response, mainly pertaining to innate immunity.^{23,24} It is a cytokine-mediated response, which leads to the deposition of neutrophils in the conjunctiva of these patients.²⁵ In both SJS and MMP, there is chronic inflammation of the ocular surface, which if not treated in time, leads to cicatrizing conjunctivitis, ultimately causing ADDE. Graft versus host disease (GVHD) is an adverse immunological event following allogeneic hematopoietic stem cell transplantation. It includes a T cell-mediated immune response which causes infiltration and inflammation of ocular structures such as the lacrimal glands and conjunctiva, leading to ADDE.^{26,27} Other acquired causes




Figure 1. Classification of aqueous deficiency dry eye disease into Sjögren's and non-Sjögren's dry eye disease. Non-Sjögren's ADDE can further be classified into congenital form such as in cases of alacrimia and acquired form seen in various auto-immune diseases and other conditions such as chemical/thermal trauma, radiation and iatrogenic damage.

of non-Sjögren's-type ADDE include ocular chemical burns, iatrogenic causes such as kerato-refractive surgeries, long-term use of topical and systemic medications and radiation therapy for head and neck cancers. Some of these conditions, like radiation therapy, cause direct damage to the lacrimal gland and the ocular surface, while others, like topical anti-glaucoma medications, cause chronic inflammation and fibrosis of the ocular surface leading to cicatrising changes.

Normal anatomy of healthy lacrimal glands

Gross and Applied Anatomy

The main lacrimal gland is located in the supero-temporal orbit, behind the orbital rim, in the lacrimal gland fossa. It extends from the lateral edge of superior rectus muscle to the frontozygomatic suture and posteriorly upto the posterior aspect of the globe.^{28,29} The lacrimal gland is seen as an almond-shaped, concavo-convex bilobed structure, divided into two lobes by the lateral horn of levator aponeurosis.²⁹⁻³² The larger lobe being the orbital lobe constitutes about 60–70% of the gland, weighs about 600–700 g and is located behind the orbital septum.^{28,31-34} Hence, the orbital lobe cannot be clinically visualized. The smaller palpebral lobe, located beneath the levator aponeurosis, is relatively flat and can be examined clinically in the superior fornix by elevation or eversion of the lateral part of the upper eyelid. It appears as a pale, pinkish, lobulated structure.^{28,32,35} Although lacrimal gland secretions are thought to mainly contribute to reflex tearing, recent reports suggest that the gland has an important role in basal tear secretion as well.³⁶⁻³⁸

There are two types of accessory lacrimal glands, the glands of Wolfring (along the tarsal border) and Krause (in the substantia propria of fornices). Glands of Wolfring are more in number, present mainly in the upper eyelid, while the Gland of Krause are larger in size. These glands, like the main lacrimal gland, are serous in nature and of eccrine variety. Parasympathetic nerves and hormones stimulate the accessory lacrimal glands, which contribute to both basal and reflex secretions.³⁹

Microscopic Structure

Microscopically, the lacrimal gland is formed of acinar epithelial cells forming lobules draining into intralobular ducts. The ducts are lined by tubular columnar cell, surrounded by myoepithelial cells.^{39,40} The intralobular ducts further merge into interlobular ducts which are lined by cuboidal cells and open in the superior and inferior fornix as 3–12 excretory ductules.³⁹⁻⁴² Recently, the role of columnar cells of ducts in the secretion of tears have also been found; however, this still requires further studies for confirmation.⁴³ The intervening interstitial connective tissue contains lymphocytes, plasma cells, mucosa-associated lymphoid tissue, fibroblasts, blood vessels and nerve fibres.^{29,32,39} Accessory lacrimal glands are similar to the main lacrimal gland microscopically.

Pathological and ultrastructural changes in lacrimal gland insufficiency

Sjögren's-Associated ADDE

In SS, the lacrimal gland undergoes progressive fibrosis and hence gradually loses its secretory function. The histological examination of the lacrimal gland has revealed acinar dilation, interstitial fibrosis, with inflammatory aggregates and atrophy with fibrosis of the gland.⁴⁴

Cicatrising Conjunctivitis-Associated ADDE

In chronic cicatrising conjunctivitis such as SJS, MMP, and chemical burns, there is an ongoing sub-conjunctival fibrosis, leading to the destruction of the lacrimal ductules, both in the palpebral and orbital lobes, which on histological examination is seen as peri-ductular, interlobar and interlobular lymphocytic infiltration and fibrosis. The acini, particularly in the orbital lobe, may remain uninvolved and retain their secretory property.^{44,45} These changes are in stark contrast to the diffuse glandular atrophy and inflammatory aggregation that is seen in SS. However, despite the relative preservation of acinar structure on histopathology, significant ultrastructural changes have also been observed, showing disturbance in productivity and secretory activity of the glands in cicatrising disease like SJS.⁴⁶

Diagnostic modalities of lacrimal gland insufficiency in ADDE

The traditional standard basic tests for the diagnosis of DED include ocular surface disease index score (OSDI), Schirmer's test, tear film height and tear film break up time (TBUT). Since this review focuses mainly on ADDE due to the lacrimal gland insufficiency, for the simplicity of discussion, the recent advances in diagnostic modalities have been classified into direct assessment of the lacrimal gland and indirect assessment of lacrimal gland insufficiency by tests based on tear film parameters.

Direct Clinical Assessment of Lacrimal Gland and Its Function

Morphometry

Clinically, on slit lamp bio-microscopy, one can assess the lacrimal gland by asking the patient to look inferonasally and lifting the upper eyelid to observe the contour, size and appearance of the palpebral lobe of the lacrimal gland. Recently a study by Singh et al. showed that in normal individuals and those with EDE, the palpebral lobe is bilaterally symmetrical, having a convex pinkish appearance. However, in cases of ADDE, the palpebral lobe size is bilaterally asymmetrical, appears flatter than normal and loses its convexity. Further, in SJS, the palpebral lobe appears smaller and there is a presence of symblepharon along with scarring.³⁵ Therefore, clinical assessment of the palpebral lobe morphology can not

only help us in diagnosing ADDE but can also help us in differentiating between the underlying etiological causes of SS and SJS.

Tear Flow Rate or Tear Secretion

Direct assessment of tear secretion (DATS) from the excretory ducts on the surface of the palpebral lobe of lacrimal lobe was first described in 1986.⁴¹ It was developed on the concept of the Seidel's test, wherein a point of leak appears as a stained flow of fluid from the focal source. Similarly, simply touching the surface of the palpebral lobe with a fluorescein strip shows the appearance of the fluid at the duct openings and its rate of flow and washout.⁴⁵ Recent studies have shown its utility in diagnosing the subtype of ADDE. Singh et al. showed the difference in the number of ductular opening and flow rate between patients of cicatrising conjunctivitis, SS and normal healthy individuals. The patients of cicatrising conjunctivitis had reduced number of ductular opening and tear flow rate as compared to the patients of SS.³⁵ Also the latency for the initiation of the tear flow and the rate of flow can be assessed. The time lag is maximum in the cicatrising group followed by the SS group. The advantage of both lacrimal gland morphometry and DATS is that they can be performed in the clinic and can aid clinicians to diagnose the subtype of DED.

Indirect Assessment of Lacrimal Gland Function (Tear Film-Based Tests)

Tear Film Breakup (TBFU) Patterns

Due to the ongoing inflammation and subsequent injury to the epithelial cells and microvilli, ocular surface shows positive staining after the topical application of fluorescein. In 2017, Yokoi et al. described the characteristic tear film breakup patterns in the various subtypes of DED, thereby proving an aid for diagnosis. They described five main patterns. As per their report, an area break of ocular surface staining points towards severe grade of ADDE, a line and dimple break indicate mild-to-moderate grade of ADDE, while a spot break determines decreased wettability of the surface.⁴⁷ Based on this pattern, DED can be classified into three independent subtypes, ADDE, decreased wettability DE (DWDE), and increased evaporation. Hence, based on the pattern, it is possible to identify which component of the tear film should be supplemented. This novel concept of a layer-by-layer diagnosis and therapy is known as "tear film-oriented diagnosis" (TFOD) and "tear film-oriented therapy" (TFOT).⁴⁸

Non-Invasive Tear Film Breakup Time (NIBUT)

Instillation of fluorescein dye is considered invasive in nature since it is associated with chances of transmission of infection; hence, a newer method known as non-invasive tear film break up time (NIBUT) was introduced.⁴⁹ It is a placido disc-based assessment of the tear film, which determines the tear film stability based on the reflection of mires from the ocular surface. The reflection gets altered when the tear film stability is

compromised.^{50,51} However, there appears to be significant difference in the values measured by TBUT and NIBUT. Average value of 3–4 NIBUT measurements provides a more accurate result, and hence the average value should be preferred over a single measurement.⁵²

Tear Meniscus Height

An objective way of assessing the tear film volume is by using the anterior segment optical coherence tomography (OCT)-based measurement of tear meniscus height (TMH). Time domain (TD-OCT) was the first OCT used for this purpose. Recent advances in spectral domain and swept source OCT have proven to be more useful, since it has the added benefit of helping in distinguishing the subgroup of DED.^{53–56} The TMH is lowest in Sjogren's syndrome, followed by non-Sjogren's ADDE.⁵⁷

Tear Strip Meniscometry

This device uses a water absorbing material similar to the Schirmer's test. It has the added benefit that it is quick, providing results within 5 seconds, and is a non-invasive method to quantify the tear volume.^{58,59}

Tear Osmolarity

In case of ADDE and EDE, due to the insufficient aqueous secretion and normal or excessive evaporation, the ocular surface remains in a chronic state of inflammation. This inflammation of ocular surface results in outpouring of inflammatory products and cytokines into the tear film. This results in increasing the tear film osmolarity. The TearLab Osmolarity System (TearLab Corporation, San Diego, California, USA) is currently the only FDA-approved commercially available in-office device for measuring tear film osmolarity.⁶⁰ It collects a sample of 50 nL of tear film and gives results within just a few seconds.⁶¹ The normal tear osmolarity has a value of 302 mOsm/L with an inter-eye difference of 8 mOsm/L.^{60,62} It is worth noting that tear osmolarity is variable depending on the time of the day, or food intake, and hence must never be considered as a static value.^{61,63}

MMP-9

Due to the evidence of increase in the pro-inflammatory cytokines such as MMP-9 in tear film of patients with DED, MMP-9 assessment has turned a useful aid for diagnosis. MMP-9 is an important endopeptidase released in the tear film during the extracellular matrix remodelling and facilitating leukocyte migration and chronic state of inflammation.^{64–66} Currently, InflammDry test (RPS, Inc., Sarasota, Florida, USA) is the only FDA-approved device for the detection of MMP-9 in patients. However, the test is non-specific for DED since the MMP-9 levels can be elevated even in other inflammatory ocular conditions such as keratitis.⁶⁷

Lactoferrin and Lysozyme

These are the normal constituents of the tear film, which are said to have a protective mechanism in the form of antibacterial, antitumor and immunomodulatory property. Lactoferrin levels in tear fluid are mainly reduced in autoimmune conditions such as SS.^{68–70}

Lacrimal Gland Imaging

CT scan, MRI scan and PET scan are some of the known modalities used for lacrimal gland imaging. Recently, the role of in vivo confocal microscopy of the lacrimal gland has been demonstrated in DED patients.⁷¹ It can assess the acinar unit density, acinar unit diameter and inflammatory cell density in both the Meibomian and lacrimal glands.⁷² This is a relatively unexplored area and there is enormous scope for further research in the field of lacrimal gland imaging.

Treatment of lacrimal gland insufficiency

Pharmacological Treatment

Lacrimal gland insufficiency manifests mainly as a reduction in the tear volume which further leads to increased tear hyperosmolarity, resulting in chronic ocular surface inflammation and disease sequelae. (Figure 2). Hence, the pharmacological treatment of ADDE includes topical tear supplements, punctal plugs, secretagogues and anti-inflammatory agents such as corticosteroids, cyclosporine and tacrolimus. There is also an important role of controlling the underlying systemic disease, particularly in immune-mediated conditions like SS and GVHD.

Tear Supplements

These have been the first line of management in patients with ADDE, regardless of the disease severity. They are known to improve the ocular surface lubrication, tear retention, tear

osmolarity and TBUT and reduce the ocular surface staining. However, there are various types of artificial tears available which differ in its constituents.^{73,74} Preservatives such as benzalkonium chloride should be avoided as it causes stinging sensation and ocular surface toxicity with long-term use. The pH of the human tear is between 6.9 and 7.5. Any compound having a higher pH causes burning sensation upon instillation. These factors further worsen the pre-existing symptoms of a patient with DED. Current recommendations are that a preservative-free compound having a lower pH should be preferred. Due to the varied compositions, the drops need to be individualized in every patient.

Secretagogues

This is a new class of drugs, which mainly acts on the receptors of the lacrimal gland and helps in increasing the tear production. This class of drugs include diquafosol tetrasodium, rebamipide, cevimeline and oral pilocarpine. Diquafosol tetrasodium is available as a 3% ophthalmic solution. It acts on P2Y2 receptors, which are found on the conjunctiva, meibomian and lacrimal gland. It increases the intracellular calcium and thereby helps in increasing the tear secretion.^{75–82} Rebamipide is a quinolinone derivative, which has a dual effect. It acts on goblet cells and increases the mucin production, thereby improving the tear film stability and epithelial regeneration. It also has an added benefit in increasing the levels of PGE2 and PGI2 and reducing the level of TNF- α , thereby preventing macrophage infiltration in the conjunctiva and reducing the amount of ocular surface inflammation and improving the tear film stability.^{82–86} Cevimeline hydrochloride is a cholinergic agonist that binds to muscarinic acetylcholine receptors, especially to the M3 receptor, producing an increase in glandular secretions. Oral pilocarpine is also known to be having similar mechanism and helps in increasing both the salivary and aqueous tear production in patients of SS.^{87–90}




Figure 2. Classification, mechanism of action and the effect on ocular surface of various topical drugs used in the treatment of ADDE. LG, lacrimal gland; PG, prostaglandins; TNF, tumor necrosis factor; M3, muscarinic receptor and ICAM-1, intercellular adhesion molecule.

Anti-inflammatory Agents

The pathogenesis of DED⁹⁰ and molecular interaction of various anti-inflammatory agents have been illustrated in **Figures 3 and 4**. Topical corticosteroids help in reducing the ocular surface inflammation by interrupting the inflammatory cycle. They mainly suppress the cytokine activation and leukocyte infiltration, thereby controlling the ongoing inflammation. However, in view of the various side effects, their continued long-term use is not recommended. Topical cyclosporine acts by inhibiting the calcineurin–phosphatase pathway via cyclophilin. It is also said to increase the goblet cell density and increase tear production. It is said to inhibit T cell activation and decrease apoptosis of epithelial and goblet cells.⁹¹ FDA has approved 0.05% cyclosporine emulsion (Restasis, Allergan Inc. Irvine, CA, USA), which has anti-inflammatory property and in addition is also known to increase the tear production.^{92–94} It is said to give better results than the use of lubricants alone in the treatment of DED.⁹⁵

Cequa (CEQUA, Sun Pharmaceutical Industries, Inc., Cranbury, NJ), a preservative-free nanomicellar form of cyclosporine, has also recently been approved by FDA. It has the advantage of better drug penetration and rapid onset of action.⁹⁶ Lifitegrast (Xiidra™, Novartis, USA) was approved in 2016 to treat DED.⁹⁷ It is a nanoparticle, which prevents the inflammatory cascade by inhibiting the binding of ICAM-1 to any integrins. It inhibits binding of leukocyte-associated antigen-1 (LFA-1) on T cells to its ligand intercellular adhesion molecule-1 (ICAM1) on antigen presenting, epithelial and vascular endothelial cells and prevents the formation of the immunological synapse that is required for full T cell activation. It is available for topical use as 5% ophthalmic solution.^{98–102} Metformin, a first-line drug used

in type 2 diabetes, has also emerged as an effective anti-inflammatory drug useful in patients of SS.¹⁰³ Multiple other treatment modalities to promote body's natural anti-inflammatory mechanism are described which include oral supplementation with Omega-3 polyunsaturated fatty acids (PUFAs), curcumin and anti-oxidants, which suppress interleukins and the inflammatory cascade.^{104,105}

Contact Lenses

Soft bandage contact lenses are known to improve the symptoms of patients with DED. Many recent studies have shown the role of scleral and PROSE (Prosthetic Replacement of Ocular Surface Ecosystem) lenses in terms of improving symptoms and visual acuity, particularly in cases with severe ADDE in patients with chronic ocular sequelae of SJS.^{106,107} These lenses are filled with fluid, which provide lubrication to the ocular surface. It improves the quality of vision as well as improved the symptoms of patients. However, the limitation is its availability, expense and need of training for its application and removal.^{108–110}

Neuronal Stimulation Therapy

Acupuncture is a form of Chinese traditional medical therapy, wherein needles are applied at specific points on the body for neuronal stimulation. It has shown to be effective in treating DED. However, further studies are needed to gauge its efficacy.^{111–113} Role of anterior ethmoidal nerve stimulation being better than lacrimal nerve stimulation for increasing the aqueous production is also known.¹¹⁴ Intranasal Tear Neurostimulator (TrueTear, Allergan plc) is a similar device




Figure 3. Mechanism of action of the anti-inflammatory agents such as steroids, cyclosporine, and Lifitegrast used in ADDE. TNF- α , tumor necrosis factor alpha; IL-, interleukin; JNK, c-Jun N-terminal kinase; ERK, extracellular signal-related kinase; APC, antigen presenting cells; MMP, matrix metalloproteinase; ICAM-1, intercellular adhesion molecule 1; LFA-1, lymphocyte function-associated antigen 1; MHCII, major histocompatibility complex II; TCR, T cell receptor; IFN- γ , interferon gamma. Adapted from Periman et al.⁹⁰




Figure 4. Site of action during the pathophysiology of ADED-induced inflammatory cascade and the molecular interaction of the various anti-inflammatory agents used in ADED. MAP-K, mitogen-activated protein kinase; JNK, c-Jun N-terminal kinase; ERK, extracellular signal-related kinase; TNF- α , tumor necrosis factor alpha; IL-, interleukin; CCL, CXCL- chemokines; ICAM-1, intercellular adhesion molecule 1; LFA-1, lymphocyte function-associated antigen 1; TGF- β , transforming growth factor beta; IFN- γ , interferon gamma and MMP, matrix metalloproteinase. Adapted from Periman et al.⁹⁰

designed to deliver microcurrents to the nasal cavity, stimulating the nasolacrimal pathway. It has recently received FDA approval to temporarily increase tear production. It has shown significant improvements in ocular dryness and discomfort compared along with a good safety profile and hence appears to be a promising new management strategy for these patients.¹¹⁵ The iTEAR®100 device is a similar device; however, unlike TrueTear, this stimulates the anterior ethmoidal nerves at the tip of the nose. Further clinical trials for its efficacy and safety are awaited.

Surgical Management

Minor Salivary Gland Transplantation (MSGT)

This surgical option has shown to be useful and effective in patients with certain types of severe ADDE. The surgery was first described in 2008.¹¹⁶ The graft tissue is obtained from under the oral labial mucosa and transplanted onto the posterior eyelid lamella. It increases the tear secretions in patients of non-Sjögren's type of ADDE.^{117–119} Recently, a modified technique has also been described, wherein the graft is taken en-bloc, retaining the overlying mucosa and submucosal tissue to act as a barrier against post-operative sub-conjunctival fibrosis. In the modified technique the graft is placed over the superior bulbar surface of the eye.¹¹⁹ Certain clinical tests have been described to help selecting the donor site and in assessing the functionality of the transplanted glands post-operatively.^{120–122} The test is similar to the DATS test, wherein a fluorescein strip is applied onto the labial mucosa, and based on the number of excretory duct openings, the donor site can be selected.^{120,121,123,124} Similarly, the test can be used to objectively assess the number of functional openings and secretory

rate in the transplanted graft on the ocular surface.¹²² It is important to remember that MSGT is not recommended in Sjögren's type of DED because the underlying disease process affects both the salivary and the lacrimal glands.

Mesenchymal Stem Cell Transplantation

Mesenchymal/stromal stem cells (MSCs) possess immunomodulatory properties and multilineage differentiation abilities. It acts by suppressing both local and systemic inflammation and inhibiting the activation and proliferation of T cells.¹²⁵ It has a potential role in treating patients with severe or refractory DED. The literature also shows its role in treating patients with steroid-resistant GVHD-induced DED.¹²⁶ Injection of MSCs derived from bone marrow or adipose tissue into the lacrimal gland has also shown efficacy.¹²⁷ Further results of randomized control trials from the same group is still awaited.

Lacrimal Gland Regeneration

Lacrimal gland regeneration using MSC, in addition to immunomodulation, is one of the main focuses of current research and is gaining popularity.^{128–131} Further advances include the development of bio-engineered lacrimal glands, which can produce tears having the protein content similar to human tears.^{132,133} Researchers have also described the establishment of 3D organoids of the human lacrimal gland, which can engraft and produce mature tear products after orthotopic transplantation in mice.¹³⁴ Organoids can be expanded over multiple months and recapitulate morphological and transcriptional features of lacrimal ducts. These exciting and novel approaches are awaiting clinical translation.

Faecal Microbial Transplant

The role and interaction of ocular-gut-microbiome have been found to have a major role to play in the pathophysiology of ocular surface diseases, especially in auto-immune conditions such as SS. Hence, supplementation with probiotics or metabolites of commensal bacteria could have a major role in future treatment directives for dry eye diseases.^{135,136} Faecal microbial transplant showed efficacy in patients of immune-mediated DED; however, further trials would be needed to prove its efficacy and safety.¹³⁷

Summary

Aqueous deficiency dry eye disease, although the less common form of DED as compared to EDE, is the more serious clinical sub-type that causes chronic ocular morbidity and is potentially blinding. The underlying cause of ADDE is the dysfunction of the lacrimal glands, which is mostly acquired and due to auto-immune diseases that either directly target the gland, like in SS, or the periglandular tissue and the secretory ductules, like in SJS. Recent advancements have improved our understanding of the pathological processes affecting the lacrimal gland, particularly in cicatrising non-Sjögren's type of ADDE. Developments in diagnostics include both simple clinical diagnostic techniques like direct assessment of lacrimal gland morphology and secretion, which can be performed in office settings without any specialized equipment and sophisticated, yet elegant and objective tear-film-based tests like osmolarity measurements. Introduction of newer therapeutic agents, such as secretagogues like diquafosol, anti-inflammatory agents like preservative free nanomicellar form of cyclosporine and lifitegrast, have expanded the treatment options and are likely to bring more relief to patients. Lastly, the role of the gut microbiome is being explored through faecal microbial transplantation in SS-associated ADDE, while the final frontier of lacrimal gland rejuvenation, repair or replacement is also being challenged through MSC therapy and bioengineered glands and organoids.

Disclosure statement

No potential conflict of interest was reported by the author(s).

Funding

This work was supported by the Hyderabad Eye Research Foundation (HERF), Hyderabad, Telangana, India. [Intra-mural grant].

ORCID

Mariya B Doctor  <http://orcid.org/0000-0003-3956-5984>
Sayan Basu  <http://orcid.org/0000-0001-5030-5003>

References

1. Donthineni PR, Kammari P, Shanbhag SS, Singh V, Das AV, Basu S. Incidence, demographics, types and risk factors of dry eye disease in India: electronic medical records driven big data analytics report I. *Ocul Surf*. 2019;17(2):250–256. doi:10.1016/j.jtos.2019.02.007.
2. Craig JP, Nichols KK, Akpek EK, et al. TFOS DEWS II definition and classification report. *Ocul Surf*. 2017;15(3):276–283. doi:10.1016/j.jtos.2017.05.008.
3. Craig JP, Nelson JD, Azar DT, et al. TFOS DEWS II report executive summary. *Ocul Surf*. 2017;15(4):802–812. doi:10.1016/j.jtos.2017.08.003.
4. Hessen M, Akpek EK. Dry eye: an inflammatory ocular disease. *J Ophthalmic Vis Res*. 2014;9:240–250.
5. Brito-Zerón P, Ramos-Casals M. Advances in the understanding and treatment of systemic complications in Sjögren's syndrome. *Curr Opin Rheumatol*. 2014;26(5):520–527. doi:10.1097/BOR.0000000000000096.
6. Wang J, Zhou L, Liu B. Update on disease pathogenesis, diagnosis, and management of primary Sjögren's syndrome. *Int J Rheum Dis*. 2020;23(6):723–727. doi:10.1111/1756-185X.13839.
7. Vitali C, Bombardieri S, Jonsson R, et al. Classification criteria for Sjögren's syndrome: a revised version of the European criteria proposed by the American-European consensus group. *Ann Rheum Dis*. 2002;61(6):554–558. doi:10.1136/ard.61.6.554.
8. Tian Y, Yang H, Liu N, Li Y, Chen J. Advances in pathogenesis of Sjögren's syndrome. *J Immunol Res*. 2021;2021:5928232. doi:10.1155/2021/5928232.
9. Lynch SV, Pedersen O. The human intestinal microbiome in health and disease. *N Engl J Med*. 2016;375(24):2369–2379. doi:10.1056/NEJMra1600266.
10. Zhong D, Wu C, Zeng X, Wang Q. The role of gut microbiota in the pathogenesis of rheumatic diseases. *Clin Rheumatol*. 2018;37(1):25–34. doi:10.1007/s10067-017-3821-4.
11. de Paiva CS, Jones DB, Stern ME, et al. Altered mucosal microbiome diversity and disease severity in Sjögren syndrome. *Sci Rep*. 2016;6(1):23561. doi:10.1038/srep23561.
12. Mandl T, Marsal J, Olsson P, Ohlsson B, Andréasson K. Severe intestinal dysbiosis is prevalent in primary Sjögren's syndrome and is associated with systemic disease activity. *Arthritis Res Ther*. 2017;19(1):237. doi:10.1186/s13075-017-1446-2.
13. Donthineni PR, Das AV, Basu S. Dry eye disease in children and adolescents in India. *Ocul Surf*. 2020 Oct;18(4):777–782. doi:10.1016/j.jtos.2020.07.019.
14. Adams J, Schaaf CP. Diagnosis and genetics of alacrima. *Clin Genet*. 2018;94(1):54–60. doi:10.1111/cge.13173.
15. Zhao Z, Allen RC. Congenital alacrima. *Orbit*. 2021;41(2):162–169. doi:10.1080/01676830.2021.1974057.
16. Gupta N, Farooqui JH, Agni M, Kumar A, Sharma M, Mathur U. Alacrima, a rare cause of pediatric dry eye. *J Aapos*. 2018;22(3):233–235. doi:10.1016/j.jaapos.2017.11.003.
17. Fraunfelder FT, Sciubba JJ, Mathers WD. The role of medications in causing dry eye. *J Ophthalmol*. 2012;2012:285851. doi:10.1155/2012/285851.
18. Singh S. Congenital alacrimia with lacrimal gland hypoplasia. *Orbit*. 2021;1. doi:10.1080/01676830.2021.1942503.
19. Bianchini JR, Macaraeg PV Jr., Lasagna L, et al. Drugs as etiologic factors in the Stevens-Johnson syndrome. *Am J Med*. 1968;44(3):390–405. doi:10.1016/0002-9343(68)90110-1.
20. Patel TK, Barvaliya MJ, Sharma D, Tripathi C. A systematic review of the drug-induced Stevens-Johnson syndrome and toxic epidermal necrolysis in Indian population. *Indian J Dermatol Venereol Leprol*. 2013;79(3):389–398. doi:10.4103/0378-6323.110749.
21. Mockenhaupt M, Viboud C, Dunant A, et al. Stevens-Johnson syndrome and toxic epidermal necrolysis: assessment of medication risks with emphasis on recently marketed drugs. The EuroSCAR-study. *J Invest Dermatol*. 2008;128(1):35–44. doi:10.1038/sj.jid.5701033.
22. McCormack JG. Mycoplasma pneumoniae and the erythema multiforme-Stevens Johnson syndrome. *J Infect*. 1981;3(1):32–36. doi:10.1016/S0163-4453(81)92236-2.

23. Schwartz RA, McDonough PH, Lee BW. Toxic epidermal necrolysis: part I. Introduction, history, classification, clinical features, systemic manifestations, etiology, and immunopathogenesis. *J Am Acad Dermatol.* 2013;69(2):173.e1-13; quiz 85-6. doi:10.1016/j.jaad.2013.05.003.
24. Williams GP, Tomlins PJ, Denniston AK, et al. Elevation of conjunctival epithelial CD45INTCD11b⁺CD16⁺CD14⁺ neutrophils in ocular Stevens- Johnson syndrome and toxic epidermal necrolysis. *Invest Ophthalmol Vis Sci.* 2013;54(7):4578-4585. doi:10.1167/iovs.13-11859.
25. Pavlos R, Mallal S, Phillips E. HLA and pharmacogenetics of drug hypersensitivity. *Pharmacogenomics.* 2012;13(11):1285-1306. doi:10.2217/pgs.12.108.
26. Carreno-Galeano JT, Dohlman TH, Kim S, Yin J, Dana R. A review of ocular graft-versus-host disease: pathophysiology, clinical presentation and management. *Ocul Immunol Inflammation.* 2021;29(6):1190-1199. doi:10.1080/09273948.2021.1939390.
27. Tung CI. Current approaches to treatment of ocular graft-versus-host disease. *Int Ophthalmol Clin.* 2017;57(2):65-88. doi:10.1097/HIO.000000000000167.
28. Lorber M. Gross characteristics of normal human lacrimal glands. *Ocul Surf.* 2007;5(1):13-22. doi:10.1016/S1542-0124(12)70049-6.
29. Eugene WR. *Wolff's Anatomy of the Eye and Orbit.* 7 ed. Philadelphia PA: WB Saunders; 1976. 222.
30. Mafee MF, Haik BG. Lacrimal gland and fossa lesions: role of computed tomography. *Radiol Clin North Am.* 1987;25:767-779.
31. Forrester JVDA, McMenamin P, Lee WR. *The Eye: Basic Sciences in Practice.* 1. Philadelphia PA: WB Saunders; 1996.
32. Snell RSLM. *Clinical Anatomy of the Eye.* 2. Malden MA: Blackwell Science; 1998.
33. S T. *Ocular Anatomy and physiology.* Boston MA: Blackwell scientific; 1993.
34. Forrester JVDA, McMenamin PG, Lee WR. *The Eye. Basic Sciences in Practice.* 2. Edinburgh: WB Saunders; 2002.
35. Singh S, Shanbhag SS, Basu S. Palpebral lobe of the human lacrimal gland: morphometric analysis in normal versus dry eyes. *Br J Ophthalmol.* 2021;105(10):1352-1357. doi:10.1136/bjophthalmol-2020-316929.
36. Jones LT. The lacrimal secretory system and its treatment. *Am J Ophthalmol.* 1966;62(1):47-60. doi:10.1016/0002-9394(66)91676-X.
37. Jordan A, Baum J. Basic tear flow Does it exist? *Ophthalmol.* 1980;87(9):920-930. doi:10.1016/S0161-6420(80)35143-9.
38. Singh S, Basu S. Effect of topical anesthesia on the secretory activity of the main lacrimal Gland. *Cornea.* 2020;39(10):e24-e5. doi:10.1097/ICO.0000000000002410.
39. Singh S, Basu S. The human lacrimal gland: historical perspectives, current understanding, and recent advances. *Curr Eye Res.* 2020;45(10):1188-1198. doi:10.1080/02713683.2020.1774065.
40. Kim EC, Doh SH, Chung SY, et al. Direct visualization of aqueous tear secretion from lacrimal gland. *Acta Ophthalmol.* 2017;95(4):e314-e22. doi:10.1111/aos.13335.
41. Bron AJ. Lacrimal streams: the demonstration of human lacrimal fluid secretion and the lacrimal ductules. *Br J Ophthalmol.* 1986;70(4):241-245. doi:10.1136/bjo.70.4.241.
42. Obata H. Anatomy and histopathology of the human lacrimal gland. *Cornea.* 2006;25(10 Suppl 1):S82-9. doi:10.1097/01.ico.0000247220.18295.d3.
43. Tóth-Molnár E, Ding C. New insight into lacrimal gland function: role of the duct epithelium in tear secretion. *Ocul Surf.* 2020;18(4):595-603. doi:10.1016/j.jtos.2020.07.002.
44. Singh S, Narang P, Vashist U, Mittal V. Histomorphological changes in lachrymal glands of patients with chronic Stevens-Johnson Syndrome. *Cornea.* 2019;38(9):e39-e40. doi:10.1097/ICO.0000000000002037.
45. Singh S, Mishra DK, Shanbhag S, et al. Lacrimal gland involvement in severe dry eyes after Stevens-Johnson Syndrome. *Ophthalmology.* 2021;128(4):621-624. doi:10.1016/j.opthta.2020.08.016.
46. Singh S, Basu S. Ultrastructural study of the lacrimal glands in severe dry eye disease following Stevens-Johnson syndrome. *Ocul Surf.* 2022;23:204-206. doi:10.1016/j.jtos.2021.10.005.
47. Yokoi N, Georgiev GA, Kato H, et al. Classification of fluorescein breakup patterns: a Novel method of differential diagnosis for dry eye. *Am J Ophthalmol.* 2017;180:72-85. doi:10.1016/j.ajo.2017.05.022.
48. Yokoi N, Georgiev GA. Tear film-oriented diagnosis and tear film-oriented therapy for dry eye based on tear film dynamics. *Invest Ophthalmol Vis Sci.* 2018;59(14):DES13-DES22. doi:10.1167/iovs.17-23700.
49. Hong J, Sun X, Wei A, et al. Assessment of tear film stability in dry eye with a newly developed keratograph. *Cornea.* 2013;32(5):716-721. doi:10.1097/ICO.0b013e3182714425.
50. Fuller DG, Potts K, Kim J. Noninvasive tear breakup times and ocular surface disease. *Optom Vis Sci.* 2013;90(10):1086-1091. doi:10.1097/OPX.0000000000000023.
51. Lee R, Yeo S, Aung HT, Tong L. Agreement of noninvasive tear break-up time measurement between Tomey RT-7000 Auto refractor-keratometer and oculus keratograph 5M. *Clin Ophthalmol.* 2016;10:1785-1790. doi:10.2147/OPTH.S110180.
52. Cho P, Douthwaite W. The relation between invasive and noninvasive tear break-up time. *Optom Vis Sci.* 1995;72(1):17-22. doi:10.1097/00006324-199501000-00004.
53. Chan HH, Zhao Y, Tun TA, Tong L. Repeatability of tear meniscus evaluation using spectral-domain Cirrus® HD-OCT and time-domain Visante® OCT. *Cont Lens Anterior Eye.* 2015;38(5):368-372. doi:10.1016/j.clae.2015.04.002.
54. Qiu X, Gong L, Sun X, Jin H. Age-related variations of human tear meniscus and diagnosis of dry eye with Fourier-domain anterior segment optical coherence tomography. *Cornea.* 2011;30(5):543-549. doi:10.1097/ICO.0b013e3181fb84ea.
55. Fukuda R, Usui T, Miyai T, Yamagami S, Amano S. Tear meniscus evaluation by anterior segment swept-source optical coherence tomography. *Am J Ophthalmol.* 2013;155(4):620-4. doi:10.1016/j.ajo.2012.11.009.
56. Akiyama R, Usui T, Yamagami S. Diagnosis of dry eye by tear meniscus measurements using anterior segment swept source optical coherence tomography. *Cornea.* 2015;34(Supplement 11):S115-20. doi:10.1097/ICO.0000000000000583.
57. Qiu X, Gong L, Lu Y, Jin H, Robitaille M. The diagnostic significance of Fourier-domain optical coherence tomography in Sjögren syndrome, aqueous tear deficiency and lipid tear deficiency patients. *Acta Ophthalmol.* 2012;90(5):e359-66. doi:10.1111/j.1755-3768.2012.02413.x.
58. Dogru M, Ishida K, Matsumoto Y, et al. Strip meniscometry: a new and simple method of tear meniscus evaluation. *Invest Ophthalmol Vis Sci.* 2006;47(5):901-1895. doi:10.1167/iovs.05-0802.
59. Ibrahim OM, Dogru M, Ward SK, et al. The efficacy, sensitivity, and specificity of strip meniscometry in conjunction with tear function tests in the assessment of tear meniscus. *Invest Ophthalmol Vis Sci.* 2011;52(5):2194-2198. doi:10.1167/iovs.10-5986.
60. Szalai E, Berta A, Szekanecz Z, Szűcs G, Módos L Jr. Evaluation of tear osmolality in non-Sjögren and Sjögren syndrome dry eye patients with the TearLab system. *Cornea.* 2012;31(8):867-871. doi:10.1097/ICO.0b013e3182532047.
61. Versura P, Profazio V, Campos EC. Performance of tear osmolality compared to previous diagnostic tests for dry eye diseases. *Curr Eye Res.* 2010;35(7):553-564. doi:10.3109/02713683.2010.484557.
62. Lemp MA, Bron AJ, Baudouin C, et al. Tear osmolality in the diagnosis and management of dry eye disease. *Am J Ophthalmol.* 2011;151(5):792-8.e1. doi:10.1016/j.ajo.2010.10.032.
63. Thulasi P, Djalilian AR. Update in current diagnostics and therapeutics of dry eye disease. *Ophthalmology.* 2017;124(11s):S27-s33. doi:10.1016/j.opthta.2017.07.022.

64. Lanza NL, McClellan AL, Batawi H, et al. Dry Eye Profiles in patients with a positive elevated surface matrix metalloproteinase 9 point-of-care test versus negative patients. *Ocul Surf.* 2016;14(2):216–223. doi:10.1016/j.jtos.2015.12.007.
65. Messmer EM, von Lindenfels V, Garbe A, Kampik A. Matrix Metalloproteinase 9 testing in dry eye disease using a commercially available point-of-care immunoassay. *Ophthalmology.* 2016;123(11):2300–2308. doi:10.1016/j.ophtha.2016.07.028.
66. Sivak JM, Fini ME. MMPs in the eye: emerging roles for matrix metalloproteinases in ocular physiology. *Prog Retin Eye Res.* 2002;21(1):1–14. doi:10.1016/S1350-9462(01)00015-5.
67. Pinto-Fraga J, Enríquez-de-Salamanca A, Calonge M, et al. Severity, therapeutic, and activity tear biomarkers in dry eye disease: an analysis from a phase III clinical trial. *Ocul Surf.* 2018;16(3):368–376. doi:10.1016/j.jtos.2018.05.001.
68. Flanagan JL, Willcox MD. Role of lactoferrin in the tear film. *Biochimie.* 2009;91(1):35–43. doi:10.1016/j.biochi.2008.07.007.
69. Danjo Y, Lee M, Horimoto K, Hamano T. Ocular surface damage and tear lactoferrin in dry eye syndrome. *Acta Ophthalmol (Copenh).* 1994;72(4):433–437. doi:10.1111/j.1755-3768.1994.tb02791.x.
70. Sonobe H, Ogawa Y, Yamada K, et al. A novel and innovative paper-based analytical device for assessing tear lactoferrin of dry eye patients. *Ocul Surf.* 2019;17(1):160–166. doi:10.1016/j.jtos.2018.11.001.
71. Kojima T, Dogru M, Kawashima M, Nakamura S, Tsubota K. Advances in the diagnosis and treatment of dry eye. *Prog Retin Eye Res.* 2020;78:100842. doi:10.1016/j.preteyeres.2020.100842.
72. Matsumoto Y, Ibrahim OMA. Application of in vivo confocal microscopy in dry eye disease. *Invest Ophthalmol Vis Sci.* 2018;59(14):Des41–des7. doi:10.1167/jovs.17-23602.
73. Lemp MA. Management of dry eye disease. *Am J Manag Care.* 2008;14:S88–101.
74. Pucker AD, Ng SM, Nichols JJ. Over the counter (OTC) artificial tear drops for dry eye syndrome. *Cochrane Database Syst Rev.* 2016;2:CD009729. doi:10.1002/14651858.CD009729.pub2.
75. Kamiya K, Nakanishi M, Ishii R, et al. Clinical evaluation of the additive effect of diquafosol tetrasodium on sodium hyaluronate monotherapy in patients with dry eye syndrome: a prospective, randomized, multicenter study. *Eye (Lond).* 2012;26(10):1363–1368. doi:10.1038/eye.2012.166.
76. Keating GM. Diquafosol ophthalmic solution 3 %: a review of its use in dry eye. *Drugs.* 2015;75(8):911–922. doi:10.1007/s40265-015-0409-7.
77. Bremond-Gignac D, Gicquel JJ, Chiambaretta F. Pharmacokinetic evaluation of diquafosol tetrasodium for the treatment of Sjögren's syndrome. *Expert Opin Drug Metab Toxicol.* 2014;10(6):905–913. doi:10.1517/17425255.2014.915026.
78. Nakamura M, Imanaka T, Sakamoto A. Diquafosol ophthalmic solution for dry eye treatment. *Adv Ther.* 2012;29(7):579–589. doi:10.1007/s12325-012-0033-9.
79. Takamura E, Tsubota K, Watanabe H, Ohashi Y. A randomised, double-masked comparison study of diquafosol versus sodium hyaluronate ophthalmic solutions in dry eye patients. *Br J Ophthalmol.* 2012;96(10):1310–1315. doi:10.1136/bjophthalmol-2011-301448.
80. Koh S, Ikeda C, Takai Y, Watanabe H, Maeda N, Nishida K. Long-term results of treatment with diquafosol ophthalmic solution for aqueous-deficient dry eye. *Jpn J Ophthalmol.* 2013;57(5):440–446. doi:10.1007/s10384-013-0251-y.
81. Carracedo G, Crooke A, Guzman-Aranguez A, de Lara Mj P, Martin-Gil A, Pintor J. The role of dinucleoside polyphosphates on the ocular surface and other eye structures. *Prog Retin Eye Res.* 2016;55:182–205. doi:10.1016/j.preteyeres.2016.07.001.
82. Fujihara T, Murakami T, Fujita H, Nakamura M, Nakata K. Improvement of corneal barrier function by the P2Y(2) agonist INS365 in a rat dry eye model. *Invest Ophthalmol Vis Sci.* 2001;42:96–100.
83. Kashima T, Itakura H, Akiyama H, Kishi S. Rebamipide ophthalmic suspension for the treatment of dry eye syndrome: a critical appraisal. *Clin Ophthalmol (Auckland, NZ).* 2014;8:1003–1010. doi:10.2147/OPTH.S40798.
84. Kinoshita S, Awamura S, Oshiden K, Nakamichi N, Suzuki H, Yokoi N. Rebamipide (OPC- 12759) in the treatment of dry eye: a randomized, double-masked, multicenter, placebo-controlled phase II study. *Ophthalmology.* 2012;119(12):2471–2478. doi:10.1016/j.ophtha.2012.06.052.
85. Tajima K, Hattori T, Takahashi H, et al. Rebamipide suppresses TNF- α production and macrophage infiltration in the conjunctiva. *Vet Ophthalmol.* 2018;21(4):347–352. doi:10.1111/vop.12510.
86. Itoh S, Itoh K, Shinohara H. Regulation of human corneal epithelial mucins by rebamipide. *Curr Eye Res.* 2014;39(2):133–141. doi:10.3109/02713683.2013.834939.
87. Fox RI. Use of cevimeline, a muscarinic M1 and M3 agonist, in the treatment of Sjögren's syndrome. *Adv Exp Med Biol.* 2002;506:1107–1116.
88. Ono M, Takamura E, Shinozaki K, et al. Therapeutic effect of cevimeline on dry eye in patients with Sjögren's syndrome: a randomized, double-blind clinical study. *Am J Ophthalmol.* 2004;138(1):6–17. doi:10.1016/j.ajo.2004.02.010.
89. Petrone D, Condemi JJ, Fife R, Gluck O, Cohen S, Dalgin P. A double-blind, randomized, placebo-controlled study of cevimeline in Sjögren's syndrome patients with xerostomia and keratoconjunctivitis sicca. *Arthritis Rheumatism.* 2002;46(3):748–754. doi:10.1002/art.510.
90. Vivino FB, Al-Hashimi I, Khan Z, et al. Pilocarpine tablets for the treatment of dry mouth and dry eye symptoms in patients with Sjögren syndrome: a randomized, placebo-controlled, fixed-dose, multicenter trial. P92-01 study group. *Arch int med.* 1999;159(2):174–181. doi:10.1001/archinte.159.2.174.
91. Periman LM, Perez VL, Saban DR, Lin MC, Neri P. The immunological basis of dry eye disease and current topical treatment options. *J Ocul Pharmacol Ther.* 2020;36(3):137–146. doi:10.1089/jop.2019.0060.
92. Leonardi A, Van Setten G, Amrane M, et al. Efficacy and safety of 0.1% cyclosporine A cationic emulsion in the treatment of severe dry eye disease: a multicenter randomized trial. *Eur J Ophthalmol.* 2016;26(4):287–296. doi:10.5301/ejo.5000779.
93. Sall K, Stevenson OD, Mundorf TK, Reis BL. Two multicenter, randomized studies of the efficacy and safety of cyclosporine ophthalmic emulsion in moderate to severe dry eye disease. CsA Phase 3 Study Group. *Ophthalmology.* 2000;107(4):631–639. doi:10.1016/S0161-6420(99)00176-1.
94. Stevenson D, Tauber J, Reis BL. Efficacy and safety of cyclosporin A ophthalmic emulsion in the treatment of moderate-to-severe dry eye disease: a dose-ranging, randomized trial. The cyclosporin A Phase 2 study group. *Ophthalmology.* 2000;107(5):967–974. doi:10.1016/S0161-6420(00)00035-X.
95. Tuan HI, Chi SC, Kang YN. An updated systematic review with meta-analysis of randomized trials on topical cyclosporin A for dry-eye disease. *Drug Des Devel Ther.* 2020;14:265–274. doi:10.2147/DDDT.S207743.
96. Mandal A, Gote V, Pal D, Ogundele A, Mitra AK. Ocular pharmacokinetics of a topical ophthalmic nanomicellar solution of cyclosporine (Cequa®) for dry eye disease. *Pharm Res.* 2019;36(2):36. doi:10.1007/s11095-018-2556-5.
97. Abidi A, Shukla P, Ahmad A. Lifitegrast: a novel drug for treatment of dry eye disease. *J Pharmacol Pharmacother.* 2016;7(4):194–198. doi:10.4103/0976-500X.195920.

98. Hsueh PY, Ju Y, Vega A, Edman MC, MacKay JA, Hamm-Alvarez SF. A multivalent ICAM-1 binding nanoparticle which inhibits ICAM-1 and LFA-1 interaction represents a new tool for the investigation of autoimmune-mediated dry eye. *Int J Mol Sci.* 2020;21(8):2758. doi:10.3390/ijms21082758.
99. Stem ME, Gao J, Morgan GA, et al. The role of ICAM-1 as a signal protein for predisposition of ocular surface inflammation. *Adv Exp Med Biol.* 2002;506(Pt B):753–759.
100. Semba CP, Gadek TR. Development of lifitegrast: a novel T-cell inhibitor for the treatment of dry eye disease. *Clin Ophthalmol (Auckland, NZ).* 2016;10:1083–1094. doi:10.2147/OPTH.S110557.
101. Donnenfeld ED, Karpecki PM, Majmudar PA, et al. Safety of lifitegrast ophthalmic solution 5.0% in patients with dry eye disease: a 1-Year, multicenter, randomized, placebo-controlled study. *Cornea.* 2016;35(6):741–748. doi:10.1097/ICO.0000000000000803.
102. Sheppard JD, Torkildsen GL, Lonsdale JD, et al. Lifitegrast ophthalmic solution 5.0% for treatment of dry eye disease: results of the OPUS-1 phase 3 study. *Ophthalmology.* 2014;121(2):475–483. doi:10.1016/j.ophtha.2013.09.015.
103. Kim J, Kim YS, Park SH. Metformin as a Treatment Strategy for Sjögren's syndrome. *Int J Mol Sci.* 2021;22:(13).
104. Sheppard JD Jr., Singh R, McClellan AJ, et al. Long-term supplementation with n-6 and n-3 PUFAs improves moderate-to-severe keratoconjunctivitis sicca: a randomized double-blind clinical trial. *Cornea.* 2013;32(10):1297–1304. doi:10.1097/ICO.0b013e318299549c.
105. Rogers NM, Kireta S, Coates PT. Curcumin induces maturation-arrested dendritic cells that expand regulatory T cells in vitro and in vivo. *Clin Exp Immunol.* 2010;162(3):460–473. doi:10.1111/j.1365-2249.2010.04232.x.
106. Basu S, Shanbhag SS, Gokani A, Kedar R, Bahuguna C, Sangwan VS. Chronic ocular sequelae of Stevens-Johnson syndrome in children: long-term impact of appropriate therapy on natural history of disease. *Am J Ophthalmol.* May 2018;189:17–28. doi:10.1016/j.ajo.2018.01.028.
107. Shanbhag SS, Shah S, Singh M, Bahuguna C, Donthinani PR, Basu S. Lid-related keratopathy in Stevens-Johnson syndrome: natural course and impact of therapeutic interventions in children and adults. *Am J Ophthalmol.* Nov 2020;219:357–365. doi:10.1016/j.ajo.2020.07.006.
108. Alipour F, Kheirkhah A, Jabarvand Behrouz M. Use of mini scleral contact lenses in moderate to severe dry eye. *Cont Lens Anterior Eye.* 2012;35(6):272–276. doi:10.1016/j.clae.2012.07.006.
109. La Porta Weber S, Becco de Souza R, Gomes J Á P, Hofling-Lima AL. The use of the esclera scleral contact lens in the treatment of moderate to severe dry eye disease. *Am J Ophthalmol.* 2016;163:167–73.e1. doi:10.1016/j.ajo.2015.11.034.
110. Bavinger JC, DeLoss K, Mian SI. Scleral lens use in dry eye syndrome. *Curr Opin Ophthalmol.* 2015;26(4):319–324. doi:10.1097/ICU.0000000000000171.
111. Na JH, Jung JH, Park JG, Song PH, Song CH. Therapeutic effects of acupuncture in typical dry eye: a systematic review and meta-analysis. *Acta Ophthalmol.* 2021;99(5):489–498. doi:10.1111/aos.14651.
112. Kim BH, Kim MH, Kang SH, Nam HJ. Optimizing acupuncture treatment for dry eye syndrome: a systematic review. *BMC Complement Altern Med.* 2018;18(1):145. doi:10.1186/s12906-018-2202-0.
113. Yang L, Yang Z, Yu H, Song H. Acupuncture therapy is more effective than artificial tears for dry eye syndrome: evidence based on a meta-analysis. *Evidence-based complementary and alternative medicine: eCAM.* 2015;2015:143858.
114. Kossler AL, Brinton M, Patel ZM, Dalal R, Ta CN, Palanker D. Chronic electrical stimulation for tear secretion: lacrimal vs. anterior ethmoid nerve. *Ocul Surf.* 2019;17(4):822–827. doi:10.1016/j.jtos.2019.08.012.
115. Senchyna M, Ousler GW, Jerkins G, et al. Effect of TrueTear™ on dry eye symptoms during exposure to a controlled adverse environment. *Invest Ophthalmol Vis Sci.* 2018;59(9):918.
116. Geerling G, Raus P, Murube J. Minor salivary gland transplantation. *Dev Ophthalmol.* 2008;41:243–254.
117. Su JZ, Wang Z, Liu XJ, Lv L, Yu GY. Use of saliva flow rate measurement in minor salivary glands autotransplantation for treatment of severe dry eye disease. *Br J Ophthalmol.* 2021;2020--317552. doi:10.1136/bjophthalmol-2020-317552.
118. Qin Y, Zhang Y, Liang Q, et al. Labial salivary gland transplantation for severe dry eye in a rhesus monkey model. *Invest Ophthalmol Vis Sci.* 2018;59(6):2478–2486. doi:10.1167/iovs.18-23966.
119. Wakamatsu TH, Sant'Anna A, Cristovam PC, Alves VAF, Wakamatsu A, Gomes JAP. Minor salivary gland transplantation for severe dry eyes. *Cornea.* 2017;36(Suppl 1):S26–s33. doi:10.1097/ICO.0000000000001358.
120. Singh S, Basu S. A novel diagnostic technique of measuring labial minor salivary gland secretions using sodium fluorescein dye: implications for patients with dry eyes. *Semin Ophthalmol.* 2022 Jan 2;37(1):111–116. doi:10.1080/08820538.2021.1926518.
121. Singh S, Basu S. Preoperative labial mucosa evaluation as a deciding tool for minor salivary gland transplantation. *Ophthalmic Plast Reconstr Surg.* 2021;37(3s):S121–s2. doi:10.1097/IOP.0000000000001842.
122. Singh S, Basu S. Functional assessment of transplanted minor salivary glands. *Cornea.* 2020;39(8):e21–e2. doi:10.1097/ICO.0000000000002359.
123. Marinho DR, Burmann TG, Kwitko S. Labial salivary gland transplantation for severe dry eye due to chemical burns and Stevens-Johnson syndrome. *Ophthalmic Plast Reconstr Surg.* 2010;26(3):182–184. doi:10.1097/IOP.0b013e3181b8c3ad.
124. Vazirani J, Bhalekar S, Amescua G, Singh S, Basu S. Minor salivary gland transplantation for severe dry eye disease due to cicatrizing conjunctivitis: multicentre long-term outcomes of a modified technique. *Br J Ophthalmol.* 2020;105(11):1485–1490. doi:10.1136/bjophthalmol-2020-316611.
125. Nauta AJ, Fibbe WE. Immunomodulatory properties of mesenchymal stromal cells. *Blood.* 2007;110(10):3499–3506. doi:10.1182/blood-2007-02-069716.
126. Weng J, He C, Lai P, et al. Mesenchymal stromal cells treatment attenuates dry eye in patients with chronic graft-versus-host disease. *Mol Ther: J Am Soc Gene Ther.* 2012;20(12):2347–2354. doi:10.1038/mt.2012.208.
127. Möller-Hansen M, Larsen AC, Toft PB, et al. Safety and feasibility of mesenchymal stem cell therapy in patients with aqueous deficient dry eye disease. *Ocul Surf.* 2021;19:43–52. doi:10.1016/j.jtos.2020.11.013.
128. Dietrich J, Schrader S. Towards lacrimal gland regeneration: current concepts and experimental approaches. *Curr Eye Res.* 2020;45(3):230–240. doi:10.1080/02713683.2019.1637438.
129. Kawakita T. Regeneration of lacrimal gland function to maintain the health of the ocular surface. *Invest Ophthalmol Vis Sci.* 2018;59(14):Des169–des73. doi:10.1167/iovs.17-23576.
130. Yao Y, Zhang Y. The lacrimal gland: development, wound repair and regeneration. *Biotechnol Lett.* 2017;39(7):939–949. doi:10.1007/s10529-017-2326-1.
131. Liu CY, Hirayama M, Ali M, Shah D, Aakalu VK. Strategies for Regenerating the Lacrimal Gland. *Current Ophthalmol Rep.* 2017;5(3):193–198. doi:10.1007/s40135-017-0142-3.
132. Hirayama M, Tsubota K, Tsuji T. Bioengineered Lacrimal Gland Organ Regeneration in Vivo. *J Funct Biomater.* 2015;6(3):634–649. doi:10.3390/jfb6030634.
133. Hirayama M, Tsubota K, Tsuji T. Generation of a bioengineered lacrimal gland by using the organ germ method. *Methods Mol Biol.* 2017;1597:153–165.

134. Bannier-Hélaouët M, Post Y, Korving J, et al. Exploring the human lacrimal gland using organoids and single-cell sequencing. *Cell Stem Cell*. 2021 Jul 1;28(7):1221–1232.e7. doi:[10.1016/j.stem.2021.02.024](https://doi.org/10.1016/j.stem.2021.02.024).
135. Pflugfelder SC, de Paiva CS. The pathophysiology of dry eye disease: what we know and future directions for research. *Ophthalmology*. 2017;124(11s):S4–s13. doi:[10.1016/j.ophttha.2017.07.010](https://doi.org/10.1016/j.ophttha.2017.07.010).
136. Trujillo-Vargas CM, Schaefer L, Alam J, Pflugfelder SC, Britton RA, de Paiva CS. The gut-eye- lacrimal gland-microbiome axis in Sjögren Syndrome. *Ocul Surf*. 2020;18(2):335–344. doi:[10.1016/j.jtos.2019.10.006](https://doi.org/10.1016/j.jtos.2019.10.006).
137. Watane A, Cavuoto KM, Rojas M, et al. Fecal microbial transplant in individuals with immune-mediated dry eye. *Am J Ophthalmol*. 2022;233:90–100. doi:[10.1016/j.ajo.2021.06.022](https://doi.org/10.1016/j.ajo.2021.06.022).



Case report

Simple oral mucosal epithelial transplantation (SOMET) for ocular surface reconstruction in Stevens-Johnson Syndrome: A case report

Mariya B. Doctor^a, Raksheeth N. Rajagopal^a, Sayan Basu^{a,b,*}^a Shantilal Shanghvi Cornea Institute, L V Prasad Eye Institute, Kallam Anji Reddy Campus, Hyderabad, Telangana, India^b Prof. Brien Holden Centre for Eye Research (BHERC), L V Prasad Eye Institute, Hyderabad 500034, Telangana, India

ARTICLE INFO

Keywords:

Stevens-Johnson Syndrome
Simple oral mucosal epithelial transplantation
Limbal stem cell deficiency
LSCD
Case report

ABSTRACT

Introduction: We report the clinical outcome of a novel surgical technique called simple oral mucosal epithelial transplantation (SOMET) for the treatment of limbal stem cell deficiency (LSCD) in a patient with Stevens-Johnson Syndrome (SJS).

Presentation of case: An eighteen-year-old girl was diagnosed as having chronic bilateral ocular sequelae of SJS. She initially underwent mucous membrane grafting (MMG) in both eyes for lid margin keratinization. Over the course of the next decade, the ocular surface cicatrization worsened in her left eye, leading to progressive symblepharon formation with total corneal conjunctivalization. She then underwent ocular surface reconstruction using bulbar MMG and SOMET. Following SOMET, the ocular surface epithelialized within 3 weeks and remained stable throughout the follow-up period. At one-year postoperatively, the visual acuity had improved from light perception to 20/250 unaided, and to 20/100 with scleral contact lens correction in the left eye.

Discussion: Simple limbal epithelial transplantation (SLET) has been a boon for the treatment of unilateral LSCD. Allogeneic SLET and kerato-limbal allografts can be useful for patients with bilateral disease, however this exposes the patients to the risks of long-term systemic immunosuppression. SOMET combines the benefits of cultivated oral mucosal epithelial transplantation (COMET) and SLET, and is an autologous and single-staged surgical alternative for patients with bilateral LSCD.

Conclusion: This case demonstrates that SOMET is a viable surgical option in cases with bilateral LSCD, eliminating the need for an allogeneic limbal graft, systemic immunosuppression, or laboratory cell culture.

1. Introduction

Stevens-Johnson Syndrome (SJS) is a rare, life-threatening, vesiculo-bullous disease of the skin and mucosa, occurring due to an immune-mediated hypersensitivity reaction to a systemic drug or infection [1]. Ocular involvement is the most common chronic sequelae of SJS, which impairs the quality of life of affected individuals [1,2]. Keratopathy and limbal stem cell deficiency (LSCD) ensue due to constant friction from lid margin keratinization, long-standing ocular surface inflammation, conjunctival cicatrization, and aqueous deficient dry eye disease, leading to sight-threatening complications [2]. Treatment of lid margin keratinization involves lid margin mucous membrane grafting (MMG) or the use of prosthetic replacement of ocular surface ecosystem (PROSE) contact lenses [3–5]. The fornix and the ocular surface can be reconstructed using amniotic membrane transplantation (AMT), ocular surface MMG, or cultivated oral mucosal epithelial transplantation

(COMET) [6,7]. Bilateral LSCD in these patients can be treated with allogeneic limbal stem cell transplantation or COMET [8]. Being an autologous procedure, COMET carries no risk of immunological rejection, improves visual acuity, ocular surface severity scoring, and helps treat LSCD [9]. However, it requires a sophisticated and expensive clinical-grade laboratory setup. Allogeneic limbal stem cell transplantation, on the other hand, requires long-term systemic immunosuppression. In this case report, we applied the concept of simple limbal epithelial transplantation (SLET) and combined it with the benefits of COMET, by replacing the limbal tissue with autologous oral mucosa, using a surgical technique termed simple oral mucosal epithelial transplantation (SOMET). This case report is as per the SCARE-2020 criteria [10].

* Corresponding author at: Prof. Brien Holden Centre for Eye Research (BHERC), L V Prasad Eye Institute, Banjara Hills, Hyderabad 500034, Telangana, India.
E-mail address: sayanbasu@lvpei.org (S. Basu).

<https://doi.org/10.1016/j.ijscr.2023.108643>

Received 5 July 2023; Received in revised form 4 August 2023; Accepted 4 August 2023

Available online 9 August 2023

2210-2612/© 2023 The Author(s). Published by Elsevier Ltd on behalf of IJS Publishing Group Ltd. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

2. Presentation of case

An eighteen-year-old girl presented to us in 2012 with complaints of photophobia, redness, and discharge in both eyes (OU) for the past three years, after an episode of multiple skin eruptions and rashes over the body, which occurred as a consequence of consuming systemic medication for fever. At presentation, she had a best corrected visual acuity (BCVA) of 20/40p in OU. There were blocked meibomian orifices, thickened and keratinized lid margins, conjunctival cicatrization, and corneal scarring with vascularization noted in OU. She was diagnosed as having ocular sequelae of SJS and was started on topical steroids, and frequent application of tear substitutes. She later underwent lid margin MMG in OU. The right eye (OD) later underwent symblepharon release, amniotic membrane grafting and cataract surgery. While OD was relatively stable (Fig. 1A), in the left eye (OS) over the period of the next ten years, she developed a progressive symblepharon covering 80–90 % of the cornea (Fig. 1B) causing her visual acuity to drop from 20/40 to light perception. At this stage we intervened surgically in OS and performed symblepharon release with ocular surface reconstruction using bulbar MMG and SOMET.

Surgical procedure (Fig. 2): under general anesthesia, after painting and draping the eye, the symblepharon was released from the cornea and a 360-degree peritomy was performed (Fig. 2A, B). A combination of sharp and blunt dissection was carried out to separate the fibrovascular pannus from the underlying corneal stroma (Fig. 2C). An amniotic membrane graft (AMG) was attached over the ocular surface with fibrin glue and perilimbal 10-0 nylon continuous sutures (Fig. 2D). The oral mucosa over the lower lip was cleaned with betadine 5 % solution. The submucosal tissue was infiltrated with 5 mL of lignocaine hydrochloride 2 % and adrenaline (1:200,000) solution. The MMG was marked with tissue markers and borders incised with number 11 blade. It was then carefully excised and placed over the surgical drape (Fig. 2E). A small triangular piece of the MMG tissue (approximately 3 mm in size) was excised and kept aside for SOMET later (Fig. 2F). The submucosal tissue was debulked, and the graft was thinned to the extent possible (Fig. 2G, H). It was divided into 2 horizontally equal parts for surface reconstruction (Fig. 2I). The grafts were attached circumferentially over the sclera around the limbus using 7-0 vicryl sutures (Fig. 2J–K). The remaining triangular tissue of MMG (Fig. 2K) was thinned and divided into 10–15 tiny pieces, which were placed over the paracentral cornea (similar to SLET), using fibrin glue, over which a soft bandage contact lens (BCL) was placed (Fig. 2L).

Post-operatively, she was followed up at 1 day, 1 week, 3 weeks, 2

months, 6 months, and 1 year. She was started on low-dose oral steroids (20 mg/day) for a week, which were tapered to 5 mg/day for a month and stopped thereafter; topical tear substitutes, and ointment of steroid and antibiotic combination applied thrice a day for 1 week and later shifted to only nightly application. The cornea had completely epithelized by 3 weeks (Fig. 3A, B) and the BCL was removed. The fornices had been restored (Fig. 3C) and the ocular surface (bulbar) MMG had also integrated well (Fig. 3D–F). We encountered no epithelial instability, or any other complication during the entire post-operative period. At the final 1-year follow-up visit, the ocular surface was stable, the cornea was completely epithelized, the SOMET grafts had completely integrated, and there was residual macular to nebular grade superficial stromal scarring with vascularization (Fig. 4A, B). The uncorrected visual acuity was 20/200 and 20/250 in OD and OS, respectively. She was improving to 20/100 in OS with PROSE contact lenses (Fig. 4C, D). The Schirmer's test value had also increased from 0 mm at 5 min pre-operatively to 6 mm post-operatively.

3. Discussion

Following SJS, patients usually present to an ophthalmologist after developing ocular symptoms such as foreign body sensation, dry eyes, or decrease in visual acuity, months to years after the acute episode [11]. The chronic stage may lead to sight-threatening complications due to lid margin keratinization, dry eye disease, and chronic ocular surface inflammation [2–6,11]. Since LSCD in SJS is often bilateral, surgical options include allogeneic limbal stem cell transplantation in the form of allo-SLET, or keratolimbal allografting (KLAL), or autologous COMET. Outcomes of limbal allografting in cases of SJS have not shown to be satisfactory due to high chances of failure and the need for long-term immunosuppression which have other potential side effects [11,12]. COMET has a success rate of about 70 % in cases of bilateral LSCD. It helps in rapid surface epithelization, reduces corneal opacification, and improves visual acuity [9,13,14]. Studies have shown that the cultured cells resemble the corneal epithelial cell having three to five cell layers, with small basal cells, flattened middle cells, and polygonal superficial cells. They express markers of corneal epithelial cell differentiation (cytokeratin 3 and connexin 43), and marker of progenitor stem cells (p63), similar to corneal epithelial cells [13–15].

COMET has the advantage of being an autologous procedure, and not requiring any systemic immunosuppression; while the disadvantages include it being a two-staged procedure, and the requirement of an expensive clinical-grade laboratory set-up for cell cultivation [8]. Based




Fig. 1. Slit lamp photographs of both eyes around ten years after of undergoing eyelid mucous membrane grafting. The right eye (A), which had also undergone symblepharon release with amniotic membrane transplantation and cataract surgery two years prior shows: thickened lid margins, integrated eyelid mucous membrane grafts in both upper and lower eyelid, with nebular grade corneal scarring. The left eye (B) shows: thickened lid margins with mucous membrane grafts in place, an extensive symblepharon covering almost the entire cornea (yellow arrow), limiting eversion of the upper lid and precluding the view of the cornea. (For interpretation of the references to colour in this figure legend, the reader is referred to the web version of this article.)




Fig. 2. Surgical technique of simple oral mucosal epithelial transplantation or SOMET (A to L). A: The symblepharon was excised (A) and the self-retaining speculum was placed to expose the cornea and ocular surface. A 360-degree peritomy was performed (B) and the fibro-vascular pannus was dissected from the underlying corneal stroma (C). An amniotic membrane graft was placed over the ocular surface using fibrin glue, along with anchoring, perilimbal, continuous sutures with 10-0 nylon (D). The oral mucous membrane graft was harvested from the lower lip (E), and a small triangular shaped tissue was excised (F, G) and kept aside, for later use as SOMET. The graft was thinned to the extent possible by excising the underlying submucosal tissue and fat (H). The tissue was then dissected into two equal parts (I). The grafts were sutured over the bulbar bare sclera, circumferentially around the limbus (J), using 7-0 vicryl sutures (K). The triangular tissue was then thinned (K) and divided into 10–15 small pieces, and placed over the cornea, using fibrin glue (L).




Fig. 3. Slit lamp photographs of the left eye three weeks after simple oral mucosal epithelial transplantation or SOMET (A to F). The tiny pieces of mucosal grafts (A, blue arrow) can be appreciated in the paracentral cornea. Fluorescein-stained image in cobalt blue filter (B) shows a completely epithelized corneal surface. Everted image of the left upper eyelid (C) reveals a deep upper fornix. Images in different gazes (D to F) showing 360-degree bulbar mucous membrane grafts, which are well taken up. (For interpretation of the references to colour in this figure legend, the reader is referred to the web version of this article.)

on the encouraging outcomes of COMET, and combining them with the principle of SLET (where autologous limbal tissue from a healthy eye is divided into multiple small pieces and placed over the AMG in the paracentral cornea for these cells to proliferate and eventually epithelize

and cover the entire corneal surface) [16,17] we decided to perform SOMET, a synthesis of both these techniques. We placed tiny pieces of the MMG, obtained from the lip mucosa, circumferentially at the paracentral cornea with the help of fibrin glue over an underlying AMG.




Fig. 4. Slit lamp photos of the left eye at the one-year follow-up visit after simple oral mucosal epithelial transplantation or SOMET (A to D). The corneal surface is stable (A), completely epithelized (B) with completely integrated SOMET grafts and macular to nebular grade superficial corneal scarring and vascularization. Images after wearing PROSE contact lenses in the left eye (C, D) which show acceptable fitting of the lenses over the reconstructed corneal and bulbar surface.

These transplants were seen as discrete tissue initially but later integrated and led to complete epithelization of the corneal surface by 3 weeks. Post-operatively, a well epithelized surface, improvement in visual acuity and Schirmer's value were noted at 1-year follow-up in our patient.

There are few reports of the oral mucosa being used as a tissue graft, without ex-vivo cultivation as in COMET, for the treatment of LSCD [18,19]. In a retrospective study done by Liu et al., an oral mucosal graft was used as a surrogate limbus in 7 eyes with severe LSCD (1 eye having LSCD due to SJS). They too simultaneously treated the symblepharon along with LSCD. Their results showed that all patients had improvement in symptoms and corneal scarring. They also demonstrated that initially, the peripheral corneal vascularization increased, which then decreased significantly after 3–6 months [18]. The first report of performing SOMET was in a rabbit model of LSCD by Inamochi et al. [19]. In this study a 3×4 mm piece of oral mucosal tissue was excised and divided into small pieces, treated with dispase II, and placed over the denuded cornea, without AMT or fibrin glue. They demonstrated that the epithelium expanded to form islands around the grafts 1 week after surgery and the corneal defect was completely epithelialized at 2 weeks, similar to the results of SLET. They went on to describe that the single surface layer of epithelial cells was CK3-positive, and basal cells were p63 positive similar to corneal epithelial cells [19]. Hence, they concluded that the direct placement of autologous oral mucosal tissue, without ex-vivo cell culture, could be a good alternative to limbal allografting in bilateral LSCD. This report is the first clinical validation of the SOMET technique proposed by Inamochi et al. The modifications from the animal study were that there was no prior treatment of the mucosal tissue with dispase, we used AMG on the cornea, and used fibrin

glue to secure the transplants, analogous to SLET for the treatment of LSCD. Here the AMG would act as a scaffold for the mucosal epithelial cells to proliferate rapidly. We also performed simultaneous bulbar surface reconstruction with MMG to address the severe symblepharon, analogous to a conjunctival autograft or MMG being performed along with SLET to address the conjunctival deficiency [17]. This approach was not only effective in anatomically restoring the ocular surface but also resulted in significant functional improvement in unaided and contact lens-corrected visual acuity.

4. Conclusion

This study highlights the potential of SOMET as a surgical alternative to allogeneic limbal transplantation and COMET for the treatment of bilateral LSCD. It combines the strengths of SLET and COMET, while avoiding the need for long-term systemic immunosuppression or laboratory cell cultivation. However, a prospective study with a larger sample size is required to validate the long-term clinical outcomes of SOMET in different cases of bilateral LSCD.

Consent

Written informed consent was obtained from the patient for publication of this case report and accompanying images. A copy of the written consent is available for review by the Editor-in-Chief of this journal on request.

Patient perspective

"I was blind and disfigured in my left eye. The surgery helped me in seeing better, but what I was most impressed with was the improvement in the cosmetic appearance of the eye. This has helped improve my self-confidence and self-esteem."

Declaration of competing interest

The authors have no conflict of interest.

Acknowledgment

The study was funded by the Hyderabad Eye Research Foundation (HERF), Hyderabad, Telangana, India.

References

- [1] S. Kohanim, S. Palioura, H.N. Saeed, E.K. Akpek, G. Amescua, S. Basu, P. H. Blomquist, C.S. Bouchard, J.K. Dart, X. Gai, J.A. Gomes, D.G. Gregory, G. Iyer, D.S. Jacobs, A.J. Johnson, S. Kinoshita, I.S. Mantagos, J.S. Mehta, V.L. Perez, S. C. Pflugfelder, V.S. Sangwan, K.C. Sippel, C. Sotozono, B. Srinivasan, D.T. Tan, R. Tandon, S.C. Tseng, M. Ueta, J. Chodosh, Stevens-Johnson Syndrome/toxic epidermal necrolysis—a comprehensive review and guide to therapy. I. Systemic disease, *Ocul. Surf.* 14 (1) (2016) 2–19, <https://doi.org/10.1016/j.jtos.2015.10.002>. Jan.
- [2] S. Kohanim, S. Palioura, H.N. Saeed, E.K. Akpek, G. Amescua, S. Basu, P. H. Blomquist, C.S. Bouchard, J.K. Dart, X. Gai, J.A. Gomes, D.G. Gregory, G. Iyer, D.S. Jacobs, A.J. Johnson, S. Kinoshita, I.S. Mantagos, J.S. Mehta, V.L. Perez, S. C. Pflugfelder, V.S. Sangwan, K.C. Sippel, C. Sotozono, B. Srinivasan, D.T. Tan, R. Tandon, S.C. Tseng, M. Ueta, J. Chodosh, Acute and chronic ophthalmic involvement in Stevens-Johnson Syndrome/toxic epidermal necrolysis — a comprehensive review and guide to therapy. II. Ophthalmic disease, *Ocul. Surf.* 14 (2) (2016) 168–188, <https://doi.org/10.1016/j.jtos.2016.02.001>. Apr.
- [3] S.S. Shanbhag, V.S. Sangwan, A. Singh, P.R. Donthineni, S. Basu, B. Srinivasan, et al., Clinical aspects of Stevens-Johnson Syndrome/toxic epidermal necrolysis with severe ocular complications in India, *Front. Med.* 8 (2021), 643955.
- [4] S.S. Shanbhag, S. Shah, M. Singh, C. Bahuguna, P.R. Donthineni, S. Basu, Lid-related keratopathy in Stevens-Johnson Syndrome: natural course and impact of therapeutic interventions in children and adults, *Am J. Ophthalmol.* 219 (2020) 357–365.
- [5] S. Basu, S.S. Shanbhag, A. Gokani, R. Kedar, C. Bahuguna, V.S. Sangwan, Chronic ocular sequelae of Stevens-Johnson Syndrome in children: long-term impact of appropriate therapy on natural history of disease, *Am J. Ophthalmol.* 189 (2018) 17–28.
- [6] G. Iyer, B. Srinivasan, S. Agarwal, S. Kamala Muralidharan, S. Arumugam, Comprehensive approach to ocular consequences of Stevens Johnson Syndrome — the aftermath of a systemic condition, *Graefes Arch. Clin. Exper. Ophthalmol.* = *Albrecht von Graefes Arch. Klin. Exper. Ophthalmol.* 252 (3) (2014) 457–467.
- [7] P. Sudana, S. Basu, S.S. Shanbhag, Oral mucous membrane grafts for total symblepharon and lid margin keratinisation post Stevens-Johnson syndrome, *BMJ Case Rep.* 13 (10) (2020).
- [8] J. Vazirani, I. Mariappan, S. Ramamurthy, S. Fatima, S. Basu, V.S. Sangwan, Surgical management of bilateral limbal stem cell deficiency, *Ocul. Surf.* 14 (3) (2016) 350–364, <https://doi.org/10.1016/j.jtos.2016.02.006>. Jul.
- [9] R. Venugopal, R. Nagpal, S. Mohanty, S. Sen, S. Kashyap, T. Agarwal, et al., Outcomes of cultivated oral mucosal epithelial transplantation in eyes with chronic Stevens-Johnson Syndrome sequelae, *Am J. Ophthalmol.* 222 (2021) 82–91.
- [10] R.A. Agha, T. Franchi, C. Sohrabi, G. Mathew, A. Kerwan, A. Thoma, et al., The SCARE 2020 guideline: updating consensus surgical CAse REport (SCARE) guidelines, *Int. J. Surg.* 84 (2020) 226–230.
- [11] R. Jain, N. Sharma, S. Basu, G. Iyer, M. Ueta, C. Sotozono, C. Kannabiran, V. M. Rathi, N. Gupta, S. Kinoshita, J.A. Gomes, J. Chodosh, V.S. Sangwan, Stevens-Johnson syndrome: the role of an ophthalmologist, *Surv. Ophthalmol.* 61 (4) (2016) 369–399, <https://doi.org/10.1016/j.survophthal.2016.01.004>. Jul–Aug.
- [12] A. Solomon, P. Ellies, D.F. Anderson, A. Touhami, M. Grueterich, E.M. Espana, et al., Long-term outcome of keratolimbal allograft with or without penetrating keratoplasty for total limbal stem cell deficiency, *Ophthalmology* 109 (6) (2002) 1159–1166.
- [13] P. Prabhasawat, P. Ekpo, M. Uprasertkul, S. Chotikavanich, N. Tesavibul, K. Pornpanich, et al., Long-term result of autologous cultivated oral mucosal epithelial transplantation for severe ocular surface disease, *Cell Tissue Bank.* 17 (3) (2016) 491–503.
- [14] K. Nishida, M. Yamato, Y. Hayashida, K. Watanabe, K. Yamamoto, E. Adachi, et al., Corneal reconstruction with tissue-engineered cell sheets composed of autologous oral mucosal epithelium, *N. Engl. J. Med.* 351 (12) (2004) 1187–1196.
- [15] S.L. Madhira, G. Vemuganti, A. Bhaduri, S. Gaddipati, V.S. Sangwan, Y. Ghanekar, Culture and characterization of oral mucosal epithelial cells on human amniotic membrane for ocular surface reconstruction, *Mol. Vis.* 14 (2008) 189–196.
- [16] V.S. Sangwan, S. Basu, S. MacNeil, D. Balasubramanian, Simple limbal epithelial transplantation (SLET): a novel surgical technique for the treatment of unilateral limbal stem cell deficiency, *Br. J. Ophthalmol.* 96 (7) (2012) 931–934.
- [17] S.S. Shanbhag, C.N. Patel, R. Goyal, P.R. Donthineni, V. Singh, S. Basu, Simple limbal epithelial transplantation (SLET): review of indications, surgical technique, mechanism, outcomes, limitations, and impact, *Indian J. Ophthalmol.* 67 (8) (2019) 1265–1277. Aug.
- [18] J. Liu, H. Sheha, Y. Fu, M. Gegengack, S.C. Tseng, Oral mucosal graft with amniotic membrane transplantation for total limbal stem cell deficiency, *Am J. Ophthalmol.* 152 (5) (2011) 739–747.e1.
- [19] A. Inamochi, A. Tomioka, K. Kitamoto, T. Miyai, T. Usui, M. Aihara, et al., Simple oral mucosal epithelial transplantation in a rabbit model, *Sci. Rep.* 9 (1) (2019) 18088.