Articles

Since 2024 Since 2023 Since 2020 Custom range...

Sort by relevance

Sort by date

Create alert

Meta-analysis of prevalence of triple-negative breast cancer and its clinical features at...

Search within citing articles

About 53 results (0.02 sec)

Advances in systemic therapies for triple negative breast cancer

RA Leon-Ferre, MP Goetz - Bmj, 2023 - bmj.com

Triple negative breast cancer (TNBC) continues to be the subtype of breast cancer with the highest rates of recurrence and mortality. The lack of expression of targetable proteins such ...

☆ Save ☑ Cite Cited by 78 Related articles All 6 versions

[нтмь] Nanomedicine tactics in cancer treatment: Challenge and hope

MS Al-Zoubi, RM Al-Zoubi - Critical Reviews in Oncology/Hematology, 2022 - Elsevier Defeating cancer is the ultimate challenge and goal of oncologists, facing various obstacles along with finding effective anti-cancer therapies and understanding drug delivery ...

☆ Save 勿 Cite Cited by 29 Related articles All 6 versions

Risk factors for the development of triple-negative breast cancer versus nontriple-negative breast cancer: a case-control study

S Nag, R Dikshit, S Desai, A Mane, S Mhatre... - Scientific Reports, 2023 - nature.com The risk factors for breast cancer have been defined in several studies but there is deficient data for specific subtypes. We report here the pathological characteristics of a breast cancer ... ☆ Save 勿 Cite Cited by 5 Related articles All 8 versions

[нтмь] Outcomes in breast cancer—does ethnicity matter?

YS Yap - ESMO open, 2023 - Elsevier

Ethnic or racial differences in breast cancer (BC) survival outcomes have been reported, but current data are largely restricted to comparisons between African Americans and non ...

☆ Save 勿 Cite Cited by 10 Related articles All 6 versions

[HTML] FOXM1 mediates GDF-15 dependent stemness and intrinsic drug resistance in breast cancer

A Modi, P Purohit, D Roy, JR Vishnoi, P Pareek... - Molecular Biology ..., 2022 - Springer Background Stemness, a key component of breast cancer (BC) heterogeneity, is responsible for chemoresistance. Growth differentiation factor-15 (GDF-15) induces drug resistance and ...

☆ Save 勿 Cite Cited by 20 Related articles All 7 versions

☆ Save 勿 Cite Cited by 15 Related articles All 6 versions

Combination of 3PO analog PFK15 and siPFKL efficiently suppresses the migration, colony formation ability, and PFK-1 activity of triple-negative breast cancers by ...

A Kashyap, SM Umar, A Dev JR... - Journal of Cellular ..., 2023 - Wiley Online Library Among all the subtypes of breast cancer, triple-negative breast cancer (TNBC) has been associated with the worst prognosis. Recently, for many solid tumors (including breast ... ☆ Save 勿 Cite Cited by 4 Related articles All 3 versions

Profile of pathogenic mutations and evaluation of germline genetic testing criteria in consecutive breast cancer patients treated at a North Indian tertiary care center

A Mittal, SVS Deo, A Gogia, A Batra, A Kumar... - Annals of Surgical ..., 2021 - Springer Background The burden of hereditary breast cancer in India is not well defined. Moreover, genetic testing criteria (National Comprehensive Cancer Network [NCCN] and ...

[нтмь] Breast cancer in young Indian women: factors, challenges in screening, and upcoming diagnostics

P Roy - Journal of Cancer Research and Clinical Oncology, 2023 - Springer Breast cancer management for young Indian women are full of challenges. The National Cancer Registry Programme (NCRP) has predicted that nearly 2, 30,000 cases of breast ... ☆ Save 勿 Cite Cited by 2 Related articles All 3 versions

Exploring the human microbiome—A step forward for precision medicine in breast cancer

A Jotshi, KK Sukla, MM Haque, C Bose... - Cancer ..., 2023 - Wiley Online Library Background The second most frequent cancer in the world and the most common malignancy in women is breast cancer. Breast cancer is a significant health concern in India ... ☆ Save 勿 Cite Cited by 3 Related articles All 5 versions

Tumor Characteristics Associated with Axillary Nodal Positivity in Triple Negative **Breast Cancer** 

N Chintapally, K Englander, J Gallagher, K Elleson... - Diseases, 2023 - mdpi.com Larger-size primary tumors are correlated with axillary metastases and worse outcomes. We evaluated the relationships among tumor size, location, and distance to nipple relative to ... ☆ Save 勿 Cite Cited by 2 Related articles All 8 versions ≫

[HTML] sciencedirect.com

[PDF] nature.com

[HTML] sciencedirect.com

[HTML] springer.com

53 citations, 1st paper is printed here

[PDF] wiley.com

[PDF] springer.com

[HTML] springer.com

Prof. L S Shashidhara

[PDF] wiley.com

[PDF] mdpi.com

Gooooogle 1 2 3 4 5 6



Q



Articles

3 results (**0.02** sec)

My profile

My library

Any time

Since 2024

Since 2023

Since 2020

Custom range...

Sort by relevance

Sort by date

Create alert

Evaluation of tumor-infiltrating lymphocytes (TILs) in molecular subtypes of an Indian cohort...

Search within citing articles

HER2-low and tumor infiltrating lymphocytes in triple-negative breast cancer: Are they connected?

X Baez-Navarro, NS van den Ende, <u>AH Nguyen</u>... - Breast Cancer ..., 2024 - Springer Most patients with triple-negative breast cancer (TNBC) are not candidates for targeted therapy, leaving chemotherapy as the primary treatment option. Recently, immunotherapy ...

☆ Save ፵ Cite Related articles All 10 versions

[нтмь] Assessment of Stromal Tumor-Infiltrating Lymphocytes in Neoadjuvant Chemotherapy for Invasive Breast Carcinoma: Predictive Insights Across Molecular ...

AA Khan, S Ahuja, K Kiruthikasri, <u>S Zaheer</u> - Pathology-Research and ..., 2024 - Elsevier Background and aims Breast cancer, a leading cause of female mortality, has prompted the widespread adoption of Neoadjuvant chemotherapy (NAC) for its potential to minimize ...

☆ Save ☑ Cite Related articles All 3 versions

[нтмь] Association of tumor-infiltrating lymphocytes with clinical outcomes in patients with triple-negative breast cancer receiving neoadjuvant chemotherapy: а ...

<u>FCA de Moraes</u>, MEC Souza, VKT Sano... - Clinical and ..., 2024 - Springer Objective Triple-negative breast cancer (TNBC) presents a clinical challenge as an aggressive tumor, correlated with unfavorable prognosis. Tumor-infiltrating lymphocytes ...

☆ Save ☑ Cite Related articles All 2 versions

Zinc transporter LIV1: A promising cell surface target for triple negative breast cancer

R Saravanan, V Balasubramanian... - Journal of Cellular ..., 2022 - Wiley Online Library Breast cancer is one of the leading causes contributing to the global cancer burden. The triple negative breast cancer (TNBC) molecular subtype accounts for the most aggressive ...

☆ Save ☑ Cite Cited by 7 Related articles All 3 versions

Analyzing the influence of IL18 in regulation of YAP1 in breast oncogenesis using cBioportal

A Rahman, <u>LS Shashidhara</u> - Cancer Reports, 2022 - Wiley Online Library Background Yes-associated protein 1 (YAP1) is responsible for tumor growth, progression and metastasis. The mechanisms controlling the generation and relative ratio of the ...

☆ Save ☑ Cite Cited by 3 Related articles All 6 versions

[PDF] springer.com

[HTML] sciencedirect.com

[HTML] springer.com

Prof. L S Shashidhara

Shout.

[PDF] wiley.com



Articles

Since 2024 Since 2023 Since 2020 Custom range...

# Sort by relevance

Sort by date

Create alert

RUNX1 and RUNX3 protect against YAP-mediated EMT, stem-ness and shorter survival outcomes...

Search within citing articles

[HTML] Lipid metabolism alteration contributes to and maintains the properties of cancer stem cells

Lipids, the basic components of the cell membrane, execute fundamental roles in almost all the cell activities including cell-cell recognition, signalling transduction and energy supplies ...

H Li, Z Feng, ML He - Theranostics, 2020 - ncbi.nlm.nih.gov

☆ Save 夘 Cite Cited by 108 Related articles All 7 versions

[HTML] RUNX1 and cancer

TC Lin - Biochimica et biophysica acta (BBA)-reviews on cancer, 2022 - Elsevier Runt-related transcription factor 1 (RUNX1) is frequently involved in the progression of acute leukemia. However, emerging and discoverable RUNX1 somatic mutations, RUNX1 ... ☆ Save ⑰ Cite Cited by 55 Related articles All 3 versions

[HTML] sciencedirect.com

Circulating IncRNA SNHG11 as a novel biomarker for early diagnosis and prognosis of colorectal cancer

W Xu, G Zhou, H Wang, Y Liu, B Chen... - ... journal of cancer, 2020 - Wiley Online Library Colorectal cancer (CRC) is the third most common cancer and the second leading cause of cancer mortality worldwide. Emerging evidence indicates that tumour cells release ... ☆ Save 夘 Cite Cited by 149 Related articles All 5 versions

[PDF] wiley.com

[HTML] nih.gov

Runx1 is a key regulator of articular cartilage homeostasis by orchestrating YAP, TGFβ, and Wnt signaling in articular cartilage formation and osteoarthritis

Y Zhang, T Zuo, A McVicar, HL Yang, YP Li, W Chen - Bone Research, 2022 - nature.com Runt-related transcription factor 1 (Runx1) plays a key role in cartilage formation, but its function in articular cartilage formation is unclear. We generated non-inducible and ... ☆ Save 夘 Cite Cited by 26 Related articles All 12 versions

[PDF] nature.com

[нтмь] Inside the stemness engine: Mechanistic links between deregulated transcription factors and stemness in cancer

EH Ervin, R French, CH Chang, S Pauklin - Seminars in cancer biology, 2022 - Elsevier Cell identity is largely determined by its transcriptional profile. In tumour, deregulation of transcription factor expression and/or activity enables cancer cell to acquire a stem-like state ... ☆ Save 勿 Cite Cited by 21 Related articles All 4 versions

[HTML] sciencedirect.com

RUNX1: an emerging therapeutic target for cardiovascular disease

A Riddell, M McBride, <u>T Braun</u>, SA Nicklin... - Cardiovascular ..., 2020 - academic.oup.com Abstract Runt-related transcription factor-1 (RUNX1), also known as acute myeloid leukaemia 1 protein (AML1), is a member of the core-binding factor family of transcription ... ☆ Save 勿 Cite Cited by 63 Related articles All 11 versions

[PDF] oup.com

The hippo kinase LATS2 controls Helicobacter pylori-induced epithelialmesenchymal transition and intestinal metaplasia in gastric mucosa

SE Molina-Castro, C Tiffon, J Giraud, H Boeuf... - Cellular and Molecular ..., 2020 - Elsevier Background & Aims Gastric carcinoma is related mostly to CagA+-Helicobacter pylori infection, which disrupts the gastric mucosa turnover and elicits an epithelial-mesenchymal ... ☆ Save 夘 Cite Cited by 54 Related articles All 12 versions

[PDF] sciencedirect.com

RUNX1-dependent mechanisms in biological control and dysregulation in cancer

D Hong, AJ Fritz, JA Gordon, CE Tye... - Journal of cellular ..., 2019 - Wiley Online Library The RUNX1 transcription factor has recently been shown to be obligatory for normal development. RUNX1 controls the expression of genes essential for proper development in ... ☆ Save 勿 Cite Cited by 65 Related articles All 8 versions

[HTML] nih.gov

RUNX1 is regulated by androgen receptor to promote cancer stem markers and chemotherapy resistance in triple negative breast cancer

NB Fernández, SM Sosa, <u>JT Roberts</u>, MS Recouvreux... - Cells, 2023 - mdpi.com Triple negative breast cancer (TNBC) is an aggressive breast cancer subtype for which no effective targeted therapies are available. Growing evidence suggests that chemotherapy ... ☆ Save 勿 Cite Cited by 10 Related articles All 16 versions ≫

[PDF] mdpi.com

[HTML] YAP1-mediated pancreatic stellate cell activation inhibits pancreatic cancer cell proliferation

Y Xiao, H Zhang, Q Ma, R Huang, J Lu, X Liang, X Liu... - Cancer letters, 2019 - Elsevier Pancreatic stellate cells (PSCs) are activated in pancreatic ductal adenocarcinoma (PDAC) and are responsible for dense desmoplastic stroma. Yes-associated protein 1 (YAP1) can ... ☆ Save 勿 Cite Cited by 52 Related articles All 4 versions

[HTML] sciencedirect.com

Goooooogle > 1 2 3 4 5 6 7

Privacy Terms Help

Prof. L S Shashidhara

Articles

Since 2024 Since 2023 Since 2020 Custom range...

Sort by relevance

Sort by date

Create alert

# Structure of the VP16 transactivator target in the Mediator

Search within citing articles

About 94 results (0.05 sec)

# The Mediator complex as a master regulator of transcription by RNA polymerase

WF Richter, S Nayak, <u>J Iwasa</u>, <u>DJ Taatjes</u> - Nature Reviews Molecular ..., 2022 - nature.com The Mediator complex, which in humans is 1.4 MDa in size and includes 26 subunits. controls many aspects of RNA polymerase II (Pol II) function. Apart from its size, a defining ... ☆ Save 夘 Cite Cited by 124 Related articles All 6 versions

### [HTML] The Mediator complex: a central integrator of transcription

BL Allen, DJ Taatjes - Nature reviews Molecular cell biology, 2015 - nature.com Abstract The RNA polymerase II (Pol II) enzyme transcribes all protein-coding and most noncoding RNA genes and is globally regulated by Mediator—a large, conformationally flexible ... ☆ Save 勿 Cite Cited by 1026 Related articles All 9 versions

### [HTML] Transcription regulation by the Mediator complex

J Soutourina - Nature reviews Molecular cell biology, 2018 - nature.com Alterations in the regulation of gene expression are frequently associated with developmental diseases or cancer. Transcription activation is a key phenomenon in the ... ☆ Save 夘 Cite Cited by 615 Related articles All 7 versions

### [нтмь] Regulation of the RNA polymerase II pre-initiation complex by its associated coactivators

S Malik, RG Roeder - Nature Reviews Genetics, 2023 - nature.com Abstract The RNA polymerase II (Pol II) pre-initiation complex (PIC) is a critical node in eukaryotic transcription regulation, and its formation is the major rate-limiting step in ... ☆ Save 勿 Cite Cited by 27 Related articles All 3 versions

#### Decreased enhancer-promoter proximity accompanying enhancer activation

NS Benabdallah, I Williamson, RS Illingworth, L Kane... - Molecular cell, 2019 - cell.com Enhancers can regulate the promoters of their target genes over very large genomic distances. It is widely assumed that mechanisms of enhancer action involve the ... ☆ Save 勿 Cite Cited by 273 Related articles All 17 versions

# [HTML] Structure and mechanism of the RNA polymerase II transcription machinery

AC Schier, DJ Taatjes - Genes & development, 2020 - genesdev.cshlp.org Abstract RNA polymerase II (Pol II) transcribes all protein-coding genes and many noncoding RNAs in eukaryotic genomes. Although Pol II is a complex, 12-subunit enzyme, it ... ☆ Save 勿 Cite Cited by 240 Related articles All 9 versions

### Structure of the human Mediator-bound transcription preinitiation complex

R Abdella, A Talyzina, S Chen, CJ Inouye, R Tjian... - Science, 2021 - science.org Eukaryotic transcription requires the assembly of a multisubunit preinitiation complex (PIC) composed of RNA polymerase II (Pol II) and the general transcription factors. The coactivator ... ☆ Save 勿 Cite Cited by 127 Related articles All 10 versions

### Structures of the human Mediator and Mediator-bound preinitiation complex

X Chen, X Yin, J Li, Z Wu, Y Qi, X Wang, W Liu, Y Xu - Science, 2021 - science.org INTRODUCTION The multisubunit Mediator binds RNA polymerase II (Pol II), transduces regulatory signals from transcription factors to Pol II, facilitates preinitiation complex (PIC) ... ☆ Save 夘 Cite Cited by 116 Related articles All 6 versions

### [HTML] Application of iterative soft thresholding for fast reconstruction of NMR data non-uniformly sampled with multidimensional Poisson Gap scheduling

SG Hyberts, AG Milbradt, AB Wagner... - Journal of biomolecular ..., 2012 - Springer The fast Fourier transformation has been the gold standard for transforming data from time to frequency domain in many spectroscopic methods, including NMR. While reliable, it has as ... ☆ Save 勿 Cite Cited by 465 Related articles All 15 versions

# The Mediator complex and transcription regulation

ZC Poss, CC Ebmeier, DJ Taatjes - Critical reviews in biochemistry ..., 2013 - Taylor & Francis The Mediator complex is a multi-subunit assembly that appears to be required for regulating expression of most RNA polymerase II (pol II) transcripts, which include protein-coding and ... ☆ Save 勿 Cite Cited by 423 Related articles All 11 versions

**IPDF1** nature.com

[HTML] nature.com

[HTML] nature.com

[HTML] nature.com

[PDF] cell.com Full View

[HTML] cshlp.org

Prof. L S Shashidhara

Shout.


[PDF] science.org

[PDF] science.org

[HTML] springer.com

[PDF] tandfonline.com

Goooooooogle > 1 2 3 4 5 6 7 8 9 10



Since 2024

Since 2023

Since 2020

Custom range...

Sort by relevance

Create alert

Sort by date

Articles

Page 2 of about 77 results (0.02 sec)

E1 ubiquitin-activating enzyme UBA-1 plays multiple roles throughout C. elegans development

Search within citing articles

[нтмь] The ubiquitin proteasome system in Caenorhabditis elegans and its regulation

N Papaevgeniou, N Chondrogianni - Redox biology, 2014 - Elsevier

Protein degradation constitutes a major cellular function that is responsible for maintenance of the normal cellular physiology either through the degradation of normal proteins or ...

☆ Save 切 Cite Cited by 100 Related articles All 13 versions

De Novo Assembly of the Manila Clam *Ruditapes philippinarum* Transcriptome Provides New Insights into Expression Bias, Mitochondrial Doubly Uniparental ...

F Ghiselli, L Milani, PL Chang... - Molecular biology ..., 2012 - academic.oup.com Males and females share the same genome, thus, phenotypic divergence requires differential gene expression and sex-specific regulation. Accordingly, the analysis of ... ☆ Save 切 Cite Cited by 116 Related articles All 14 versions

Synaptic protein degradation controls sexually dimorphic circuits through regulation of DCC/UNC-40

Y Salzberg, V Pechuk, A Gat, H Setty, S Sela... - Current biology, 2020 - cell.com
Sexually dimorphic circuits underlie behavioral differences between the sexes, yet the
molecular mechanisms involved in their formation are poorly understood. We show here that ...

☆ Save 切 Cite Cited by 25 Related articles All 12 versions

Ubiquitination and the proteasome as drug targets in trypanosomatid diseases MJ Bijlmakers - Frontiers in Chemistry, 2021 - frontiersin.org

The eukaryotic pathogens Trypanosoma brucei, Trypanosoma cruzi and Leishmania are responsible for debilitating diseases that affect millions of people worldwide. The numbers of ... 

☆ Save 切り Cite Cited by 33 Related articles All 7 versions ≫

The Caenorhabditis elegans Kinesin-3 Motor UNC-104/KIF1A Is Degraded upon Loss of Specific Binding to Cargo

J Kumar, BC Choudhary, R Metpally, Q Zheng... - PLoS ..., 2010 - journals.plos.org
UNC-104/KIF1A is a Kinesin-3 motor that transports synaptic vesicles from the cell body
towards the synapse by binding to PI (4, 5) P2 through its PH domain. The fate of the motor ...

☆ Save 切り Cite Cited by 93 Related articles All 21 versions ≫

The amazing ubiquitin-proteasome system: structural components and implication in aging

EN Tsakiri, IP Trougakos - International review of cell and molecular ..., 2015 - Elsevier Proteome quality control (PQC) is critical for the maintenance of cellular functionality and it is assured by the curating activity of the proteostasis network (PN). PN is constituted of several ... ☆ Save 切り Cite Cited by 75 Related articles All 4 versions

[HTML] The Machado-Joseph disease deubiquitylase ATX-3 couples longevity and proteostasis

K Kuhlbrodt, PC Janiesch, <u>É Kevei</u>, <u>A Segref</u>... - Nature cell ..., 2011 - nature.com
Protein ubiquitylation is a key post-translational control mechanism contributing to different physiological processes, such as signal transduction and ageing. The size and linkage of a ...

☆ Save ワワ Cite Cited by 106 Related articles All 9 versions

The conceivable functions of protein ubiquitination and deubiquitination in reproduction

J Wang, Q Zhou, J Ding, T Yin, P Ye... - Frontiers in Physiology, 2022 - frontiersin.org

Protein ubiquitination with general existence in virtually all eukaryotic cells serves as a significant post-translational modification of cellular proteins, which leads to the degradation ...

☆ Save 切 Cite Cited by 11 Related articles All 4 versions ≫

F-box protein FBXB-65 regulates anterograde transport of the kinesin-3 motor UNC-104 through a PTM near its cargo-binding PH domain

V Sabharwal, SPP Boyanapalli, <u>A Shee</u>... - Journal of cell ..., 2024 - journals.biologists.com Axonal transport in neurons is essential for cargo movement between the cell body and synapses. Caenorhabditis elegans UNC-104 and its homolog KIF1A are kinesin-3 motors ... ☆ Save 切り Cite Cited by 1 Related articles All 5 versions

Adipose gene expression profiles reveal insights into the adaptation of northern Eurasian semi-domestic reindeer (*Rangifer tarandus*)

M Weldenegodguad, K Pokharel, L Niiranen... - Communications ..., 2021 - nature.com Reindeer (Rangifer tarandus) are semi-domesticated animals adapted to the challenging conditions of northern Eurasia. Adipose tissues play a crucial role in northern animals by ... ☆ Save ∇∇ Cite Cited by 15 Related articles All 19 versions

Goooooogle >
Previous 1 2 3 4 5 6 7 8 Nex

[HTML] sciencedirect.com

[PDF] oup.com

[PDF] cell.com Full View

[PDF] frontiersin.org

[PDF] plos.org

Prof. L S Shashidhara

[PDF] frontiersin.org

[HTML] nature.com

[PDF] biologists.com

[PDF] nature.com

Articles

Since 2024

Since 2023

Since 2020 Custom range...

Sort by relevance Sort by date

Create alert

Regulation of sperm gene expression by the GATA factor ELT-1

Search within citing articles

About 25 results (0.02 sec)

Meiotic Development in Caenorhabditis elegans

DY Lui, MP Colaiácovo - Germ Cell Development in C. elegans, 2013 - Springer Caenorhabditis elegans has become a powerful experimental organism with which to study meiotic processes that promote the accurate segregation of chromosomes during the ...

☆ Save 勿 Cite Cited by 133 Related articles All 10 versions

[HTML] The regulation of spermatogenesis and sperm function in nematodes

RE Ellis, GM Stanfield - Seminars in cell & developmental biology, 2014 - Elsevier In the nematode C. elegans, both males and self-fertile hermaphrodites produce sperm. As a result, researchers have been able to use a broad range of genetic and genomic techniques ... ☆ Save 夘 Cite Cited by 103 Related articles All 8 versions

A spatial and temporal map of C. elegans gene expression

WC Spencer, G Zeller, JD Watson, SR Henz... - Genome ..., 2011 - genome.cshlp.org The C. elegans genome has been completely sequenced, and the developmental anatomy of this model organism is described at single-cell resolution. Here we utilize strategies that ... ☆ Save 勿 Cite Cited by 335 Related articles All 10 versions

Spermatogenesis

DS Chu, DC Shakes - Germ cell development in C. elegans, 2013 - Springer During spermatogenesis, pluripotent germ cells differentiate to become efficient delivery vehicles to the oocyte of paternal DNA. Though male and female germ cells both undergo ...

☆ Save 夘 Cite Cited by 109 Related articles All 7 versions

The conserved molting/circadian rhythm regulator NHR-23/NR1F1 serves as an essential co-regulator of *C. elegans* spermatogenesis

JM Ragle, AL Aita, KN Morrison... - ..., 2020 - journals.biologists.com

In sexually reproducing metazoans, spermatogenesis is the process by which uncommitted germ cells give rise to haploid sperm. Work in model systems has revealed mechanisms ... ☆ Save 夘 Cite Cited by 27 Related articles All 10 versions

Characterisation of *Caenorhabditis elegans*sperm transcriptome and proteome

X Ma, Y Zhu, C Li, P Xue, Y Zhao, S Chen, F Yang... - Bmc Genomics, 2014 - Springer Background Although sperm is transcriptionally and translationally quiescent, complex populations of RNAs, including mRNAs and non-coding RNAs, exist in sperm. Previous ...

☆ Save 勿 Cite Cited by 57 Related articles All 15 versions

[нтмь] Transcriptional regulation of gene expression in C. elegans

V Reinke, M Krause, P Okkema - WormBook: the online review of C ..., 2013 - ncbi.nlm.nih.gov Protein coding gene sequences are converted to mRNA by the highly regulated process of transcription. The precise temporal and spatial control of transcription for many genes is an ...

☆ Save 勿 Cite Cited by 57 Related articles All 7 versions

Sex determination in nematode germ cells

RE Ellis - Sexual Development, 2023 - karger.com Background: Animal germ cells differentiate as sperm or as oocytes. These sexual fates are controlled by complex regulatory pathways to ensure that the proper gametes are made at ...

☆ Save 勿 Cite Cited by 6 Related articles All 8 versions

NHR-23 and SPE-44 regulate distinct sets of genes during Caenorhabditis *elegans* spermatogenesis

JM Ragle, KN Morrison, AA Vo, ZE Johnson... - G3, 2022 - academic.oup.com Spermatogenesis is the process through which mature male gametes are formed and is necessary for the transmission of genetic information. While much work has established how ...

☆ Save 勿 Cite Cited by 6 Related articles All 15 versions

SPE-44 implements sperm cell fate

M Kulkarni, DC Shakes, K Guevel, HE Smith - PLoS genetics, 2012 - journals.plos.org The sperm/oocyte decision in the hermaphrodite germline of Caenorhabditis elegans provides a powerful model for the characterization of stem cell fate specification and ...

☆ Save 勿 Cite Cited by 42 Related articles All 21 versions ≫

[HTML] nih.gov

[HTML] sciencedirect.com

[PDF] cshlp.org Free from Publisher

[HTML] nih.gov

[PDF] biologists.com

[PDF] springer.com

Prof. L S Shashidhara

[HTML] nih.gov

[PDF] karger.com

[PDF] oup.com **Full View** 

[PDF] plos.org

Gooogle > 1 2 3

Next