






## ARTICLE


<https://doi.org/10.1038/s41467-020-17325-y>

OPEN

# A rare variant of African ancestry activates 8q24 lncRNA hub by modulating cancer associated enhancer


Kaivalya Walavalkar <sup>1</sup>, Bharath Saravanan<sup>1,2</sup>, Anurag Kumar Singh<sup>1</sup>, Ranveer Singh Jayani <sup>3</sup>, Ashwin Nair<sup>1,2</sup>, Umer Farooq<sup>1,4</sup>, Zubairul Islam<sup>1,2</sup>, Deepanshu Soota<sup>1</sup>, Rajat Mann<sup>1</sup>, Padubidri V. Shivaprasad <sup>1</sup>, Matthew L. Freedman <sup>5,6,7</sup>, Radhakrishnan Sabarinathan<sup>1</sup>, Christopher A. Haiman<sup>8</sup> & Dimple Notani <sup>1</sup>✉

Genetic variation at the 8q24 locus is linked with the greater susceptibility to prostate cancer in men of African ancestry. One such African ancestry specific rare variant, rs72725854 (A>G/T) (~6% allele frequency) has been associated with a ~2-fold increase in prostate cancer risk. However, the functional relevance of this variant is unknown. Here we show that the variant rs72725854 is present in a prostate cancer-specific enhancer at 8q24 locus. Chromatin-conformation capture and dCas9 mediated enhancer blocking establish a direct regulatory link between this enhancer and lncRNAs PCAT1, PRNCR1 and PVT1. The risk allele ('T') is associated with higher expression of PCAT1, PVT1 and c-myc in prostate tumors. Further, enhancer with the risk allele gains response to androgen stimulation by recruiting the transcription factor SPDEF whereas, non-risk alleles remain non-responsive. Elevated expression of these lncRNAs and c-myc in risk allele carriers may explain their greater susceptibility to prostate cancer.

<sup>1</sup>Genetics and Development, National Centre for Biological Sciences, Tata Institute of Fundamental Research, Bangalore, Karnataka 560065, India. <sup>2</sup>Sastra Deemed University, Thanjavur, Tamil Nadu 613401, India. <sup>3</sup>Howard Hughes Medical Institute, Department of Medicine, University of California San Diego, 9500 Gilman Drive, La Jolla, CA 92037, USA. <sup>4</sup>Trans-Disciplinary University, IVRI road, Bangalore, Tamil Nadu 560064 Karnataka, India. <sup>5</sup>Department of Medical Oncology, Dana-Farber Cancer Institute, Boston, MA 02215, USA. <sup>6</sup>The Eli and Edythe L. Broad Institute, Cambridge, MA 02142, USA. <sup>7</sup>Centre for Functional Cancer Epigenetics, Dana-Farber Cancer Institute, Boston, MA 02215, USA. <sup>8</sup>Department of Preventive Medicine, Keck School of Medicine, University of Southern California, Los Angeles, CA 90007, USA. ✉email: [dnotani@ncbs.res.in](mailto:dnotani@ncbs.res.in)

# An interdependent network of functional enhancers regulates transcription and EZH2 loading at the *INK4a/ARF* locus

## Graphical abstract



## Authors

Umer Farooq, Bharath Saravanan, Zubairul Islam, ..., Sweetly Meel, Sudha Swaminathan, Dimple Notani

## Correspondence

dnotani@ncbs.res.in

## In brief

A dense enhancer cluster adjacent to the *INK4a/ARF* locus is the most reproducible GWAS hotspot. Using a series of enhancer deletions, Farooq et al. show that only a subset of enhancers loop with promoters. These interacting enhancers form a single functional unit that relies completely on each enhancer for the gene regulation.

## Highlights

- Only a few enhancers from the dense multi-enhancer cluster regulate the *INK4a/ARF* locus
- Functional enhancers are not defined by high levels of H3K27ac or eRNAs
- Deletion of a single functional enhancer renders the entire SE non-functional
- Enhancer activation prevents EZH2 loading onto the *INK4a/ARF* promoters

