

Comparison of dietary proportions for sources Z and G and Enteromorpha 1.00 -0.75 -Proportion ... 0.25 -0.00 -Z and G Enteromorpha Source

Comparison of dietary proportions for sources Z and G and Enteromorpha 1.00 -0.75 -Proportion ... 0.25 -0.00 -Z and G Enteromorpha Source

Comparison of dietary proportions between sources 0.75 -Proportion -0.25 -0.00 -Grass Zostera U.lactuca Enteromorpha Source

Comparison of dietary proportions between sources 0.75 -Proportion -0.25 -0.00 -Grass Enteromorpha Zostera U.lactuca Source

Comparison of dietary proportions between sources 8.0 0.6 Proportion 9.9 0.2 -0.0 Grass Zostera U.lactuca Enteromorpha Source

Comparison of dietary proportions between sources 0.75 -Proportion 0.50 0.25 -0.00 -Grass Zostera U.lactuca Enteromorpha Source

Comparison of dietary proportions between sources 0.75 -Proportion 0.500 J 0.25 -0.00 -Grass Zostera U.lactuca Enteromorpha Source

Comparison of dietary proportions between sources 1.00 -0.75 -Proportion ... 0.25 -0.00 -Grass Zostera U.lactuca Enteromorpha Source


Comparison of dietary proportions between sources 1.00 -0.75 -Proportion ... 0.25 -0.00 -Grass Zostera U.lactuca Enteromorpha Source

Comparison of dietary proportions between sources 1.00 -0.75 -Proportion 0.25 -0.00 -Grass Zostera U.lactuca Enteromorpha Source

Comparison of dietary proportions between sources 1.00 -0.75 -Proportion ... 0.25 -0.00 -Grass Zostera U.lactuca Enteromorpha Source

Comparison of dietary proportions between sources 1.00 -0.75 -Proportion -0.25 -0.00 -Grass U.lactuca Zostera Enteromorpha Source


Comparison of dietary proportions for source Zostera 0.75 -Proportion ... 0.25 -0.00 -Period 5 Period 1 Period 2 Period 3 Period 4 Period 6 Period 7 Period 8 Group



Comparison of dietary proportions for source Zostera 0.75 -Proportion 0.25 -0.00 -Period 5 Period 1 Period 2 Period 3 Period 4 Period 6 Period 7 Period 8 Group


Comparison of dietary proportions for source Zostera 0.75 -0.50 -Proportion 0.25 -0.00 -Period 5 Period 8 Period 2 Period 3 Period 4 Period 6 Period 7 Group


Comparison of dietary proportions for source Zostera 1.00 -0.75 -Proportion 0.25 -0.00 -Period 5 Period 1 Period 2 Period 3 Period 4 Period 6 Period 7 Period 8 Group





Comparison of dietary proportions for source Zostera 1.00 -0.75 -Proportion 0.25 -0.00 -Period 5 Period 1 Period 2 Period 3 Period 4 Period 6 Period 7 Period 8 Group


Comparison of dietary proportions for source Zostera 0.75 -0.50 -Proportion 0.25 -0.00 -Period 5 Period 8 Period 2 Period 3 Period 4 Period 6 Period 7 Group











Tracers plot 15.0 -Mixtures Period 1 Mixtures Period 2 12.5 -Mixtures Period 3 Mixtures Period 4 Mixtures Period 5 d15N_PI Mixtures Period 6 X Mixtures Period 7 Mixtures Period 8 10.0 -0 Zostera × 0 Grass U.lactuca Enteromorpha 7.5 -+ -30 -25 -20 -15 -10 d13C_PI

