



**HKU  
Med**

**LKS Faculty of Medicine**  
**School of Nursing**  
香港大學護理學院

---

**NURS2600**  
**NURSING OF ADULT I**

# **Cardiovascular System III**

## **Coronary Artery Diseases (CAD)**

---

**DR. JOANNA KO**  
**LECTURER**

**TEL: +852 3917 6630 / EMAIL: [KOJOANNA@HKU.HK](mailto:KOJOANNA@HKU.HK)**


# Learning Outcomes

---

After this lecture, students will be able to:

- Understand the definition and pathophysiology of CAD
- Identify different CAD such as atherosclerosis, angina pectoris and acute coronary syndrome (ACS)/acute myocardial infarction (AMI)
- Familiarize with the risk factors and clinical manifestations of ACS/AMI
- Understand the medical and surgical management of treating ACS/AMI
- Demonstrate the nursing process to manage patients with ACS/AMI

# Anatomy of Coronary Arteries



# Coronary Artery Disease (CAD)

---


## Coronary artery disease

- I. Atherosclerosis
- II. Angina Pectoris
- III. Acute Coronary Syndrome / Acute Myocardial Infarction (AMI)

Major Coronary Arteries that supply the  with O<sub>2</sub> and nutrient become damaged or diseased or containing deposits (plaque) in these coronary arteries and inflammation are usually to blame for coronary artery disease.

# I. Atherosclerosis

## Atherosclerosis





# Pathophysiology of Atherosclerosis


Deposition of fats, cholesterol, calcium, and other substances in of endothelium of coronary arteries

The loss of arterial elasticity due to vessel thickening and stiffening

Accumulation of lipid complexes triggers the inflammatory response of immune system

Macrophages are drawn to the site to scavenge the materials

Macrophages fill with lipids to form atherosclerotic plaque (+/-rupture)




# Consequence of Atherosclerosis

---

➤ Coronary artery stenosis

➤ Myocardial ischemia

➔ Acute coronary syndrome/myocardial infarction



# Risk Factors of Atherosclerosis

---

## Non-modifiable

- Family history of CAD
- Increasing age (> 45 years for men; > 55 years for women)
- Gender (men develop CAD at an earlier age than women)
- Race (higher incidence in African Americans than in Caucasians)

## Modifiable

- Hyperlipidemia (LDL: harmful effect on the arterial wall)
- Cigarette smoking, tobacco use
- Hypertension
- Diabetes
- Metabolic syndrome
- Obesity
- Physical inactivity




## II. Angina Pectoris

- A medical term for chest pain or discomfort due to coronary heart disease
- The heart muscle demand > supply


### Types of Angina Pectoris

- **Stable Angina:** predictable and consistent pain that occurs on exertion and is relieved by rest and / or nitroglycerin (TNG)
- **Unstable angina:** Due to rupture of plaque and symptoms increase in frequency and severity; it **may not be relieved with rest or nitroglycerin (TNG)**



# III. Acute Coronary Syndrome

- An umbrella term for conditions caused by sudden blockage of the blood supply to the heart:
  - Unstable angina
  - ECG change
  - Elevation of cardiac enzymes




<http://medicalnotesonline.blogspot.com/2011/01/cardiology-myocardial-infarction-and.html>

# Cardiac Markers

- Cardiac markers are used in the diagnosis and risk stratification of patients with chest pain and suspected acute coronary syndrome (ACS).
- Substances that are released into the blood when the heart is damaged or stressed

Cardiac markers	Descriptions	Indications
Cardiac troponin	<ul style="list-style-type: none"><li>Regulatory protein complex</li><li>Two cardiac-specific isoforms: T and I</li></ul>	<ul style="list-style-type: none"><li>Injury to heart</li><li>Released within 3-4 hours</li></ul>
Creatine Kinase (CK)	<ul style="list-style-type: none"><li>Enzyme</li><li>CK-MM, found mostly in skeletal muscles</li><li>CK-MB, found mostly in the heart muscles</li><li>CK-BB, found mostly in brain tissues</li></ul>	<ul style="list-style-type: none"><li>Injury to skeletal muscle and/or heart cells</li><li>3 to 6 hours after injury, peaks in 18 to 24 hours</li></ul>
Myoglobin	<ul style="list-style-type: none"><li>Heme protein that helps transport oxygen</li><li>Found in cardiac and skeletal muscle</li></ul>	<ul style="list-style-type: none"><li>Injury to skeletal muscle and/or heart cells</li><li>not specific in indicating cardiac event</li></ul>

# Cardiac Markers



<https://www.biomerieux-diagnostics.com/vidasr-acute-coronary-syndrome-ac-s-panel>

OPEN ACCESS  
RESEARCH ARTICLE

PDF/EPUB

Tools Share

# High-sensitive Troponin I (HsTnI)

## Jump to

Abstract  
Methods  
Results  
Discussion  
Sources of Funding  
Disclosures  
Footnotes  
References  
eLetters

## High-Sensitivity Troponin I Levels and Coronary Artery Disease Severity, Progression, and Long-Term Outcomes

Ayman Samman Tahhan, Pratik Sandesara, Salim S. Hayek, Muhammad Hammadah, Ayman Alkhoder, Heval M. Kelli, Matthew Topel, Wesley T. O'Neal, Nima Ghasemzadeh, Yi-An Ko, Mohamad Mazen Gafeer, Naser Abdelhadi, Fahad Choudhary, Keyur Patel, Agim Beshiri, Gillian Murtagh, Jonathan Kim, Peter Wilson, Leslee Shaw, Viola Vaccarino, ... [See all authors](#)

Originally published 21 Feb 2018 | <https://doi.org/10.1161/JAHA.117.007914> | Journal of the American Heart Association. 2018;7:e007914

### Abstract

#### Background

The associations between high-sensitivity troponin I (hsTnI) levels and coronary artery disease (CAD) severity and progression remain unclear. We investigated whether there is an association between hsTnI and angiographic severity and progression of CAD and whether the predictive value of hsTnI level for incident cardiovascular outcomes is independent of CAD severity.

#### Methods and Results

In 3087 patients (aged  $63 \pm 12$  years, 64% men) undergoing cardiac catheterization without evidence of acute myocardial infarction, the severity of CAD was calculated by the number of major coronary arteries with  $\geq 50\%$  stenosis and the Gensini score. CAD progression was assessed in a subset of 717 patients who had undergone  $\geq 2$  coronary angiograms  $> 3$  months before enrollment. Patients were followed up for incident all-cause mortality and incident cardiovascular events. Of the total population, 11% had normal angiograms, 23% had nonobstructive CAD, 20% had 1-vessel CAD, 20% had 2-vessel CAD, and 26% had 3-vessel CAD. After adjusting for age, sex, race, body mass index, smoking, hypertension, diabetes mellitus history, and renal function, hsTnI levels were independently associated with the severity of CAD measured by the Gensini score (log 2  $\beta = 0.31$ ; 95% confidence interval, 0.18–0.44;  $P < 0.001$ ) and with CAD progression (log 2  $\beta = 0.36$ ; 95% confidence interval, 0.14–0.58;  $P = 0.001$ ). hsTnI level was also a significant predictor of incident death, cardiovascular death, myocardial infarction, revascularization, and cardiac hospitalizations, independent of the aforementioned covariates and CAD severity.

#### Conclusions

Higher hsTnI levels are associated with the underlying burden of coronary atherosclerosis, more rapid progression of CAD, and higher risk of all-cause mortality and incident cardiovascular events. Whether more aggressive treatment aimed at reducing hsTnI levels can modulate disease progression requires further investigation.

# Point-of-Care Testing for HsTnI

High-sensitive  
troponin I reading  
for only 3  
minutes





<https://blog.hightechcampus.com/htce/siemens-healthineers-bringing-point-of-care-testing-closer-to-the-patient>



# Myocardial Infarction


- **Non-ST elevated MI (NSTEMI)**
  - Usually have **transient** thrombosis or **incomplete** coronary occlusion
  - **ST-segment depression** or **T wave inversion**; Usually do not develop pathological Q wave
- **ST elevated MI (STEMI)**
  - Tend to have **more extensive** MI that is associated with **prolonged and complete** coronary occlusion
  - **ST-segment elevation** in 2 or more contiguous leads; Pathological Q wave usually seen





*acute*


ST elevation MI



Non-ST elevation MI

# Pathological Q Wave

- During MI, Q wave can be normal or abnormal
- Abnormal Q wave usually follows ST elevation
- Wider and deeper than normal Q
- Presence of pathological Q can represent old infarct
- Indication of myocardial necrosis



# Myocardial Infarction (MI) – Clinical Manifestations

---

- Chest pain
  - Sudden onset, not relieved by rest or medications
    - Location: substernal, retrosternal, epigastric area
    - May radiate like angina
    - Quality: heaviness, crushing pressure, burning, constriction
    - More severe
    - Timing: last longer than 15-20 mins
- Dyspnoea
- Sweating, weakness
- Nausea & vomiting
- Cool, pale and moist skin
- ↑ HR; ↑ RR
- Severe anxiety: has a sense of impending doom and death
- ± loss of consciousness and even sudden death
- *Approximately 1/3 of all myocardial infarctions are silent, without chest pain or other symptoms*



# Myocardial Infarction (MI) – Complications

---

- Arrhythmias
  - Premature ventricular contraction (PVC)
  - Ventricular tachycardia (VT) or ventricular fibrillation (VF)
- Cardiogenic shock
- Infarct extension
- Heart failure
- Structural defect
- Pericarditis

# Assessment of ACS

---

- History taking: past medical history, family history, medication history of antiplatelet or cholesterol lowering drugs
- Physical examination: chest pain, heart murmurs
- Obtain 12-lead ECG to be read within 10 minutes
- Monitor patient's vital signs and connect patient to cardiac monitor
- Obtain laboratory blood specimens of cardiac biomarkers
- Obtain other diagnostics to clarify the diagnosis:
  - Echocardiogram
  - Cardiac catheterization





# COLDSPA

---

## **COLDSPAA** **Symptom Analysis Mnemonic**

<i>Mnemonic</i>	<i>Question</i>
<b>C</b> haracter	Describe the sign or symptom (feeling, appearance, sound, smell or taste if applicable)
<b>O</b> nset	When did it begin?
<b>L</b> ocation	Where is it? Does it radiate? Does it occur anywhere else?
<b>D</b> uration	How long does it last? Does it recur?
<b>S</b> everity	How bad is it? How much does it bother you?
<b>P</b> attern	What makes it better or worse?
<b>A</b> ssociated factors / how it <b>A</b> ffects the pt	What other symptoms occur with it? How does it affect you?

Table 2-3 taken from:  
Lewis, P and Foley, D 2011, *Weber & Kelly's Health Assessment in Nursing*, Lippincott Williams & Wilkins Pty Ltd, a division of Wolters Kluwer Health, NSW.

# Chest Pain – PQRST Assessment

	Factor	Questions to ask patient
P	Precipitating events/Provocation	What events or activities precipitated the pain E.g. exercise
Q	Quality of pain	What does the pain feel like (e.g. pressure, dull, aching, tight, squeezing, heaviness)? Dull pain: MI; Sharp pain: ? Aortic dissection
R	Radiating of pain	Where is the pain located? Any radiation? MI: pain commonly radiate to left shoulder
S	Severity of pain	On a scale of 0 to 10 with 0 indicating no pain and 10 being the most severe pain you would imagine, what no. would you give the pain?
T	Timing	When did the pain begin? Has the pain changed since this time? Have you had pain like this before?










# Implications of Chest Pain

---

CARDIAC	NON-CARDIAC
<ul style="list-style-type: none"><li>• Acute coronary syndrome</li><li>• Aortic dissection</li><li>• Pericarditis</li><li>• Myocarditis</li><li>• Valvular disease</li></ul>	<ul style="list-style-type: none"><li>• Gastrointestinal Esophageal spasm or reflux Peptic ulcer</li><li>• Pulmonary Pneumonia Pulmonary embolism Pneumothorax</li><li>• Neurological (nerve root pain, herpes zoster)</li><li>• Musculoskeletal (e.g. osteochondritis)</li></ul>

**Atypical** presentations of ACS (e.g. epigastric pain, upper back pain or dizziness) are frequent in younger (25-40 years) and older (>75 years) patients and in women. Asymptomatic myocardial ischemia (**silent ischemia**) is particularly common among diabetics.


## CENTRAL ILLUSTRATION: Patient Assessment With Suspected ACS

	Likelihood of myocardial infarction (MI)				
	LOW				HIGH
I. Clinical setting Symptoms and vital signs					 CPR/shock
II. Electro-cardiogram (ECG)	 Normal ECG	 ST depression (mild)	 ST depression	 ST elevation	
III. Troponin level at 0h		—	—/+	+	++ +++
IV. Troponin change (within 1, 2 or 3h)		—	—/+	+	++ If any of the above, consider direct rule-in
Triage decision	Rule-out MI		Observe		Rule-in MI
Differential diagnosis	Noncardiac		Unstable angina	Other cardiac	NSTEMI STEMI

Twerenbold, R. et al. J Am Coll Cardiol. 2017;70(8):996-1012.

# AHA Guideline (2021)

<https://www.ahajournals.org/doi/10.1161/CIR.0000000000001029>




# Diagnostic Investigations of ACS

## 1. Echocardiography/Transthoracic Echocardiography

- Transmission of high-frequency sound waves through the chest wall & the recording of the return signals
- Determine cardiac output (ejection fraction), ventricular wall motion, chamber size & the etiology of heart murmurs
- Evaluate the function of heart valves, including prosthetic heart valves



<https://www.specialistcardiology.com.au/tests/echocardiography/transthoracic-echocardiogram-echo-tte>



[https://www.researchgate.net/figure/Transthoracic-echocardiography-Trans-thoracic-echocardiography-in-A-parasternal\\_fig1\\_318844962](https://www.researchgate.net/figure/Transthoracic-echocardiography-Trans-thoracic-echocardiography-in-A-parasternal_fig1_318844962)




# Diagnostic Investigations of ACS (Cont'd)

---

## 2. Cardiac angiogram

- An invasive test performed to look for narrowings or blockages in the blood vessels that supply the heart
- It is the gold standard diagnostic test for CAD (Bluemke et al., 2008)



<https://unitedcardiology.com.au/tests-procedures/coronary-angiography-angioplasty-and-stenting/>

# Medical Management of ACS

---

## ❖ Goals:

- ✓ Minimize myocardial damage
- ✓ Preserve myocardial function
- ✓ Prevent complications

# Medical Management Options

---

- Evaluate for indication for reperfusion therapy:
  - Reestablish coronary flow
  - **Percutaneous Coronary Intervention (PCI)**: Door-to-balloon inflation time is 90 minutes
  - **Thrombolytic (fibrinolytic) therapy**: Door-to-needle time is 30 minutes
  - **Coronary Artery Bypass Grafting**
- Continue therapy as indicated:
  - IV heparin, low-molecular-weight heparin
  - Antiplatelet: Aspirin, Clopidogrel (Plavix), Ticagrelor (Brilinta)
  - Beta-blocker (Metoprolol, Atenolol)
  - ACEI (Enalapril, Lisinopril)
  - Statin (Simvastatin, Atorvastatin)
  - Nitroglycerin

# Updated Medical Treatment for ACS

- ❖ In 1990, **MONA** was given a Class 1 recommendation for almost all patients with myocardial infarction
- ❖ MONA is no longer as beneficial to the cardiac patient
- ❖ **Morphine** had been found to potentially mask ischemic symptoms
- ❖ **Oxygen** was not found to make a difference in the patients with normal SpO<sub>2</sub>
- ❖ Gouda et al. (2016): oxygen could have paradoxical effects of increased coronary vascular resistance
- ❖ Morphine and oxygen **increased mortality** in patients with ACS

Cureus

Open Access Review  
Article

DOI: 10.7759/cureus.2114

## Morphine, Oxygen, Nitrates, and Mortality Reducing Pharmacological Treatment for Acute Coronary Syndrome: An Evidence- based Review

José Nunes de Alencar Neto <sup>1</sup>

1. Departamento De Cardiologia, Hospital São Paulo

✉ Corresponding author: José Nunes de Alencar Neto, josenunesalencar@gmail.com  
Disclosures can be found in Additional Information at the end of the article

### Abstract

Since it was first reported in 1912, acute coronary syndrome (ACS) has become the leading cause of death in the Western world. Several improvements that have been made over the years in the pharmacological treatment of ACS have reduced the relative risk of death due to myocardial infarction from 35-45% previously to approximately 5.5% at present. Universities, websites, and educational videos commonly use a mnemonic for morphine, oxygen, nitrates, and aspirin (MONA) to refer to the adjuvant treatment used for the management of ACS.

We review the scientific data pertaining to treatment strategies for the management of ACS and discuss whether MONA remains relevant in the present scenario.

While using morphine and oxygen is associated with risks such as higher mortality and increase in the size of the infarct, respectively, several available drugs such as fibrinolytics, anticoagulants, beta-blockers, renin-angiotensin-aldosterone system inhibitors, P2Y<sub>12</sub> inhibitors, and statins are known to be useful to treat ACS.

MONA should be viewed as an obsolete teaching and learning aid, and therefore we recommend that its use be discontinued for the management of ACS.

Categories: Cardiology

Keywords: cardiology, adjuvant treatment, myocardial infarction, acute coronary syndrome, morphine, aspirin, oxygen, nitrate

### Introduction And Background

Acute coronary syndrome (ACS) is the leading cause of death in the Western world. In the United States, approximately one in seven deaths is secondary to coronary heart disease. It is estimated that more than one million Americans experience a myocardial infarction (MI) every year, which in effect means that one American experiences an MI every 24 seconds [1].

Diagnostic and treatment modalities have vastly evolved over the years. Since this disease entity was first reported by Herrick in 1912 [2], adjuvant and pharmacological modalities for the management of ACS have evolved greatly and resulted in better pain relief and mortality

Received 11/22/2017  
Review began 11/28/2017  
Review ended 01/20/2018  
Published 01/25/2018

© Copyright 2018  
Alencar Neto. This is an open  
access article distributed under the  
terms of the Creative Commons  
Attribution License CC-BY 3.0.,  
which permits unrestricted use,  
distribution, and reproduction in any  
medium, provided the original  
author and source are credited.

# Nitroglycerin

- Treat episodes of angina (chest pain) in people who have coronary artery disease
- Spray or sublingual vasodilator

Chest pain

Take 1<sup>st</sup> dose of TNG

Not relieved: take 2<sup>nd</sup> dose after 5 minutes

Not relieved: take 3<sup>rd</sup> dose after another 5 minutes


Not relieved: seek medical help!!!



<https://www.indiamart.com/proddetail/nitroglycerin-sublingual-tablet-usp-0-5-mg-16880074548.html>

# Percutaneous Coronary Intervention (PCI)

- Invasive interventional procedures to treat angina and acute MI
- A guide wire is inserted through radial or femoral artery
- Relieves narrowing and obstruction of the arteries by compressing the atheroma
- Improve blood flow → Allows more blood and oxygen delivered to the heart muscle
- Includes:
  - ✧ Percutaneous transluminal coronary angioplasty (PCI)
  - ✧ Intracoronary stent implantation
  - ✧ *Atherectomy*: a non-surgical procedure that inserts a tiny device on the end of a catheter inside the artery and the plaque is "shaved" away
  - ✧ *Brachytherapy*: Transluminal laser revascularization




<https://www.indiamart.com/proddetail/angiography-surgery-19187188848.html>



# Intra-Coronary Stent Implantation

- Drug-eluting stents are placed within the artery
- Stents help prevent
  - Abrupt closure of arteries shortly after PCI
  - Re-stenosis
  - Formation of thrombi
  - Scar tissue within the coronary artery lesion
  - Decrease the risks of artery spasm after PCI



<https://rodnieoro.com/home/percutaneous-coronary-intervention-angioplasty>

**Double anti-platelet therapy:** Patients with coronary artery stents are usually maintained on full doses of **aspirin + anti-platelet drug for at least 1 year**

# Complications of PCI

---

- Acute arrhythmias
- Dissection, perforation of the artery
- Acute MI
- Bleeding, infection and pain from puncture site
- Allergic reaction to the dye used
- Damage to the kidneys caused by the dye used
- Blood clots that can trigger strokes, heart attacks or other serious problems

# Nursing Interventions Before PCI

---


- Instruct the patient to fast (usually for 8 to 12 hours) before the procedure
- Inform the patient about the expected duration of the procedure & advise that it will involve lying on a hard table for < 2 hours
- Reassure the patient that IV medications are given to maintain comfort
- Explain that palpitation may be felt in the chest because of extra heartbeats that almost always occur
- Tell the patient that the injection of a contrast agent may produce a flushed feeling throughout the body
- Encourage the patient to express fears and anxieties

# Nursing Interventions After PCI

---

- Observe the catheter access site for bleeding or hematoma formation
- Assess peripheral pulses and skin temperature in the affected extremity (dorsalis pedis & posterior tibial pulses in the lower extremity, **capillary refill** in the upper extremity)
- BP & HR Q1H X 4 then Q4H until discharge
- Assess the patient for affected extremity pain, numbness, or tingling sensations that may indicate arterial insufficiency
- Screen carefully for arrhythmias by observing the cardiac monitor
- Femoral artery access: the head of the bed elevated no greater than 30 degrees, bed rest >6 hours
- Radial artery access: bed rest 2-3 hours
- Observe for any oozing from the hemostasis band or pressure dressing
- No blood taking and blood pressure monitoring on the affect extremeity
- Administer analgesic medication as prescribed for discomfort

# Hemostasis After PCI



<https://www.hightidehealth.com/sureseal-bandages.html>

Source from: <https://thoracickey.com/arterial-and-venous-access-2/>

## Thrombolytic Therapy

---

- Administer intravenously or directly into the coronary artery in the cardiac catheterization lab

Example: *Streptokinase, recombinant tissue plasminogen activator (rt-PA)*

- Thrombolysis to allow reperfusion
- Cardiac catheterization and other invasive procedures might be required if the thrombus is unable to dissolve
- Should not be used in patient with risk of bleeding



# Thrombolytic Therapy

---

## ❖ Indications:

- Chest pain for longer than 20 mins, unrelieved by nitroglycerin
- ST-segment elevation in at least two leads that face the same site of the heart
- 6-hours from onset of pain

## ❖ Absolute Contraindications:

- Active bleeding
- Known bleeding disorder
- History of hemorrhagic stroke
- History of intracranial vessel malformation
- Recent major surgery or trauma
- Uncontrolled hypertension
- Pregnancy


# Coronary Artery Bypass Grafting (CABG)

- A procedure of bypass a blocked section of a coronary artery and to deliver oxygen to the heart


A blood vessel from another part of the body (e.g. **saphenous vein**, **left internal mammary artery-LIMA**) is grafted and bypass the obstruction

- Indications:
  - Alleviation of angina that cannot be controlled with medication or PCI
  - For **multi-vessels CAD**
  - Treatment for complication from an unsuccessful PCI

## Coronary artery bypass surgery



# Coronary Artery Bypass Grafting



# Nursing Diagnoses of ACS

---

- ✓ *Decreased cardiac output* related to altered contractility and altered heart rate and rhythm
- ✓ *Acute pain* related to an imbalance between myocardial oxygen supply and demand
- ✓ *Anxiety* related to perceived threats of death, pain and/or possible lifestyle changes
- ✓ *Activity intolerance* related to general weakness secondary to decreased cardiac output and poor lung and tissue perfusion
- ✓ *Risk of ineffective peripheral tissue perfusion* related to decreased cardiac output
- ✓ *Ineffective self-health maintenance* related lack of knowledge of disease process, risk factor reduction, rehabilitation, medication

# Nursing Interventions of ACS

Nursing Diagnoses	Nursing interventions
Release pain	<ul style="list-style-type: none"><li>➤ Assess and monitor pain severity</li><li>➤ Obtain 12-lead ECG during pain episode</li><li>➤ Administer medications to release the pain</li></ul>
Promote adequate tissue perfusion	<ul style="list-style-type: none"><li>➤ Advice the patient to stop all activities</li><li>➤ Administer oxygen as ordered</li><li>➤ Administer thrombolytic drugs if applicable</li></ul>
Reduce anxiety	<ul style="list-style-type: none"><li>➤ Assess and monitor verbal &amp; non-verbal signs of anxiety</li><li>➤ Encourage verbalization of feelings, perception and fear</li><li>➤ Provide factual information concerning diagnosis, treatment and prognosis</li></ul>
Balance physical activity with energy conservative activities	<ul style="list-style-type: none"><li>➤ Assess patient's difficulty to perform ADL</li><li>➤ Monitor patient's response to cardiac medication (BP &amp; P)</li><li>➤ Arrange exercise &amp; rest period to avoid fatigue</li></ul>

---

# **Cardiac Rehabilitation**

---




# Cardiac Rehabilitation

---

- A process aiming to reduce cardiovascular risk factors and improve lifestyle
  - For secondary prevention of cardiovascular complications such as mortality, heart failure, myocardial infarction, coronary artery disease
- I. Exercise counseling and training
  - II. Education for heart-healthy living
  - III. Counseling to reduce stress

# Local Cardiac Rehabilitation Programme



# Education on Healthy Living

---

- Cardiac risk factor modification
  - a. smoking cessation
  - b. low fat diet
  - c. exercise
  - d. weight reduction
- Compliance of prescribed medication regime
- Self-care of chest pain (e.g taking TNG, take rest, seek emergency medical care)

# References

---

- American Heart Association. (2016). Guideline for the Management of Heart. *Advanced Cardiovascular Life Support Provider Manual*. American Heart Association.
- Carpenito-Moyet, L. J. (Ed.). (2017). *Nursing diagnosis: Application to clinical practice (15<sup>th</sup> ed.)*. Philadelphia: Wolters Kluwer.
- Centers for Disease Control and Prevention. (2015). *Coronary Artery Disease (CAD)*. Retrieved from [https://www.cdc.gov/heartdisease/coronary\\_ad.htm](https://www.cdc.gov/heartdisease/coronary_ad.htm)
- Hinkle, J. L., & Cheever, K. H. (2014). *Brunner & Suddarth's textbook of medical-surgical nursing (13<sup>th</sup> ed.)*. Philadelphia: Lippincott Williams & Wilkins.
- Lewis, S. L., Bucher, L., Heitkemper, M. M., Harding, M. M., Kwong, J., & Roberts, D. (2016). *Medical-surgical nursing: assessment and management of clinical problems (9<sup>th</sup> ed.)*. St. Louis, Missouri: Elsevier Health Sciences.
- Smeltzer, S.C, Bare, B.G. Hinkle, J.L. & Cheever, K.H. (2014). *Brunner & Suddarth's Textbook of Medical & Surgical Nursing*. (13<sup>th</sup> ed.). Philadelphia: Lippincott Williams & Wilkins.

---

# Thank You

---