



**HKU
Med**

LKS Faculty of Medicine
School of Biomedical Sciences
香港大學生物醫學學院

Enrichment Course in Biology

BIO17-20 Body defense – A brief introduction to Immunology

Vera SF Chan, Assistant Professor

Sections covered in this lecture

- I. a general concept of body defense
- II. introduction to key immune components
- III. introduction to two major types of immune response
- IV. examples of coordinated immune response in health and diseases

BIO17 Learning outcome



HKU
Med

At the end of section I, you should be able to

- describe **three levels** of body defense
- list the characteristics of the **first line of defense**

The immune system is an army of body guards to defend against invading pathogens!



HKU
Med




Our body is equipped with three lines of defense against infection



HKU
Med

Invading
pathogens



1st : **Barriers** – physical, chemical and microbiological components that prevent entry

2nd : **Non-specific immune responses** that rapidly remove invading pathogens

3rd : **Specific immune responses** that target and “memorize” specifically to the invading pathogens and mount specific actions

Barriers: first line of defense



HKU
Med




Figure 2-4 Immunobiology, 6/e. (© Garland Science 200


Immune system: second and third lines of defense



HKU
Med



Lymphoid organs




First line defense in skin



- Outmost epidermis is keratinized – layers of dead cornified cells form a physical barrier that prevent pathogen entry

- Fatty acids secreted by sebaceous gland is antiseptic
- Dermicidin, an anti-microbial peptide, produced by sweat gland suppress microbial growth





Barriers in respiratory tract



HKU
Med

- Ciliated mucus secreting cells:
 1. Slimy mucus traps microbes
 2. Cilia beat towards upper respiratory tract to remove trapped microbes
- Tight junction formed a seal between epithelial cells to prevent entry of microbes into the blood stream
- Defensins, antimicrobial proteins, secreted by respiratory epithelial cells to suppress microbial growth




Barriers in gastrointestinal (GI) tract




HKU
Med


- Saliva contain enzymes that digest microbes
- Gastric and intestinal juices contain digestive enzymes
- Acidic pH in stomach kills microbes
- Antimicrobial defensins are secreted by intestinal epithelial cells
- Symbiotic microbes (normal gut flora) suppress growth of invading pathogens



Other barriers



- Tear /lacrimal gland – produce tears that
 - 1) contain enzymes that digest microbes
 - 2) wash away trapped microbes or dust particles in the eyes and nose



- Acidic secretion in vagina that suppress microbial growth
- Normal vagina flora/microbiota

First line of defense: summary



HKU
Med

The barriers can be:

- **Mechanical/physical**

- Keratinization of skin
- Hairs, cilia and mucus
- Epithelial cells are joined by tight junctions

- **Chemical**

- Acidic pH (skin, stomach, vagina)
- Enzymes: lysozymes (in saliva, sweat, tears), pepsin (gut)
- Fatty acids (skin)
- Antibacterial peptides – e.g. defensins (intestine, lung) and dermcidin (sweat glands)

- **Microbial**

- normal flora (in skin, gut and vagina) keep microbiological balance by competition for nutrients or produce substances toxic to other microbes

BIO18 Learning outcome



HKU
Med

At the end section II, you should be able to

- list major components of the immune system

Common terminology in immunology




HKU
Med

- **Immune responses** – cellular and molecular events that defend the host against pathogens or adverse events
- **Antigen** – any molecules/ cellular structures that induce immune responses; usually from foreign material but may derive from our own tissues (self/auto-antigen).
- **Immunogenicity** – ability of a substance to induce either humoral and/or cell-mediated immune responses
- **Antigen receptors** – specific proteins on cell surface of lymphocytes to interact with antigens and transduce intracellular signals for cell activation
- **Antibody** – also known as immunoglobulin, a protein produced by immune cells (plasma cells) to bind to specific antigens.
- **Humoral immunity** – antibody-mediated immune responses
- **Cell-mediated immunity** – the immune effector functions are carried out by activated immune cells

Immunology made easy



HKU
Med



Our immune system is designed to detect and act upon:

1. Self vs non-self
e.g. infection,
transplant rejection
2. Danger signals
e.g. cancer cells,
tissue damages

Components of immune system



HKU
Med


The immune system includes the body parts that help in the **recognition and destruction/removal** of foreign/harmful materials.

- (i) organs and cells of immune system
- (ii) Non-specific /innate immunity
- (iii) Specific / adaptive immunity

Lymphoid organs/tissues



HKU
Med




(I) Primary lymphoid organs – sites of leukocyte development from precursor cells

- **Bone marrow** – source of haematopoietic stem cells (blood cell precursors)
- Thymus

(II) Secondary lymphoid organs – sites of leukocytes interaction, activation and differentiation

- Spleen
- Lymph nodes
- Tissue associated lymphoid tissues (e.g. Tonsils, Peyer's patches)

An overview of cells in the immune system




Common cell types in the blood

lymphocyte (20-40%)

- T cells – for adaptive immunity
- B cells - for adaptive immunity
- NK cells –killer cells, innate response

Monocyte (3-10%)

- Phagocytic,
- can turn into macrophages in tissues



← **Erythrocyte**

Polymorphonuclear (PMN) leukocyte (45-65%)

- also refer as granulocytes
- most abundant granulocytes in blood is neutrophils
- phagocytic cells

The human immune system

- The **second and third** lines of defense are both part of our active immune system
- All immune cell populations are developed from haematopoietic (from Greek, means “blood making”) stem cells (HSC) of the bone marrow. Cord blood is also another source of hematopoietic stem cells.
- HSC can be differentiated into the major players in the immune system (**granulocytes, monocytes, lymphocytes, dendritic cells**) as well as cells not involved in immune action, such as erythrocytes (red blood cells) and megakaryocytes (for blood clotting).
- White blood cells (or called **leukocytes**) in the body are the immune cell populations that guard us against infections or danger signals

The human immune system



- Immune cells circulate in the body via blood vessels and lymphatics and take residence in **lymphoid organs** (e.g. spleen, lymph nodes) as well as any tissue organs (e.g. tissue macrophages)
- Some immune cells mainly perform non-specific immune actions (**innate immunity**) e.g. macrophages, granulocytes, monocytes, natural killer (NK) cells, dendritic cells
- and some immune cells specialize in antigen-specific immune responses (**adaptive immunity**) e.g. B cells and T cells
- Each cell type is specialized to carry out different functions and **act synergistically** to mediate protection

BIO19 Learning outcome



HKU
Med

At the end of section III, you should be able to

- Understand the major differences between **innate** and **adaptive** immunity
- Give examples of innate and adaptive immunity

Non-specific immunity (innate immunity)



HKU
Med


- If microbes evade first line of defense, the innate immune response will be ready to take action
- Innate responses **pre-exist** in all individuals, and is ready to act **rapidly** (within minutes or hours) upon exposure to microbes
- The action is **non-specific** because it responds in similar fashion to a broad category of micro-organisms, and there is **no enhancement of responses in subsequent re-exposure** to the same pathogens i.e. no immunological memory of pathogen
- Innate immune functions can be carried out **by immune cells** directly or by the **macro-molecules** (humoral factors) they produced.
- Examples of immune cells for innate responses: macrophages, neutrophils, natural killer (NK) cells, mast cells, eosinophils, basophils and dendritic cells

Examples of non-specific immune function -1



HKU
Med

- Direct killing of microbes by **phagocytosis** – a mechanism of “eat and digest” of microbes or cell debris
- Phagocytes (e.g. macrophages, neutrophils) can engulf bacteria and digest them intracellularly



A bacterium being enclosed by a macrophage

- Natural killer (NK) cells can kill infected or abnormal cells in close contact
- Kill by **releasing toxic granules** to penetrate and lyse the target cells directly




A target cell being attacked by two NK cells

Examples of non-specific immune function - 2



Complement cascade

- **Complement proteins** are a family of **serum proteins** that elicit non-specific immune protection
- A **chain reaction** will be initiated by infectious microbes to produce several active complement components (e.g. C3b, C3a, C5a, MAC) which effect in various immune responses



Specific immunity (adaptive/acquired immunity)



HKU
Med


- Specific immunity is **not inborn** but need to be “**acquired**” upon exposure to microbes
- Specific immune response takes days to weeks to develop
- The response is **antigen-specific** as it only recognizes the invading pathogens that initiate the response. It does not cross react with other pathogens, even different strain of the same species.
- Specific immunity has “**immunological memory**” – it gives better and faster responses in the subsequent re-exposure of the same pathogens
- Two major cell types for specific immune responses: **B cells and T cells**

T and B cells have different mechanisms of effector action




HKU
Med

B cells – humoral response




T cells – cellular response




Two arms of adaptive immunity

A. Humoral immunity – B cells




- B cells bear specific antigen receptors (B cell receptor, BCR) on cell surface for pathogens/antigen recognition
- Once activated by pathogens, B cells become plasma cells that secrete specific antibodies
- Antibodies are proteins that interact with antigens directly
- Antibodies are antigen-specific, only respond to the initial activating antigen



- antibodies can inactivate microbes to prevent further cell invasion
- neutralize toxins and infectious particles,
- facilitate response of other immune cells



Two arms of adaptive immunity

B. Cell-mediated immunity – T cells



Cytotoxic T cells

- Carry out direct killing by releasing toxic granules to the infected cells during cell-cell contact



Helper T cells

- Provide “help” for other immune cells (e.g. B cells, NK cells etc.) through secretion of cytokines (immune cells growth factors and function modulators) or by cell-cell contact

- T cells also bear **specific antigen recognition receptors** (T cell receptor, TCR) on the cell surface
- two major subsets of T cells with distinct functions

Summary:

A brief overview of immune system



HKU
Med

1st line: Characteristics of barriers

- Mechanical, chemical, microbial

2nd line: Innate Immune defense

- characteristics
 - inborn, not antigen specific, quick response and short duration, no immunological memory
- examples of innate defense
 - Cellular: phagocytosis, natural killer cytotoxicity
 - Macro-molecules: complements

3rd line: Specific/adaptive immune defense

- Characteristics
 - acquired during life, antigen-specific in action, takes longer to initiate, develop immunological memory
- examples of adaptive response
 - Humoral: antibodies (B cells)
 - Cell-mediated: T cells

This is a very brief overview and many details will be covered in the relevant immunology lectures in the curriculum

BIO20 Learning outcome



HKU
Med

Key learning points:

- a basic understanding on the coordinated events in the induction of acute inflammation
- Some examples to illustrate the role of immune responses in health and diseases

Inflammation




HKU
Med

Inflammation - local accumulation of fluid, plasma proteins, and white blood cells that is initiated by physical injury, infection, or a local immune response.



- acute local Inflammation: coordinated actions of innate immune responses
- features of acute inflammation: redness, heat, swelling and pain
- These cardinal signs appear rapidly after injury

Key steps in inflammation induction



Step 1 – Tissue Injury

- e.g. caused by bacterial infection, physical cut or burn
- leads to bleeding, cell death and/or microbial invasion which activate resident immune cells
- Local resident immune cells (e.g. macrophages and mast cells) have damage or pathogen-sensing receptors
 - activated and release cytokines and chemokines

<http://iahealth.net/inflammation/>


<https://www.youtube.com/watch?v=iZYLelJwe4w>

Key steps in inflammation induction

Step 2: Release of chemical mediators

Chemical mediators include:

- **Histamine and prostaglandins** (lipid mediators) – induce local vascular responses
- **Inflammatory cytokines** – molecules that modulate the actions or growth of immune cells to elicit inflammation responses
- **chemokines** – molecules that attract other immune cells to the site of inflammation



Key steps in inflammation induction



Step 3: effector outcomes caused by the chemical mediators

(a) vasodilation

- Dilation of blood vessels or capillaries
→ increased blood flow to the affected area
→ **red** and **hot** (cardinal signs)
- Heat increases the metabolic rate of cells
→ promote healing



(b) Increased capillary permeability

- Fluid leaks into tissues forming an inflammatory exudate containing proteins and cells
- develop **swelling** which exerts pressure and cause **pain**

Key steps in inflammation induction



(c) Chemotaxis

- **chemokines** attract the migration of other immune cells into the affected site
- blood leukocytes (mainly neutrophils and monocytes) migrate to the injured area
- recruited **neutrophils and resident macrophages are phagocytic**, thus facilitate the killing and removal of invading micro-organisms
- activated neutrophils and macrophages further **produce cytokines** that suppress growth of microbes, activate immune cells and induce tissue repairs
- **adaptive immune responses develop later** and act upon it if the infection cannot be resolved by the innate responses.

Vaccination – harnessing adaptive immunity



HKU
Med




Vaccination – development of immunological memory for pathogens

APPENDIX. Hong Kong Childhood Immunisation Programme

Age	Immunisation RECOMMENDED
Newborn	B. C. G. vaccine Hepatitis B vaccine — first dose
1 Month	Hepatitis B vaccine — second dose
2 Months	DTaP-IPV vaccine — first dose Pneumococcal vaccine — first dose
4 Months	DTaP-IPV vaccine — second dose Pneumococcal vaccine — second dose
6 Months	DTaP-IPV vaccine — third dose Pneumococcal vaccine — third dose Hepatitis B vaccine — third dose
1 Year	MMR (Measles, Mumps & Rubella) vaccine — first dose Pneumococcal vaccine — booster dose Varicella vaccine — first dose
1.5 Years	DTaP-IPV vaccine — booster dose
Primary 1	MMRV (Measles, Mumps, Rubella & Varicella) vaccine — second dose DTaP-IPV vaccine — booster dose
Primary 6	dTaP-IPV vaccine — booster dose

Note: DTaP-IPV vaccine: Diphtheria, Tetanus, acellular Pertussis & Inactivated Poliovirus vaccine; dTaP-IPV vaccine: Diphtheria, Tetanus, acellular Pertussis (reduced dose) & Inactivated Poliovirus vaccine



Allergy: over-reaction of immune responses



Hypersensitivity 4


- **Hypersensitive immune reactions** against environmental substances
- involve **antibody-mediated** and/or **cell-mediated mechanisms** to produce distinct clinical manifestations
- clinical manifestations can be divided into:
 - (i) IgE-mediated allergy e.g. allergic rhinitis (hay fever), allergic asthma, food/drug allergy
 - (ii) Cell-mediated allergy e.g. allergic contact dermatitis (allergy induced by poison Ivy)

Diseases of the immune system



HKU
Med

- *Autoimmune diseases (AID)* arise when **chronic inflammatory** response **against self antigen** is causing tissue damages or malfunctioning of organ systems




Lupus



Rheumatoid arthritis

Body Parts That Can Be Affected by Autoimmune Diseases



Diseases of the immune system



(SCID, severe combined immunodeficiency)

Immunodeficiency

- defects (congenital or acquired) leading to reduced or absence of specific immune components
- recurrent infections – ↑ susceptibility, severity and duration
- ↑ **Opportunistic infections** – caused by microbes that normally do not easily cause serious disease in healthy individuals e.g. candidiasis, streptococcus, HSV etc.
- ↑ **malignancies** incidence



**HKU
Med**

**LKS Faculty of Medicine
The University of Hong Kong
香港大學李嘉誠醫學院**