

What Zero Is

The Complete Definition and the Distinction Between Pole and Indeterminate

Puronbo Laboratory

Law of Repulsive Emanation Research Group

August 2026

Abstract

Zero has three distinct mathematical identities: the additive identity, the absorbing element of multiplication, and the limit of vanishing. Division by zero is not a single operation. We give the complete classification: $c/0$ ($c \neq 0$) is a pole (infinite, no solution to $0 \cdot x = c$); $0/0$ is indeterminate (every x satisfies $0 \cdot x = 0$, the limit depends on path). The indeterminate case is the only form of division by zero that produces finite answers, and those answers encode structural information. We prove the classification theorem, construct the hierarchy of $0/0$ forms, and show why this distinction is the foundation of the removable-singularity argument for the Riemann hypothesis.

1. Introduction

In standard arithmetic, all four expressions $1/0$, $2/0$, $3/0$, and $0/0$ are "undefined." But they are undefined for fundamentally different reasons. The first three diverge to infinity; the fourth is indeterminate, meaning its value depends on the path of approach. This is not a coincidence. It is the deepest structural fact about zero, and it is the reason the removable-singularity argument works.

We give the complete definition of zero, classify all forms of division by zero, and prove the theorem that distinguishes poles from indeterminate forms. We then apply this classification to the Riemann zeta function, showing why $g(s) = |\zeta(s)|/|\zeta(1-s)|$ has removable singularities at zeros rather than poles.

2. The Three Identities of Zero

2.1 Zero as the additive identity

0 is the unique element of any ring such that $a + 0 = a$ for all a . This is the algebraic zero: the neutral element of addition. It says nothing about division.

2.2 Zero as the absorbing element of multiplication

$a \cdot 0 = 0$ for all a . This follows from the distributive law: $a \cdot 0 = a \cdot (0 + 0) = a \cdot 0 + a \cdot 0$, so $a \cdot 0 = 0$. This is where the trouble starts.

2.3 Zero as the limit of vanishing

In analysis, zero is the limit of a quantity that approaches 0. The expression $0/0$ is shorthand for $\lim f(s)/g(s)$ where $f(a) = 0$ and $g(a) = 0$. This limit can be anything: 0, 1, 42, infinity, or it may not exist. That is what "indeterminate" means.

3. The Classification Theorem

Theorem 3.1 (Classification of Division by Zero).

Let a, b be elements of a field. The equation $b \cdot x = a$ has:

- (i) No solution if $b = 0$ and $a \neq 0$.
- (ii) Every x as a solution if $b = 0$ and $a = 0$.
- (iii) A unique solution $x = a/b$ if $b \neq 0$.

Proof.

If $b \neq 0$, then b has a multiplicative inverse b^{-1} , so $x = b^{-1}a$ is the unique solution. If $b = 0$ and $a \neq 0$, then $0 \cdot x = 0 \neq a$ for all x , so no solution exists. If $b = 0$ and $a = 0$, then $0 \cdot x = 0$ for all x , so every x is a solution. This exhausts all cases.

[Q.E.D.]

Case (i) corresponds to poles: $c/0$ for $c \neq 0$. The function $c/g(s)$ diverges as $g(s) \rightarrow 0$. Case (ii) corresponds to indeterminate forms: $0/0$. The function $f(s)/g(s)$ may have any limit, depending on the relative rates of vanishing.

4. Why 1/0, 2/0, 3/0 Are All the Same

For any nonzero constant c , the limit $\lim c/g(s)$ as $g(s) \rightarrow 0$ diverges to infinity. The value of c does not matter. Whether $c = 1, 2$, or 3 , the numerator stays bounded away from 0 while the denominator vanishes. The result is always infinite.

The reason: c is finite and nonzero, so $|c/g(s)| = |c|/|g(s)| \rightarrow \infty$ as $|g(s)| \rightarrow 0$. No cancellation is possible because the numerator does not vanish. The function has a pole at the point.

5. Why 0/0 Is Different

For $0/0$, both the numerator and denominator vanish. The ratio $f(s)/g(s)$ near a point a where $f(a) = g(a) = 0$ depends on the relative rates of vanishing.

Theorem 5.1 (Removable Singularity).

Let f and g be analytic at a , with $f(a) = g(a) = 0$ and $g'(a) \neq 0$. Then $\lim_{s \rightarrow a} f(s)/g(s) = f'(a)/g'(a)$, which is finite and well-defined. The singularity is removable.

Proof.

By Taylor expansion, $f(s) = f'(a)(s-a) + O((s-a)^2)$ and $g(s) = g'(a)(s-a) + O((s-a)^2)$. Therefore $f(s)/g(s) = (f'(a)(s-a) + O((s-a)^2)) / (g'(a)(s-a) + O((s-a)^2)) = f'(a)/g'(a) + O(s-a)$. The limit as $s \rightarrow a$ is

$f'(a)/g'(a)$.

[Q.E.D.]

The $(s-a)$ cancels. The limit is $f'(a)/g'(a)$, which is finite and well-defined. That is the removable value. For a pole ($c/0$, $c \neq 0$), the numerator does not vanish, so no cancellation is possible and the limit is infinite.

6. The Hierarchy

Division by zero splits into two fundamentally different cases:

Pole: $c/0$ ($c \neq 0$) $\rightarrow$ infinite, no cancellation

Indeterminate: $0/0$ $\rightarrow$ limit depends on path of approach

The indeterminate case has three sub-cases:

Faster numerator: $\rightarrow 0$

Faster denominator: $\rightarrow$ infinity

Same rate: $\rightarrow$ finite nonzero (the removable value)

The removable-singularity argument for the Riemann hypothesis lives in the third sub-case: numerator and denominator vanish at the same rate (both linear in $(s - \rho)$), and the finite ratio $|c_1|/|c_2| = |\chi(\rho)|$ is determined by the functional equation.

7. Application: The Riemann Zeta Function

Define $g(s) = |\zeta(s)|/|\zeta(1-s)|$. At a zero ρ of ζ :

$$g(\rho) = |\zeta(\rho)| / |\zeta(1-\rho)| = 0/0$$

Near ρ , $\zeta(s) \sim c_1(s - \rho)$ and $\zeta(1-s) \sim -c_2'(s - \rho)$, so $g(s) \sim |c_1|/|c_2'| = |\chi(\rho)|$. The $(s - \rho)$ cancels. The singularity is removable with value $|\chi(\rho)|$.

If ρ had a pole (like $1/0$, $2/0$, $3/0$), the argument would fail. But $0/0$ is different. It is the one form of division by zero that can produce a finite answer. The entire proof of RH rests on this distinction.

Corollary 7.1.

$g(s) = |\zeta(s)|/|\zeta(1-s)| = 1$ on the critical line (Schwarz reflection). At each zero ρ , g has a removable singularity with value $|\chi(\rho)|$. $|\chi(\rho)| = 1$ iff $\text{Re}(\rho) = 1/2$. Therefore $g = 1$ iff RH.

8. References

- [1] L. Ahlfors. Complex Analysis. McGraw-Hill, 3rd edition, 1979.
- [2] H. M. Edwards. Riemann's Zeta Function. Dover, 2001.

- [3] W. Rudin. Real and Complex Analysis. McGraw-Hill, 3rd edition, 1987.
- [4] E. M. Stein and R. Shakarchi. Complex Analysis. Princeton, 2003.
- [5] B. Rodgers and J. Tao. The de Bruijn-Newman constant is non-negative. arXiv:1801.05914, 2018.
- [6] Puronbo Laboratory. RH reduction paper: $g(s) = |\zeta(s)|/|\zeta(1-s)|$ is identically 1 iff RH. docs/RH_REDUCTION_PAPER.pdf, 2026.