


Question: $T = 9.5$. If $\log_2 x^T - \log_4 x = \log_8 x^k$ is an identity for all $x > 0$, compute the value of k .

Construction-based method



[DECOMPOSE] Break down the problem into subquestions: 1. What is ... 2. How to ...

[COT] 1. First, ... The equation given is: $\log_2(x^T) - \log_4(x) = \log_8(x^k)$...

[VERIFY] Verify the answer ...


[POT] ... Therefore, $k=27$ is the right answer.

Final Answer:

27

[Template]
Reasoning
Meta-Thinking

R1-like method



Okay, so I need to ... Hmm, let's break this down step by step.

... Given that $k = 3 \cdot (9.5 - 0.5) = 3 \cdot (9) = 27$

Wait, hold on. Let me double-check my steps.


Therefore, $k=27$ is the right answer. I think that's solid. ... So I'm confident ...

Final Answer is

27

Reasoning
Meta-Thinking

ReMA: Reinforced Meta-thinking Agents



Meta-Thinking

1. Let me rewrite each term using the natural logarithm ($\ln$).

2. Starting with the left-hand side: 1. $(\log_2 x^T)$: This can be written as $\frac{\ln x^T}{\ln 2}$...

3. Since this equation must hold for all $x > 0$, we can divide both sides by $\frac{\ln x}{\ln 2}$...

4. Solve for k : $k = 9 \cdot 3 = 27$...

5. Wait, let me double-check this step to ensure there are no mistakes...[FINISH]

6. The final answer is

27

Reasoning
Meta-Thinking

Reasoning steps

