

SOLUTIONS MANUAL FOR

---

Heat Conduction  
Fourth Edition

---

by

Yemen Yener  
Sadik Kakac



CRC Press  
Taylor & Francis Group



# SOLUTIONS MANUAL FOR

---

## Heat Conduction

Fourth Edition

---

\_\_\_\_\_ by \_\_\_\_\_

Yemen Yener  
Sadik Kakac



CRC Press

Taylor & Francis Group

Boca Raton London New York

---

CRC Press is an imprint of the  
Taylor & Francis Group, an **informa** business

CRC Press  
Taylor & Francis Group  
6000 Broken Sound Parkway NW, Suite 300  
Boca Raton, FL 33487-2742

© 2012 by Taylor & Francis Group, LLC  
CRC Press is an imprint of Taylor & Francis Group, an Informa business

No claim to original U.S. Government works

Printed in the United States of America on acid-free paper  
Version Date: 20111102

International Standard Book Number: 978-1-4665-0784-5 (Paperback)

This book contains information obtained from authentic and highly regarded sources. Reasonable efforts have been made to publish reliable data and information, but the author and publisher cannot assume responsibility for the validity of all materials or the consequences of their use. The authors and publishers have attempted to trace the copyright holders of all material reproduced in this publication and apologize to copyright holders if permission to publish in this form has not been obtained. If any copyright material has not been acknowledged please write and let us know so we may rectify in any future reprint.

Except as permitted under U.S. Copyright Law, no part of this book may be reprinted, reproduced, transmitted, or utilized in any form by any electronic, mechanical, or other means, now known or hereafter invented, including photocopying, microfilming, and recording, or in any information storage or retrieval system, without written permission from the publishers.

For permission to photocopy or use material electronically from this work, please access [www.copyright.com](http://www.copyright.com) (<http://www.copyright.com/>) or contact the Copyright Clearance Center, Inc. (CCC), 222 Rosewood Drive, Danvers, MA 01923, 978-750-8400. CCC is a not-for-profit organization that provides licenses and registration for a variety of users. For organizations that have been granted a photocopy license by the CCC, a separate system of payment has been arranged.

**Trademark Notice:** Product or corporate names may be trademarks or registered trademarks, and are used only for identification and explanation without intent to infringe.

Visit the Taylor & Francis Web site at  
<http://www.taylorandfrancis.com>

and the CRC Press Web site at  
<http://www.crcpress.com>

# CHAPTER 1

## FOUNDATIONS OF HEAT TRANSFER

---

PROB. 1.1: Since the flow is steady, the first law (1.31) reduces to

$$\int_{c.s.} \rho (h + \frac{1}{2} V^2 + gz) \vec{V} \cdot \hat{n} dA = q_{c.s.} - \dot{W}_{shear} - \dot{W}_{shaft} \int_{c.v.} \dot{q}_e d\sigma \quad (1)$$

Assume that all the properties are constant over cross-sections ① and ②. Thus,

$$\int_{c.s.} \rho (h + \frac{1}{2} V^2 + gz) \vec{V} \cdot \hat{n} dA = \rho_2 (h_2 + \frac{1}{2} V_2^2 + gz_2) V_2 A_2 - \rho_1 (h_1 + \frac{1}{2} V_1^2 + gz_1) V_1 A_1$$

The continuity condition yields

$$\rho_1 A_1 V_1 = \rho_2 A_2 V_2 = \dot{m}$$

Thus,

$$\int_{c.s.} \rho (h + \frac{1}{2} V^2 + gz) \vec{V} \cdot \hat{n} dA = \dot{m} [(h_2 + \frac{1}{2} V_2^2 + gz_2) - (h_1 + \frac{1}{2} V_1^2 + gz_1)]$$

Furthermore, if it is further assumed that the flow is frictionless, then  $\dot{W}_{shear} = 0$ . In addition,

$$q_{c.s.} = q, \quad \dot{W}_{shaft} = \dot{W} \quad \text{and} \quad \dot{q}_e = 0$$


Therefore, the steady-flow energy equation (1) can be rewritten as

---


$$h_1 + \frac{1}{2} V_1^2 + gz_1 + \frac{q}{\dot{m}} = h_2 + \frac{1}{2} V_2^2 + gz_2 + \frac{\dot{W}}{\dot{m}}$$


---

PROB. 1.2:



Since  $\vec{V} = 0$ ,  $\dot{W}_{\text{shear}} = \dot{W}_{\text{shaft}} = 0$ , Eq. (1.30) reduces to

$$\int_V g \frac{\partial e}{\partial t} dv = \dot{q}_s + \int_V \dot{q}_e dv \quad \dots \quad (1)$$

$$\int_V g \frac{\partial u}{\partial t} dv = \int_V g c \frac{\partial T}{\partial t} dv \quad \dots \quad (2)$$

for solids  $du = c dT$ ,  $c = \text{specific heat}$

Furthermore,

$$\dot{q}_s = \int_S \vec{q} \cdot \hat{n} ds = \int_V \vec{V} \cdot \vec{q}'' dv \quad \dots \quad (3)$$

Here,  $\vec{q}''$  is the heat flux vector at  $dS$ , where  $\hat{n}$  represents the outward-drawn unit vector normal to  $dS$ . Then, by substituting Eqs.(2) and (3) into (1), we obtain

$$\int_V \left[ -\vec{V} \cdot \vec{q}'' + \dot{q}_e - g c \frac{\partial T}{\partial t} \right] dv = 0$$

Since this volume integral vanishes for every volume element  $V$ , its integrand must vanish everywhere, thus yielding

$$-\vec{V} \cdot \vec{q}'' + \dot{q}_e = g c \frac{\partial T}{\partial t}$$

PROB. 1.3: Under the conditions stated

the 1st law gives:


$$q(x) = q(x+\Delta x)$$

As  $\Delta x \rightarrow 0$ ,

$$q(x+\Delta x) = q(x) + \frac{dq}{dx} \Delta x$$

Hence, as  $\Delta x \rightarrow 0$ , the first law reduces to

$$\frac{dq}{dx} = 0 \text{ or } q(x) = \text{const.}$$



PROB. 1.4:

$$\frac{dq}{dx} = 0 \quad \begin{matrix} \uparrow \\ \text{First law} \end{matrix} \quad \& \quad q = -kA \frac{dT}{dx} \quad \begin{matrix} \uparrow \\ \text{Fourier's law} \end{matrix} \Rightarrow \underbrace{\frac{d}{dx} \left( -kA \frac{dT}{dx} \right)}_{\frac{d^2T}{dx^2}} = 0$$

$$\frac{d^2T}{dx^2} = 0 \Rightarrow T(x) = Ax + B$$

$$\left. \begin{array}{l} T(0) = T_1 \Rightarrow B = T_1 \\ T(L) = T_2 \Rightarrow A = \frac{T_2 - T_1}{L} \end{array} \right\} \quad \therefore T(x) = T_1 - \frac{T_1 - T_2}{L} x$$

PROB. 1.5: The first law gives (see Prob. 1.3)


$$q = -kA \frac{dT}{dx} = \text{Const.} \Rightarrow k \frac{dT}{dx} = -\frac{q}{A}$$

$$\text{or } k_0(1+\beta T) dT = -\frac{q}{A} dx$$

$$\therefore [T(x_1) - T(x_2)] + \beta \frac{1}{2} [T^2(x_1) - T^2(x_2)] = -\frac{q}{Ak_0} (x_1 - x_2)$$

$$T_3 - T_1 + \frac{\beta}{2} (T_3^2 - T_1^2) = -\frac{qL}{Ak_0} \quad \text{--- (1)}$$

$$T_2 - T_3 + \frac{\beta}{2} (T_2^2 - T_3^2) = -\frac{qL}{Ak_0} \quad \text{--- (2)}$$




These two equations yield:

$$\left. \begin{aligned} \beta &= -2 \frac{T_1 + T_2 - 2T_3}{T_1^2 + T_2^2 - 2T_3^2} \\ k_0 &= \frac{2(\varrho/A) \times L}{(T_2 - T_1) \left[ 1 + \frac{\beta}{2}(T_1 + T_2) \right]} \end{aligned} \right\} \Rightarrow k(T) = k_0 [1 + \beta T]$$


---

PROB. 1.6:


$$\begin{aligned} dT &= \frac{\partial T}{\partial x} dx + \frac{\partial T}{\partial y} dy + \frac{\partial T}{\partial z} dz \\ &= \vec{\nabla} T \cdot (\overset{i}{dx} + \overset{j}{dy} + \overset{k}{dz}) \\ &= \vec{\nabla} T \cdot d\vec{r} \end{aligned}$$



On the  $T = \text{Const.}$  surface  $dT = 0$ . Thus,  $\vec{\nabla} T$  is a vector normal to all such  $d\vec{r}$ . On the other hand,  $d\vec{r}$  is a vector tangent to the isothermal surface passing through P. Therefore,  $\vec{\nabla} T$  is a vector normal to the isothermal surface.


Again,  $dT = \vec{\nabla} T \cdot d\vec{r}$

In order for  $dT > 0$ ,  $\cos \theta > 0$ ; that is,  $\vec{\nabla} T$  has to point in the direction of increasing temperature as shown.



PROB. 1.7:

$$\vec{\nabla}T \cdot \hat{s} = \left( \hat{i} \frac{\partial T}{\partial x} + \hat{j} \frac{\partial T}{\partial y} + \hat{k} \frac{\partial T}{\partial z} \right) \cdot \left( \hat{i} \cos\alpha + \hat{j} \cos\beta + \hat{k} \cos\gamma \right)$$



$$= \frac{\partial T}{\partial x} \underbrace{\cos\alpha}_{dx/ds} + \frac{\partial T}{\partial y} \underbrace{\cos\beta}_{dy/ds} + \frac{\partial T}{\partial z} \underbrace{\cos\gamma}_{dz/ds}$$

$$= \frac{\partial T}{\partial s}$$

PROB. 1.8:

$$T(^{\circ}\text{C}) = 150 - 400x^2$$


$$q''_1 = -k \left( \frac{\partial T}{\partial x} \right)_{x=0} = -k (-800x)_{x=0}$$

$$= 0 \text{ W/m}^2$$

$$q''_2 = -k \left( \frac{\partial T}{\partial x} \right)_{x=L} = -k (-800L)$$

$$= 40 \times 800 \times 0.25$$

$$= 8,000 \text{ W/m}^2$$




PROB. 1.9:

$$q_1 = -kA \left( \frac{\partial T}{\partial x} \right)_{x=0} \text{ and } q_2 = -kA \left( \frac{\partial T}{\partial x} \right)_{x=L}$$

$$-\left( \frac{\partial T}{\partial x} \right)_{x=0} < -\left( \frac{\partial T}{\partial x} \right)_{x=L}$$

$$q_1 < q_2$$

$\therefore$  The wall is being cooled at  $t_1$ .



PROB. 1.10:


$$h = \frac{-k \left( \frac{\partial T}{\partial y} \right)_{y=0}}{T_w - T_\infty} = \frac{-(0.62) \frac{W}{m \cdot K} \times (-80 \times 800) \frac{K}{m}}{(100 - 20) K}$$

$$= 496 \frac{W}{m^2 \cdot K}$$



---

PROB. 1.11:

Energy balance yields



$$1,500 \text{ W/m}^2$$



$$\alpha_s (1,500) \frac{W}{m^2} \times (D \times L) m^2$$

$$= \epsilon (\pi D L) m^2 \times \sigma (T_e^4) K^4$$

$\uparrow$   
 $5.67 \times 10^{-8} \text{ W/(m}^2 \cdot \text{K}^4)$

L = Length of cylinder


$$T_e = \frac{1500}{\pi \times 5.67 \times 10^{-8}} = 303 \text{ K} = 30^\circ \text{C.}$$

## CHAPTER 2

### GENERAL HEAT CONDUCTION EQUATION

PROB. 2.1:

- Homogeneous and isotropic solid, i.e.,  $k = \text{const.}$
- Let there be no heat sources or sinks, i.e.,  $\dot{q} = 0$ .
- Assume the density  $\rho$  and specific heat  $c$  are independent of time  $t$ .



$$\left\{ \begin{array}{l} \text{Net rate of heat conduction} \\ \text{into the C.V. in the } x\text{-direction} \end{array} \right\} = q_{x1x} - q_{x1x+\Delta x} = q_{x1\Delta x} - \left\{ q_{x1x} + \frac{\partial q_{x1x}}{\partial x} \Delta x \right\}$$

$$= - \frac{\partial q_{x1x}}{\partial x} \Delta x$$

where

$$q_{x1x} = -k (\Delta y \Delta z) \frac{\partial T}{\partial x} \quad \leftarrow \begin{array}{l} \text{Rate of heat conduction} \\ \text{in the } x\text{-direction} \\ \text{at } x \text{ into the C.V.} \end{array}$$

$$\therefore q_{x1x} - q_{x1x+\Delta x} = k \frac{\partial^2 T}{\partial x^2} \Delta V, \quad \Delta V = \Delta x \Delta y \Delta z$$

Similarly,


$$q_{y1y} - q_{y1y+\Delta y} = k \frac{\partial^2 T}{\partial y^2} \Delta V \quad \text{and} \quad q_{z1z} - q_{z1z+\Delta z} = k \frac{\partial^2 T}{\partial z^2} \Delta V$$

Thus, energy balance on the C.V. gives

$$k \left\{ \frac{\partial^2 T}{\partial x^2} + \frac{\partial^2 T}{\partial y^2} + \frac{\partial^2 T}{\partial z^2} \right\} \Delta V = \left\{ \begin{array}{l} \text{Net rate of increase} \\ \text{of internal energy} \end{array} \right\} = \frac{\partial}{\partial t} \left\{ T c \rho \Delta V \right\}$$

$$\therefore \boxed{\frac{\partial^2 T}{\partial x^2} + \frac{\partial^2 T}{\partial y^2} + \frac{\partial^2 T}{\partial z^2} = \frac{1}{\alpha} \frac{\partial T}{\partial t}}, \quad \alpha = \frac{k}{\rho c}$$

PROB. 2.2: (a)



$$q_{tr1r} - q_{tr1r+\Delta r} = - \frac{\partial q_{tr1r}}{\partial r} \Delta r = \frac{\partial}{\partial r} \left[ k r \Delta \phi \Delta z \frac{\partial T}{\partial r} \right] \Delta r = \frac{k}{r} \frac{\partial}{\partial r} \left( r \frac{\partial T}{\partial r} \right) \Delta V$$

$$q_{phi1phi} - q_{phi1phi+\Delta phi} = - \frac{\partial q_{phi1phi}}{\partial (r\phi)} r \Delta \phi = \frac{\partial}{\partial \phi} \left[ k \Delta r \Delta z \frac{\partial T}{\partial (r\phi)} \right] \Delta \phi = \frac{k}{r^2} \frac{\partial^2 T}{\partial \phi^2} \Delta V$$

$$q_{z1z} - q_{z1z+\Delta z} = - \frac{\partial q_{z1z}}{\partial z} \Delta z = \frac{\partial}{\partial z} \left[ k r \Delta \phi \Delta r \frac{\partial T}{\partial z} \right] \Delta z = k \frac{\partial^2 T}{\partial z^2} \Delta V$$

$$\therefore k \left\{ \frac{1}{r} \frac{\partial}{\partial r} \left( r \frac{\partial T}{\partial r} \right) + \frac{1}{r^2} \frac{\partial^2 T}{\partial \phi^2} + \frac{\partial^2 T}{\partial z^2} \right\} \Delta V = \underbrace{\frac{\partial}{\partial t} \left\{ T c_p \Delta V \right\}}_{\text{Net rate of increase of internal energy in c.v.}}$$

Net rate of heat conduction into c.v.

Net rate of increase of internal energy in c.v.

$$\boxed{\frac{1}{r} \frac{\partial}{\partial r} \left( r \frac{\partial T}{\partial r} \right) + \frac{1}{r^2} \frac{\partial^2 T}{\partial \phi^2} + \frac{\partial^2 T}{\partial z^2} = \frac{1}{\alpha} \frac{\partial T}{\partial t}}, \alpha = \frac{k}{\rho c}$$

PROB. 2.3: General heat conduction equation in the rectangular coordinates:

$$\frac{\partial^2 T}{\partial x^2} + \frac{\partial^2 T}{\partial y^2} + \frac{\partial^2 T}{\partial z^2} + \frac{q}{k} = \frac{1}{\alpha} \frac{\partial T}{\partial t}$$

a) The cylindrical coordinates:

$$\left. \begin{array}{l} x = r \cos \phi \\ y = r \sin \phi \\ z = z \end{array} \right\} \quad \left. \begin{array}{l} r = (x^2 + y^2)^{1/2} \\ \phi = \tan^{-1}(\frac{y}{x}) \\ z = z \end{array} \right.$$

Thus,

$$\begin{aligned} \frac{\partial T}{\partial x} &= \frac{\partial T}{\partial r} \frac{\partial r}{\partial x} + \frac{\partial T}{\partial \phi} \frac{\partial \phi}{\partial x} = \cos \phi \frac{\partial T}{\partial r} - \frac{\sin \phi}{r} \frac{\partial T}{\partial \phi} \\ \frac{\partial^2 T}{\partial x^2} &= \frac{\partial}{\partial x} \left( \frac{\partial T}{\partial x} \right) = \frac{\partial}{\partial r} \left( \frac{\partial T}{\partial x} \right) \frac{\partial r}{\partial x} + \frac{\partial}{\partial \phi} \left( \frac{\partial T}{\partial x} \right) \frac{\partial \phi}{\partial x} \\ &= \cos^2 \phi \frac{\partial^2 T}{\partial r^2} + 2 \frac{\cos \phi \sin \phi}{r^2} \frac{\partial T}{\partial \phi} - 2 \frac{\sin \phi \cos \phi}{r} \frac{\partial^2 T}{\partial r \partial \phi} \\ &\quad + \frac{\sin^2 \phi}{r} \frac{\partial^2 T}{\partial \phi^2} + \frac{\sin^2 \phi}{r^2} \frac{\partial^2 T}{\partial \phi^2} \end{aligned}$$

Similarly,

$$\begin{aligned} \frac{\partial T}{\partial y} &= \sin \phi \frac{\partial T}{\partial r} + \frac{\cos \phi}{r} \frac{\partial T}{\partial \phi} \\ \frac{\partial^2 T}{\partial y^2} &= \sin^2 \phi \frac{\partial^2 T}{\partial r^2} + 2 \frac{\sin \phi \cos \phi}{r} \frac{\partial^2 T}{\partial \phi \partial r} - 2 \frac{\sin \phi \cos \phi}{r^2} \frac{\partial^2 T}{\partial \phi^2} \\ &\quad + \frac{\cos^2 \phi}{r} \frac{\partial^2 T}{\partial r^2} + \frac{\cos^2 \phi}{r^2} \frac{\partial^2 T}{\partial \phi^2} \end{aligned}$$

In addition, in both coordinate systems:  $\frac{\partial^2 T}{\partial z^2} = \frac{\partial^2 T}{\partial z^2}$

Therefore,

$$\frac{\partial^2 T}{\partial x^2} + \frac{\partial^2 T}{\partial y^2} + \frac{\partial^2 T}{\partial z^2} = \frac{\partial^2 T}{\partial r^2} + \frac{1}{r} \frac{\partial T}{\partial r} + \frac{1}{r^2} \frac{\partial^2 T}{\partial \phi^2} + \frac{\partial^2 T}{\partial z^2}$$

and the general heat conduction equation in the cylindrical coordinate is

$$\frac{\partial^2 T}{\partial r^2} + \frac{1}{r} \frac{\partial T}{\partial r} + \frac{1}{r^2} \frac{\partial^2 T}{\partial \phi^2} + \frac{\partial^2 T}{\partial z^2} + \frac{q}{k} = \frac{1}{\alpha} \frac{\partial T}{\partial t}$$

b) The spherical coordinates:

$$\left. \begin{array}{l} x = r \sin \theta \cos \phi \\ y = r \sin \theta \sin \phi \\ z = r \cos \theta \end{array} \right\} \quad \begin{array}{l} r = \sqrt{x^2 + y^2 + z^2} \\ \phi = \tan^{-1} \left( \frac{y}{x} \right) \\ \theta = \cos^{-1} \frac{z}{\sqrt{x^2 + y^2 + z^2}} \end{array}$$

$$\begin{aligned} \frac{\partial T}{\partial x} &= \frac{\partial T}{\partial r} \frac{\partial r}{\partial x} + \frac{\partial T}{\partial \phi} \frac{\partial \phi}{\partial x} + \frac{\partial T}{\partial \theta} \frac{\partial \theta}{\partial x} \\ &= \sin \theta \cos \phi \frac{\partial T}{\partial r} - \frac{\sin \phi}{r \sin \theta} \frac{\partial T}{\partial \phi} + \frac{\cos \theta \cos \phi}{r} \frac{\partial T}{\partial \theta} \end{aligned}$$

$$\begin{aligned} \frac{\partial^2 T}{\partial x^2} &= \frac{\partial}{\partial r} \left( \frac{\partial T}{\partial x} \right) \frac{\partial r}{\partial x} + \frac{\partial}{\partial \phi} \left( \frac{\partial T}{\partial x} \right) \frac{\partial \phi}{\partial x} + \frac{\partial}{\partial \theta} \left( \frac{\partial T}{\partial x} \right) \frac{\partial \theta}{\partial x} \\ &= \sin^2 \theta \cos^2 \phi \frac{\partial^2 T}{\partial r^2} + 2 \frac{\cos \phi \sin \phi}{r^2 \sin^2 \theta} \frac{\partial T}{\partial \phi} - 2 \frac{\sin \phi \cos \phi}{r} \frac{\partial^2 T}{\partial r \partial \phi} \\ &\quad + \frac{\cos \theta}{r^2} \left( -2 \sin \theta \cos^2 \phi + \frac{\sin^2 \phi}{\sin \theta} \right) \frac{\partial T}{\partial \theta} + 2 \frac{\sin \theta \cos \theta \cos^2 \phi}{r} \frac{\partial^2 T}{\partial r \partial \theta} \\ &\quad + \left( \frac{\cos^2 \theta \cos^2 \phi}{r} + \frac{\sin^2 \phi}{r} \right) \frac{\partial T}{\partial r} + \frac{\sin^2 \phi}{r^2 \sin^2 \theta} \frac{\partial^2 T}{\partial \phi^2} \\ &\quad - 2 \frac{\cos \theta \sin \phi \cos \phi}{r^2 \sin \theta} \frac{\partial^2 T}{\partial \phi \partial \theta} + \frac{\cos^2 \theta \cos^2 \phi}{r^2} \frac{\partial^2 T}{\partial \theta^2} \end{aligned}$$

Similarly,

$$\begin{aligned} \frac{\partial T}{\partial y} &= \frac{\partial T}{\partial r} \frac{\partial r}{\partial y} + \frac{\partial T}{\partial \phi} \frac{\partial \phi}{\partial y} + \frac{\partial T}{\partial \theta} \frac{\partial \theta}{\partial y} \\ &= \sin \theta \sin \phi \frac{\partial T}{\partial r} + \frac{\cos \phi}{r \sin \theta} \frac{\partial T}{\partial \phi} + \frac{\cos \theta \sin \phi}{r} \frac{\partial T}{\partial \theta} \end{aligned}$$

$$\begin{aligned}
 \frac{\partial^2 T}{\partial y^2} &= \sin^2 \theta \sin^2 \phi \frac{\partial^2 T}{\partial r^2} - 2 \frac{\sin \phi \cos \phi}{r^2 \sin^2 \theta} + 2 \frac{\sin \phi \cos \phi}{r} \frac{\partial^2 T}{\partial r \partial \theta} \\
 &+ \frac{\cos \theta}{r^2} \left( -2 \sin \theta \sin^2 \phi + \frac{\cos^2 \phi}{\sin \theta} \right) \frac{\partial T}{\partial \theta} + 2 \frac{\sin \theta \cos \theta \sin^2 \phi}{r} \frac{\partial^2 T}{\partial r \partial \theta} \\
 &+ \frac{\cos^2 \theta \sin^2 \phi + \cos^2 \phi}{r} \frac{\partial T}{\partial r} + \frac{\cos^2 \phi}{r^2 \sin^2 \theta} \frac{\partial^2 T}{\partial \theta^2} \\
 &+ 2 \frac{\cos \theta \sin \phi \cos \phi}{r^2 \sin \theta} \frac{\partial^2 T}{\partial \phi \partial \theta} + \frac{\cos^2 \theta \sin^2 \phi}{r^2} \frac{\partial^2 T}{\partial \phi^2}
 \end{aligned}$$

and

$$\frac{\partial T}{\partial z} = \cos \frac{\partial T}{\partial r} - \frac{\sin \theta}{r} \frac{\partial T}{\partial \theta}$$

$$\begin{aligned}
 \frac{\partial^2 T}{\partial z^2} &= \cos^2 \theta \frac{\partial^2 T}{\partial r^2} + 2 \frac{\sin \theta \cos \theta}{r^2} \frac{\partial T}{\partial \theta} - 2 \frac{\sin \theta \cos \theta}{r} \frac{\partial^2 T}{\partial r \partial \theta} \\
 &+ \frac{\sin^2 \theta}{r} \frac{\partial^2 T}{\partial r^2} + \frac{\sin^2 \theta}{r^2} \frac{\partial^2 T}{\partial \theta^2}
 \end{aligned}$$

Thus,

$$\begin{aligned}
 \frac{\partial^2 T}{\partial x^2} + \frac{\partial^2 T}{\partial y^2} + \frac{\partial^2 T}{\partial z^2} &= \underbrace{\frac{1}{r^2} \frac{\partial}{\partial r} \left( r^2 \frac{\partial T}{\partial r} \right)}_{\frac{\partial^2 T}{\partial r^2} + \frac{2}{r} \frac{\partial T}{\partial r}} + \underbrace{\frac{1}{r^2 \sin \theta} \left( \sin \theta \frac{\partial T}{\partial \theta} \right)}_{\frac{\cos \theta}{r^2 \sin \theta} \frac{\partial T}{\partial \theta} + \frac{1}{r^2} \frac{\partial^2 T}{\partial \theta^2}} \\
 &+ \frac{1}{r^2 \sin^2 \theta} \frac{\partial^2 T}{\partial \phi^2}
 \end{aligned}$$

and the general heat conduction equation in the spherical coordinates is


$$\frac{1}{r^2} \frac{\partial}{\partial r} \left( r^2 \frac{\partial T}{\partial r} \right) + \frac{1}{r^2 \sin \theta} \left( \sin \theta \frac{\partial T}{\partial \theta} \right) + \frac{1}{r^2 \sin^2 \theta} \frac{\partial^2 T}{\partial \phi^2} + \frac{q}{k} = \frac{1}{\alpha} \frac{\partial T}{\partial t}$$

PROB. 2.4:

B.C. at  $x=0$ :

$$-k \left( \frac{\partial T}{\partial x} \right)_{x=0} = h [T_\infty - T]_{x=0} + q''_s$$

$$\boxed{\left[ -k \frac{\partial T}{\partial x} + h T \right]_{x=0} = h T_\infty + q''_s}$$




PROB. 2.5:

1) Since the contact is perfect,


$$T_1|_{\vec{r}_s} = T_2|_{\vec{r}_s}$$

2)

$$-k_1 \frac{\partial T_1}{\partial n} \Big|_{n=0} + \mu p V = -k_2 \frac{\partial T_2}{\partial n} \Big|_{n=0}$$



PROB. 2.6:



The temperature distribution will be one-dimensional, that is,  $T=T(r,t)$ . Thus, the formulation of the problem for  $T(r,t)$  can be given as


$$\text{Diff. Eq.} \rightarrow \frac{\partial^2 T}{\partial r^2} + \frac{2}{r} \frac{\partial T}{\partial r} = \frac{1}{\alpha} \frac{\partial T}{\partial t}, \quad \alpha = \frac{k}{gc}$$

$$\text{Initial Cond.} \rightarrow T(r,0) = T_i$$

$$\text{Two boundary condns.} \rightarrow T(0,t) = \text{finite}; \quad -k \frac{\partial T}{\partial r} \Big|_{r=r_0} = h [T(r_0,t) - T_\infty]$$

PROB. 2.7: Because of the symmetry in the  $\phi$ -direction, under steady-state conditions  $T = T(r, z)$ . In addition, the temperature of the lower surface will be equal to the temperature of the boiling water,  $T_f$  (due to very large  $h$ ). Thus, the formulation is given by


$$\frac{\partial^2 T}{\partial r^2} + \frac{1}{r} \frac{\partial T}{\partial r} + \frac{\partial^2 T}{\partial z^2} = 0$$



$$T(0, z) = \text{finite}; -k \left( \frac{\partial T}{\partial r} \right)_{r=r_0} = h [T(r_0, z) - T_\infty]$$

$$T(r, 0) = T_f; \left( \frac{\partial T}{\partial z} \right)_{z=H} = 0$$

PROB. 2.8:



$$k_x = 2 k_y$$

For unit depth:

$$\begin{aligned} q_{x|x} - q_{x|x+\Delta x} &= - \frac{\partial q_{x|x}}{\partial x} \Delta x = \frac{\partial}{\partial x} \left( k_x \Delta y \frac{\partial T}{\partial x} \right) \Delta x \\ &= k_x \frac{\partial^2 T}{\partial x^2} \Delta x \Delta y \end{aligned}$$

and

$$\begin{aligned} q_{y1y} - q_{y1y+\Delta y} &= - \frac{\partial q_{y1y}}{\partial y} \Delta y = \frac{\partial}{\partial y} \left( k_y \Delta x \frac{\partial T}{\partial y} \right) \Delta y \\ &= k_y \frac{\partial^2 T}{\partial y^2} \Delta x \Delta y \end{aligned}$$

Thus, under steady-state conditions and in the absence of heat sources or sinks, energy balance on the c.v. shown yields

$$k_x \frac{\partial^2 T}{\partial x^2} + k_y \frac{\partial^2 T}{\partial y^2} = 0 \quad \text{or} \quad 2 \frac{\partial^2 T}{\partial x^2} + \frac{\partial^2 T}{\partial y^2} = 0$$

Now, the formulation of the problem for  $T=T(x,y)$  can be written as

$$2 \frac{\partial^2 T}{\partial x^2} + \frac{\partial^2 T}{\partial y^2} = 0$$

$$T(0,y) = T_1 ; \quad T(a,y) = T_1$$

$$T(x,0) = T_1 ; \quad T(x,b) = T_2$$

If we introduce a new temperature function as

$$\theta(x,y) = T(x,y) - T_1$$


then the formulation can be rewritten as

$$2 \frac{\partial^2 \theta}{\partial x^2} + \frac{\partial^2 \theta}{\partial y^2} = 0$$

$$\theta(0,y) = 0 ; \quad \theta(a,y) = 0$$

$$\theta(x,0) = 0 ; \quad \theta(x,b) = T_2 - T_1$$

PROB. 2.9:



Formulation of the problem for  $T(x,t)$  can be written as

$$\text{D.E.} \rightarrow \frac{\partial^2 T}{\partial x^2} + \frac{\dot{q}_0}{k} e^{-\beta t} = \frac{1}{\alpha} \frac{\partial T}{\partial t}, \quad \alpha = \frac{k}{gc}$$

$$\text{I.C.} \rightarrow T(x,0) = T_i'$$

$$\text{B.C.'s} \rightarrow \frac{\partial T(0,t)}{\partial x} = 0; \quad T(L,t) = T_i'$$

If we define a new temperature function as

$$\theta(x,t) = T(x,t) - T_i'$$

then the formulation can be rewritten as

$$\frac{\partial^2 \theta}{\partial x^2} + \frac{\dot{q}_0}{k} e^{-\beta t} = \frac{1}{\alpha} \frac{\partial \theta}{\partial t}$$

$$\theta(x,0) = 0$$

$$\frac{\partial \theta(0,t)}{\partial x} = 0; \quad \theta(L,t) = 0$$

PROB. 2.10:

$$\left\{ \begin{array}{l} \frac{\partial^2 T}{\partial x^2} = \frac{1}{\alpha} \frac{\partial T}{\partial t} \\ T(x, 0) = T_i(x) \\ T(0, t) = T_1; T(L, t) = T_2 \end{array} \right.$$

Assume that there are more than one solution. Let  $T^{(1)}(x, t)$  and  $T^{(2)}(x, t)$  be two different solutions. They both satisfy the formulation of the problem:

$$\frac{\partial^2 T^{(1)}}{\partial x^2} = \frac{1}{\alpha} \frac{\partial T^{(1)}}{\partial t}$$

$$T^{(1)}(x, 0) = T_i(x)$$

$$T^{(1)}(0, t) = T_1; T^{(1)}(L, t) = T_2$$

$$\frac{\partial^2 T^{(2)}}{\partial x^2} = \frac{1}{\alpha} \frac{\partial T^{(2)}}{\partial t}$$

$$T^{(2)}(x, 0) = T_i(x)$$

$$T^{(2)}(0, t) = T_1; T^{(2)}(L, t) = T_2$$

Define a new function  $\phi = T^{(1)} - T^{(2)}$ . Since the problem is linear,  $\phi$  will satisfy the following problem:

$$\frac{\partial^2 \phi}{\partial x^2} = \frac{1}{\alpha} \frac{\partial \phi}{\partial t}$$

$$\phi(x, 0) = 0$$

$$\phi(0, t) = 0; \phi(L, t) = 0$$

Physical interpretation: The solution of this problem,  $\phi(x)$ , represents the unsteady-state temperature distribution in a slab with zero initial condition and zero boundary conditions. Hence, it will always be zero, which implies  $T^{(1)} = T^{(2)}$ .

Mathematical interpretation: Multiply the differential equation by  $\phi$  and then integrate over  $x$  from 0 to L:

$$\underbrace{\int_0^L \phi \frac{\partial^2 \phi}{\partial x^2} dx}_{\phi \frac{\partial \phi}{\partial x} \Big|_0^L - \int_0^L \left(\frac{\partial \phi}{\partial x}\right)^2 dx} = \underbrace{\frac{1}{\alpha} \int_0^L \phi \frac{\partial \phi}{\partial t} dx}_{\frac{1}{2\alpha} \frac{d}{dt} \int_0^L \phi^2 dx}$$

$$-\int_0^L \left(\frac{\partial \phi}{\partial x}\right)^2 dx = \frac{1}{2\alpha} \frac{d}{dt} \int_0^L \phi^2 dx$$

Now, integrate this result over t from  $t=0$  to any  $t$ :

$$\underbrace{- \int_0^t \left[ \int_0^L \left(\frac{\partial \phi}{\partial x}\right)^2 dx \right] dt}_{\text{Always negative}} = \frac{1}{2\alpha} \underbrace{\left\{ \int_0^L \phi^2(x,t) dx - \int_0^L \phi^2(x,0) dx \right\}}_{\text{Always positive}}$$


Hence, the left - and right-hand sides can be equal to each other only if they are both zero.

$$\therefore \int_0^L \phi^2(x,t) dx = 0$$

This, in turn, implies

$$\phi(x,t) \equiv 0 \quad \text{in } 0 < x < L.$$

PROB. 2.11:



$$\frac{\partial^2 T}{\partial x^2} + \frac{\partial^2 T}{\partial y^2} = 0$$

$$\left. \frac{\partial T}{\partial x} \right|_{x=0} = 0, \quad -k \left. \frac{\partial T}{\partial x} \right|_{x=a} = q''_1$$

$$\left. \frac{\partial T}{\partial y} \right|_{y=0} = 0, \quad k \left. \frac{\partial T}{\partial y} \right|_{y=b} = q''_2$$

a) Under steady-state conditions:

$$q''_2 \cdot a = q''_1 \cdot b$$

$$\therefore \underline{q''_1 = \frac{a}{b} q''_2}$$

b) Note that if  $T(x,y)$  is a solution, then  $T(x,y) + C$  is also a solution, where  $C$  is any constant. This can be proved by direct substitution in the formulation of the problem. That is, let  $\theta = T + c$ . Then,

$$\frac{\partial^2 \theta}{\partial x^2} + \frac{\partial^2 \theta}{\partial y^2} = 0$$

$$\left. \frac{\partial \theta}{\partial x} \right|_{x=0} = 0, \quad -k \left. \frac{\partial \theta}{\partial x} \right|_{x=a} = q''_1$$

$$\left. \frac{\partial \theta}{\partial y} \right|_{y=0} = 0, \quad +k \left. \frac{\partial \theta}{\partial y} \right|_{y=b} = q''_2$$

Thus, both  $T(x,y)$  and  $\theta(x,y)$  satisfy the same problem.

PROB. 2.12:


$$\left\{ \begin{array}{l} \frac{d}{dx} \left[ k(T) \frac{dT}{dx} \right] + \dot{q} = 0 \\ \frac{dT(0)}{dx} = 0 \\ T(L) = T_w \end{array} \right\} \rightarrow \theta(x) = \frac{1}{k_w} \int_{T_w}^{T(x)} k(T') dT' \rightarrow \left\{ \begin{array}{l} \frac{d^2 \theta}{dx^2} + \frac{\dot{q}}{k_w} = 0 \\ \frac{d\theta(0)}{dx} = 0 \\ \theta(L) = 0 \end{array} \right\}$$

↑  
Linear in  $\theta(x)$ .

provided that  $\dot{q}$  is not a non-linear function of the temperature  $T$ .

PROB. 2.13:

$$-k \frac{\partial \theta}{\partial n} \Big|_{n=0} = h [T(\vec{r}_s, t) - T_\infty] \quad ①$$



Let

$$\theta(\vec{r}, t) = \frac{1}{k_R} \int_{T_R}^{T(\vec{r}, t)} k(T') dT' , \quad k_R = k(T_R)$$

Then, the boundary condition ① would be rewritten as

$$-k_R \frac{\partial \theta}{\partial n} \Big|_{n=0} = h^*(\vec{r}_s, t) \theta(\vec{r}_s, t) \quad ②$$

provided that the heat transfer coefficient  $h$  in ① is given by

$$h = \frac{h^*(\vec{r}_s, t)}{k_R [T(\vec{r}_s, t) - T_{\infty}]} \int_{T_R}^{T(\vec{r}_s, t)} k(T) dT' \quad ③$$


where  $h^*(\vec{r}_s, t)$  is any function of  $\vec{r}_s$  and  $t$ .

Note that (a) the condition ② is a linear and homogeneous boundary condition of the third kind, and (b) for this transformation to be possible the heat transfer coefficient  $h$  has to be a function of the surface temperature  $T(\vec{r}_s, t)$  as specified by the relation ③.

# CHAPTER 3

## ONE-DIMENSIONAL STEADY-STATE HEAT CONDUCTION

PROB. 3.1:



$$\left. \begin{aligned} \frac{680 - 120}{R_1 + R_2} &= q_1 \\ \frac{700 - 490}{R_1 + R_2} &= q_2 \end{aligned} \right\} \quad \frac{q_2}{q_1} = \frac{210}{560} = 0.375 \text{ or } 37.5\%$$

PROB. 3.2: a) Under steady-state conditions,

$$q = -k 2\pi r L \frac{dT}{dr} = \text{const.}, \quad k = 0.137 + 0.002T(\text{°C})$$



$$\int_{T_0}^{T_i} k(T) dT = \frac{q}{2\pi L} \ln \frac{r_0}{r_i} \quad [k] = \frac{\text{W}}{\text{m} \cdot \text{K}}$$

$$0.137(500 - 80) + \frac{0.002}{2} (500^2 - 80^2) = \frac{q}{2\pi L} \ln \frac{25}{12.5}$$

$$12 \text{ W/(m}^2 \cdot \text{K}) \quad \therefore \quad \frac{q}{L} = 1,311.5 \frac{\text{W}}{\text{m}}$$

$$\text{b)} \quad \frac{q}{L} = 1,311.5 = h \times 2\pi r_0 \times (80 - T_\infty) \quad \Rightarrow \quad T_\infty = 10.4^\circ\text{C}$$

PROB. 3,3:



$$R_i' = \frac{1}{h_i \pi ID} = \frac{1}{4500 \times \pi \times 10.126 \times 0.0254} = 2.75 \times 10^{-4} \frac{\text{m} \cdot \text{K}}{\text{W}}$$

$$R_t = \frac{1}{2\pi k} \ln \frac{OD}{ID} = \frac{1}{2\pi \times 40} \ln \frac{12.75}{10.126} = 9.17 \times 10^{-4} \frac{\text{m} \cdot \text{K}}{\text{W}}$$

$$R_o = \frac{1}{h_o \pi OD} = \frac{1}{12 \times \pi \times 2 \times r_2} = \frac{0.01326}{r_2}$$

$$R_1 = \frac{1}{2\pi k_1} \ln \frac{r_1}{(OD/2)} = 0.7958 [1.8206 + \ln r_1]; R_2 = \frac{1}{2\pi k_2} \ln \frac{r_2}{r_1} = 2.04 \ln \frac{r_2}{r_1}$$

$$\left. \begin{aligned} \frac{400 - 300}{R_i' + R_t + R_1} &= \frac{48 - 30}{R_o} \rightarrow \ln r_1 = \frac{0.0926}{r_2} - 1.816 \\ \frac{300 - 48}{R_2} &= \frac{48 - 30}{R_o} \rightarrow \ln \frac{r_2}{r_1} = \frac{0.091}{r_2} \end{aligned} \right\} \Rightarrow \boxed{\ln r_2 = \frac{0.1836}{r_2} - 1.816}$$

Assume  $r_2$  (m)  $\rightarrow \ln r_2 \rightarrow$  find  $r_2$  (m)

0.35	-1.2914	0.275
0.275	-1.1483	0.317
0.317	-1.237	0.29
0.3	-1.204	0.2999

High-temp. insulation

thickness:  $r_1 - \frac{OD}{2} = 0.058 \text{ m}$

Magnesia insulation thickness

$$\therefore r_2 = 0.3 \text{ m} \rightarrow r_1 = 0.22 \text{ m}$$

$$r_2 - r_1 = 0.3 - 0.22 = 0.08 \text{ m}$$

PROB. 3.4:

$$q = \frac{T_i - T_o}{\sum R_{th}}$$

$$\sum R_{th} = \frac{1}{4\pi} \left[ \frac{1}{r_1^2 h_i} + \frac{r_2 - r_1}{r_2 r_1 k_1} + \frac{r - r_2}{r r_2 k_2} + \frac{1}{r^2 h_o} \right]$$

$$\frac{d}{dr} (\sum R_{th}) = \frac{1}{4\pi r^2} \left[ \frac{1}{k_2} - \frac{2}{h_o r} \right] = 0$$

This equation is satisfied when  $r = \frac{2k_2}{h_o}$ .

This value of  $r$  is termed the critical radius of insulation for a sphere, i.e.,  $r_{cr} = 2k_2/h_o$ . One more differentiation yields

$$\frac{d^2}{dr^2} (\sum R_{th}) \Big|_{r=r_{cr}} = \frac{h_o^3}{32\pi k_2^4} > 0$$

Hence, at  $r=r_{cr} = 2k_2/h_o$ , the total thermal resistance is minimum and therefore  $q$  is maximum.

---


PROB. 3.5: (Refer to Fig. 3.11 in text)

Since  $r_{cr} = \frac{0.151}{6} = 0.0252 \text{ m} = 2.52 \text{ cm} > r_2 = 0.5 \text{ in} = 1.27 \text{ cm}$ , adding insulation will increase the heat loss until  $r$  becomes equal to  $r_{cr}$ , and then the heat loss will start decreasing.

$$\begin{aligned} \text{Heat loss without insulation: } q_1 &= \frac{T_f - T_\infty}{R_1} \\ \text{Heat loss with insulation: } q_2 &= \frac{T_f - T_\infty}{R_2} \end{aligned} \quad \left. \begin{array}{l} \text{If } q_1 = q_2, \text{ then} \\ R_1 = R_2 \end{array} \right\}$$

$$\therefore \frac{1}{h r_2} = \frac{1}{k_2} \ln \frac{r}{r_2} + \frac{1}{h r}$$

This relation is satisfied when  $r = 6.1 \text{ cm}$ . This is the value of  $r$  at which the loss with and without the insulation will be the same.




PROB. 3.6: Formulation of the problem:

$$\frac{d^2T}{dx^2} + \frac{\dot{q}}{k} = 0$$

$$\frac{dT(0)}{dx} = 0$$

$$-k \frac{dT(L)}{dx} = h [T(L) - T_{\infty}]$$



Sol. to DE:  $T(x) = -\frac{\dot{q}x^2}{2k} + C_1 x + C_2$

B.C. at  $x=0 \Rightarrow C_1 = 0$  and B.C. at  $x=L \Rightarrow C_2 = T_{\infty} + \frac{\dot{q}L^2}{2k} + \frac{\dot{q}L}{h}$

$$\therefore T(x) = T_{\infty} + \frac{\dot{q}L^2}{2k} \left[ 1 - \left( \frac{x}{L} \right)^2 \right] + \frac{\dot{q}L}{h}$$

Heat flux from the surface at  $x=L$ :  $q''_L = \dot{q}L$

PROB. 3.7:  $L=3$  in,  $\dot{q}=5 \times 10^4$  W/m<sup>3</sup>,  $T_{\infty}=30^\circ\text{C}$ ,  $h=600$  W/(m<sup>2</sup>·K), and  $k=17$  W/(m·K). From Prob. 3.6, the max. temp. in the wall will be given by

$$\begin{aligned} T_{\max} &= T_{\infty} + \frac{\dot{q}L^2}{2k} + \frac{\dot{q}L}{h} = T_{\infty} + \frac{\dot{q}L^2}{2k} \left( 1 + 2 \frac{k}{hL} \right) \\ &= 30 + \frac{5 \times 10^4 \times 3 \times 2.54 \times 10^{-2}}{2 \times 17} \left( 1 + 2 \times \frac{17}{3 \times 2.54 \times 10^{-2} \times 600} \right) \\ &= 225.4^\circ\text{C} \end{aligned}$$

PROB. 3.8: Formulation of the problem:

$$\left\{ \begin{array}{l} \frac{d^2T}{dx^2} + \frac{\dot{q}}{k} = 0 \\ T(0) = T_1; \quad T(L) = T_2 \end{array} \right\} \quad \text{SOL. to DE: } T(x) = -\frac{\dot{q}x^2}{2k} + C_1 x + C_2$$

$$\frac{T(0) = T_1 \Rightarrow C_2 = T_1 \text{ & } T(L) = T_2 \Rightarrow C_1 = \frac{T_2 - T_1}{L} + \frac{\dot{q}L}{2k}}{T(x) - T_1 = \frac{T_2 - T_1}{L} x + \frac{\dot{q}x}{2k} [L - x]}$$

PROB. 3.9:

Formulation  $\left\{ \begin{array}{l} \frac{d^2T}{dr^2} + \frac{1}{r} \frac{dT}{dr} + \frac{\dot{q}}{k} = 0 \\ k \frac{dT(r_i)}{dr} = h [T(r_i) - T_f] \\ \frac{dT(r_0)}{dr} = 0 \end{array} \right\}$

Sol. to D.E.

$$T(r) = -\frac{\dot{q}r^2}{4k} + c_1 \ln r + c_2$$

B.C. at  $r_0 \Rightarrow c_1 = \frac{\dot{q}r_0^2}{2k}$

B.C. at  $r_i$   $\Rightarrow c_2 = T_f + \frac{\dot{q}r_i^2}{4k} \left\{ 1 - 2 \left( \frac{r_0}{r_i} \right)^2 \ln r_i + \frac{2}{Bi} \left[ 1 - \left( \frac{r_0}{r_i} \right)^2 \right] \right\}, Bi = \frac{h r_i}{k}$

Substituting  $c_1$  &  $c_2$  into the solution of DE one gets the temp. dist.

Rate of Heat Transfer to the gas / length =  $\pi (r_0^2 - r_i^2) \dot{q}_f$

PROB. 3.10:

Formulation  $\left\{ \begin{array}{l} \frac{d^2T}{dr^2} + \frac{1}{r} \frac{dT}{dr} + \frac{\dot{q}}{k} = 0 \\ -k \frac{dT(r_0)}{dr} = h_2 [T(r_0) - T_{f2}] \\ k \frac{T(r_i)}{dr} = h_1 [T(r_i) - T_{f1}] \end{array} \right\}$

Sol. to DE:

$$T(r) = -\frac{\dot{q}r^2}{4k} + c_1 \ln r + c_2$$

B.C.'s yield:

$$c_1 = \frac{1}{\Delta_1} \left\{ \frac{\dot{q}}{2} \left[ \frac{r_0}{h_2} + \frac{r_i}{h_1} \right] + \frac{\dot{q}}{4k} [r_0^2 - r_i^2] + (T_{f2} - T_{f1}) \right\}$$

$$c_2 = -\frac{\Delta_2}{\Delta_1} \left\{ \frac{\dot{q}}{2} \left[ \frac{r_0}{h_2} + \frac{r_i}{h_1} \right] + \frac{\dot{q}}{4k} [r_0^2 - r_i^2] + (T_{f2} - T_{f1}) \right\} + \frac{\dot{q}r_0}{2h_2} + \frac{\dot{q}r_i^2}{4k} + T_{f2}$$

where

$$\Delta_1 = k \left( \frac{1}{r_0 h_2} + \frac{1}{r_i h_1} \right) + \ln \frac{r_0}{r_i} \quad \& \quad \Delta_2 = \frac{k}{r_0 h_2} + \ln r_0$$

Substituting  $c_1$  &  $c_2$  into the solution of DE gives the temp. dist.

Then, one would obtain the surface temperatures from

$$T(r_0) = -\frac{\dot{q}r_0^2}{4k} + c_1 \ln r_0 + c_2 \quad \& \quad T(r_i) = -\frac{\dot{q}r_i^2}{4k} + c_1 \ln r_i + c_2$$

Max. Temp.?  $\frac{dT}{dr} = -\frac{\dot{q}r}{2k} + \frac{c_1}{r} = 0 \Rightarrow r^* = \sqrt{2k c_1 / \dot{q}}$

If  $r_i \leq r^* \leq r_0$  then  $T_{max} = T(r^*)$

If  $r^* < r_i$  then  $T_{max} = T(r_i)$

If  $r^* > r_0$  then  $T_{max} = T(r_0)$

PROB. 3.11:

Formulation of the problem

$$\left\{ \begin{array}{l} \frac{d^2T}{dx^2} + \frac{\dot{q}_o}{k} \left[ 1 - \left( \frac{x}{L} \right)^2 \right] = 0 \\ \frac{dT(0)}{dx} = 0 \\ T(L) = T_w \end{array} \right\} \rightarrow \begin{array}{l} T(x) = -\frac{\dot{q}_o}{k} \left[ \frac{x^2}{2} - \frac{x^4}{12L^2} \right] + c_1 x + c_2 \\ c_1 = 0 \\ c_2 = T_w + \frac{5}{12} \frac{\dot{q}_o L^2}{k} \end{array}$$

$$\therefore T(x) - T_w = \frac{\dot{q}_o L^2}{2k} \left[ \left( \frac{x}{L} \right)^2 - \frac{1}{6} \left( \frac{x}{L} \right)^4 + \frac{5}{6} \right]$$


---

PROB. 3.12:

Formulation of the problem

$$\left\{ \begin{array}{l} \frac{d^2T}{dr^2} + \frac{1}{r} \frac{dT}{dr} + \frac{\dot{q}_o}{k} \left( 1 - \frac{r}{r_o} \right) = 0 \\ T(0) = \text{finite} \\ T(r_o) = T_w \end{array} \right\} \rightarrow \begin{array}{l} T(r) = -\frac{\dot{q}_o}{k} \left( \frac{r^2}{4} - \frac{r^3}{9r_o} \right) + c_1 \ln r + c_2 \\ c_1 = 0 \\ c_2 = T_w + \frac{\dot{q}_o r_o^2}{k} \frac{5}{36} \end{array}$$


$$\therefore \frac{T(r) - T_w}{\dot{q}_o r_o^2 / k} = \frac{5}{36} - \left( \frac{r}{r_o} \right)^2 \left( \frac{1}{4} - \frac{1}{9} \frac{r}{r_o} \right)$$


---

PROB. 3.13: Formulation of the problem:

$$\left\{ \begin{array}{l} \frac{d^2T}{dr^2} + \frac{1}{r} \frac{dT}{dr} + \frac{\dot{q}}{k} = 0 \\ T(0) = \text{finite} \\ -k \frac{dT(r_o)}{dr} = h [T(r_o) - T_\infty] \end{array} \right\}$$

Sol. to DE:  $T(r) = -\frac{\dot{q} r^2}{4k} + c_1 \ln r + c_2$



B.C. at  $r=0 \Rightarrow c_1=0$  & B.C. at  $r=r_o \Rightarrow c_2 = T_\infty + \frac{\dot{q} r_o}{2h} + \frac{\dot{q} r_o^2}{4k}$

$$\therefore T(r) = T_\infty + \frac{\dot{q} r_o^2}{4k} \left[ 1 - \left( \frac{r}{r_o} \right)^2 \right] + \frac{\dot{q} r_o}{2h}$$

PRDB. 3.14: Assumptions:

1. steady-state.
2.  $k_1$  &  $k_2$  are constant.
3. One-dimensional.
4. Perfect contact at the interface.

Formulation:

$$\frac{d^2T_1}{dr^2} + \frac{1}{r} \frac{dT_1}{dr} + \frac{\dot{q}_o}{k} \left[ 1 - \left( \frac{r}{r_i} \right)^2 \right] = 0, \quad 0 \leq r < r_i$$

$$\frac{d^2T_2}{dr^2} + \frac{1}{r} \frac{dT_2}{dr} = 0, \quad r_i < r < r_2$$

$$\frac{dT_1(0)}{dr} = 0 \quad \text{or} \quad T_1(0) = \text{finite}$$

$$-k_1 \frac{dT_1(r_i)}{dr} = -k_2 \frac{dT_2(r_i)}{dr}$$

$$T_1(r_i) = T_2(r_i) \quad \leftarrow \text{Assuming perfect contact.}$$

$$-k_2 \frac{dT_2(r_2)}{dr} = h [T_2(r_2) - T_f]$$

Solutions to DE's:

$$T_1(r) = -\frac{\dot{q}_o r}{4k_1} \left[ 1 - \left( \frac{r}{2r_i} \right)^2 \right] + C_1 \ln r + C_2 \quad \& \quad T_2(r) = C_3 \ln r + C_4$$

Boundary conditions give:


$$C_1 = 0, \quad C_2 = \frac{3\dot{q}_o r_i^2}{16k_1} + \frac{\dot{q}_o r_i^2}{4k_2} \left[ \ln \frac{r_2}{r_i} + \frac{k_2}{r_2 h} \right] + T_f$$

$$C_3 = -\frac{\dot{q}_o r_i^2}{4k_2}, \quad C_4 = \frac{\dot{q}_o r_i^2}{4k_2} \left[ \frac{k_2}{r_2 h} + \ln r_2 \right] + T_f$$

Thus,

$$T_1(r) - T_f = -\frac{\dot{q}_o r^2}{4k_1} \left[ 1 - \left( \frac{r}{2r_i} \right)^2 \right] + \frac{3}{16} \frac{\dot{q}_o r_i^2}{k_1} + \frac{\dot{q}_o r_i^2}{4k_2} \left[ \ln \frac{r_2}{r_i} + \frac{k_2}{r_2 h} \right], \quad 0 \leq r \leq r_i$$

$$T_2(r) - T_f = \frac{\dot{q}_o r_i^2}{4k_2} \left[ \frac{k_2}{r_2 h} - \ln \left( \frac{r}{r_2} \right) \right], \quad r_i \leq r \leq r_2$$




PROB. 3.15: (a) An energy balance on the system shown yields

$$q_f(x) = q_f(x+\Delta x) + q_{\text{conv.}}$$

As  $\Delta x \rightarrow 0$

$$= q_f(x) + \frac{dq}{dx} \Delta x + P \Delta x h [T(x) - T_\infty]$$



A = cross-section area

P = perimeter

$$\left. \begin{aligned} \frac{dq}{dx} + hP [T(x) - T_\infty] &= 0 \\ q_f &= -kA \frac{dT}{dx} \end{aligned} \right\} \Rightarrow \frac{d^2T}{dx^2} - \frac{hP}{kA} [T(x) - T_\infty] = 0$$

$$\text{Let } \left\{ \begin{aligned} \theta(x) &= T(x) - T_\infty \\ m^2 &= \frac{hP}{kA} \end{aligned} \right\} \rightarrow \frac{d^2\theta}{dx^2} - m^2 \theta = 0$$

$$\therefore \theta(x) = C_1 \sinh mx + C_2 \cosh mx$$

$$T(0) = T_1 \Rightarrow \theta(0) = T_1 - T_\infty = \theta_1 \Rightarrow C_2 = \theta_1$$

$$T(L) = T_2 \Rightarrow \theta(L) = T_2 - T_\infty = \theta_2 \Rightarrow C_1 = \frac{\theta_2 - \theta_1 \cosh mL}{\sinh mL}$$

$$\therefore \theta(x) = \frac{\theta_2 - \theta_1 \cosh mL}{\sinh mL} \sinh mx + \theta_1 \cosh mx$$

$$\boxed{\theta(x) = \theta_2 \frac{\sinh mx}{\sinh mL} + \theta_1 \frac{\sinh [m(L-x)]}{\sinh mL}}$$


b)

$$q_L = -kA \frac{dT(0)}{dx} + kA \frac{dT(L)}{dx} = kA \left[ \frac{dT(L)}{dx} - \frac{dT(0)}{dx} \right]$$

$$= kAm (\theta_2 + \theta_1) \left[ \coth mL - \frac{1}{\sinh mL} \right]$$

$$\boxed{q_L = (\theta_2 + \theta_1) \sqrt{hPKA} \frac{\cosh mL - 1}{\sinh mL}}$$

PROB. 3.16:



From Eq. (3.132) in text:

$$\frac{T_{(L)} - T_{\infty}}{T_b - T_{\infty}} = \frac{149 - T_{\infty}}{65 - T_{\infty}} = \frac{1}{\cosh mL + N \sinh mL}$$

where

$$m = \sqrt{\frac{hP}{kA}}, \quad P = \pi \times 0.5 \text{ in} \quad \& \quad A = \frac{\pi}{4} (0.5^2 - 0.3^2) = \pi \times 0.04 \text{ in}^2$$

$$\frac{A}{P} = \frac{\pi \times 0.04}{\pi \times 0.5} = 0.8 \text{ in} = 0.8 \times 2.54 \times 10^{-2} = 0.02032 \text{ m}$$

$$m = \sqrt{\frac{100}{26 \times 0.02032}} = 13.76 \text{ m}^{-1} \Rightarrow mL = 13.76 \times 2 \times 2.54 \times 10^{-2} = 0.7$$

$$N = \frac{h}{mk} = \frac{100}{13.76 \times 26} = 0.28$$

$$\frac{149 - T_{\infty}}{65 - T_{\infty}} = \frac{1}{\cosh(0.7) + 0.28 \sinh(0.7)} = 0.682$$

$$149 - T_{\infty} = 44.33 - 0.682 \times T_{\infty}$$




$$\underline{T_{\infty} = 32.9^{\circ}\text{C}}$$

PROB. 3.17: For steady-state problems with no internal energy generation Eq. (2.16) reduces to

$$\nabla^2 T = 0$$

In rectangular coordinates,

$$\frac{\partial^2 \theta}{\partial x^2} + \frac{\partial^2 \theta}{\partial y^2} + \frac{\partial^2 \theta}{\partial z^2} = 0, \quad \theta = T - T_{\infty}$$



Integrate over  $y$  from  $y=0$  to  $y=l$ :

$$\int_{y=0}^l \left\{ \frac{\partial^2 \theta}{\partial x^2} + \frac{\partial^2 \theta}{\partial y^2} + \frac{\partial^2 \theta}{\partial z^2} \right\} dy = \int_{y=0}^l \left\{ \frac{\partial^2 \theta}{\partial x^2} + \frac{\partial^2 \theta}{\partial z^2} \right\} dy + \left. \frac{\partial \theta}{\partial y} \right|_{y=0}^{y=l} = 0$$

Integrate over  $z$  from  $z=0$  to  $z=\delta$ :

$$\int_{z=0}^{\delta} \int_{y=0}^l \frac{\partial^2 \theta}{\partial x^2} dy dz + \underbrace{\int_{y=0}^l \left. \frac{\partial \theta}{\partial z} \right|_{z=0}^{z=\delta} dy + \int_{z=0}^{\delta} \left. \frac{\partial \theta}{\partial y} \right|_{y=0}^{y=l} dz = 0$$

$$-\frac{h}{k} \left\{ \int_{y=0}^l [\theta(x, y, \delta) + \theta(x, y, 0)] dy + \int_{z=0}^{\delta} [\theta(x, \delta, z) + \theta(x, 0, z)] dz \right\}$$


For one-dimensional extended surfaces  $\theta(x, y, z) \cong \theta(x)$ . Thus,

$$(S \cdot l) \frac{d^2 \theta}{dx^2} - \frac{h}{k} 2(l + \delta) \theta(x) = 0$$

$$\boxed{\frac{d^2 \theta}{dx^2} - \frac{hP}{kA} \theta(x) = 0}$$

$$A = \delta \cdot l, \quad P = 2(l + \delta)$$

PROB. 3.18:



For unit depth:

$$q'_f = \sqrt{2hkL} \theta_b \frac{I_1(2m\sqrt{L})}{I_0(2m\sqrt{L})}$$

$$m = \sqrt{\frac{2hL}{kb}} = \sqrt{\frac{2 \times 250 \times 0.1}{50 \times 0.02}} = 7.07 \text{ m}^{1/2}$$

$$2m\sqrt{L} = 2 \times 7.07 \sqrt{0.1} = 4.47$$

$$\therefore q'_f = \sqrt{2 \times 250 \times 50 \times 0.02 \times (200 - 40)} \times \frac{I_1(4.47)}{I_0(4.47)} = \underline{3147 \text{ W/m}}$$

Alternative solution:

$$q'_f = h A_f \eta_f (T_\infty - T_b)$$

Refer to Fig. 3.25:

$$L_c = 0.1 \text{ m} \quad \& \quad A_m = 0.1 \times \frac{0.02}{2} = 1 \times 10^{-3} \text{ m}^2$$

$$L_c^{3/2} \left( \frac{h}{kA_m} \right)^{1/2} = (0.1)^{3/2} \left( \frac{250}{50 \times 10^{-3}} \right)^{1/2} = 2.24 \Rightarrow \eta_f \approx 41\%$$

$$\therefore q'_f = 250 \times 2 \times [0.1^2 + 0.01^2]^{1/2} \times 0.41 \times (200 - 40) \approx \underline{3280 \text{ W/m}}$$

PROB. 3.19:

Pipe: ID = 1 in

Fins: L = 0.75 in


Pipe & Fins:

$$t = 0.25 \text{ cm} \text{ (assumed)}$$

$$k = 41 \text{ W/(m.K)}$$

$$h_i = 500 \text{ W/(m}^2\text{.K)}$$

$$h_o = 12 \text{ W/(m}^2\text{.K)}$$



Heat transfer rate from hot water to air per unit length of the tube:

$$q' = \frac{95 - 15}{\frac{1}{h_i A_i} + \frac{1}{2\pi k} \ln \frac{OD}{ID} + \frac{1}{8 \sqrt{h_o P k A} \tanh mL + h_o A_{unf}}}$$

where

$$A_i = \pi ID = \pi \times 2.54 \times 10^{-2} = 7.98 \times 10^{-2} \text{ m}^2, P = 2 \text{ m}, A = 0.25 \times 10^{-2} \text{ m}^2$$

$$m = \sqrt{\frac{h_o P}{k A}} = \left( \frac{12 \times 2}{41 \times 0.25 \times 10^{-2}} \right)^{1/2} = 15.3 \text{ m}^{-1}$$

$$mL = 15.3 \times 0.75 \times 2.54 \times 10^{-2} = 0.2915, \tanh mL = 0.2835$$


$$A_{unf} = [\pi \times (\underbrace{2.54 + 0.5}_{OD}) - 8 \times 0.25] \times 10^{-2} = 7.55 \times 10^{-2} \text{ m}^2$$

$$\frac{80}{q'} = \frac{100}{500 \times 7.98} + \frac{1}{2\pi \times 41} \ln \frac{3.04}{2.54} + \frac{1}{8 \sqrt{12 \times 2 \times 41 \times 0.25 \times 10^{-2}} \times 0.2835 + 12 \times 7.55 \times 10^{-2}}$$

$$\Rightarrow \underline{q' = 320.2 \text{ W/m}}$$

PROB. 3.20: a) Differential equation:

Method - I: Control-volume approach.



$$q(x) = Tc \rho VA + \left( -kA \frac{dT}{dx} \right)$$

P = Perimeter  
A = Cross-section area

$$q(x+\Delta x) = q(x) + \frac{dq}{dx} \Delta x$$

↑ As  $\Delta x \rightarrow 0$

Energy (thermal) balance on the C.V. gives

$$q(x) = q(x+\Delta x) + hP\Delta x [T(x) - T_\infty]$$


$$\frac{dq}{dx} + hP [T(x) - T_\infty] = 0$$

$$\rho c V A \frac{dT}{dx} + \frac{d}{dx} \left( -kA \frac{dT}{dx} \right) + hP [T(x) - T_\infty] = 0$$

$$\frac{d^2T}{dx^2} - \frac{V}{\alpha} \frac{dT}{dx} - \frac{hP}{KA} [T(x) - T_\infty] = 0$$

$\alpha = \frac{k}{\rho c}$

Method - II: System approach.



Energy balance on the system:

$$\cancel{q(x)} = \underbrace{q(x+\Delta x)}_{\cancel{q(x)} + \frac{dq}{dx} \Delta x} + hP\Delta x [T(x) - T_\infty] + \left\{ \begin{array}{l} \text{Rate of change} \\ \text{of internal energy} \\ \text{of the system} \end{array} \right\}$$

$$\frac{dq}{dx} + hP [T - T_\infty] + gCAV \frac{dT}{dx} = 0$$

$$\frac{d}{dt} (gCA \Delta x T) = gCA \Delta x \frac{dT}{dx} \frac{dx}{dt}$$

In this case:  $q(x) = -kA \frac{dT}{dx}$

$$\therefore \boxed{\frac{d^2T}{dx^2} - \frac{V}{\alpha} \frac{dT}{dx} - \frac{hP}{kA} [T - T_\infty] = 0}$$

Boundary conditions:  $T(0) = T_0$  and  $T(L) = T_L$ .

b) Let  $\theta(x) = T(x) - T_\infty$ .

$$\boxed{\frac{d^2\theta}{dx^2} - \frac{V}{\alpha} \frac{d\theta}{dx} - m^2 \theta = 0} \quad m^2 = \frac{hP}{kA}$$

$$\theta(0) = T_0 - T_\infty; \quad \theta(L) = T_L - T_\infty$$

Solution is given by

$$\boxed{\theta(x) = T(x) - T_\infty = e^{ax} \left\{ \frac{\bar{e}^{aL}(T_L - T_\infty) \sinh bx + (T_0 - T_\infty) \sinh b(L-x)}{\sinh bL} \right\}}$$

where

$$a = \frac{V}{2\alpha} \quad \text{and} \quad b = \sqrt{a^2 + m^2}$$

PROB. 3.21: From Prob. 3.20,

$$\Theta(x) = e^{ax} \left[ \frac{e^{-aL}(T_L - T_\infty) \sinh bx + (T_0 - T_\infty) \sinh b(L-x)}{\sinh bL} \right]$$

where

$$a = \frac{V}{2\alpha}, \quad b = \sqrt{a^2 + m^2}, \quad m^2 = \frac{hP}{kA}.$$

In this solution,  $V$  and  $L$  are considered to be known. On the other hand, if we are seeking a relationship between  $V$  and  $L$  (and the other parameters  $T_\infty, h, P, A, k, g, c, T_0$ , and  $T_L$ ), then  $V$  becomes an unknown for given  $L$  (and the other parameters). Therefore, to obtain such a relationship, a third condition has to be imposed on this solution. Since at the spool the temperature is  $T_L$ , it can be assumed that

$$\frac{dT(L)}{dx} \approx 0 \rightarrow \frac{d\Theta(L)}{dx} \approx 0 \quad \text{which yields} \rightarrow \boxed{\frac{aL}{(T_L - T_\infty) \sinh bL} = \frac{a}{b}}$$

This is a transcendental equation for  $V$ , which can be solved, for example, numerically to find  $V$  as a function of the parameters of the problem.

An Approximate Solution for  $V$ : Neglect heat conduction along the wire.

Then,

$$\left. \begin{aligned} \frac{d\Theta}{dx} + \frac{hP}{gcAV} \Theta &= 0 \\ \Theta(0) &= T_0 - T_\infty \end{aligned} \right\} \quad \begin{aligned} \Theta(x) &= \Theta(0) \exp \left\{ -\frac{hP}{gcAV} x \right\} \\ \Theta(L) &= T_L - T_\infty = (T_0 - T_\infty) e^{-\frac{hP}{gcAV} L} \end{aligned}$$

$$\therefore V = \frac{hPL}{gcA} \left\{ \ln \frac{T_0 - T_\infty}{T_L - T_\infty} \right\}^{-1}$$

PROB. 3.22: Assumptions: 1) Steady-state

(i.e.,  $\omega = \text{const.}$ ); 2)  $h = \text{const.}$ ;

3)  $T_{\infty} = \text{Const.}$ ; 4)  $k = \text{const.}$  (given);


5)  $\rho r = \text{const.}$  (given); 6)  $\mu = \text{const.}$ ;

7) Since  $\delta \ll R$ , the temp. distribution can be assumed to be one-dimensional;

that is,  $T = T(r)$ ; 8) Negligible heat loss from the edges at  $r = R$ ;

that is

$$\frac{dT(R)}{dr} \approx 0$$



An energy balance on the system shown yields

$$q(r) + \mu \rho r \omega 2\pi r \Delta r = \underbrace{q(r+\Delta r)}_{q(r) + \frac{dq}{dr} \Delta r} + 2h 2\pi r \Delta r [T(r) - T_{\infty}]$$

$$q(r) = -k 2\pi r (2\delta) \frac{dT}{dr} \rightarrow \theta(r) = T(r) - T_{\infty} \quad \text{Const.}$$

$$r^2 \frac{d^2 \theta}{dr^2} + r \frac{d\theta}{dr} - m^2 r^2 \theta(r) = -\frac{\mu(\rho r) \omega}{2k\delta} r^2, \quad m^2 = \frac{h}{k\delta}$$

$$\therefore \theta(r) = A I_0(mr) + B K_0(mr) + \underbrace{\frac{\mu(\rho r) \omega}{2h}}_{\text{Particular solution}}$$

Boundary conditions:

$$T(0) = \text{finite} \rightarrow \theta(0) = \text{finite} \rightarrow B = 0$$

$$\frac{dT(R)}{dr} = 0 \rightarrow \frac{d\theta(R)}{dr} = 0 \rightarrow A = 0$$

Thus,

$$\underline{\underline{\theta(r) = T(r) - T_{\infty} = \frac{\mu \rho \omega}{2h} r}}$$

PROB. 2.23: If  $p = \text{const.}$ , then the formulation of the problem becomes  
(See Prob. 2.22)

$$r^2 \frac{d^2\theta}{dr^2} + r \frac{d\theta}{dr} - m^2 r^2 \theta(r) = \alpha r^3, \quad \theta(r) = T(r) - T_\infty, \quad m^2 = \frac{h}{k\delta} \quad \& \quad \alpha = -\frac{\mu p \omega}{2k\delta}$$

with the same boundary conditions as in Prob. 2.22.

$$\theta(r) = C_1 I_0(mr) + C_2 K_0(mr) + \theta_p(r)$$

$$\text{Let } \theta_p(r) = A(r) I_0(mr) + B(r) K_0(mr)$$

$$\frac{d\theta_p}{dr} = \underbrace{\frac{dA}{dr} I_0(mr) + \frac{dB}{dr} K_0(mr)}_{\text{Set } = 0} + A(r) m I_1(mr) - B(r) m K_1(mr)$$

$$\frac{dA}{dr} I_0(mr) + \frac{dB}{dr} K_0(mr) = 0 \quad — (1)$$

$$\begin{aligned} \frac{d^2\theta_p}{dr^2} &= \frac{dA}{dr} m I_1(mr) - \frac{dB}{dr} m K_1(mr) + A(r) m [m I_0(mr) - \frac{1}{r} I_1(mr)] \\ &\quad - B(r) m [-m K_0(mr) - \frac{1}{r} K_1(mr)] \end{aligned}$$

Now, substitute  $\theta_p$ ,  $\frac{d\theta_p}{dr}$  and  $\frac{d^2\theta_p}{dr^2}$  into the differential equation:

$$\begin{aligned} &r^2 \left[ \frac{dA}{dr} m I_1(mr) - \frac{dB}{dr} m K_1(mr) + A(r) m I_0(mr) - \cancel{\frac{A(r)m}{r} I_1(mr)} \right] \\ &+ B(r) m^2 K_0(mr) + \cancel{\frac{B(r)m}{r} K_1(mr)} + r \left[ A(r) m I_1(mr) - \cancel{B(r)m K_1(mr)} \right] \\ &- m^2 r^2 \left[ A(r) I_0(mr) + B(r) K_0(mr) \right] = \alpha r^3 \end{aligned}$$

$$\frac{dA}{dr} I_1(mr) - \frac{dB}{dr} K_1(mr) = \frac{\alpha}{m} r \quad — (2)$$

From Eqs. (1) and (2), we obtain

$$\frac{dA}{dr} = \frac{\alpha}{m} \frac{r K_0(mr)}{\Delta(mr)} \quad \text{and} \quad \frac{dB}{dr} = -\frac{\alpha}{m} \frac{r I_0(mr)}{\Delta(mr)}$$

where

$$\Delta(mr) = I_1(mr)K_0(mr) + K_1(mr)I_0(mr) = \frac{1}{mr}$$

See Hildebrand, Adv. Calculus for Applications, 2<sup>nd</sup> ed., p. 178, Prob. 44.4c.

Thus,

$$\frac{dA}{dr} = \alpha r^2 K_0(mr) \quad \text{and} \quad \frac{dB}{dr} = -\alpha r^2 I_0(mr)$$

$$A(r) = \alpha \int_0^r K_0(m\bar{s}) \bar{s}^2 d\bar{s}, \quad B(r) = -\alpha \int_0^r I_0(m\bar{s}) \bar{s}^2 d\bar{s} \leftarrow \begin{array}{l} \text{Note that we} \\ \text{let } A(0) = B(0) = 0 \end{array}$$

(Why?)

The solution can now be written as

$$\theta(r) = C_1 I_0(mr) + C_2 K_0(mr) + \alpha \int_0^r [I_0(m\bar{s}) K_0(m\bar{s}) - I_0(m\bar{s}) K_0(m\bar{s})] \bar{s}^2 d\bar{s} \quad (3)$$

$$\theta(0) = \text{finite} \implies C_2 = 0 \quad (4)$$

$$\begin{aligned} \frac{d\theta(R)}{dr} &= C_1 m I_1(mR) + \alpha \left[ I_0(mR) K_0(mR) - I_0(mR) K_0(mR) \right] R^2 \\ &\quad + \alpha m \int_0^R [I_1(m\bar{s}) K_0(m\bar{s}) + I_0(m\bar{s}) K_1(m\bar{s})] \bar{s}^2 d\bar{s} = 0 \\ \therefore C_1 &= -\frac{\alpha}{I_1(mR)} \int_0^R [I_1(m\bar{s}) K_0(m\bar{s}) + I_0(m\bar{s}) K_1(m\bar{s})] \bar{s}^2 d\bar{s} \end{aligned} \quad (5)$$

Now, substitute  $C_1$  and  $C_2$  from (5) and (4) into (3) to obtain the final form of the solution  $\theta(r)$ .

PROB. 3.24: (a) Differential equation:


$$\frac{d}{dz} \left[ A(z) \frac{d\theta}{dz} \right] - \frac{h P(z)}{k} \theta(z) = 0$$

where

$$\theta(z) = T(z) - T_{\infty}$$

$$A(z) = \pi r^2 = \pi \left( \frac{r_0}{L} z \right)^2 = \pi \left( \frac{r_0}{L} \right)^2 z^2$$

$$P(z) = 2\pi \frac{r_0}{L} z$$



$$z^2 \frac{d^2\theta}{dz^2} + 2z \frac{d\theta}{dz} - m^2 z \theta = 0 , \quad m^2 = \frac{2hL}{kr_0}$$

Solution of this equation is

$$\theta(z) = \frac{-1/2}{z} [ A I_1(2mz^{1/2}) + B K_1(2mz^{1/2}) ]$$

Boundary Conditions:

$$\theta(0) = \text{finite} \Rightarrow B = 0^*$$

$$\theta(L) = T_w - T_{\infty} = \theta_w \Rightarrow A = \frac{\sqrt{L} \theta_w}{I_1(2m\sqrt{L})}$$

$$\therefore \frac{\theta(z)}{\theta_w} = \frac{T(z) - T_{\infty}}{T_w - T_{\infty}} = \sqrt{\frac{L}{z}} \frac{I_1(2m\sqrt{z})}{I_1(2m\sqrt{L})}$$

$$\begin{aligned} b) Q_L &= +k \pi r_0^2 \frac{dT(L)}{dz} = k \pi r_0^2 \frac{d\theta(L)}{dz} = \frac{k \pi r_0^2 \theta_w \sqrt{L}}{I_1(2m\sqrt{L})} \frac{d}{dz} \left\{ \frac{I_1(2m\sqrt{z})}{\sqrt{z}} \right\}_{z=L} \\ &= \frac{\pi r_0^2 k \theta_w \sqrt{L}}{I_1(2m\sqrt{L})} \left\{ \frac{m I_2(2m\sqrt{z})}{z} \right\}_{z=L} = \frac{\pi r_0^2 k (T_w - T_{\infty})}{\sqrt{L}} \frac{m I_2(2m\sqrt{L})}{I_1(2m\sqrt{L})} \end{aligned}$$

\* Because,  $\lim_{z \rightarrow 0} \left\{ \frac{1}{\sqrt{z}} I_1(2m\sqrt{z}) \right\} \rightarrow 0$  and  $\lim_{z \rightarrow \infty} \left\{ \frac{1}{\sqrt{z}} K_1(2m\sqrt{z}) \right\} \rightarrow \infty$

PROB. 3.25: The fin equation:

$$\frac{d}{dx} \left[ A(x) \frac{d\theta}{dx} \right] - \frac{hP}{k} \theta(x) = 0$$

where

and  $\theta(x) = T(x) - T_{\infty}$

$$\begin{cases} A(x) = 2b(x/L)^2 \\ P(x) = 2 \end{cases}$$

For unit depth



$$\frac{d}{dx} \left[ x^2 \frac{d\theta}{dx} \right] - m^2 \theta(x) = 0, \quad m^2 = \frac{hL^2}{b^2 k}$$

↓ Let  $z = \ln x$  or  $x = e^z$

$$\frac{d^2\theta}{dz^2} + \frac{d\theta}{dz} - m^2 \theta(z) = 0 \rightarrow \theta(z) = e^{\frac{1}{2}z} [c_1 e^{-az} + c_2 e^{az}]$$

$$a = \frac{1}{2} \sqrt{1+4m^2}$$

$$\therefore \theta(x) = x^{\frac{1}{2}} [c_1 x^{-a} + c_2 x^a]$$


At  $x=0$ :  $\theta = \text{finite} \Rightarrow c_1 = 0$  (Note  $a > \frac{1}{2}$ )

At  $x=L$ :  $\theta = T_0 - T_{\infty} \Rightarrow c_2 = \frac{T_0 - T_{\infty}}{L^{a-\frac{1}{2}}}$

$$\therefore \frac{T(x) - T_{\infty}}{T_0 - T_{\infty}} = \left( \frac{x}{L} \right)^{a-\frac{1}{2}}$$


Rate of heat loss per unit depth:

$$q'_{\text{loss}} = k \cdot 2b \left( \frac{dT}{dx} \right)_{x=L} = \dots = \frac{2kb}{L} (T_0 - T_{\infty}) (a - \frac{1}{2})$$



PROB. 3.26:

Under steady-state conditions the rod will have a uniform temperature, say  $T_s$ . Thus,



$$(P\mu)(2\pi r_1)(r_1 \omega) = \frac{T_s - T_\infty}{\frac{1}{2\pi k} \ln \frac{r_2}{r_1} + \frac{1}{2\pi r_2 h}}$$

Note that we have taken the interface temperature =  $T_s$ . (Perfect Contact)

$$\therefore T_s = T_\infty + P\mu r_1^2 \omega \left\{ \frac{1}{k} \ln \frac{r_2}{r_1} + \frac{1}{r_2 h} \right\}$$


$$\frac{dT_s}{dr_2} = P\mu r_1^2 \omega \left\{ \frac{1}{kr_2} - \frac{1}{r_2^2 h} \right\} = 0 \Rightarrow r_2 = \frac{k}{h}$$

Thus, the rod (and therefore the interface) temperature will be the lowest for  $r_2 = \frac{k}{h}$ , provided that  $r_1 < \frac{k}{h}$ .

$$\text{Check: } \frac{d^2 T_s}{dr_2^2} \Big|_{r_2 = \frac{k}{h}} = ( ) \left\{ -\frac{1}{kr_2^2} + \frac{2}{hr_2^3} \right\} \Big|_{r_2 = \frac{k}{h}} = ( ) \frac{h^2}{k^3} > 0$$



Thus, at  $r_2 = \frac{k}{h}$ ,  $T_s$  is a minimum, if  $r_1 < \frac{k}{h}$ .

What happens if  $r_1 > \frac{k}{h}$ ? In that case, for  $T_s$  to be minimum  $r_2 \sim r_1$ . See the plot below:



PROB. 3.27:

- Because of symmetry, consider only one half of the system.
- Since the shaft is thin,  $T = T(x)$ .
- Since the sleeve is of negligible thickness, its temperature at any  $x$  will be the same as that of the shaft's surface at the same  $x$ .
- Assume  $k = \text{const.}$



When  $0 < x < L/2$ :

$$q''_w = \mu P r_0 \omega - h [T(x) - T_{\infty}] \quad \rightarrow \quad \frac{d^2\theta}{dx^2} - m_1^2 \theta(x) = -\frac{\mu P r_0 \omega P}{k A}$$

Because of the fact that the sleeve has negligible thickness.

$$\text{where } \theta(x) = T(x) - T_{\infty}, \quad m_1^2 = \frac{h_1 P}{k A}$$

$$\theta(x) = C_1 \sinh m_1 x + C_2 \cosh m_1 x + \theta_p, \quad \theta_p = \frac{\mu P r_0 \omega P}{k A} \frac{KA}{h_1 P} = \frac{\mu P r_0 \omega}{h_1}$$

$$\text{At } x=0: \quad \frac{d\theta}{dx} = 0 \quad \Rightarrow \quad \frac{d\theta}{dx} = 0 \Rightarrow C_1 = 0$$

$$\theta(x) = C_2 \cosh m_1 x + \frac{\mu P r_0 \omega}{h_1} \quad \text{--- ①}$$

When  $\frac{L}{2} < x < L$ :  $q''_w = h_2 [T_\infty - T(x)]$

$$\therefore \frac{d^2\theta}{dx^2} - m_2^2 \theta(x) = 0, \quad m_2^2 = \frac{h_2 P}{kA}$$

$$\theta(x) = C_3 \sinh m_2(L-x) + C_4 \cosh m_2(L-x)$$

$$\text{At } x=L: \quad \frac{dT}{dx} = \frac{d\theta}{dx} = 0 \Rightarrow C_3 = 0$$

$$\theta(x) = C_4 \cosh m_2(L-x) \quad \dots \quad (2)$$

At  $x = L/2$ : From Eqs. (1) & (2) we obtain

$$C_2 \cosh m_1 \frac{L}{2} + \frac{\mu P_f \omega}{h_1} = C_4 \cosh m_2 \frac{L}{2} \quad (\text{continuity of temp.})$$

$$C_2 m_1 \sinh m_1 \frac{L}{2} = -C_4 m_2 \sinh m_2 \frac{L}{2} \quad (\text{continuity of heat flux})$$

$$C_1 = -\frac{\mu P_f \omega}{h_1} \frac{m_2 \sinh m_2 \frac{L}{2}}{m_1 \sinh m_1 \frac{L}{2} \cosh m_2 \frac{L}{2} + m_2 \cosh m_1 \frac{L}{2} \sinh m_2 \frac{L}{2}}$$

$$C_2 = \frac{\mu P_f \omega}{h_1} \frac{m_1 \sinh m_1 \frac{L}{2}}{m_1 \sinh m_1 \frac{L}{2} \cosh m_2 \frac{L}{2} + m_2 \cosh m_1 \frac{L}{2} \sinh m_2 \frac{L}{2}}$$

Thus,


$$T(x) - T_\infty = \frac{\mu P_f \omega}{h_1} \left[ 1 - \frac{m_2 \sinh m_2 \frac{L}{2} \cosh m_1 x}{m_1 \sinh m_1 \frac{L}{2} \cosh m_2 \frac{L}{2} + m_2 \cosh m_1 \frac{L}{2} \sinh m_2 \frac{L}{2}} \right]$$

when  $0 < x < L/2$


$$T(x) - T_\infty = \frac{\mu P_f \omega}{h_1} \frac{m_1 \sinh m_1 \frac{L}{2} \cosh m_2 (L-x)}{m_1 \sinh m_1 \frac{L}{2} \cosh m_2 \frac{L}{2} + m_2 \cosh m_1 \frac{L}{2} \sinh m_2 \frac{L}{2}}$$

when  $L/2 < x < L$

PROB. 3.28:



Consider a section of the rod:



A thermal energy balance on the section (c.v.) shown gives

$$\begin{aligned} q_f(x) + \pi r_0^2 \Delta x \dot{q}_f &= q_f(x + \Delta x) + q_{\text{conv.}} \\ &\downarrow \text{As } \Delta x \rightarrow 0 \\ &= q_f(x) + \frac{dq_f}{dx} \Delta x + h 2\pi r_0 \Delta x [T(x) - T_\infty] \end{aligned}$$

$$\frac{dq_f}{dx} + 2\pi r_0 h [T(x) - T_\infty] = \pi r_0^2 \dot{q}_f$$

$$q_f(x) = -k \pi r_0^2 \frac{dT}{dx}$$

Assuming  $k = \text{const.}$ ,  $\rightarrow -k \pi r_0^2 \frac{d^2 T}{dx^2} + 2\pi r_0 h [T(x) - T_\infty] = \pi r_0^2 \dot{q}_f$

$$\therefore \frac{d^2 \theta}{dx^2} - m^2 \theta(x) = -\frac{\dot{q}_f}{k}, \quad m = \frac{2h}{r_0 k}$$

$$\theta(x) = T(x) - T_\infty$$

$0 < x < L/2$ :  $\dot{q} \neq 0$

$$\theta_1(x) = A \sinh mx + B \cosh mx + \frac{\dot{q}}{m^2 k}$$

$L/2 < x < L$ :  $\dot{q} = 0$

$$\theta_2(x) = C \sinh m(L-x) + D \cosh m(L-x)$$

$$\text{At } x=0: \frac{d\theta_1}{dx} = 0 \Rightarrow A=0$$

$$\text{At } x=L: \frac{d\theta_2}{dx} = 0 \Rightarrow C=0$$

$$\text{At } x=\frac{L}{2}: \theta_1 = \theta_2 \Rightarrow B \cosh m \frac{L}{2} + \frac{\dot{q}}{m^2 k} = D \cosh m \frac{L}{2}$$

$$\frac{d\theta_1}{dx} = \frac{d\theta_2}{dx} \Rightarrow B m \sinh m \frac{L}{2} = -D m \sinh m \frac{L}{2}$$

$$\therefore B = -D$$

$$2B \cosh m \frac{L}{2} = -\frac{\dot{q}}{m^2 k}$$


$$\therefore B = -D = -\frac{\dot{q}}{2m^2 k} \frac{1}{\cosh m \frac{L}{2}}$$

Thus,

$$T(x) - T_\infty = \frac{\dot{q}}{m^2 k} \left[ 1 - \frac{1}{2} \frac{\cosh mx}{\cosh m \frac{L}{2}} \right], \quad 0 \leq x \leq \frac{L}{2}$$

$$T(x) - T_\infty = \frac{\dot{q}}{2m^2 k} \frac{\cosh m(L-x)}{\cosh m \frac{L}{2}}, \quad \frac{L}{2} \leq x \leq L$$

PROB. 3.29:



Consider a section of the metal strip:

$$\frac{q(x) \rightarrow T(x) \rightarrow q(x+\Delta x)}{x \rightarrow | \Delta x |} \quad q_s$$

Energy (thermal) balance on the c.v. shown per unit depth gives

$$\left. \begin{aligned} q(x) &= q(x+\Delta x) + q_s \\ &\downarrow \text{As } \Delta x \rightarrow 0 \\ &= q(x) + \frac{dq}{dx} \Delta x + q_s \end{aligned} \right\} \quad \text{where} \quad q_s = \begin{cases} 2h(T-T_\infty)\Delta x - q''_0\Delta x, & 0 < x < w_1 \\ 2h(T-T_\infty)\Delta x, & w_1 < x < w_2 \end{cases}$$

and

$$q(x) = -k(d \cdot 1) \frac{dT}{dx}$$



$$\frac{d^2T}{dx^2} - \frac{q_s}{k \cdot d} = 0$$

$$0 \leq x \leq w_1 : T_1(x) \equiv T(x).$$

$$\frac{d^2T}{dx^2} - m^2(T_1 - T_\infty) = \frac{q''_0}{kd}, \quad m^2 = \frac{2h}{kd}$$

$$\therefore T_1(x) - T_\infty = C_1 \cosh(mx) + C_2 \sinh(mx) + \frac{q''_0}{2h}$$

$$\omega_1 \leq x \leq \omega_2 : T_2(x) = T(x)$$

$$\frac{d^2 T_2}{dx^2} - m^2 (T_2 - T_\infty) = 0$$

$$\therefore T_2(x) = C_3 \cosh m(\omega_1 - x) + C_4 \sinh m(\omega_2 - x)$$

$$\text{At } x=0: \frac{dT_1}{dx} = 0 \Rightarrow C_2 = 0$$

$$\text{At } x=\omega_2: \frac{dT_2}{dx} = 0 \Rightarrow C_4 = 0$$

$$\text{At } x=\omega_1: T_1 = T_2 \Rightarrow C_1 \cosh(m\omega_1) + \frac{q''_0}{2h} = C_3 \cosh m(\omega_2 - \omega_1)$$

$$\frac{dT_1}{dx} = \frac{dT_2}{dx} \Rightarrow C_1 m \sinh(m\omega_1) = C_3 (-m) \sinh m(\omega_2 - \omega_1)$$



$$C_1 = -\frac{q''_0}{2h} \frac{\sinh m(\omega_2 - \omega_1)}{\sinh(m\omega_2)}$$


$$C_3 = \frac{q''_0}{2h} \frac{\sinh(m\omega_1)}{\sinh(m\omega_2)}$$

Thus,


$$T(x) - T_\infty = \frac{q''_0}{2h} \left[ 1 - \frac{\sinh m(\omega_2 - \omega_1)}{\sinh(m\omega_2)} \cosh(mx) \right], \quad 0 \leq x \leq \omega_1$$

$$T(x) - T_\infty = \frac{q''_0}{2h} \frac{\sinh(m\omega_1)}{\sinh(m\omega_2)} \cosh m(\omega_2 - x), \quad \omega_1 \leq x \leq \omega_2$$

PROB. 3.30:



Formulation (by control volume approach):



$$q(x) = q(x+\Delta x) + q_{\text{conv.}} \quad \begin{matrix} \text{Thermal energy balance} \\ \text{for unit depth.} \end{matrix}$$

As  $\Delta x \rightarrow 0$

$$= q(x) + \frac{dq}{dx} \Delta x + 2 \Delta x h [T(x) - T_f]$$

$$\frac{dq}{dx} + 2h [T(x) - T_f] = 0$$

where

$$q(x) = C_p r_0 \delta T(x) + (-k \delta \frac{dT}{dx})$$

$$\therefore \frac{d^2T}{dx^2} - 2a \frac{dT}{dx} - m^2 [T(x) - T_f] = 0, \quad m^2 = \frac{2h}{k\delta}, \quad a = \frac{1}{2} \frac{r_0 \omega}{\alpha}, \quad \alpha = \frac{k}{C_p c}$$

When  $0 < x < L (= \pi r_0)$  Let  $T(x) = T_i(x)$

$$\frac{d^2T_i}{dx^2} - 2a \frac{dT_i}{dx} - m_1^2 [T_i(x) - T_g] = 0, \quad m_1^2 = \frac{2h_1}{k\delta}$$

$$\therefore T_i(x) - T_g = e^{ax} [A \sinh b_1(L-x) + B \cosh b_1(L-x)], \quad b_1 = \sqrt{a^2 + m_1^2}$$

When  $L (= \pi r_0) < x < 2L$  Let  $T(x) = T_2(x)$

$$\frac{d^2 T_2}{dx^2} - 2\alpha \frac{dT_2}{dx} - m_2^2 [T_2(x) - T_\infty] = 0, \quad m_2^2 = \frac{2h_2}{k\delta}$$

$$\therefore T_2(x) - T_\infty = e^{\alpha x} [C \sinh b_2(L-x) + D \cosh b_2(L-x)], \quad b_2 = \sqrt{\alpha^2 + m_2^2}$$

Apply B.C.'s:

$$\text{At } x=L: \quad T_1(L) = T_2(L) \quad \Rightarrow \quad D-B = e^{-\alpha L} (T_g - T_\infty) \quad -(1)$$

$$\frac{dT_1}{dx} = \frac{dT_2}{dx} \quad \Rightarrow \quad \alpha(D-B) = Cb_2 - Ab_1 \quad -(2)$$

$$\text{At } x=0 \& 2L: \quad T_1(0) = T_2(2L)$$


$$\Rightarrow A \sinh b_1 L + B \cosh b_1 L + (T_g - T_\infty) = e^{2\alpha L} [-C \sinh b_2 L + D \cosh b_2 L] \quad -(3)$$

$$\frac{dT_1(0)}{dx} = \frac{dT_2(L)}{dx}$$

$$\begin{aligned} \Rightarrow \alpha [A \sinh b_1 L + B \cosh b_1 L] + [-Ab_1 \cosh b_1 L - Bb_1 \sinh b_1 L] \\ = \alpha e^{2L} [-C \sinh b_2 L + D \cosh b_2 L] \\ + \alpha^2 e^{2L} [-Cb_2 \cosh b_2 L - b_2 D \sinh b_2 L] \quad -(4) \end{aligned}$$

Solution of the set of four algebraic equations (1), (2), (3) and (4) yields the unknown constants  $A, B, C$ , and  $D$  in the solutions for  $T_1(x) \& T_2(x)$ .

PROB. 3.31:



Under steady-state conditions, a thermal energy balance on the system shown gives

$$q(r) + q_{f_o} + 2q_{conv.} = q(r + \Delta r) \quad (1)$$

As  $\Delta r \rightarrow 0$ ,

$$q(r + \Delta r) = q(r) + \frac{dq}{dr} \Delta r \quad \text{with } q(r) = -k 2\pi r \delta \frac{dT}{dr}$$

$$q_{f_o} = A 2\pi r \Delta r q''_o, \quad A = \begin{cases} 1, & 0 < r < r_o \\ 0, & r_o < r < R \end{cases}$$

$$q_{conv.} = 2\pi r \Delta r h [T_\infty - T(r)]$$

Thus, assuming  $k = \text{const.}$ , Eq.(1) reduces to

$$\frac{1}{r} \frac{d}{dr} \left( r \frac{d\theta}{dr} \right) - m^2 \theta(r) = - \frac{A q''_o}{K \delta} \quad (2)$$

$$\text{where } \theta(r) = T(r) - T_\infty \quad \text{and } m^2 = \frac{2h}{K\delta}$$

Solution of Eq.(2),

$$\theta(r) = \begin{cases} \theta_1(r) = C_1 I_0(mr_o) + C_2 K_0(mr) + \frac{q''_o}{2h}, & 0 < r < r_o \\ \theta_2(r) = C_3 I_0(mr_o) + C_4 K_0(mr), & r_o < r < R \end{cases}$$

Boundary conditions:

$$\textcircled{1} \quad T(0) = \text{finite} \rightarrow \theta_1(0) = \text{finite} \Rightarrow C_2 = 0$$

$$\textcircled{2} \quad \frac{dT(R)}{dr} = 0 \rightarrow \frac{d\theta(R)}{dr} = 0 \leftarrow \text{Negligible heat loss from edge}$$

$$\Rightarrow C_3 m I_1(mR) - C_4 m K_1(mR) = 0 \Rightarrow C_4 = C_3 \frac{I_1(mR)}{K_1(mR)}$$

$$\text{Thus, } \theta_2(r) = C_3 \left[ I_0(mr) + \frac{I_1(mR)}{K_1(mR)} K_0(mr) \right]$$

$$\textcircled{3} \quad \theta_1(r_0) = \theta_2(r_0) \leftarrow \text{Continuity of temperature at } r=r_0.$$

$$C_1 I_0(mr_0) + \frac{Q''_0}{2h} = C_3 \left[ I_0(mr_0) + \frac{I_1(mR)}{K_1(mR)} K_0(mr_0) \right] \quad (3)$$

$$\textcircled{4} \quad \frac{d\theta_1(r_0)}{dr} = \frac{d\theta_2(r_0)}{dr} \leftarrow \text{Continuity of heat flux at } r=r_0.$$

$$C_1 m I_1(mr_0) = C_3 \left[ m I_1(mr_0) - \frac{I_1(mR)}{K_1(mR)} m K_1(mR) \right] \quad (4)$$

Solutions of Eqs. (3) & (4) give

$$C_1 = \frac{\Delta_1}{\Delta_2} \frac{Q''_0}{2h} \quad \text{and} \quad C_3 = \frac{K_1(mR)}{\Delta_2} \frac{Q''_0}{2h} \Rightarrow C_4 = \frac{I_1(mR)}{\Delta_2} \frac{Q''_0}{2h}$$

where


$$\Delta_1 = \frac{1}{I_1(mr_0)} \left[ I_1(mr_0) K_1(mR) - I_1(mR) K_1(mr_0) \right]$$

$$\Delta_2 = \frac{I_1(mR)}{I_1(mr_0)} \left[ I_1(mr_0) K_0(mr_0) + I_0(mr_0) K_1(mr_0) \right]$$


Thus,

$$T(r) - T_{\infty} = \begin{cases} \theta_1(r) = \frac{Q''_0}{2h} \left[ 1 + \frac{\Delta_1}{\Delta_2} I_0(mr) \right], & 0 < r < r_0 \\ \theta_2(r) = \frac{Q''_0}{2h} \frac{1}{\Delta_2} \left[ K_1(mR) I_0(mr) + I_1(mR) K_0(mr) \right], & r_0 < r < R \end{cases}$$

PROB. 3.32:



Consider a section of the rod as shown:



Thermal energy balance on the section shown:

$$\cancel{\dot{q}_{f_z}(z)} + \dot{q}(z) \pi r_0^2 \Delta z = \cancel{\dot{q}_{f_z}(z)} + \frac{d\dot{q}_z}{dz} \Delta z + h 2\pi r_0 \Delta z [T(z) - T_\infty]$$

$$\left. \begin{aligned} \frac{d\dot{q}_z}{dz} + h 2\pi r_0 [T(z) - T_\infty] &= \dot{q}(z) \pi r_0^2 \\ \dot{q}_z &= -k \pi r_0^2 \frac{dT}{dz} \end{aligned} \right\}$$

$$\left[ \begin{aligned} \frac{d^2\theta}{dz^2} - m^2 \theta(z) &= -\frac{\dot{q}_0}{k} e^{-\beta z}, & \theta(z) &= T(z) - T_\infty \\ \theta(0) &= T_0 - T_\infty, & \theta(\infty) &= 0 \end{aligned} \right. \quad \left. \begin{aligned} m^2 &= \frac{2h}{kr_0} \end{aligned} \right]$$

$$\theta(z) = A e^{-mz} + B e^{mz} + \theta_p(z)$$

$$\text{Particular solution: } \theta_p(z) = \begin{cases} \frac{\dot{q}_0}{k(m^2 - \beta^2)} e^{-\beta z}, & \beta \neq m \\ \frac{\dot{q}_0}{2k\beta} z e^{-\beta z}, & \beta = m \end{cases}$$

$$\theta(\infty) = 0 \Rightarrow B = 0$$

$$\theta(0) = T_0 - T_{\infty} \Rightarrow A = \begin{cases} (T_0 - T_{\infty}) - \frac{\dot{q}_0}{k(m^2 - \beta^2)}, & \beta \neq m \\ (T_0 - T_{\infty}), & \beta = m \end{cases}$$


---

$$\therefore \theta(z) = T(z) - T_{\infty} = \begin{cases} (T_0 - T_{\infty}) e^{-mz} + \frac{\dot{q}_0}{k(m^2 - \beta^2)} [e^{-\beta z} - e^{-mz}], & \beta \neq m \\ \left[ T_0 - T_{\infty} + \frac{\dot{q}_0}{2k\beta} z \right] e^{-\beta z}, & \beta = m \end{cases}$$


---

# CHAPTER 4

## THE STURM-LIOUVILLE THEORY AND FOURIER EXPANSIONS

---

PROB. 4.1:  $y(x) = A [B_{12} y_1(x) - B_{11} y_2(x)]$

$$B_{12} = \alpha_1 y_2(a) + \beta_1 \frac{dy_2(a)}{dx}, \quad B_{11} = \alpha_1 y_1(a) + \beta_1 \frac{dy_1(a)}{dx}$$

B.C. (4.3b):  $\alpha_1 y(a) + \beta_1 \frac{dy(a)}{dx} = 0$

$$\alpha_1 A [B_{12} y_1(a) - B_{11} y_2(a)] + \beta_1 A [B_{12} \frac{dy_1(a)}{dx} - B_{11} \frac{dy_2(a)}{dx}] = 0$$

$$\underbrace{\left[ \alpha_1 y_1(a) + \beta_1 \frac{dy_1(a)}{dx} \right]}_{B_{11}} B_{12} - \underbrace{\left[ \alpha_1 y_2(a) + \beta_1 \frac{dy_2(a)}{dx} \right]}_{B_{12}} B_{11} = 0$$

$$B_{11} B_{12} - B_{12} B_{11} = 0 \quad \checkmark$$

B.C. (4.3c):  $\alpha_2 y(b) + \beta_2 \frac{dy(b)}{dx} = 0$

$$\alpha_2 A [B_{12} y_1(b) - B_{11} y_2(b)] + \beta_2 A [B_{12} \frac{dy_1(b)}{dx} - B_{11} \frac{dy_2(b)}{dx}] = 0$$

$$\underbrace{\left[ \alpha_2 y_1(b) + \beta_2 \frac{dy_1(b)}{dx} \right]}_{B_{21}} B_{12} - \underbrace{\left[ \alpha_2 y_2(b) + \beta_2 \frac{dy_2(b)}{dx} \right]}_{B_{22}} B_{11} = 0$$

$$B_{21} B_{12} - B_{22} B_{11} = 0 \quad \text{per Eq. (4.7b)}$$

Following the same procedure, it can also be shown that Eq.(4.10) satisfies the B.C.'s (4.3 b,c).

PROB. 4.2:

$$\frac{d^2y}{dx^2} - \lambda^2 y = 0 \rightarrow y(x) = C_1 \sinh \lambda x + C_2 \cosh \lambda x$$

$y(0) = 0 \text{ and } y(L) = 0$

$$y(0) = 0 \Rightarrow C_2 = 0$$

$$y(L) = 0 \Rightarrow C_1 \sinh \lambda L = 0$$

$$C_1 \sinh \lambda L = 0 \quad \begin{cases} C_1 = 0 \rightarrow \text{Trivial solution} \\ \sinh \lambda L = 0 \Rightarrow \lambda L = 0 \Rightarrow \lambda = 0 \end{cases}$$

Note that  $\lambda = 0$  is the only real root of  $\sinh \lambda L = 0$

PROB. 4.3:

$$A_0; A_1 + A_2 x; A_3 + A_4 x + A_5 x^2$$

$$\int_0^1 A_0 (A_1 + A_2 x) dx = 0 \rightarrow A_1 + \frac{1}{2} A_2 = 0 \quad ①$$

$$\int_0^1 A_0 (A_3 + A_4 x + A_5 x^2) dx = 0 \rightarrow A_3 + \frac{1}{2} A_4 + \frac{1}{3} A_5 = 0 \quad ②$$

$$\int_0^1 (A_1 + A_2 x)(A_3 + A_4 x + A_5 x^2) dx = 0$$

$$\rightarrow A_1 A_3 + \frac{1}{2} (A_1 A_4 + A_2 A_3) + \frac{1}{3} (A_1 A_5 + A_2 A_4) + \frac{1}{4} A_2 A_5 = 0 \quad ③$$

There are 6 constants to be determined, but there are only 3 equations. Hence, 3 of the constants can be taken arbitrarily, say,  $A_0 = 1$ ,  $A_1 = 1$  &  $A_3 = 1$ . Thus, we obtain

$$A_2 = -2, A_4 = -6 \text{ and } A_5 = 6$$

Hence, the orthogonal set would be

$$\{1, 1-2x, 1-6x+6x^2\}$$

PROB. 4.4:

$$a_0(x) \frac{d^2y}{dx^2} + a_1(x) \frac{dy}{dx} + [a_2(x) + \lambda a_3(x)] y = 0$$

$$\underbrace{\frac{d^2y}{dx^2} + \frac{a_1}{a_0} \frac{dy}{dx}}_{\frac{1}{P(x)} \frac{d}{dx} \left[ \bar{P}(x) \frac{dy}{dx} \right]} + \left[ \frac{a_2}{a_0} + \lambda \frac{a_3}{a_0} \right] y = 0$$

$$\therefore \frac{1}{\bar{P}(x)} \frac{d\bar{P}}{dx} = \frac{a_1}{a_0} \rightarrow \frac{d\bar{P}}{\bar{P}} = \frac{a_1}{a_0} dx \rightarrow \ln \bar{P}(x) = \int \frac{a_1}{a_0} dx + \ln C$$

$$\bar{P}(x) = C_1 e^{\int \frac{a_1}{a_0} dx} = C_1 P(x); \quad P(x) = e^{\int \frac{a_1}{a_0} dx}$$

$$\frac{1}{C_1 P(x)} \frac{d}{dx} \left[ C_1 P(x) \frac{dy}{dx} \right] = \frac{1}{P(x)} \frac{d}{dx} \left[ P(x) \frac{dy}{dx} \right]$$

$$\frac{d}{dx} \left[ P(x) \frac{dy}{dx} \right] + \left[ \underbrace{\frac{a_2}{a_0} P(x)}_{q(x)} + \lambda \underbrace{\frac{a_3}{a_0} P(x)}_{w(x)} \right] y = 0$$

PROB. 4.5:

$$\boxed{\begin{aligned} \frac{d^2y}{dx^2} + \lambda^2 y &= 0 \\ \frac{dy(0)}{dx} &= 0, \quad \frac{dy(L)}{dx} = 0 \end{aligned}}$$

$$y(x) = A \sin \lambda x + B \cos \lambda x$$

$$\frac{dy}{dx} = A \lambda \cos \lambda x - B \lambda \sin \lambda x$$

$$\frac{dy(0)}{dx} = 0 \Rightarrow A = 0$$

$$\frac{dy(L)}{dx} = 0 \Rightarrow B \lambda \sin \lambda L = 0 \Rightarrow \sin \lambda L = 0 \Rightarrow \lambda_n = \frac{n\pi}{L}, n=0,1,2,\dots$$

$$\therefore \phi_n(x) = \cos \lambda_n x; \quad \lambda_n = \frac{n\pi}{L}, n=0,1,2,\dots$$

PROB. 4.6: (a)

$$\frac{d^2y}{dx^2} + \lambda^2 y = 0 \rightarrow y(x) = C_1 \cos \lambda x + C_2 \sin \lambda x$$

$$\frac{dy(0)}{dx} = 0, \alpha y(L) + \beta \frac{dy(L)}{dx} = 0 \quad \frac{dy(0)}{dx} = 0 \Rightarrow C_2 = 0$$

$$\therefore \phi(x) = \cos \lambda x$$

$$\alpha y(L) + \beta \frac{dy(L)}{dx} = 0 \Rightarrow \alpha \cancel{\phi} \cos \lambda L + \beta (-\cancel{\phi} \lambda \sin \lambda L) = 0$$

$$\lambda \tan \lambda L = H, \quad H = \frac{\alpha}{\beta}$$

$$\therefore \phi_n(x) = \cos \lambda_n x$$

and  $\lambda_n$ 's are the positive roots of  $\lambda \tan \lambda L = H$ .

(b)

$$f(x) = \sum_{n=1}^{\infty} A_n \cos \lambda_n x, \quad 0 < x < L$$

where

$$A_n = \frac{1}{N_n} \int_0^L f(x) \cos \lambda_n x \, dx$$

with

$$N_n = \int_0^L \cos^2 \lambda_n x \, dx = \frac{1}{2\lambda_n} [ \lambda_n L + \sin \lambda_n L \cos \lambda_n L ]$$

PROB. 4.7: (a)

$$\left. \begin{array}{l} \frac{d^2y}{dx^2} + \lambda^2 y = 0 \\ y(0) = 0 \\ y(L) - L \frac{dy(L)}{dx} = 0 \end{array} \right\} \text{Note that } \lambda = 0 \text{ is an eigenvalue.}$$

$$y_0(x) = a_0 x + b_0$$

$$y_0(0) = 0 \rightarrow b_0 = 0$$

$$y_0(L) - L \frac{dy_0(L)}{dx} = 0 \rightarrow a_0 L - L a_0 = 0 \quad \checkmark$$

$$\therefore y_0(x) = a_0 \phi_0(x), \quad \phi_0(x) = x$$

When  $\lambda \neq 0$ ,

$$y_n(x) = a_n \sin \lambda_n x + b_n \cos \lambda_n x$$

$$y_n(0) = 0 \Rightarrow b_n = 0$$

$$y_n(L) - L \frac{dy_n(L)}{dx} = 0 \Rightarrow \sin \lambda_n L - L \lambda_n \cos \lambda_n L = 0$$

$$\tan \lambda_n L = \lambda_n L$$

↓

Thus, the eigenfunctions are

$$\lambda_n, n=1, 2, 3, \dots$$

$$\phi_n(x) = \begin{cases} x, & n=0 \\ \sin \lambda_n x, & n \neq 0 \end{cases}$$

and the eigenvalues are the zeros of

$$\tan \lambda_n L = \lambda_n L, n=0, 1, 2, \dots$$

(b)

$$f(x) = A_0 x + \sum_{n=1}^{\infty} A_n \sin \lambda_n x, 0 < x < L$$

$$A_0 = \frac{1}{N_0} \int_0^L f(x) x \, dx, \quad N_0 = \int_0^L x^2 \, dx = \frac{L}{3}$$

$$A_n = \frac{1}{N_n} \int_0^L f(x) \sin \lambda_n x \, dx, \quad N_n = \int_0^L \sin^2 \lambda_n x \, dx \\ = \dots = \frac{L}{2} \sin^2 \lambda_n L$$

Thus,

$$A_0 = \frac{3}{L} \int_0^L f(x) x \, dx, \quad n=0$$

$$A_n = \frac{2}{L \sin^2 \lambda_n L} \int_0^L f(x) \sin \lambda_n x \, dx, \quad n=1, 2, 3, \dots$$

PROB. 4.8:

$$f(x) = \begin{cases} 1, & x < \frac{L}{2} \\ 0, & x > \frac{L}{2} \end{cases}$$

$$f(x) = \sum_{n=1}^{\infty} A_n \sin \frac{n\pi}{L} x, \quad 0 < x < L$$

$$\begin{aligned} A_n &= \frac{2}{L} \int_0^L f(x) \sin \frac{n\pi}{L} x \, dx = \frac{2}{L} \int_0^{L/2} \sin \frac{n\pi}{L} x \, dx \\ &= \frac{2}{L} \left[ -\frac{L}{n\pi} \cos \frac{n\pi}{L} x \right]_{x=0}^{x=L/2} = \frac{2}{n\pi} \left[ 1 - \cos \frac{n\pi}{2} \right] \end{aligned}$$

$$\therefore f(x) = \frac{2}{\pi} \sum_{n=1}^{\infty} \frac{1}{n} \left[ 1 - \cos \frac{n\pi}{2} \right] \sin \frac{n\pi}{L} x, \quad 0 < x < L$$


---

PROB. 4.9:

$$f(x) = A_0 + \sum_{n=1}^{\infty} A_n \cos \frac{n\pi}{L} x, \quad 0 < x < L$$

$$A_0 = \frac{1}{L} \int_0^L f(x) \, dx = \frac{1}{L} \int_0^{L/2} \, dx = \frac{1}{2}$$

$$\begin{aligned} A_n &= \frac{2}{L} \int_0^L f(x) \cos \frac{n\pi}{L} x \, dx \\ &= \frac{2}{L} \int_0^{L/2} \cos \frac{n\pi}{L} x \, dx = \frac{2}{n\pi} \sin \frac{n\pi}{2} \end{aligned}$$

$$\therefore f(x) = \frac{1}{2} + \frac{2}{\pi} \sum_{n=1}^{\infty} \frac{1}{n} \sin \frac{n\pi}{2} \cdot \cos \frac{n\pi}{L} x, \quad 0 < x < L$$

PROB. 4.10: Since  $\{\phi_n(x); n=0,1,2,\dots\}$  is the complete set of all the characteristic functions of the Sturm-Liouville problem (4.13), and  $f(x)$  is any piecewise differentiable function in the interval  $(a,b)$ ,

$$f(x) = \sum_{n=0}^{\infty} A_n \phi_n(x)$$

is a valid expansion in  $(a,b)$ , where

$$A_n = \frac{1}{N_n} \int_a^b f(x) \phi_n(x) w(x) dx \quad \text{with} \quad N_n = \int_a^b \phi_n^2(x) w(x) dx$$

and  $w(x)$  is the weight function. Multiply both sides of this expansion by  $f(x)w(x)$  and integrate the resultant relation over  $(a,b)$ :

$$\begin{aligned} \int_a^b [f(x)]^2 w(x) dx &= \int_a^b f(x) \left\{ \sum_{n=0}^{\infty} A_n \phi_n(x) \right\} w(x) dx \\ &= \sum_{n=0}^{\infty} A_n \underbrace{\int_a^b f(x) \phi_n(x) w(x) dx}_{N_n A_n} = \sum_{n=0}^{\infty} N_n A_n^2 \end{aligned}$$

PROB. 4.11: (a)

$$1 = \sum_{n=1}^{\infty} A_n \sin \frac{n\pi}{L} x$$

$$\therefore A_n = \frac{2}{L} \int_0^L 1 \cdot \sin \frac{n\pi}{L} x dx = -\frac{2}{n\pi} \cos \frac{n\pi}{L} x \Big|_0^L = -\frac{2}{n\pi} [(-1)^n - 1]$$

$$(b) \quad \underbrace{\int_0^L 1^2 dx}_L = \sum_{n=1}^{\infty} \underbrace{\frac{L}{2} \left\{ -\frac{2}{n\pi} [(-1)^n - 1] \right\}^2}_{\frac{4L}{\pi^2} \sum_{n=1}^{\infty} \frac{1 - (-1)^n}{n^2}} \Rightarrow \frac{\pi^2}{4} = \sum_{n=1}^{\infty} \frac{1 - (-1)^n}{n^2}$$

PROB. 4.12:

$$f(x) = B_0 + \sum_{n=1}^{\infty} \left[ A_n \sin \frac{n\pi}{L} x + B_n \cos \frac{n\pi}{L} x \right], \quad -L < x < L$$

$$B_0 = \frac{1}{2L} \int_{-L}^L f(x) dx = \frac{1}{2L} \int_{-L}^{L/2} dx = \frac{3}{4}$$

$$A_n = \frac{1}{L} \int_{-L}^L f(x) \sin \frac{n\pi}{L} x dx, \quad n=1, 2, 3, \dots$$

$$= \frac{1}{L} \int_{-L}^{L/2} \sin \frac{n\pi}{L} x dx = \frac{1}{n\pi} \underbrace{[\cos(n\pi) - \cos(\frac{n\pi}{2})]}_{(-1)^n}$$

$$B_n = \frac{1}{L} \int_{-L}^L f(x) \cos \frac{n\pi}{L} x dx, \quad n=1, 2, 3, \dots$$

$$= \frac{1}{L} \int_{-L}^{L/2} \cos \frac{n\pi}{L} x dx = \frac{1}{n\pi} \sin \frac{n\pi}{2}$$

$$\therefore f(x) = \frac{3}{4} + \frac{1}{\pi} \sum_{n=1}^{\infty} \frac{1}{n} \left\{ \left[ (-1)^n - \cos \frac{n\pi}{2} \right] \sin \frac{n\pi}{L} x + \sin \frac{n\pi}{2} \cos \frac{n\pi}{L} x \right\}$$

PROB. 4.13:

$$\frac{d^2y}{dx^2} + \lambda^2 y = 0 \rightarrow y(x) = A \sin \lambda x + B \cos \lambda x$$

$$y(-L) = y(L) \rightarrow \left\{ 2A \sin \lambda L = 0 \right\}$$

$$\frac{dy(-L)}{dx} = \frac{dy(L)}{dx} \rightarrow \left\{ 2B \lambda \sin \lambda L = 0 \right\}$$

↓

$$\lambda_n = \frac{n\pi}{L}, \quad n=0, 1, 2, \dots$$

Note that corresponding to each eigenvalue  $\lambda_n$  there are two eigenfunctions:

$$\phi_n(x) = \sin \lambda_n x \text{ and } \psi_n(x) = \cos \lambda_n x, n=0, 1, 2, \dots$$

(b) Since the above eigenvalue problem is a Sturm-Liouville system (with periodic boundary conditions), the set

$$\{ \sin \lambda_n x, \cos \lambda_n x; n=0, 1, 2, \dots \}$$

is a complete orthogonal set over the interval  $(-L, L)$  with respect to  $w(x)=1$ . Thus, an arbitrary function  $f(x)$ , piecewise differentiable over  $(-L, L)$ , can be expanded as

$$\begin{aligned} f(x) &= \sum_{n=0}^{\infty} (a_n \sin \lambda_n x + b_n \cos \lambda_n x) \\ &= b_0 + \sum_{n=1}^{\infty} (a_n \sin \lambda_n x + b_n \cos \lambda_n x), \quad -L < x < L \end{aligned}$$

where expansion coefficients can then be calculated as

$$a_n = \frac{\int_{-L}^L f(x) \sin \lambda_n x dx}{\int_{-L}^L \sin^2 \lambda_n x dx} = \frac{1}{L} \int_{-L}^L f(x) \sin \lambda_n x dx \quad n=1, 2, 3, \dots$$

$$b_0 = \frac{\int_{-L}^L f(x) dx}{\int_{-L}^L dx} = \frac{1}{2L} \int_{-L}^L f(x) dx$$

$$b_n = \frac{\int_{-L}^L f(x) \cos \lambda_n x dx}{\int_{-L}^L \cos^2 \lambda_n x dx} = \frac{1}{L} \int_{-L}^L f(x) \cos \lambda_n x dx \quad n=1, 2, 3, \dots$$

Here, it should be noted that, due to orthogonality,

$$\int_{-L}^L \sin \lambda_n x \cos \lambda_m x dx = 0; n, m = 0, 1, 2, 3, \dots$$

$$\int_{-L}^L \sin \lambda_n x \sin \lambda_m x dx = 0, \quad m \neq n$$

and

$$\int_{-L}^L \cos \lambda_n x \cos \lambda_m x dx = 0, \quad m \neq n$$

PROB. 4.14:

$$r = \sum_{n=1}^{\infty} A_n J_0(\lambda_n r), \quad 0 < r < r_0$$

where  $\lambda_n$ 's are the positive roots of  $J_0(\lambda_n r_0) = 0$ .

$$\begin{aligned} A_n &= \frac{1}{N_n} \int_0^{r_0} r J_0(\lambda_n r) r dr, \quad N_n = \frac{r_0^2}{2} J_1^2(\lambda_n r_0) \\ &\quad \vdots \\ &= \frac{1}{N_n} \frac{1}{\lambda_n} \left\{ r_0^2 J_1(\lambda_n r_0) - \frac{1}{\lambda_n} \int_0^{r_0} J_0(\lambda_n r) dr \right\} \end{aligned}$$

↑  
Table 4.2

Note that  $\int_0^{r_0} J_0(\lambda_n r) dr$  cannot be further simplified.

The function,

$$\int_0^r J_0(\lambda_n r) dr$$

is a tabulated function.

PROB. 4.15:

$$r^2 \frac{d^2 R}{dr^2} + r \frac{dR}{dr} - (\lambda^2 r^2 + \nu^2) R = 0 \rightarrow R(r) = A I_\nu(\lambda r) + B K_\nu(\lambda r)$$

$$R(0) = \text{finite} \rightarrow B = 0$$

$$R(r_0) = 0 \rightarrow A I_\nu(\lambda r_0) = 0$$

If  $\nu = 0$ ,  $I_0(\lambda r_0) = 0$  has no real root.

In this case,  $A = 0$ . But this leads to  $R(r) = 0$

If  $\nu > 0$ ,  $I_\nu(\lambda r_0) = 0 \Rightarrow \lambda = 0$ . But this also leads to trivial solution  $R(r) = 0$ .

---

PROB. 4.16: (a)

$$\left\{ \begin{array}{l} \frac{d^2 R}{dr^2} + \frac{1}{r} \frac{dR}{dr} + \lambda^2 R = 0 \\ R(a) = 0, \quad R(b) = 0 \end{array} \right\}$$

$$R(r) = C_1 J_0(\lambda r) + C_2 Y_0(\lambda r)$$

$$C_1 J_0(\lambda a) + C_2 Y_0(\lambda a) = 0$$

$$C_1 J_0(\lambda b) + C_2 Y_0(\lambda b) = 0$$

In order to have non-trivial solution for  $C_1$  and  $C_2$ ,

$$\begin{vmatrix} J_0(\lambda a) & Y_0(\lambda a) \\ J_0(\lambda b) & Y_0(\lambda b) \end{vmatrix} = 0 \quad \text{or} \quad J_0(\lambda a) Y_0(\lambda b) - J_0(\lambda b) Y_0(\lambda a) = 0$$

$\Downarrow$

$$\lambda_n, \quad n = 1, 2, \dots$$

Then, the algebraic equations for  $C_1$  and  $C_2$  become identical, yielding

$$C_1 = -C_2 \frac{Y_0(\lambda_n a)}{J_0(\lambda_n a)} = -C_2 \frac{Y_0(\lambda_n b)}{J_0(\lambda_n b)}$$

Thus,

$$\begin{aligned} R_n(r) &= C_1 \left[ J_0(\lambda_n r) + \frac{C_2}{C_1} Y_0(\lambda_n r) \right] \\ &= C_1 \left[ J_0(\lambda_n r) - \frac{J_0(\lambda_n b)}{Y_0(\lambda_n b)} Y_0(\lambda_n r) \right] \\ &= \underbrace{\frac{C_1}{Y_0(\lambda_n b)}}_{\text{Arbitrary Constant}} \left[ J_0(\lambda_n r) Y_0(\lambda_n b) - J_0(\lambda_n b) Y_0(\lambda_n r) \right] \end{aligned}$$

---

$\therefore$  Eigenfunctions :  $\underline{\phi_n(r) = J_0(\lambda_n r) Y_0(\lambda_n b) - J_0(\lambda_n b) Y_0(\lambda_n r)}$

---

(b)

$$f(r) = \sum_{n=1}^{\infty} A_n \phi_n(r), \quad a < r < b$$

$$\therefore A_n = \frac{1}{N_n} \int_a^b f(r) \phi_n(r) r dr$$

where

$$N_n = \int_a^b \phi_n^2(r) r dr$$

$$\begin{aligned} &\vdots \\ &= \frac{2}{\pi^2 \lambda_n^2} \left[ 1 - \frac{J_0^2(\lambda_n b)}{J_0^2(\lambda_n a)} \right] \end{aligned}$$

# CHAPTER 5

## STEADY TWO- AND THREE-DIMENSIONAL HEAT CONDUCTION


---

PROB. 5.1: Solution by superposition:

$$\begin{array}{c}
 \text{Diagram showing superposition of four terms:} \\
 \text{Term 1: } \boxed{\begin{matrix} 100^\circ & \\ 100 & 100 \\ 100 & \end{matrix}} = 0 \\
 \text{Term 2: } \boxed{\begin{matrix} 100 & \\ T & 0 \\ 0 & \end{matrix}} + 0 \\
 \text{Term 3: } \boxed{\begin{matrix} 0 & \\ T & 100 \\ 0 & \end{matrix}} + 0 \\
 \text{Term 4: } \boxed{\begin{matrix} 0 & \\ T & 100 \\ 0 & \end{matrix}} + 0
 \end{array}$$

$$4T = 100 \Rightarrow T = 25^\circ C$$

Solution by using Eq. (5.19):




$$\begin{aligned}
 T\left(\frac{a}{2}, \frac{a}{2}\right) &= \frac{200}{\pi} \sum_{n=1}^{\infty} \frac{[1 - (-1)^n]}{n} \frac{\sin \frac{n\pi}{a} \frac{a}{2} \sinh \frac{n\pi}{a} \frac{a}{2}}{\sinh \frac{n\pi}{a} a} \\
 &= \frac{200}{\pi} \left\{ 2 \frac{\sinh \frac{\pi}{2}}{\sinh \pi} - \frac{2}{3} \frac{\sinh \frac{3\pi}{2}}{\sinh 3\pi} \right. \\
 &\quad \left. + \frac{2}{5} \frac{\sinh \frac{5\pi}{2}}{\sinh 5\pi} - \frac{2}{7} \frac{\sinh \frac{7\pi}{2}}{\sinh 7\pi} \dots \right\}
 \end{aligned}$$

$$\begin{aligned}
 T\left(\frac{a}{2}, \frac{a}{2}\right) &= \frac{400}{\pi} (0.199268 - 0.002994 + 0.000078 \\
 &\quad - 2 \times 10^{-6} + \dots)
 \end{aligned}$$

$$\approx 25.00006^\circ C = 25^\circ C$$

PROB. 5.2:



$$\frac{T(x,y)}{T_0} = \frac{2}{\pi} \sum_{n=1}^{\infty} \frac{[1-(-1)^n]}{n} \frac{\sin \frac{n\pi}{a} x \sinh \frac{n\pi}{a} (b-y)}{\sinh \frac{n\pi}{a} b}$$

$$\frac{x}{a} = 0.25, \quad \frac{y}{b} = 0.75 \quad \& \quad b = 2a$$

$$\frac{T(x,y)}{T_0} = \frac{2}{\pi} \sum_{n=1}^{\infty} \frac{[1-(-1)^n]}{n} \frac{\sin \frac{n\pi}{4} \sinh \frac{n\pi}{2}}{\sinh 2n\pi}$$

$$\begin{aligned} \frac{T(x,y)}{T_0} &= \frac{2}{\pi} \left\{ 2 \frac{\sin \frac{\pi}{4} \sinh \frac{\pi}{2}}{\sinh 2\pi} + \frac{2}{3} \frac{\sin \frac{3\pi}{4} \sinh \frac{3\pi}{2}}{\sinh 6\pi} \right. \\ &\quad \left. + \frac{2}{5} \frac{\sin \frac{5\pi}{4} \sinh \frac{5\pi}{2}}{\sinh 10\pi} + \frac{2}{7} \frac{\sin \frac{7\pi}{4} \sinh \frac{7\pi}{2}}{\sinh 14\pi} + \dots \right\} \end{aligned}$$

$$= \frac{4}{\pi} \left[ 6.0777 \times 10^{-3} + 1.7086 \times 10^{-7} - 8.274 \times 10^{-12} \right. \\ \left. - 4.769 \times 10^{-16} + \dots \right]$$

$$\therefore \frac{T(x,y)}{T_0} \approx 7.74 \times 10^{-3}$$

PROB. 5.3: Formulation of the problem:

$$\left\{ \begin{array}{l} \frac{\partial^2 T}{\partial x^2} + \frac{\partial^2 T}{\partial y^2} = 0 \\ T(0,y) = 0; \quad T(a,y) = f(y) \\ T(x,0) = 0; \quad T(x,b) = 0 \end{array} \right\}$$

Let  $T(x,y) = X(x) \cdot Y(y)$

Then,

$$\left\{ \begin{array}{l} \frac{d^2Y}{dy^2} + \lambda^2 Y = 0 \\ Y(0) = Y(b) = 0 \end{array} \right\} \Rightarrow Y_n(y) = A_n \sin \lambda_n y \quad \Rightarrow \left\{ \begin{array}{l} \frac{d^2X_n}{dx^2} - \lambda_n^2 X_n = 0 \\ X_n(0) = 0 \end{array} \right\}$$

↓

Thus,

$$X_n(x) = B_n \sinh \lambda_n x$$

$$T(x,y) = \sum_{n=1}^{\infty} a_n \sinh \lambda_n x \sin \lambda_n y$$


↓

$$f(y) = \sum_{n=1}^{\infty} \underbrace{a_n \sin \lambda_n a}_{b_n} \sin \lambda_n y \quad \Rightarrow b_n = \frac{2}{b} \int_0^b f(y) \sin \lambda_n y dy$$


---


$$\therefore T(x,y) = \frac{2}{b} \sum_{n=1}^{\infty} \frac{\sinh \frac{n\pi}{b} x \cdot \sin \frac{n\pi}{b} y}{\sinh \frac{n\pi}{b} a} \cdot \int_0^b f(y') \sin \frac{n\pi}{b} y' dy'$$

PROB. 5.4:



$$\eta = a - x$$

From Prob. 5.3:


$$\frac{1}{b} \sum_{n=1}^{\infty} \frac{\sinh \frac{n\pi}{b} \eta \cdot \sin \frac{n\pi}{b} y}{\sinh \frac{n\pi}{b} a} \cdot \int_0^b g(y') \sin \frac{n\pi}{b} y' dy'$$


---


$$\therefore T(x,y) = \frac{2}{b} \sum_{n=1}^{\infty} \frac{\sinh \frac{n\pi}{b} (a-x) \sin \frac{n\pi}{b} y}{\sinh \frac{n\pi}{b} a} \cdot \int_0^b g(y') \sin \frac{n\pi}{b} y' dy'$$

PROB. 5.5:

$$T(x,y) = \frac{2}{\pi} \left\{ T_1 \sum_{n=1}^{\infty} \frac{[1-(-1)^n]}{n} \frac{\sin \frac{n\pi}{a} x \sinh \frac{n\pi}{a} y}{\sinh \frac{n\pi}{a} b} \right. \\ + T_2 \sum_{n=1}^{\infty} \frac{[1-(-1)^n]}{n} \frac{\sin \frac{n\pi}{a} x \sinh \frac{n\pi}{a} (b-y)}{\sinh \frac{n\pi}{a} b} \\ + T_3 \sum_{n=1}^{\infty} \frac{[1-(-1)^n]}{n} \frac{\sinh \frac{n\pi}{b} x \sin \frac{n\pi}{b} y}{\sinh \frac{n\pi}{b} a} \\ \left. + T_4 \sum_{n=1}^{\infty} \frac{[1-(-1)^n]}{n} \frac{\sinh \frac{n\pi}{b} (a-x) \sin \frac{n\pi}{b} y}{\sinh \frac{n\pi}{b} a} \right\}$$



Assume  $k=\text{const.}$

$$T(x,y) = \frac{2}{\pi} \sum_{n=1}^{\infty} \frac{[1-(-1)^n]}{n} \left\{ \frac{\sin \frac{n\pi}{a} x}{\sinh \frac{n\pi}{a} b} [T_1 \sinh \frac{n\pi}{a} y + T_2 \sinh \frac{n\pi}{a} (b-y)] \right. \\ \left. + \frac{\sin \frac{n\pi}{b} y}{\sinh \frac{n\pi}{b} a} [T_3 \sinh \frac{n\pi}{b} x + T_4 \sinh \frac{n\pi}{b} (a-x)] \right\}$$

PROB. 5.6:


$$\text{Let } \theta(x,y) = T(x,y) - T_1.$$

Then the formulation of the problem  
is given by

$$\frac{\partial^2 \theta}{\partial x^2} + \frac{\partial^2 \theta}{\partial y^2} = 0$$

$$\frac{\partial \theta(0,y)}{\partial x} = 0; \quad \theta(a,y) = 0$$

$$\frac{\partial \theta(x,0)}{\partial y} = 0; \quad \theta(x,b) = T_2 - T_1 \equiv \theta_0$$



$$\text{Let } \theta(x,y) = X(x)Y(y)$$

Thus,

$$\left\{ \begin{array}{l} \frac{d^2X}{dx^2} + \lambda_n^2 X = 0 \\ \frac{dX(0)}{dx} = 0 \\ X(a) = 0 \end{array} \right\} \Rightarrow X_n(x) = A_n \cos \lambda_n x \quad \Rightarrow \left\{ \begin{array}{l} \frac{d^2Y_n}{dy^2} - \lambda_n^2 Y_n = 0 \\ \frac{dY_n(0)}{dy} = 0 \end{array} \right\}$$

$\Downarrow$

Then,

$$\Theta(x, y) = \sum_{n=0}^{\infty} a_n \cos \lambda_n x \cosh \lambda_n y$$

$Y_n(y) = B_n \cosh \lambda_n y$

$\Downarrow$

$$\theta_0 = \sum_{n=0}^{\infty} a_n \cosh \lambda_n b \cos \lambda_n x$$

$\Downarrow$

$$a_n = \frac{\theta_0}{\cosh \lambda_n b} \frac{\int_0^a \cos \lambda_n x dx}{\int_0^a \cosh^2 \lambda_n x dx} = \dots = \frac{(-1)^n 4 \theta_0}{(2n+1)\pi \cosh \lambda_n b}$$

$$\therefore \frac{\Theta(x, y)}{\theta_0} = \frac{T(x, y) - T_1}{T_2 - T_1} = \frac{4}{\pi} \sum_{n=0}^{\infty} \frac{(-1)^n}{(2n+1)} \cos \frac{2n+1}{a} \frac{\pi}{2} x \cdot \frac{\cosh \frac{2n+1}{a} \frac{\pi}{2} y}{\cosh \frac{2n+1}{a} \frac{\pi}{2} b}$$


PROB. 5.7: a) Formulation of the problem

in terms of  $\theta(x, y) = T(x, y) - T_\infty$ :

$$\frac{\partial^2 \theta}{\partial x^2} + \frac{\partial^2 \theta}{\partial y^2} = 0$$

$$\left. \frac{\partial \theta}{\partial x} \right|_{x=0} = 0, \left[ k \frac{\partial \theta}{\partial x} + h \theta(x, y) \right]_{x=a} = 0$$

$$\theta(x, 0) = T_g - T_\infty, \left[ k \frac{\partial \theta}{\partial y} + h \theta(x, y) \right]_{y=b} = 0$$



$$\text{Let } \Theta(x, y) = X(x) \cdot Y(y)$$

Then,

$$\left\{ \begin{array}{l} \frac{d^2X}{dx^2} + \lambda^2 X = 0 \\ \frac{dX(0)}{dx} = 0 \\ k \frac{dX(a)}{dx} + h X(a) = 0 \end{array} \right\} \Rightarrow \begin{array}{l} \frac{d^2Y_n}{dy^2} - \lambda_n^2 Y_n = 0 \\ k \frac{dY_n(b)}{dy} + h Y_n(b) = 0 \end{array}$$

↓

$$X_n(x) = A_n \cos \lambda_n x$$

where  $\lambda_n$  are the roots of

$$(\lambda a) \tan(\lambda a) = Bi, \quad Bi = \frac{ha}{k}$$

↓

$$\lambda_n, \quad n=1, 2, 3, \dots$$

$$Y_n(y) = B_n \left\{ \sinh \lambda_n (b-y) + \frac{\lambda_n a}{Bi} \cosh \lambda_n (b-y) \right\}$$

Thus,

$$\Theta(x, y) = \sum_{n=1}^{\infty} a_n \cos \lambda_n x \cdot \left\{ \sinh \lambda_n (b-y) + \frac{\lambda_n a}{Bi} \cosh \lambda_n (b-y) \right\}$$

$$\Rightarrow T_g - T_{\infty} = \sum_{n=1}^{\infty} a_n \cos \lambda_n x \cdot \left\{ \sinh \lambda_n b + \frac{\lambda_n a}{Bi} \cosh \lambda_n b \right\}$$

$$\therefore a_n \left\{ \sinh \lambda_n b + \frac{\lambda_n a}{Bi} \cosh \lambda_n b \right\} = \frac{1}{N_n} \int_0^a (T_g - T_{\infty}) \cos \lambda_n x dx = \frac{(T_g - T_{\infty})}{N_n} \frac{\sin \lambda_n a}{\lambda_n}$$

$$\text{where } N_n = \frac{1}{2\lambda_n} [\lambda_n a + \sin \lambda_n a \cos \lambda_n a] \quad (\text{From Table 4.1})$$

$$a_n = 2(T_g - T_{\infty}) \frac{\sin \lambda_n a}{[\lambda_n a + \sin \lambda_n a \cos \lambda_n a][\sinh \lambda_n b + \frac{\lambda_n a}{Bi} \cosh \lambda_n b]}$$

$$\therefore \frac{T(x,y) - T_{\infty}}{T_g - T_{\infty}} = 2 \sum_{n=1}^{\infty} \frac{\sin \lambda_n a \cos \lambda_n x}{[\lambda_n a + \sin \lambda_n a \cos \lambda_n a]} \cdot \frac{\sinh \lambda_n (b-y) + \frac{\lambda_n a}{Bi} \cosh \lambda_n (b-y)}{[\sinh \lambda_n b + \frac{\lambda_n a}{Bi} \cosh \lambda_n b]}$$

b) The rate of heat loss per unit depth can be calculated as

$$q'_L = 2 \int_0^a \left\{ -k \frac{\partial T}{\partial y} \Big|_{y=0} dx \right\}$$

$$= 4k(T_g - T_{\infty}) \sum_{n=1}^{\infty} \left\{ \frac{\sin^2 \lambda_n a}{[\lambda_n a + \sin \lambda_n a \cos \lambda_n a]} \frac{\cosh \lambda_n b + \frac{\lambda_n a}{Bi} \sinh \lambda_n b}{\sinh \lambda_n b + \frac{\lambda_n a}{Bi} \cosh \lambda_n b} \right\}$$

PROB. 5.B: Formulation of the problem in terms of  $\theta(x,y) = T(x,y) - T_1$ ,  
(see Prob. 2.8):

$$2 \frac{\partial^2 \theta}{\partial x^2} + \frac{\partial^2 \theta}{\partial y^2} = 0$$

$$\theta(0,y) = 0; \quad \theta(a,y) = 0$$

$$\theta(x,0) = 0; \quad \theta(x,b) = T_2 - T_1$$

Let  $\eta = \sqrt{2}y$ . Then, the formulation becomes:

$$\frac{\partial^2 \theta}{\partial x^2} + \frac{\partial^2 \theta}{\partial \eta^2} = 0$$

$$\theta(0,\eta) = 0; \quad \theta(a,\eta) = 0$$


$$\theta(x,0) = 0; \quad \theta(x,\sqrt{2}b) = T_2 - T_1$$

Now, the solution can be written as [see Eq. (5.19)]:

$$\frac{T(x,y) - T_1}{T_2 - T_1} = \frac{2}{\pi} \sum_{n=1}^{\infty} \frac{[1 - (-1)^n]}{n} \frac{\sin \frac{n\pi}{a} x \sinh \frac{n\pi}{a} \sqrt{2}y}{\sinh \frac{n\pi}{a} \sqrt{2}b}$$

PROB. 5.9: Formulation:

$$\left\{ \begin{array}{l} \frac{\partial^2 T}{\partial x^2} + \frac{\partial^2 T}{\partial y^2} = 0 \\ \frac{\partial T(0,y)}{\partial x} = 0; \quad T(a,y) = 0 \\ T(x,0) = f(x); \quad T(x,\infty) = 0 \end{array} \right\}$$



Let  $T(x,y) = X(x) \cdot Y(y)$ . Then,

$$\left\{ \begin{array}{l} \frac{d^2 X}{dx^2} + \lambda^2 X = 0 \\ \frac{dX(0)}{dx} = 0 \\ X(a) = 0 \end{array} \right\} \Rightarrow X_n(x) = A_n \cos \lambda_n x \quad \lambda_n = \frac{(2n+1)\pi}{a}, \quad n=0, 1, 2, \dots$$

and

$$\left\{ \begin{array}{l} \frac{d^2 Y_n}{dy^2} - \lambda_n^2 Y_n = 0 \\ Y_n(\infty) = 0 \end{array} \right\} \Rightarrow Y_n(y) = B_n e^{-\lambda_n y}$$

Thus,

$$T(x,y) = \sum_{n=0}^{\infty} \underbrace{A_n B_n}_{a_n} \cos \lambda_n x e^{-\lambda_n y}$$

$$f(x) = \sum_{n=0}^{\infty} a_n \cos \lambda_n x \quad \rightarrow a_n = \frac{2}{a} \int_0^a f(x) \cos \lambda_n x dx$$

---


$$\therefore T(x,y) = \frac{2}{a} \sum_{n=0}^{\infty} \left\{ \int_0^a f(x') \cos \lambda_n x' dx' \right\} \cdot \cos \lambda_n x e^{-\lambda_n y}$$



---

The rate of heat dissipated from the surface at  $x=a$  per unit depth can be calculated as

$$q'_L = \int_0^a \left\{ -k \left( \frac{\partial T}{\partial y} \right)_{y=0} \right\} dx = \frac{2k}{a} \sum_{n=0}^{\infty} (-1)^n \int_0^a f(x) \cos \lambda_n x dx$$


---

PROB. 5.10:




Hence, the solution can be written as

$$T(x,y) - T_1 = \frac{2}{\pi} (T_2 - T_1) \left\{ \sum_{n=1}^{\infty} \frac{[1 - (-1)^n]}{n} \frac{\sin \frac{n\pi}{a} x \times \sinh \frac{n\pi}{a} (a-y)}{\sinh n\pi} \right. \\ \left. + \sum_{n=1}^{\infty} \frac{[1 - (-1)^n]}{n} \frac{\sinh \frac{n\pi}{a} x \times \sin \frac{n\pi}{a} y}{\sinh n\pi} \right\}$$

Or,

$$\frac{T(x,y) - T_1}{T_2 - T_1} = \frac{2}{\pi} \sum_{n=1}^{\infty} \frac{[1 - (-1)^n]}{n \sinh n\pi} \left\{ \sin \frac{n\pi}{a} x \times \sinh \frac{n\pi}{a} (a-y) \right. \\ \left. + \sinh \frac{n\pi}{a} x \times \sin \frac{n\pi}{a} y \right\}$$

PROB. 5.11:



$$T(x,y) - T_1 = \frac{2}{\pi} \left\{ - (T_2 - T_1) \sum_{n=1}^{\infty} \frac{[1 - (-1)^n]}{n} \frac{\sinh \frac{n\pi}{a} x \times \sin \frac{n\pi}{a} y}{\sinh n\pi} \right. \\ \left. + (T_2 - T_1) \sum_{n=1}^{\infty} \frac{[1 - (-1)^n]}{n} \frac{\sin \frac{n\pi}{a} x \times \sinh \frac{n\pi}{a} (a-y)}{\sinh n\pi} \right\}$$


$$\frac{T(x,y) - T_1}{T_2 - T_1} = \frac{2}{\pi} \sum_{n=1}^{\infty} \frac{[1 - (-1)^n]}{n \sinh n\pi} \left[ \sin \frac{n\pi}{a} x \times \sinh \frac{n\pi}{a} (a-y) \right. \\ \left. - \sinh \frac{n\pi}{a} x \times \sin \frac{n\pi}{a} y \right]$$

PROB. 5.12: Formulation of the problem:

$$\frac{\partial^2 T}{\partial x^2} + \frac{\partial^2 T}{\partial y^2} + \frac{\dot{q}(x)}{k} = 0$$

$$\frac{\partial T(0,y)}{\partial x} = 0, \quad T(a,y) = 0$$

$$\frac{\partial T(x,0)}{\partial y} = 0, \quad T(x,b) = 0$$



a) Let  $T(x,y) = \psi(x,y) + \phi(x)$ . Then, the problem separates into two as:

$$\frac{\partial^2 \psi}{\partial x^2} + \frac{\partial^2 \psi}{\partial y^2} = 0$$

$$\frac{\partial \psi(0,y)}{\partial x} = 0, \quad \psi(a,y) = 0$$

$$\frac{\partial \psi(x,0)}{\partial y} = 0, \quad \psi(x,b) = -\phi(x)$$

$$\frac{d^2 \phi}{dx^2} + \frac{\dot{q}(x)}{k} = 0$$

$$\frac{d\phi(0)}{dx} = 0, \quad \phi(a) = 0$$

---


$$\phi(x) = \frac{1}{k} \int_{x_0}^a \int_{x''}^x \dot{q}(x'') dx'' dx'$$


---

The solution to  $\psi(x,y)$  problem (See Prob. 5.6):

$$\psi(x,y) = \sum_{n=0}^{\infty} A_n \cos \frac{(2n+1)\pi}{2a} x \times \cosh \frac{(2n+1)\pi}{2a} y$$

$$-\phi(x) = \sum_{n=0}^{\infty} A_n \cos \frac{(2n+1)\pi}{2a} x \times \cosh \frac{(2n+1)\pi}{2a} b$$

$$\therefore A_n \cosh \frac{(2n+1)\pi}{2a} b = -\frac{2}{a} \int_0^a \phi(x) \cos \frac{(2n+1)\pi}{2a} x dx$$

Thus, the solution is given by

$$T(x,y) = \frac{1}{k} \int_x^a \int_0^{x'} \dot{q}(x'') dx'' dx' - \frac{2}{ak} \sum_{n=0}^{\infty} \frac{\cos \frac{(2n+1)\pi}{2a} x \cosh \frac{(2n+1)\pi}{2a} y}{\cosh \frac{(2n+1)}{2a} b} \int_0^a \int_0^{x''} \dot{q}(x'') dx'' dx' dx''$$

b) Let

$$T(x,y) = \sum_{n=0}^{\infty} A_n(y) \cos \frac{(2n+1)\pi}{2a} x, \quad \frac{dA_n(0)}{dy} = A_n(b) = 0$$

and

$$\frac{\dot{q}(x)}{k} = \sum_{n=0}^{\infty} B_n \cos \frac{(2n+1)\pi}{2a} x$$

*Known constants*  $\rightarrow B_n = \frac{2}{ak} \int_0^a \dot{q}(x) \cos \frac{(2n+1)\pi}{2a} x dx, n=0,1,2,\dots$

Note that the expansion given for  $T(x,y)$  satisfies all the boundary conditions of the problem. Now, substitute both expansions into the differential equation to obtain

$$\sum_{n=0}^{\infty} \left[ \frac{d^2 A_n}{dy^2} - \lambda_n^2 A_n(y) + B_n \right] \cos \lambda_n x = 0 \quad \text{with } \lambda_n = \frac{(2n+1)\pi}{2a}, n=0,1,2,\dots$$

$\downarrow$

$$\left\{ \begin{array}{l} \frac{d^2 A_n}{dy^2} - \lambda_n^2 A_n(y) = -B_n \\ \frac{dA_n(0)}{dy} = 0; A_n(b) = 0 \end{array} \right\} \rightarrow A_n(y) = C_n \sinh \lambda_n y + D_n \cosh \lambda_n y + \frac{B_n}{\lambda_n^2}$$

$$\frac{dA_n(0)}{dy} = 0 \Rightarrow C_n = 0$$

$$A_n(b) = 0 \Rightarrow D_n = -\frac{B_n}{\lambda_n^2 \cosh \lambda_n b}$$

$$\therefore A_n(y) = \frac{B_n}{\lambda_n^2} \left[ 1 - \frac{\cosh \lambda_n y}{\cosh \lambda_n b} \right]$$

Thus,

$$T(x, y) = \frac{2}{ak} \sum_{n=0}^{\infty} \frac{\cos \lambda_n x}{\lambda_n^2} \left[ 1 - \frac{\cosh \lambda_n y}{\cosh \lambda_n b} \right] \cdot \int_0^a \dot{q}(x') \cos \lambda_n x' dx'$$

c) Let

$$T(x, y) = \sum_{m=0}^{\infty} \sum_{n=0}^{\infty} A_{mn} \cos \frac{(2m+1)\pi}{2a} x \cdot \cos \frac{(2n+1)\pi}{2b} y$$

satisfies all  
the boundary  
conditions

and

$$\begin{aligned} \frac{\dot{q}(x)}{k} &= \sum_{m=0}^{\infty} \sum_{n=0}^{\infty} B_{mn} \cos \frac{(2m+1)\pi}{2a} x \cdot \cos \frac{(2n+1)\pi}{2b} y \\ \downarrow & \\ B_{mn} &= \frac{4}{abk} \int_0^b \int_0^a \dot{q}(x) \cos \beta_m x \cos \lambda_n y dx dy \\ &= \frac{4}{abk} \frac{2b}{(2n+1)\pi} (-1)^n \int_0^a \dot{q}(x) \cos \beta_m x dx \end{aligned}$$

$$\begin{aligned} \beta_m &= \frac{(2m+1)\pi}{2a} \\ \lambda_n &= \frac{(2n+1)\pi}{2b} \end{aligned}$$

known constants

Substituting both expansions into the differential equation, we obtain

$$\sum_{m=0}^{\infty} \sum_{n=0}^{\infty} \left\{ [-\beta_m^2 - \lambda_n^2] A_{mn} + B_{mn} \right\} \cdot \cos \beta_m x \cdot \cos \lambda_n y = 0$$

$$A_{mn} = \frac{B_{mn}}{\beta_m^2 + \lambda_n^2}$$

Thus,

$$T(x, y) = \frac{8}{ak\pi} \sum_{m=0}^{\infty} \sum_{n=0}^{\infty} \frac{(-1)^n}{(2n+1)} \frac{\cos \beta_m x \cdot \cos \lambda_n y}{\beta_m^2 + \lambda_n^2} \cdot \int_0^a \dot{q}(x') \cos \beta_m x' dx'$$

PROB. 5.13:

$$\frac{\partial^2 T}{\partial x^2} + \frac{\partial^2 T}{\partial y^2} + \frac{q(x,y)}{k} = 0$$

$$\frac{\partial T(0,y)}{\partial x} = 0; \quad T(a,y) = 0$$

$$\frac{\partial T(x,0)}{\partial y} = 0; \quad T(x,b) = 0$$

Let

$$T(x,y) = \sum_{m=0}^{\infty} \sum_{n=0}^{\infty} A_{mn} \cos \frac{(2m+1)\pi}{2a} x \cos \frac{(2n+1)\pi}{2b} y$$

and

$$\frac{q(x,y)}{k} = \sum_{m=0}^{\infty} \sum_{n=0}^{\infty} B_{mn} \cos \frac{(2m+1)\pi}{2a} x \cos \frac{(2n+1)\pi}{2b} y$$

$$\therefore B_{mn} = \frac{4}{abk} \int_0^a \int_0^b q(x,y) \cos \lambda_m x \cos \beta_n y dx dy$$

$$\text{with } \lambda_m = \frac{(2m+1)\pi}{2a} \quad \text{and } \beta_n = \frac{(2n+1)\pi}{2b}$$

Substitute both expansions into the DE and obtain

$$\therefore A_{mn} = \frac{B_{mn}}{\lambda_m^2 + \beta_n^2}$$

Thus,

$$T(x,y) = \frac{4}{abk} \sum_{m=0}^{\infty} \sum_{n=0}^{\infty} \frac{\cos \lambda_m x \cos \beta_n y}{\lambda_m^2 + \beta_n^2} \int_0^a \int_0^b q(x',y') \cos \lambda_m x' \cos \beta_n y' dx' dy'$$

PROB. 5.14: Obtain the solution from Eq. (5.108) by replacing  $F(z)$  by  $T_0$ :

$$\frac{T(r,z)}{T_0} = \frac{2}{\pi} \sum_{n=1}^{\infty} \frac{[1 - (-1)^n]}{n} \frac{J_0\left(\frac{n\pi}{L}r\right) \sin \frac{n\pi}{L}z}{J_0\left(\frac{n\pi}{L}r_0\right)}$$

PROB. 5.15: Make use of Eq. (5.98) and replace  $z$  by  $(L-z)$  to obtain

$$T(r,z) = \frac{2}{r_0^2} \sum_{n=1}^{\infty} \frac{J_0(\lambda_n r)}{J_1^2(\lambda_n r_0)} \frac{\sinh \lambda_n z}{\sinh \lambda_n L} \int_0^{r_0} f(r') J_0(\lambda_n r') r' dr'$$

where  $\lambda_n$ 's are the positive roots of  $J_0(\lambda r_0) = 0$ .

PROB. 5.16: a) Obtain the solution by superposition by the use of Eqs.(5.98) and (5.108) and the result of Prob. 5.15:

$$\begin{aligned} T(r,z) &= \frac{2}{L} \sum_{n=1}^{\infty} \frac{J_0\left(\frac{n\pi}{L}r\right)}{J_0\left(\frac{n\pi}{L}r_0\right)} \sin \frac{n\pi}{L}z \cdot \int_0^L f_1(z') \sin \frac{n\pi}{L}z' dz' \\ &\quad + \frac{2}{r_0^2} \sum_{n=1}^{\infty} \frac{J_0(\lambda_n r)}{J_1^2(\lambda_n r_0) \sinh \lambda_n L} \left\{ \sinh \lambda_n (L-z) \int_0^{r_0} f_2(r') J_0(\lambda_n r') r' dr' \right. \\ &\quad \left. + \sinh \lambda_n z \int_0^{r_0} f_3(r') J_0(\lambda_n r') r' dr' \right\} \end{aligned}$$

where  $\lambda_n$ 's are the positive roots of  $J_0(\lambda r_0) = 0$

b) If  $f_1(z) = T_0$  and  $f_2(r) = f_3(r) = T_1$ , then the solution can be written as

$$\frac{T(r,z) - T_1}{T_0 - T_1} = \frac{2}{\pi} \sum_{n=1}^{\infty} \frac{[1 - (-1)^n]}{n} \frac{J_0\left(\frac{n\pi}{L}r\right) \sin \frac{n\pi}{L}z}{J_0\left(\frac{n\pi}{L}r_0\right)}$$

or

$$\frac{T(r,z) - T_0}{T_1 - T_0} = \frac{2}{r_0} \sum_{n=1}^{\infty} \frac{1}{\lambda_n} \frac{J_0(\lambda_n r)}{J_1(\lambda_n r_0)} \frac{\sinh \lambda_n(L-z) + \sinh \lambda_n z}{\sinh \lambda_n L}$$

where Eq. (5.99) has been used.

---

PROB. 5.17: Formulation in terms of  $\theta(r,z) = T(r,z) - T_\infty$ :

$$\left\{ \begin{array}{l} \frac{\partial^2 \theta}{\partial r^2} + \frac{1}{r} \frac{\partial \theta}{\partial r} + \frac{\partial^2 \theta}{\partial z^2} = 0 \\ \theta(0,z) = \text{finite}; \quad \frac{\partial \theta(r_0, z)}{\partial r} + \frac{h}{k} \theta(r_0, z) = 0 \\ \theta(r, 0) = f(r) - T_\infty; \quad \frac{\partial \theta(r, L)}{\partial z} + \frac{h}{k} \theta(r, L) = 0 \end{array} \right\}$$

Let  $\theta(r,z) = R(r) \cdot Z(z)$ . Then,

$$\theta(r,z) = \sum_{n=1}^{\infty} a_n J_0(\lambda_n r) \left[ \sinh \lambda_n(L-z) + \frac{\lambda_n k}{h} \cosh \lambda_n L \right]$$

where  $\lambda_n$  are the positive roots of  $\frac{J_1(\lambda r_0)}{J_0(\lambda r_0)} = \frac{h \lambda_n}{k}$ . The boundary condition at  $z=0$  yields

$$f(r) - T_\infty = \sum_{n=1}^{\infty} a_n J_0(\lambda_n r) \left[ \sinh \lambda_n L + \frac{\lambda_n k}{h} \cosh \lambda_n L \right]$$

$$\Rightarrow a_n \left[ \sinh \lambda_n L + \frac{\lambda_n k}{h} \cosh \lambda_n L \right] = \frac{1}{N_n} \int_0^{r_0} [f(r) - T_\infty] J_0(\lambda_n r) r dr$$

$$\text{where } N_n = \frac{r_0^2}{2} \left[ 1 + \frac{h^2}{k^2 \lambda_n^2} \right] J_0^2(\lambda_n r_0) = \frac{r_0^2}{2} \left[ J_0^2(\lambda_n r_0) + J_1^2(\lambda_n r_0) \right]$$

Thus,

$$\theta(r,z) = \frac{2}{r_0^2} \sum_{n=1}^{\infty} \frac{J_0(\lambda_n r) \left[ \sinh \lambda_n(L-z) + \frac{\lambda_n k}{h} \cosh \lambda_n L \right]}{\left[ J_0^2(\lambda_n r_0) + J_1^2(\lambda_n r_0) \right] \left[ \sinh \lambda_n L + \frac{\lambda_n k}{h} \cosh \lambda_n L \right]} \int_0^{r_0} [f(r) - T_\infty] J_0(\lambda_n r) r dr$$

PROB. 5.18: Formulation of the problem :

$$\frac{\partial^2 T}{\partial r^2} + \frac{1}{r} \frac{\partial T}{\partial r} + \frac{\partial^2 T}{\partial z^2} = 0$$

$$T(0, z) = \text{finite}; \quad T(r_0, z) = T_0 + \sin \frac{\pi z}{L}$$

$$T(r, 0) = T_0; \quad T(r, L) = T_0$$

$$\text{Let } \theta(r, z) = T(r, z) - T_0 = R(r) \cdot Z(z)$$

Then,

$$\left. \begin{array}{l} \frac{d^2 Z}{dz^2} + \lambda^2 Z = 0 \\ Z(0) = 0; \quad Z(L) = 0 \end{array} \right\} \Rightarrow Z_n(z) = A_n \sin \frac{n\pi}{L} z, \quad n=1, 2, 3, \dots$$

and

$$\left. \begin{array}{l} \frac{d^2 R_n}{dr^2} + \frac{1}{r} \frac{dR_n}{dr} - \lambda_n^2 R_n = 0 \\ R_n(0) = \text{finite} \end{array} \right\} \Rightarrow R_n(r) = B_n I_0(\lambda_n r)$$

Thus,

$$\theta(r, z) = \sum_{n=1}^{\infty} C_n I_0(\lambda_n r) \sin \frac{n\pi}{L} z$$

$$\downarrow$$

$$\sin \frac{\pi z}{L} = \sum_{n=1}^{\infty} C_n I_0(\lambda_n r_0) \sin \frac{n\pi}{L} z$$

$$\therefore C_1 = \frac{1}{I_0(\lambda_n r_0)}, \quad \text{and} \quad C_n = 0, \quad n=2, 3, \dots$$

Then, the solution becomes


$$\theta(r, z) = T(r, z) - T_0 = \frac{I_0(\frac{\pi r}{L})}{I_0(\frac{\pi r_0}{L})} \sin \frac{\pi z}{L}$$

PROB. 5.19: The temperature distribution is, in fact, one-dimensional, that is,

$$T = T(z)$$

because, if  $\frac{\partial T}{\partial r}$  is zero at  $r=r_0$  then under steady-state conditions it will be zero for all values of  $r$  in the rod. Therefore,

$$\underline{T(z) = T_1 + \frac{T_2 - T_1}{H} z}$$



Alternative Method of Solution with the assumption of  $T(r,z)$ :  
Formulation of the problem in terms of  $\theta(r,z) = T(r,z) - T_1$ :

$$\frac{\partial^2 \theta}{\partial r^2} + \frac{1}{r} \frac{\partial \theta}{\partial r} + \frac{\partial^2 \theta}{\partial z^2} = 0$$

$$\theta(0,z) = \text{finite}; \quad \frac{\partial \theta(r_0, z)}{\partial r} = 0$$

$$\theta(r,0) = 0; \quad \theta(r,H) = T_2 - T_1$$

$$\text{Let } \theta(r,z) = R(r) \cdot Z(z)$$

$$\left\{ \begin{array}{l} r^2 \frac{d^2 R}{dr^2} + r \frac{dR}{dr} + \lambda^2 r^2 R = 0 \\ R(0) \neq 0 \\ \frac{dR(r_0)}{dr} = 0 \end{array} \right\} \quad \begin{array}{l} R_n(r) = A_n J_0(\lambda_n r), \quad n=0,1,2,\dots \\ \text{where } \lambda_n \text{'s are the positive roots of} \\ \frac{d}{dr} [J_0(\lambda r)] = 0 \text{ or } J_1(\lambda r) = 0 \end{array}$$

Note that  $\lambda_0 = 0$  is a characteristic value.

Corresponding to each  $\lambda_n$ ,

$$\left\{ \begin{array}{l} \frac{d^2 z_n}{dz^2} - \lambda_n^2 z_n = 0 \\ z_n(0) = 0 \end{array} \right\} \Rightarrow \left\{ \begin{array}{l} z_0(z) = B_0 z \text{ for } \lambda_0 = 0 \\ z_n(z) = B_n \sinh \lambda_n z, n=1,2,\dots \end{array} \right.$$

Hence, the product solution can be constructed as

$$\Theta(r,z) = \underbrace{C_0}_{A_0 B_0} z + \sum_{n=1}^{\infty} \underbrace{A_n B_n}_{C_n} J_0(\lambda_n r) \sinh \lambda_n z$$

Imposing the B.C. at  $z=H$ :

$$T_2 - T_1 = C_0 H + \sum_{n=1}^{\infty} \underbrace{C_n \sinh \lambda_n H}_{b_n} J_0(\lambda_n r) \quad \leftarrow \text{Fourier-Bessel expansion of } T_2 - T_1.$$

$$\therefore b_0 = \frac{2}{r_0^2} \int_0^{r_0} (T_2 - T_1) r dr = T_2 - T_1 \rightarrow C_0 = \frac{T_2 - T_1}{H}$$

$$b_n = \frac{1}{N_n} \int_0^{r_0} (T_2 - T_1) J_0(\lambda_n r) r dr = \underbrace{\frac{T_2 - T_1}{N_n} \int_0^{r_0} J_0(\lambda_n r) r dr}_0 = 0$$

$$\Rightarrow C_n = 0, n=1,2,3,\dots$$

Because

Hence,

$$\Theta(r,z) = \frac{T_2 - T_1}{H} z$$

or


$\{1, J_0(\lambda_n r); n=1,2,3,\dots\}$   
is an orthogonal set (complete).

$$\boxed{T(r,z) - T_1 = \frac{T_2 - T_1}{H} z}$$

PROB. 5.20: a) Formulation of the problem

in terms of  $\theta(r, z) = T(r, z) - T_0$ :

$$\frac{\partial^2 \theta}{\partial r^2} + \frac{1}{r} \frac{\partial \theta}{\partial r} + \frac{\partial^2 \theta}{\partial z^2} = 0$$



$$\theta(0, z) = \text{finite}; \quad -k \frac{\partial \theta}{\partial r} \Big|_{r=r_0} = h [\theta(r_0, z) - \theta_\infty], \quad \theta_\infty = T_\infty - T_0$$

$$\theta(r, 0) = 0; \quad \theta(r, L) = 0$$

Let  $\theta(r, z) = R(r) \cdot Z(z)$  to obtain

$$\left\{ \begin{array}{l} \frac{d^2 Z}{dz^2} + \lambda^2 Z = 0 \\ Z(0) = 0 \\ Z(L) = 0 \end{array} \right\} \Rightarrow Z_n(z) = A_n \sin \lambda_n z \quad \text{and} \quad \left\{ \begin{array}{l} r^2 \frac{d^2 R_n}{dr^2} + r \frac{d R_n}{dr} - \lambda_n^2 r^2 R_n = 0 \\ R_n(0) = \text{finite} \end{array} \right\}$$

$$R_n(r) = B_n I_0(\lambda_n r)$$

$$\therefore \theta(r, z) = \sum_{n=1}^{\infty} a_n I_0(\lambda_n r) \sin \lambda_n z$$

$$\text{B.C. at } r=r_0 \Rightarrow h \theta_\infty = \sum_{n=1}^{\infty} [h I_0(\lambda_n r_0) + k \lambda_n I_1(\lambda_n r_0)] a_n \sin \lambda_n z$$

$$a_n = \frac{2\theta_\infty}{n\pi} \frac{[1 - (-1)^n]}{I_0(\lambda_n r_0) + \frac{n\pi}{B_i} I_1(\lambda_n r_0)}, \quad B_i = \frac{hL}{k}$$

$$\therefore \frac{\theta(r, z)}{\theta_\infty} = \frac{T(r, z) - T_0}{T_\infty - T_0} = 2 \frac{B_i}{\pi} \sum_{n=1}^{\infty} \frac{[1 - (-1)^n]}{n} \frac{I_0(\lambda_n r) \sin \lambda_n z}{B_i I_0(\lambda_n r_0) + n\pi I_1(\lambda_n r_0)}$$

b)

$$Q_{\text{loss}} = 2 \times \int_0^{r_0} \left\{ -k 2\pi r dr \frac{\partial T}{\partial z} \Big|_{z=0} \right\} dr = 8\pi \frac{k}{L} B_i \theta_\infty \sum_{n=1}^{\infty} \frac{[(-1)^n - 1] \int_0^{r_0} I_0(\lambda_n r) r dr}{B_i I_0(\lambda_n r_0) + n\pi I_1(\lambda_n r_0)}$$

$$= 8k B_i \theta_\infty r_0 \sum_{n=1}^{\infty} \frac{[(-1)^n - 1]}{n} \frac{I_1(\lambda_n r_0)}{B_i I_0(\lambda_n r_0) + n\pi I_1(\lambda_n r_0)}$$


PROB. 5.21: Formulation in terms of

$$\theta(r, z) = T(r, z) - T_0 \text{ is:}$$

$$\frac{\partial^2 \theta}{\partial r^2} + \frac{1}{r} \frac{\partial \theta}{\partial r} + \frac{\partial^2 \theta}{\partial z^2} + \frac{\dot{q}}{k} = 0$$

$$\theta(r_0, z) = \theta(r, 0) = \theta(r, L) = 0$$

$$\text{Let } \theta(r, z) = \psi(r, z) + \phi(z)$$



$$\frac{\partial^2 \psi}{\partial r^2} + \frac{1}{r} \frac{\partial \psi}{\partial r} + \frac{\partial^2 \psi}{\partial z^2} = 0$$

$$\psi(r_0, z) = -\phi(z)$$

$$\psi(r, 0) = \psi(r, L) = 0$$

$$\frac{\partial^2 \phi}{\partial z^2} + \frac{\dot{q}}{k} = 0$$

$$\phi(0) = 0$$

$$\phi(L) = 0$$

From

Eq. (5.108)

$$\phi(z) = \frac{\dot{q}L^2}{2k} \left[ \frac{z}{L} - \left( \frac{z}{L} \right)^2 \right]$$

$$\psi(r, z) = \frac{2}{L} \sum_{n=1}^{\infty} \frac{I_0\left(\frac{n\pi r}{L}\right)}{I_0\left(\frac{n\pi r_0}{L}\right)} \sin \frac{n\pi}{L} z \cdot \int_0^L \left\{ -\frac{\dot{q}L^2}{2k} \left[ \frac{z}{L} - \left( \frac{z}{L} \right)^2 \right] \right\} \sin \frac{n\pi z'}{L} dz'$$

$$= -\frac{2\dot{q}L^2}{\pi^3 k} \sum_{n=1}^{\infty} \frac{[1 - (-1)^n]}{n^3} \frac{I_0\left(\frac{n\pi r}{L}\right)}{I_0\left(\frac{n\pi r_0}{L}\right)} \sin \frac{n\pi}{L} z$$

Thus,

$$T(r, z) - T_0 = \frac{\dot{q}L^2}{2k} \left[ \frac{z}{L} - \left( \frac{z}{L} \right)^2 \right]$$

$$- \frac{2\dot{q}L^2}{\pi^3 k} \sum_{n=1}^{\infty} \frac{[1 - (-1)^n]}{n^3} \frac{I_0\left(\frac{n\pi r}{L}\right)}{I_0\left(\frac{n\pi r_0}{L}\right)} \sin \frac{n\pi}{L} z$$

PROB. 5.22: Formulation in terms of  $\theta(r,z) = T(r,z) - T_0$ :

$$\left\{ \begin{array}{l} \frac{\partial^2 \theta}{\partial r^2} + \frac{1}{r} \frac{\partial \theta}{\partial r} + \frac{\partial^2 \theta}{\partial z^2} + \frac{\dot{q}(r,z)}{k} = 0 \\ \theta(r_0, z) = \theta(r, 0) = \theta(r, L) = 0 \end{array} \right\}$$

Let

$$\theta(r,z) = \sum_{m=1}^{\infty} \sum_{n=1}^{\infty} A_{mn} J_0(\lambda_n r) \sin \frac{m\pi}{L} z$$

where  $\lambda_n$  are the positive roots of  $J_0(\lambda_n r_0) = 0$ .

Also,

$$\frac{\dot{q}(r,z)}{k} = \sum_{m=1}^{\infty} \sum_{n=1}^{\infty} B_{mn} J_0(\lambda_n r) \sin \frac{m\pi}{L} z$$

where

$$B_{mn} = \frac{2}{k L N_n} \int_0^{r_0} \int_0^L \dot{q}(r,z) \sin \frac{m\pi}{L} z J_0(\lambda_n r) r dr dz$$

with

$$N_n = \frac{r_0^2}{2} J_1^2(\lambda_n r_0)$$

Note that the expansion for  $\theta(r,z)$  satisfies all the boundary conditions. Now, substitute both expansions into the DE and obtain

$$A_{mn} = \frac{B_{mn}}{\lambda_n^2 + \left(\frac{m\pi}{L}\right)^2}$$

Thus,

---


$$T(r,z) - T_0 = \frac{4}{L r_0^2 k} \sum_{m=1}^{\infty} \sum_{n=1}^{\infty} \frac{J_0(\lambda_n r) \sin \frac{m\pi}{L} z}{J_1^2(\lambda_n r_0) [\lambda_n^2 + \left(\frac{m\pi}{L}\right)^2]} \int_0^L \int_0^{r_0} \dot{q}(r',z') \sin \frac{m\pi}{L} z' J_0(\lambda_n r') r' dr' dz'$$


---

PROB. 5.23: From Prob. 2.11, the formulation of the problem:

$$\frac{\partial^2 T}{\partial x^2} + \frac{\partial^2 T}{\partial y^2} = 0$$

$$\left. \frac{\partial T}{\partial x} \right|_{x=0} = 0, \quad -k \left. \frac{\partial T}{\partial x} \right|_{x=a} = q_1''$$

$$\left. \frac{\partial T}{\partial y} \right|_{y=0} = 0, \quad k \left. \frac{\partial T}{\partial y} \right|_{y=b} = q_2'' = \frac{b}{a} q_1''$$

Let  $T(x,y) = \psi(x,y) + \phi(x) + \Omega(y)$

$\frac{\partial^2 \psi}{\partial x^2} + \frac{\partial^2 \psi}{\partial y^2} = 0$ $\left. \frac{\partial \psi}{\partial x} \right _{x=0} = 0, -k \left. \frac{\partial \psi}{\partial x} \right _{x=a} = 0$ $\left. \frac{\partial \psi}{\partial y} \right _{y=0} = 0, k \left. \frac{\partial \psi}{\partial y} \right _{y=b} = 0$ $\downarrow$ By inspection $\psi(x,y) = \text{const.} = A$	$\frac{d^2 \phi}{dx^2} + \frac{d^2 \Omega}{dy^2} = 0$ $\frac{d^2 \phi}{dx^2} = -\frac{d^2 \Omega}{dy^2} = \text{const.} = C$ $\frac{d\phi(0)}{dx} = 0, -k \frac{d\phi(a)}{dx} = q_1''$ $\frac{d\Omega(0)}{dy} = 0, k \frac{d\Omega(b)}{dy} = \frac{b}{a} q_1''$ $\downarrow$ $\phi(x) = B - \frac{q_1''}{2ak} x^2$ $\Omega(y) = D + \frac{q_1''}{2ak} y^2$
---	--

$\therefore T(x,y) = \boxed{A + \frac{q_1''}{2ak} (y^2 - x^2)}$

Any arbitrary constant =  $A + B + D$ .

PROB. 5.24: a) Formulation:

$$\frac{\partial^2 T}{\partial x^2} + \frac{\partial^2 T}{\partial y^2} + \frac{\dot{q}}{k} = 0$$

$$\left. \frac{\partial T}{\partial x} \right|_{x=0} = 0; \quad -k \left. \frac{\partial T}{\partial x} \right|_{x=a} = q_1''$$

$$\left. \frac{\partial T}{\partial y} \right|_{y=0} = 0; \quad k \left. \frac{\partial T}{\partial y} \right|_{y=b} = q_2''$$

b) Under steady-state conditions,

$$q_2'' \times a + \dot{q} \times (a \times b) = q_1'' \times b \Rightarrow \underline{q_2' = b \times \left( \frac{1}{a} q_1'' - \dot{q} \right)}$$

c) Note that if  $T(x,y)$  is a solution, then  $T(x,y) + C$  is also a solution, where  $C$  is any constant. This can be proved by direct substitution (see Prob. 2.11).

d) Let  $T(x,y) = \psi(x,y) + \phi(x) + \Omega(y)$ . This leads to

$$\frac{\partial^2 \psi}{\partial x^2} + \frac{\partial^2 \psi}{\partial y^2} = 0$$

$$\left. \frac{\partial \psi}{\partial x} \right|_{x=0} = 0; \quad -k \left. \frac{\partial \psi}{\partial x} \right|_{x=a} = 0$$

$$\left. \frac{\partial \psi}{\partial y} \right|_{y=0} = 0; \quad k \left. \frac{\partial \psi}{\partial y} \right|_{y=b} = 0$$

By inspection

$$\psi(x,y) = A = \text{constant}$$

and

$$\frac{d^2 \phi}{dx^2} = -\frac{d^2 \Omega}{dy^2} - \frac{\dot{q}}{k} = \text{const.} = C$$

$$\frac{d \phi(0)}{dx} = 0; \quad -k \frac{d \phi(a)}{dx} = q_1''$$

$$\frac{d \Omega(0)}{dy} = 0; \quad k \frac{d \Omega(b)}{dy} = q_2''$$

↓

$$\phi(x) = B - \frac{q_1''}{2ak} x^2$$

$$\Omega(y) = D + \frac{q_2''}{2bk} y^2$$

Thus,

$$\underline{T(x,y) = a + \frac{1}{2k} \left( \frac{q_2''}{b} y^2 - \frac{q_1''}{a} x^2 \right)}$$

A + B + D ← Arbitrary constant


PROB. 5.25: Formulation of  
the problem:

$$\frac{\partial^2 T}{\partial r^2} + \frac{1}{r} \frac{\partial T}{\partial r} + \frac{\partial^2 T}{\partial z^2} = 0$$

$$T(0, z) = \text{finite}; -k \frac{\partial T}{\partial r} \Big|_{r=r_0} = q''_{f_2}$$

$$\frac{\partial T}{\partial z} \Big|_{z=0} = 0$$

$$-k \frac{\partial T}{\partial z} \Big|_{z=H} = \begin{cases} -q''_{f_1}, & 0 < r < r_1 \\ 0, & r_1 < r < r_0 \end{cases}$$



Note that  $q''_{f_1} \pi r_1^2 = q''_{f_2} 2\pi r_0 H \Rightarrow q''_{f_1} = \frac{2r_0 H}{r_1^2} q''_{f_2}$  *{This condition has to be satisfied to have the steady-state condition.}*

a) If  $T$  is a solution of the problem, then

$T+C$  will also be a solution, where  $C$  is any constant. This can be proved by direct substitution into the formulation.

b)

$$\text{Let } T(r, z) = \psi(r, z) + \phi(r) + \Omega(z)$$

$$\frac{\partial^2 \psi}{\partial r^2} + \frac{1}{r} \frac{\partial \psi}{\partial r} + \frac{\partial^2 \psi}{\partial z^2} = 0$$

$$\psi(0, z) = \text{finite}; \quad \frac{\partial \psi}{\partial r} \Big|_{r=r_0} = 0$$

$$\frac{\partial \psi}{\partial z} \Big|_{z=0} = 0$$




$$\frac{d^2 \phi}{dr^2} + \frac{1}{r} \frac{d \phi}{dr} = -\frac{d^2 \Omega}{dz^2} = \text{const.} = -\frac{q''}{k}$$

$$\phi(0) = \text{finite}; \quad -k \frac{d \phi(r_0)}{dr} = q''_{f_2}$$

$$\frac{d \Omega(H)}{dz} = 0$$

$$-k \frac{\partial T}{\partial z} \Big|_{z=H} = \begin{cases} -q''_{f_2} + q''_{f_3}, & 0 < r < r_1 \\ q''_{f_3}, & r_1 < r < r_0 \end{cases} \quad -k \frac{d \Omega(H)}{dz} = -q''_{f_3}$$

← see next page ↑



$\Downarrow$

$$q''_{f3} \pi r_0^2 = \dot{q} \pi r_0^2 H$$

$$q''_{f3} = H \dot{q} = \frac{2H}{r_0} q''$$

$\Downarrow$

$$\dot{q} \pi r_0^2 H = q''_{f2} 2\pi r_0 H$$

$$\dot{q} = \frac{2}{r_0} q''_{f2}$$

$\Downarrow$

$$\dot{q} = \frac{2}{r_0} q''_{f2}$$

check

$\Downarrow$

$$q''_{f3} = \frac{2H}{r_0} \frac{r_1^2}{2\pi r_0 H} q''$$

$$q''_{f3} = \left(\frac{r_1}{r_0}\right)^2 q''$$

$$\begin{aligned} \pi r_1^2 (q''_{f1} - q''_{f3}) &= q''_{f3} \pi (r_0^2 - r_1^2) \\ r_1^2 \left[ \left(\frac{r_0}{r_1}\right)^2 - 1 \right] q''_{f3} &= q''_{f3} (r_0^2 - r_1^2) \end{aligned}$$

$$\therefore r_0^2 - r_1^2 = r_0^2 - r_1^2 \quad \checkmark$$

Solution to  $\Omega(z)$  problem:

$$\Omega(z) = \frac{q_i''}{2kH} \left(\frac{r_i}{r_0}\right)^2 z^2 + A$$

↑ Any arbitrary constant

Solution to  $\phi(r)$  problem:

$$\phi(r) = -\frac{q_i''}{4kH} \left(\frac{r_i}{r_0}\right)^2 r^2 + B$$

↑ Any arbitrary constant

Solution to  $\psi(r,z)$  problem:

$$\text{Let } \psi(r,z) = R(r) \cdot Z(z)$$



$$r^2 \frac{d^2 R}{dr^2} + r \frac{dR}{dr} + \lambda^2 r^2 R = 0$$

$$R(0) = \text{finite}$$

$$\frac{dR(0)}{dr} = 0$$



$$R_n(r) = \begin{cases} C_0 \\ C_n J_0(\lambda_n r), n=1,2,3,\dots \end{cases}$$

$$\frac{d^2 Z_n}{dz^2} - \lambda_n^2 Z_n = 0$$

$$\frac{dZ_n(0)}{dz} = 0$$



$$Z_n(z) = \begin{cases} D_0 \\ D_n \cosh \lambda_n z \end{cases}$$

where  $\lambda_n$  are the roots of  $J_0(\lambda_n r_0) = 0$

$$\therefore \psi(r,z) = b_0 + \sum_{n=1}^{\infty} a_n J_0(\lambda_n r) \cosh \lambda_n z$$

and

$$b_0 + A + B$$

$$T(r,z) = a_0 + \frac{q_i''}{2kH} \left(\frac{r_i}{r_0}\right)^2 \left(z^2 - \frac{1}{2} r^2\right) + \sum_{n=1}^{\infty} a_n J_0(\lambda_n r) \cosh \lambda_n z$$

$$\begin{aligned}
 k \frac{\partial T}{\partial z} \Big|_{z=H} &= \underbrace{q''_1 \left( \frac{r_1}{r_0} \right)^2}_{C_0} + \sum_{n=1}^{\infty} \underbrace{(k a_n \lambda_n \sinh \lambda_n H)}_{C_n} J_0(\lambda_n r) \\
 &= C_0 + \sum_{n=1}^{\infty} C_n J_0(\lambda_n r) \quad \leftarrow \text{Valid expansion in } (0, r_0)
 \end{aligned}$$

$$C_0 = \frac{2}{r_0^2} \int_0^{r_0} k \frac{\partial T}{\partial z} \Big|_{z=H} \cdot r \cdot dr = \frac{2}{r_0^2} \int_0^{r_1} q''_1 r dr = \left( \frac{r_1}{r_0} \right)^2 q''_1$$

$$C_n = \frac{1}{N_n} \int_0^{r_0} k \frac{\partial T}{\partial z} \Big|_{z=H} J_0(\lambda_n r) \cdot r \cdot dr$$

Where  $N_n = \frac{r_0^2}{2} J_0^2(\lambda_n r_0)$   $\leftarrow$  From Table 4.2

$$\begin{aligned}
 &= \frac{q''_1}{N_n} \int_0^{r_1} J_0(\lambda_n r) r dr = \frac{q''_1}{N_n} \frac{1}{\lambda_n} \int_0^{r_1} \frac{d}{dr} [r J_1(\lambda_n r)] dr \\
 &= \frac{2}{r_0^2} q''_1 \frac{r_1 J_1(\lambda_n r_1)}{\lambda_n J_0^2(\lambda_n r_0)}
 \end{aligned}$$

$$\begin{aligned}
 \therefore T(r, z) &= A_0 + \frac{q''_1}{2kH} \left( \frac{r_1}{r_0} \right)^2 \left\{ z^2 - \frac{1}{2} r^2 \right. \\
 &\quad \left. + 4 \frac{H}{r_1} \sum_{n=1}^{\infty} \frac{J_0(\lambda_n r) J_1(\lambda_n r_1) \cosh \lambda_n z}{\lambda_n^2 J_0^2(\lambda_n r_0) \sinh \lambda_n H} \right\}
 \end{aligned}$$

Any arbitrary constant.

PROB. 5.26: a) Formulation in terms of  $\theta(r, z) = T(r, z) - T_1$ :

$$\left\{ \begin{array}{l} \frac{\partial^2 \theta}{\partial r^2} + \frac{1}{r} \frac{\partial \theta}{\partial r} + \frac{\partial^2 \theta}{\partial z^2} = 0 \\ \theta(0, z) = \text{finite}; \quad \theta(r_0, z) = 0 \\ \theta(r, 0) = f(r) - T_1; \quad \theta(r, \infty) = 0 \end{array} \right\}$$

Let  $\theta(r, z) = R(r) \cdot Z(z)$ . Then,

$$\theta(r, z) = \sum_{n=1}^{\infty} c_n e^{\lambda_n z} J_0(\lambda_n r), \quad \text{where } \lambda_n \text{ are the positive roots of } J_0(\lambda_n r_0) = 0$$

$$f(r) - T_1 = \sum_{n=1}^{\infty} c_n J_0(\lambda_n r)$$

$$c_n = \frac{1}{N_n} \int_0^{r_0} [f(r) - T_1] J_0(\lambda_n r) r dr, \quad N_n = \frac{r_0^2}{2} J_1^2(\lambda_n r_0)$$

$$\therefore T(r, z) - T_1 = \frac{2}{r_0^2} \sum_{n=1}^{\infty} \frac{J_0(\lambda_n r)}{J_1^2(\lambda_n r_0)} e^{\lambda_n z} \int_0^{r_0} [f(r') - T_1] J_0(\lambda_n r') r' dr'$$

b) If  $f(r) = T_0 = \text{const.}$ , then

$$\int_0^{r_0} [f(r') - T_1] J_0(\lambda_n r') r' dr' = \dots = \frac{T_0 - T_1}{\lambda_n} r_0 J_1(\lambda_n r_0)$$

$$\therefore \frac{T(r, z) - T_1}{T_0 - T_1} = \frac{2}{r_0} \sum_{n=1}^{\infty} \frac{J_0(\lambda_n r)}{\lambda_n J_1(\lambda_n r_0)} e^{-\lambda_n z}$$

c)

$$q_{T_2}(0) = \int_0^{r_0} \left\{ -k 2\pi r \left( \frac{\partial T}{\partial z} \right)_{z=0} dr \right\} = \dots = 4\pi k (T_0 - T_1) \sum_{n=1}^{\infty} \frac{1}{\lambda_n}$$

PROB. 5.27: a) Formulation in terms of  $\theta(r, z) = T(r, z) - T_{\infty}$ :

$$\frac{\partial^2 \theta}{\partial r^2} + \frac{1}{r} \frac{\partial \theta}{\partial r} + \frac{\partial^2 \theta}{\partial z^2} = 0$$

$$\theta(0, z) = \text{finite}; \quad -k \frac{\partial \theta}{\partial r} \Big|_{r=r_0} = h \theta(r_0, z)$$

$$\theta(r, 0) = T_0 - T_{\infty}; \quad \theta(r, \infty) = 0$$

$$\text{Let } \theta(r, z) = R(r) Z(z) \Rightarrow \theta(r, z) = \sum_{n=1}^{\infty} a_n J_0(\lambda_n r) e^{-\lambda_n z}$$

where  $\lambda_n$  are the positive zeros of

$$k \lambda_n J_1(\lambda_n r_0) = h J_0(\lambda_n r_0) \Rightarrow \frac{J_1(\lambda_n r_0)}{J_0(\lambda_n r_0)} = \frac{Bi}{\lambda_n r_0}, \quad Bi = \frac{h r_0}{k}$$

Now,

$$\theta(r, 0) = T_0 - T_{\infty} \Rightarrow T_0 - T_{\infty} = \sum_{n=1}^{\infty} a_n J_0(\lambda_n r)$$

Thus,

$$a_n = \frac{1}{N_n} \int_0^{r_0} (T_0 - T_{\infty}) J_0(\lambda_n r) r dr = \frac{T_0 - T_{\infty}}{N_n} \frac{r_0}{\lambda_n} J_1(\lambda_n r_0)$$

where (from Table 4.2),

$$N_n = \frac{r_0^2}{2} \left[ 1 + \frac{1}{\lambda_n^2} \left( \frac{h}{k} \right)^2 \right] J_0^2(\lambda_n r_0) = \frac{r_0^2}{2} \left[ J_0^2(\lambda_n r_0) + J_1^2(\lambda_n r_0) \right]$$

$$\therefore \theta(r, z) = T(r, z) - T_{\infty} = \frac{2}{r_0} (T_0 - T_{\infty}) \sum_{n=1}^{\infty} \frac{J_1(\lambda_n r_0) J_0(\lambda_n r)}{\lambda_n [J_0^2(\lambda_n r_0) + J_1^2(\lambda_n r_0)]} e^{-\lambda_n z}$$

b) The rate of heat loss from the rod,

$$q_{\text{loss}} = \int_0^{r_0} \left\{ -k 2\pi r dr \frac{\partial T(r, 0)}{\partial z} \right\} = \dots = 4\pi k (T_0 - T_{\infty}) \sum_{n=1}^{\infty} \frac{J_1^2(\lambda_n r_0)}{\lambda_n [J_0^2(\lambda_n r_0) + J_1^2(\lambda_n r_0)]}$$


PROB. 5.28: Formulation of the problem

in terms of  $\theta = T - T_i$ :

$$\frac{\partial^2 \theta}{\partial r^2} + \frac{1}{r} \frac{\partial \theta}{\partial r} + \frac{\partial^2 \theta}{\partial z^2} = 0$$

$$\theta(0, z) \neq \infty; \quad \theta(r_0, z) = 0$$

$$-k \frac{\partial \theta}{\partial z} \Big|_{z=0} = q''_w; \quad \theta(r, \infty) = 0$$



$$\text{Let } \theta(r, z) = R(r) \cdot Z(z)$$

$$\therefore \theta(r, z) = \sum_{n=1}^{\infty} C_n e^{-\lambda_n z} J_0(\lambda_n r)$$

where  $\lambda_n$ 's are the positive roots of  $J_0(\lambda_n r_0) = 0$ .

$$\text{B.C. at } z=0 \Rightarrow -\frac{q''_w}{k} = -\sum_{n=1}^{\infty} C_n \lambda_n J_0(\lambda_n r_0)$$

Thus,

$$\begin{aligned} C_n \lambda_n &= \frac{1}{N_n} \int_0^{r_0} \frac{q''_w}{k} J_0(\lambda_n r) r dr, \quad N_n = \frac{r_0^2}{2} J_1^2(\lambda_n r_0) \\ &= \frac{q''_w}{k} \frac{2}{\lambda_n r_0} \frac{1}{J_1(\lambda_n r_0)} \end{aligned}$$

$$\therefore T(r, z) - T_i = \frac{2q''_w}{kr_0} \sum_{n=1}^{\infty} \frac{1}{\lambda_n^2} \frac{J_0(\lambda_n r)}{J_1(\lambda_n r_0)} e^{-\lambda_n z}$$


PROB. 5.29: Formulation in terms of

$$\theta(r, z) = T(r, z) - T_{\infty} :$$

$$\frac{\partial^2 \theta}{\partial r^2} + \frac{1}{r} \frac{\partial \theta}{\partial r} + \frac{\partial^2 \theta}{\partial z^2} = 0$$

$$\theta(0, z) \neq \infty ; -k \left. \frac{\partial \theta}{\partial r} \right|_{r=r_0} = h \theta(r_0, z)$$

$$-k \left. \frac{\partial \theta}{\partial z} \right|_{z=0} = q''_{tw} ; \quad \theta(r, \infty) = 0$$



$$\text{Let } \theta(r, z) = R(r) \cdot Z(z)$$

$$\therefore \theta(r, z) = \sum_{n=1}^{\infty} C_n J_0(\lambda_n r) e^{-\lambda_n z}$$

where  $\lambda_n$  are the positive zeros of

$$k \lambda_n J_1(\lambda_n r_0) = h J_0(\lambda_n r_0) \Rightarrow \frac{J_1(\lambda_n r_0)}{J_0(\lambda_n r_0)} = \frac{Bi}{\lambda_n r_0}, \quad Bi = \frac{h r_0}{k}$$

$$\therefore -\frac{q''_{tw}}{k} = -\sum_{n=1}^{\infty} C_n \lambda_n J_0(\lambda_n r_0) \quad \leftarrow \text{From B.C. at } z=0.$$

Thus,

$$\begin{aligned} C_n \lambda_n &= \frac{1}{N_n} \int_0^{r_0} \frac{q''_{tw}}{k} J_0(\lambda_n r) r dr, \\ &= \frac{1}{N_n} \frac{q''_{tw}}{k} \frac{r_0}{\lambda_1} J_1(\lambda_1 r_0) \end{aligned}$$

where, from Table 4.2,


$$N_n = \frac{r_0^2}{2} \left[ + + \frac{1}{\lambda_n^2} \left( \frac{h}{k} \right)^2 \right] J_0^2(\lambda_n r_0) = \frac{r_0^2}{2} \left[ J_0^2(\lambda_n r_0) + J_1^2(\lambda_n r_0) \right]$$

---


$$\therefore \theta(r, z) = T(r, z) - T_{\infty} = \frac{2q''_{tw}}{kr_0} \sum_{n=1}^{\infty} \frac{J_1(\lambda_n r_0) J_0(\lambda_n r)}{\lambda_n^2 [J_0^2(\lambda_n r_0) + J_1^2(\lambda_n r_0)]} e^{-\lambda_n z}$$


---

PROB. 5.30:



Formulation in terms of  $\theta(x,y) = T(x,y) - T_0$ :

$$\frac{\partial^2 \theta}{\partial x^2} + \frac{\partial^2 \theta}{\partial y^2} = 0$$

$$\begin{aligned}\theta(0,y) &= 0; & \theta(L,y) &= (T_1 - T_0) e^{-y/H} \\ \theta(x,0) &= T_1 - T_0; & \lim_{y \rightarrow \infty} \theta(x,y) &= 0\end{aligned}$$

$$\text{Let } \theta(x,y) = \psi(x,y) + e^{-y/H} \phi(x)$$

$$\frac{\partial^2 \psi}{\partial x^2} + \frac{\partial^2 \psi}{\partial y^2} = 0 \quad \leftarrow \quad \frac{\partial^2 \phi}{\partial x^2} + \phi(x) = 0 \quad \rightarrow$$

$$\psi(0,y) = 0; \psi(L,y) = 0$$

$$\psi(x,0) = (T_1 - T_0) - \phi(x)$$

$$\lim_{y \rightarrow \infty} \psi(x,y) = 0$$

$$\Downarrow \psi(x,y) = X(x)Y(y)$$

$$\phi(0) = 0$$

$$\phi(L) = T_1 - T_0$$

$$\Downarrow \underline{\underline{\phi(x) = (T_1 - T_0) \frac{\sin x}{\sin L}}}$$

$$\left\{ \begin{array}{l} \frac{d^2 X}{dx^2} + \lambda_n^2 X = 0 \\ X(0) = 0 \\ X(L) = 0 \end{array} \right\} \Rightarrow X_n(x) = A_n \sin \lambda_n x$$

$$\lambda_n = \frac{n\pi}{L}, n = 1, 2, 3, \dots$$

and

$$\left\{ \begin{array}{l} \frac{d^2 Y_n}{dy^2} - \lambda_n^2 Y_n = 0 \\ Y_n(\infty) = 0 \end{array} \right\} \Rightarrow Y_n(y) = B_n e^{-\lambda_n y}$$

$$\therefore \underline{\underline{\psi(x,y) = \sum_{n=1}^{\infty} a_n \sin \lambda_n x e^{-\lambda_n y}}}$$

$$\psi(x,0) = (T_1 - T_0) \left[ 1 - \frac{\sin x}{\sin L} \right] = \sum_{n=1}^{\infty} a_n \sin \lambda_n x$$

$$a_n = \frac{2}{L} (T_1 - T_0) \int_0^L \left[ 1 - \frac{\sin x}{\sin L} \right] \sin \lambda_n x dx \\ = 2 (T_1 - T_0) \left\{ \frac{1 - (-1)^n}{n\pi} - \frac{(-1)^n n\pi}{L^2 - (n\pi)^2} \right\}$$

Thus,

$$\frac{T(x,y)-T_0}{T_1 - T_0} = e^{-y} \frac{\sin x}{\sin L} + 2 \sum_{n=1}^{\infty} \left\{ \frac{1 - (-1)^n}{n\pi} - \frac{(-1)^n n\pi}{L^2 - (n\pi)^2} \right\} \sin \frac{n\pi}{L} x e^{-\frac{n\pi}{L} y}$$


PROB. 5.31: Formulation in terms of

$$\Theta(r,z) = T(r,z) - T_0 :$$

$$\frac{\partial^2 \Theta}{\partial r^2} + \frac{1}{r} \frac{\partial \Theta}{\partial r} + \frac{\partial^2 \Theta}{\partial z^2} = 0$$

$$\Theta(0,z) = \text{finite}; \quad \Theta(r_0, z) = (T_1 - T_0) e^z$$

$$\Theta(r,0) = (T_1 - T_0); \quad \lim_{z \rightarrow \infty} \Theta(r,z) = 0$$



$$\text{Let } \Theta(r,z) = \psi(r,z) + e^z \phi(r)$$

$$\frac{\partial^2 \psi}{\partial r^2} + \frac{1}{r} \frac{\partial \psi}{\partial r} + \frac{\partial^2 \psi}{\partial z^2} = 0$$

$$\frac{d^2 \phi}{dr^2} + \frac{1}{r} \frac{d\phi}{dr} + \phi(r) = 0$$

$$\psi(0,z) = \text{finite}; \quad \psi(r_0, z) = 0$$

$$\phi(0) = \text{finite}$$

$$\psi(r,0) = (T_1 - T_0) - \phi(r)$$

$$\phi(r_0) = T_1 - T_0$$

$$\lim_{z \rightarrow \infty} \psi(r,z) = 0$$

$$\frac{\phi(r)}{(T_1 - T_0) \frac{J_0(r)}{J_0(r_0)}} =$$

If we let  $\psi(r, z) = R(r) \cdot Z(z)$ , then

$$\left\{ \begin{array}{l} \frac{d^2 R}{dr^2} + \frac{1}{r} \frac{dR}{dr} + \lambda^2 R = 0 \\ R(0) = \text{finite} \\ R(r_0) = 0 \end{array} \right\} \Rightarrow \begin{aligned} R_n(r) &= A_n J_0(\lambda_n r) \\ \text{where } \lambda_n &\text{'s are the positive} \\ \text{zeros of } J_0(\lambda_n r_0) &= 0 \end{aligned}$$

and

$$\left\{ \begin{array}{l} \frac{d^2 Z_n}{dz^2} - \lambda_n^2 Z_n = 0 \\ Z_n(\infty) = 0 \end{array} \right\} \Rightarrow Z_n(z) = B_n e^{-\lambda_n z}$$

Thus,

$$\psi(r, z) = \sum_{n=1}^{\infty} a_n e^{\lambda_n z} J_0(\lambda_n r)$$

$$\Downarrow$$

$$\psi(r, 0) = (T_1 - T_0) \left[ 1 - \frac{J_0(r)}{J_0(r_0)} \right] = \sum_{n=1}^{\infty} a_n J_0(\lambda_n r)$$

$$a_n = \frac{1}{N_n} (T_1 - T_0) \int_0^{r_0} \left[ 1 - \frac{J_0(r)}{J_0(r_0)} \right] J_0(\lambda_n r) r dr$$

$$\therefore N_n = \frac{r_0^2}{2} J_1^2(\lambda_n r_0)$$

$$= \frac{2}{r_0} \frac{T_1 - T_0}{J_1(\lambda_n r_0)} \frac{1}{\lambda_n (1 - \lambda_n^2)}$$

$$\therefore \frac{T(r, z) - T_0}{T_1 - T_0} = e^z \frac{J_0(r)}{J_0(r_0)} + \frac{2}{r_0} \sum_{n=1}^{\infty} \frac{1}{\lambda_n (1 - \lambda_n^2)} \frac{J_0(\lambda_n r)}{J_1(\lambda_n r_0)} e^{-\lambda_n z}$$

PROB. 5.32: Formulation in terms

of  $\Theta(r, z) = T(r, z) - T_0$ :

$$\frac{\partial^2 \Theta}{\partial r^2} + \frac{1}{r} \frac{\partial \Theta}{\partial r} + \frac{\partial^2 \Theta}{\partial z^2} + \frac{\dot{q}}{k} = 0$$

$$\left. \frac{\partial \Theta}{\partial r} \right|_{r=r_1} = 0; \quad \Theta(r_2, z) = 0$$

$$\Theta(r, 0) = 0; \quad \Theta(r, L) = 0$$

Let

$$\Theta(r, z) = \psi(r, z) + \phi(z)$$

$$\frac{\partial^2 \psi}{\partial r^2} + \frac{1}{r} \frac{\partial \psi}{\partial r} + \frac{\partial^2 \psi}{\partial z^2} = 0$$

$$\left. \frac{\partial \psi}{\partial r} \right|_{r=r_1} = 0, \quad \psi(r_2, z) = -\phi(z)$$


$$\psi(r, 0) = 0, \quad \psi(r, L) = 0$$

$$\Downarrow \psi(r, z) = R(r)Z(z)$$

$$\left\{ \begin{array}{l} \frac{d^2 Z}{dz^2} + \lambda^2 Z = 0 \\ Z(0) = 0 \\ Z(L) = 0 \end{array} \right. \Rightarrow$$

$$Z_n(z) = A_n \sin \lambda_n z$$

$$\lambda_n = \frac{n\pi}{L}, \quad n=1, 2, 3, \dots$$



$$\frac{d^2 \phi}{dz^2} + \frac{\dot{q}}{k} = 0$$

$$\phi(0) = 0$$

$$\phi(L) = 0$$

---


$$\phi(z) = \frac{\dot{q}L^2}{2k} \left[ \frac{z}{L} - \left( \frac{z}{L} \right)^2 \right]$$

Corresponding to each  $\lambda_n$ ,

$$r^2 \frac{d^2 R_n}{dr^2} + r \frac{dR_n}{dr} - \lambda_n^2 r^2 R_n = 0$$

$$\therefore R_n(r) = B_n I_0(\lambda_n r) + C_n K_0(\lambda_n r)$$

$$\frac{\partial \psi}{\partial r} \Big|_{r=r_1} = 0 \rightarrow \frac{dR_n}{dr} \Big|_{r=r_1} = 0 \rightarrow \underbrace{B_n \lambda_n I_1(\lambda_n r_1) - C_n \lambda_n K_1(\lambda_n r_1)}_{C_n = B_n \frac{I_1(\lambda_n r_1)}{K_1(\lambda_n r_1)}} = 0$$

$$C_n = B_n \frac{I_1(\lambda_n r_1)}{K_1(\lambda_n r_1)}$$

$$\therefore R_n(r) = \frac{B_n}{K_1(\lambda_n r_1)} [I_0(\lambda_n r) K_1(\lambda_n r_1) + I_1(\lambda_n r_1) K_0(\lambda_n r)]$$

Hence, the solution for  $\psi(r, z)$  is

$$\psi(r, z) = \sum_{n=1}^{\infty} a_n [I_0(\lambda_n r) K_1(\lambda_n r_1) + I_1(\lambda_n r_1) K_0(\lambda_n r)] \sin \lambda_n z$$

$$\therefore -\dot{\phi}(z) = -\frac{\dot{q}L^2}{2k} \left[ \frac{z}{L} - \left( \frac{z}{L} \right)^2 \right] = \sum_{n=1}^{\infty} b_n \sin \lambda_n z$$

where

$$b_n = a_n [I_0(\lambda_n r_2) K_1(\lambda_n r_1) + I_1(\lambda_n r_1) K_0(\lambda_n r_2)] = \frac{2}{L} \int_0^L [-\dot{\phi}(z)] \sin \lambda_n z dz$$

$$= -\frac{\dot{q}L}{k} \int_0^L \left[ \frac{z}{L} - \left( \frac{z}{L} \right)^2 \right] \sin \lambda_n z dz = -\frac{2\dot{q}L^2}{k} \cdot \frac{1 - (-1)^n}{(n\pi)^3}$$

$$\therefore \psi(r, z) = -\frac{2\dot{q}L^2}{k\pi^3} \sum_{n=1}^{\infty} \frac{[1 - (-1)^n]}{n^3} \frac{[I_0(\lambda_n r) K_1(\lambda_n r_1) + I_1(\lambda_n r_1) K_0(\lambda_n r)]}{[I_0(\lambda_n r_2) K_1(\lambda_n r_1) + I_1(\lambda_n r_1) K_0(\lambda_n r_2)]} \sin \lambda_n z$$

Thus,

---



$$T(r, z) - T_0 = \frac{\dot{q}L^2}{2k} \left[ \frac{z}{L} - \left( \frac{z}{L} \right)^2 \right]$$

$$-\frac{2\dot{q}L^2}{k\pi^3} \sum_{n=1}^{\infty} \frac{[1 - (-1)^n]}{n^3} \frac{[I_0(\lambda_n r) K_1(\lambda_n r_1) + I_1(\lambda_n r_1) K_0(\lambda_n r)]}{[I_0(\lambda_n r_2) K_1(\lambda_n r_1) + I_1(\lambda_n r_1) K_0(\lambda_n r_2)]} \sin \lambda_n z$$


---

PRDB. 5.33:

- Because of symmetry, consider only one half of the rod.
- Assume  $k = \text{constant}$ .



Formulation in terms of  $\theta(r, z) = T(r, z) - T_\infty$ :

$$\left. \begin{aligned} & \frac{\partial^2 \theta}{\partial r^2} + \frac{1}{r} \frac{\partial \theta}{\partial r} + \frac{\partial^2 \theta}{\partial z^2} + \frac{\dot{Q}(z)}{k} = 0 \\ & \theta(0, z) = \text{finite}; \quad -k \frac{\partial \theta}{\partial r} \Big|_{r=r_0} = h \theta(r_0, z) \\ & \frac{\partial \theta}{\partial z} \Big|_{z=0} = \frac{\partial \theta}{\partial z} \Big|_{z=L} = 0 \end{aligned} \right\} \quad \begin{aligned} & \text{where} \\ & \dot{Q}(z) = \begin{cases} \dot{q}, & 0 < z < \frac{L}{2} \\ 0, & \frac{L}{2} < z < L \end{cases} \end{aligned}$$

Expand both  $\theta(r, z)$  and  $\dot{Q}(z)/k$  into Fourier series as follows:

$$\theta(r, z) = \sum_{n=0}^{\infty} A_n(r) \cos \lambda_n z = A_0(r) + \sum_{n=1}^{\infty} A_n(r) \cos \lambda_n z, \quad 0 < z < L$$

and

$$\frac{\dot{Q}(z)}{k} = \sum_{n=0}^{\infty} B_n \cos \lambda_n z = B_0 + \sum_{n=1}^{\infty} B_n \cos \lambda_n z, \quad 0 < z < L$$

where  $\lambda_n = \frac{n\pi}{L}$ ,  $n=0, 1, 2, \dots$ , and  $\cos \lambda_n z$  are the eigenvalues and eigenfunctions of

$$\left. \begin{aligned} & \frac{d^2 z}{dz^2} + \lambda_n^2 z = 0 \\ & \frac{dz(0)}{dz} = \frac{dz(L)}{dz} = 0 \end{aligned} \right\}$$

Therefore,

$$B_0 = \frac{1}{L} \int_0^L \frac{\dot{Q}(z)}{k} dz = \frac{\dot{q}}{Lk} \int_0^{L/2} dz = \frac{\dot{q}}{2k}$$

$$B_n = \frac{2}{L} \int_0^L \frac{\dot{Q}(z)}{k} \cos \lambda_n z dz = \frac{2\dot{q}}{Lk} \int_0^{L/2} \cos \lambda_n z dz = \frac{2\dot{q}}{Lk\lambda_n} \sin \lambda_n \frac{L}{2}$$

Here we note that the expansion for  $\theta(r, z)$  satisfies the two

boundary conditions in  $z$ -direction. Now, substitute both expansions into the heat conduction equation:

$$\frac{1}{r} \frac{\partial}{\partial r} \left[ r \frac{\partial}{\partial r} \sum_{n=0}^{\infty} A_n(r) \cos \lambda_n z \right] + \frac{\partial^2}{\partial z^2} \sum_{n=0}^{\infty} A_n(r) \cos \lambda_n z + \sum_{n=0}^{\infty} B_n \cos \lambda_n z = 0$$

$$\sum_{n=0}^{\infty} \left[ \frac{1}{r} \frac{d}{dr} \left( r \frac{dA_n}{dr} \right) - \lambda_n^2 A_n(r) + B_n \right] \cos \lambda_n z = 0$$

Thus,

$$\frac{1}{r} \frac{d}{dr} \left( r \frac{dA_0}{dr} \right) = -B_0$$

$$\frac{1}{r} \frac{d}{dr} \left( r \frac{dA_n}{dr} \right) - \lambda_n^2 A_n(r) = -B_n, \quad n=1,2,3,\dots$$

Also, the boundary conditions in  $r$ -direction yield:

$$\Theta(0, z) = \text{finite} \Rightarrow \sum_{n=0}^{\infty} A_n(0) \cos \lambda_n z = \text{finite}$$

$$\therefore A_n(0) = \frac{1}{N_n} \int_0^L (\text{finite}) \cos \lambda_n z dz = \text{finite}$$

and

$$-k \left( \frac{\partial \theta}{\partial r} \right)_{r=r_0} = h \theta(r_0, z) \Rightarrow \sum_{n=0}^{\infty} \left[ k \frac{dA_n(r_0)}{dr} + h A_n(r_0) \right] \cos \lambda_n z = 0$$

$$\therefore k \frac{dA_n(r_0)}{dr} + h A_n(r_0) = 0, \quad n=0,1,2,\dots$$

Thus, when  $n=0$ ,

$$\left. \begin{aligned} \frac{1}{r} \frac{d}{dr} \left( r \frac{dA_0}{dr} \right) &= -B_0 \\ A_0(0) &= \text{finite} \\ k \frac{dA_0(r_0)}{dr} + h A_0(r_0) &= 0 \end{aligned} \right\} \Rightarrow A_0(r) = \frac{B_0 r_0^2}{4} \left[ 1 + \frac{2}{Bl} - \left( \frac{r}{r_0} \right)^2 \right]$$

where  $Bl = \frac{hr_0}{k}$

When  $n=1, 2, 3, \dots$ ,

$$\frac{1}{r} \frac{d}{dr} \left( r \frac{dA_n}{dr} \right) - \lambda_n^2 A_n(r) = -B_n$$

$$\therefore A_n(r) = C_n I_0(\lambda_n r) + D_n K_0(\lambda_n r) + \frac{B_n}{\lambda_n^2}$$

$$A_n(0) = \text{finite} \Rightarrow D_n = 0$$

$$k \frac{A_n(r_0)}{dr} + h A_n(r_0) = 0 \Rightarrow k \lambda_n C_n I_1(\lambda_n r_0) + h \left[ C_n I_0(\lambda_n r_0) + \frac{B_n}{\lambda_n^2} \right] = 0$$

$$\therefore C_n = - \frac{1}{\lambda_n^2} \frac{h B_n}{k \lambda_n I_1(\lambda_n r_0) + h I_0(\lambda_n r_0)}$$

Thus,

$$A_n(r) = \frac{B_n}{\lambda_n^2} \left[ 1 - \frac{h}{k \lambda_n I_1(\lambda_n r_0) + h I_0(\lambda_n r_0)} \right], n=1, 2, 3, \dots$$


and the solution is given by

$$\Theta(r, z) = T(r, z) - T_{\infty} = \frac{\dot{q} r_0^2}{8k} \left[ 1 + \frac{2k}{hr_0} - \left( \frac{r}{r_0} \right)^2 \right]$$

$$+ \frac{2\dot{q}}{Lk} \sum_{n=1}^{\infty} \frac{\sin \lambda_n L}{\lambda_n^3} \left[ 1 - \frac{h}{k \lambda_n I_1(\lambda_n r_0) + h I_0(\lambda_n r_0)} \right] \cos \lambda_n z$$

PROB. 5.34:

- Because of symmetry, consider only one half of the system.
- Since the sleeve is of negligible thickness, its temperature at any  $z$  will be the same as the shaft's surface temperature.
- Assume constant thermal conductivity.



Formulation in terms of  $\theta(r, z) = T(r, z) - T_{\infty}$ :

$$\frac{\partial^2 \theta}{\partial r^2} + \frac{1}{r} \frac{\partial \theta}{\partial r} + \frac{\partial^2 \theta}{\partial z^2} = 0$$

$$\left. \frac{\partial \theta}{\partial z} \right|_{z=0} = \left. \frac{\partial \theta}{\partial z} \right|_{z=L} = 0$$

$$\theta(0, z) = \text{finite}; \quad k \left. \frac{\partial \theta}{\partial r} \right|_{r=r_0} = q''_{\text{fw}}(z) = \begin{cases} \mu P \omega r_0 - h_1 \theta(r_0, z), & 0 < z < L/2 \\ -h_2 \theta(r_0, z), & L/2 < z < L \end{cases}$$

$$\text{Let } \theta(r, z) = R(r)Z(z)$$



$$\left\{ \begin{array}{l} \frac{d^2 Z}{dz^2} + \lambda_n^2 Z = 0 \\ \frac{dZ(0)}{dz} = \frac{dZ(L)}{dz} = 0 \end{array} \right\} \Rightarrow \text{where } Z_n(z) = A_n \cos \lambda_n z$$

$$\lambda_n = \frac{n\pi}{L}, \quad n = 0, 1, 2, \dots$$

and

$$\left. \begin{aligned} \frac{d^2 R_n}{dr^2} + \frac{1}{r} \frac{dR_n}{dr} - \lambda_n^2 R_n = 0 \\ R_n(0) = \text{finite} \end{aligned} \right\} \Rightarrow R_n(r) = B_n I_0(\lambda_n r)$$

Thus,

$$\Theta(r, z) = \sum_{n=0}^{\infty} a_n I_0(\lambda_n r) \cos \lambda_n z = a_0 + \sum_{n=1}^{\infty} a_n I_0(\lambda_n r) \cos \lambda_n z \quad (1)$$

In order to find the expansion coefficients  $a_n$ , we now apply the boundary condition at  $r=r_0$ :

$$k \sum_{n=1}^{\infty} a_n \lambda_n I_1(\lambda_n r_0) \cos \lambda_n z = q''_w(z)$$

Next,

$$\int_0^L \cos \lambda_n z \left\{ k \sum_{n=1}^{\infty} a_n \lambda_n I_1(\lambda_n r_0) \cos \lambda_n z = q''_w(z) \right\} dz$$
$$= \int_0^L \cos^2 \lambda_n z dz = \int_0^L q''_w(z) \cos \lambda_n z dz$$
$$k a_n \lambda_n I_1(\lambda_n r_0) \left( \frac{L}{2} \right) = \boxed{d_n}, \quad n=0, 1, 2, \dots \quad (2)$$

where

$$\begin{aligned} d_n &= \mu \rho \omega r_0 \int_0^{L/2} \cos \lambda_n z dz - h_1 \int_0^{L/2} \Theta(r_0, z) \cos \lambda_n z dz - h_2 \int_{L/2}^L \Theta(r_0, z) \cos \lambda_n z dz \\ &= Q_n - (h_1 - h_2) \sum_{m=0}^{\infty} a_m I_0(\lambda_m r_0) D_{mn} \end{aligned} \quad (3)$$

Here,  $Q_n = \mu \rho \omega r_0 D_{0n}$ ,  $n=0, 1, 2, \dots$

and

$$D_{mn} = \int_0^L \cos \lambda_m z \cos \lambda_n z dz = - \int_{-L/2}^{L/2} \cos \lambda_m z \cos \lambda_n z dz$$

which yields

$$D_{00} = \frac{L}{2}, \quad D_{mm} = D_{nn} = \frac{L}{4}, \quad m, n = 1, 2, 3, \dots$$

$$D_{0n} = \frac{1}{\lambda_n} \sin \lambda_n \frac{L}{2}, \quad D_{m0} = \frac{1}{\lambda_m} \sin \lambda_m \frac{L}{2}$$

$$D_{mn} = \frac{\lambda_m^2 D_{m0} \cos \lambda_n \frac{L}{2} - \lambda_n^2 D_{0n} \cos \lambda_m \frac{L}{2}}{\lambda_m^2 - \lambda_n^2}, \quad m \neq n$$

Now, substituting ③ into ② gives

$$k \lambda_n I_1(\lambda_n r_0) \frac{L}{2} a_n + (h_1 - h_2) \sum_{m=0}^{\infty} I_0(\lambda_m r_0) D_{mn} a_m = Q_n$$

which can be rewritten as

$$\sum_{m=0}^{\infty} C_{mn} a_m = Q_n, \quad n = 0, 1, 2, \dots \quad (4)$$

where

$$C_{mn} = k \lambda_m I_1(\lambda_m r_0) D_{00} \delta_{mn} + (h_1 - h_2) I_0(\lambda_m r_0) D_{mn}, \quad \delta_{mn} = \begin{cases} 1, & m=n \\ 0, & m \neq n \end{cases}$$

Equation ④ is a set of (infinite) simultaneous algebraic equations for the coefficients of the solution ①. Of course, the solution of the first  $N+1$  equations ( $m, n = 0, 1, 2, \dots, N$ ) gives  $a'_n$ 's of


$$\theta(r, z) \approx \sum_{n=0}^N a'_n I_0(\lambda_n r) \cos \lambda_n z$$

PROB. 5.35: a) Formulation:

$$\frac{\partial^2 T}{\partial r^2} + \frac{1}{r} \frac{\partial T}{\partial r} + \frac{1}{r^2} \frac{\partial^2 T}{\partial \phi^2} = 0$$

$$T(0, \phi) = 0, T(r_0, \phi) = f(\phi)$$

$$T(r, 0) = 0, T(r, \alpha) = 0$$



$$\text{Let } T(r, \phi) = R(r) \cdot \psi(\phi)$$

$$\left\{ \begin{array}{l} \frac{d^2 \psi}{d\phi^2} + \lambda^2 \psi = 0 \\ \psi(0) = 0 \\ \psi(\alpha) = 0 \end{array} \right\} \Rightarrow \underbrace{\psi_n = A_n \sin \lambda_n \phi}_{\lambda_n = \frac{n\pi}{\alpha}, n=1, 2, \dots} \Rightarrow \left\{ \begin{array}{l} r^2 \frac{d^2 R_n}{dr^2} + r \frac{dR_n}{dr} - \lambda_n^2 R_n = 0 \\ R_n(0) = 0 \end{array} \right\}$$

↓

$$\underbrace{R_n = B_n r^{\lambda_n}}$$

$$\therefore T(r, \phi) = \sum_{n=1}^{\infty} a_n r^{\lambda_n} \sin \lambda_n \phi$$

$$\Rightarrow f(\phi) = \sum_{n=1}^{\infty} a_n r_0^{\lambda_n} \sin \lambda_n \phi \Rightarrow a_n r_0^{\lambda_n} = \frac{2}{\alpha} \int_0^{\alpha} f(\phi') \sin \lambda_n \phi' d\phi'$$

$$T(r, \phi) = \frac{2}{\alpha} \sum_{n=1}^{\infty} \left(\frac{r}{r_0}\right)^{\lambda_n} \sin \lambda_n \phi \cdot \int_0^{\alpha} f(\phi') \sin \lambda_n \phi' d\phi'$$

b)  $f(\phi) = T_1 = \text{constant.}$


$$T(r, \phi) = \frac{2T_1}{\alpha} \sum_{n=1}^{\infty} \frac{[1 - (-1)^n]}{n} \left(\frac{r}{r_0}\right)^{\lambda_n} \sin \lambda_n \phi$$

PROB. 5.36:

$$\frac{\partial^2 T}{\partial r^2} + \frac{1}{r} \frac{\partial T}{\partial r} + \frac{1}{r^2} \frac{\partial^2 T}{\partial \phi^2} = 0$$

$$T(0, \phi) = \text{finite}, T(r_0, \phi) = 0$$

$$T(r, 0) = T_0, T(r, \alpha) = 0$$



$$\text{Let } T(r, \phi) = \psi(r, \phi) + \Omega(\phi)$$

$$\frac{\partial^2 \psi}{\partial r^2} + \frac{1}{r} \frac{\partial \psi}{\partial r} + \frac{1}{r^2} \frac{\partial^2 \psi}{\partial \phi^2} = 0$$

$$\psi(0, \phi) = \text{finite} - \Omega(\phi) = \underbrace{\text{finite}}_0$$

$$\psi(r_0, \phi) = -\Omega(\phi)$$

$$\Omega(r, 0) = 0, \Omega(r, \alpha) = 0$$

$$\frac{d^2 \Omega}{d \phi^2} = 0$$

$$\Omega(0) = T_0, \Omega(\alpha) = 0$$

$$\underline{\underline{\Omega(\phi) = T_0 \left(1 - \frac{\phi}{\alpha}\right)}}$$


↓ See Prob. 5.28.

$$\psi(r, \phi) = \frac{2}{\alpha} \sum_{n=1}^{\infty} \left(\frac{r}{r_0}\right)^{\frac{n\pi}{\alpha}} \sin \frac{n\pi}{\alpha} \phi \cdot \int_0^\alpha \underbrace{\psi(r_0, \phi') \sin \frac{n\pi}{\alpha} \phi' d\phi'}_{-\Omega(\phi')}$$

$$= -\frac{2T_0}{\pi} \sum_{n=1}^{\infty} \frac{1}{n} \left(\frac{r}{r_0}\right)^{\frac{n\pi}{\alpha}} \sin \frac{n\pi}{\alpha} \phi$$

$$\therefore T(r, \phi) = T_0 \left(1 - \frac{\phi}{\alpha}\right) - \frac{2T_0}{\pi} \sum_{n=1}^{\infty} \frac{1}{n} \left(\frac{r}{r_0}\right)^{\frac{n\pi}{\alpha}} \sin \frac{n\pi}{\alpha} \phi$$

PROB. 5.37:



Formulation in terms of  $\theta(r, \phi) = T(r, \phi) - T_\infty$ :

$$\frac{\partial^2 \theta}{\partial r^2} + \frac{1}{r} \frac{\partial \theta}{\partial r} + \frac{1}{r^2} \frac{\partial^2 \theta}{\partial \phi^2} = 0$$

$$\theta(0, \phi) = 0; \quad k \frac{\partial \theta}{\partial r} \Big|_{r=r_0} = q''_s \cos \phi - h \theta(r_0, \phi)$$

$$\frac{\partial \theta}{\partial \phi} \Big|_{\phi=0} = 0; \quad \theta(r, \alpha) = 0$$

$$\text{Let } \theta(r, \phi) = R(r) \cdot \psi(\phi)$$

$$\left\{ \begin{array}{l} \frac{d^2 \psi}{d \phi^2} + \lambda^2 \psi = 0 \\ \frac{d \psi(0)}{d \phi} = 0 \\ \psi(\alpha) = 0 \end{array} \right\} \Rightarrow \psi_n = A_n \cos \lambda_n \phi \quad \left\{ \begin{array}{l} \lambda_n = \frac{2n-1}{\alpha} \frac{\pi}{2}, \quad n=1, 2, \dots \\ R_n(0) = 0 \end{array} \right.$$

$$R_n(r) = B_n r^{\lambda_n}$$

$$\therefore \theta(r, \phi) = \sum_{n=1}^{\infty} a_n r^{\lambda_n} \cos \lambda_n \phi$$

Applying the B.C. at  $r=r_0$  gives

$$\begin{aligned} q'' \cos \phi &= \sum_{n=1}^{\infty} a_n [h r_0^{\lambda_n} + k \lambda_n r_0^{\lambda_n - 1}] \cos \lambda_n \phi \\ &= \sum_{n=1}^{\infty} a_n h r_0^{\lambda_n} \left[ 1 + \frac{\lambda_n}{Bi} \right] \cos \lambda_n \phi, \quad Bi = \frac{hr_0}{k} \end{aligned}$$

$$\begin{aligned} \therefore a_n h r_0^{\lambda_n} \left[ 1 + \frac{\lambda_n}{Bi} \right] &= \frac{2q''}{\alpha} \int_0^\alpha \cos \phi \cos \lambda_n \phi d\phi \\ &\vdots \\ &= q'' (-1)^n \cos \alpha \frac{(2n-1)\pi}{1-\lambda_n^2} \end{aligned}$$

Thus,

$$\theta(r, \phi) = \frac{q'' \pi \cos \alpha}{h} \sum_{n=1}^{\infty} \left( \frac{r}{r_0} \right)^{\lambda_n} \frac{(-1)^n (2n-1)}{1-\lambda_n^2} \frac{\cos \lambda_n \phi}{\left[ 1 + \frac{\lambda_n}{Bi} \right]}$$

or

$$T(r, \phi) - T_\infty = \frac{q'' \pi r_0 \cos \alpha}{k} \sum_{n=1}^{\infty} (-1)^n \left( \frac{r}{r_0} \right)^{\lambda_n} \frac{2n-1}{1-\lambda_n^2} \frac{\cos \lambda_n \phi}{[Bi + \lambda_n]}$$

PROB. 5.38: Formulation of the problem:


$$\frac{\partial^2 T}{\partial r^2} + \frac{1}{r} \frac{\partial T}{\partial r} + \frac{1}{r^2} \frac{\partial^2 T}{\partial \phi^2} = 0$$

$$T(r_0, \phi) = f(\phi)$$

$$T(r, \phi) \rightarrow \text{finite} \quad r \rightarrow \infty$$

$$T(r, \phi) = T(r, \phi + 2\pi)$$

$$\frac{\partial T(r, \phi)}{\partial \phi} = \frac{\partial T(r, \phi + 2\pi)}{\partial \phi}$$



Seeking a product solution of the form  $T(r, \phi) = R(r)\Phi(\phi)$  yields the general solution

$$T(r, \phi) = (A_1 r^\lambda + A_2 r^{-\lambda}) (B_1 \sin \lambda \phi + B_2 \cos \lambda \phi)$$

for the differential equation, where the sign of the separation constant is consistent with the fact that the  $\phi$ -direction is the homogeneous direction for  $T(r, t)$ . On the other hand, imposing the two boundary conditions in the  $\phi$ -direction gives

$$\lambda_n = n, \quad n = 0, 1, 2, \dots$$

Furthermore,  $T(\infty, \phi) = \text{finite}$  gives  $A_1 = 0$ . Thus, the solution of the problem can be written as

$$T(r, \phi) = \sum_{n=0}^{\infty} r^{-n} (a_n \sin n\phi + b_n \cos n\phi)$$

The boundary condition  $T(r_0, \phi) = f(\phi)$  yields

$$f(\phi) = b_0 + \sum_{n=1}^{\infty} r_0^{-n} (a_n \sin n\phi + b_n \cos n\phi)$$

Thus,

$$b_0 = \frac{1}{2\pi} \int_0^{2\pi} f(\phi) d\phi$$

$$a_n = \frac{r_0^n}{\pi} \int_0^{2\pi} f(\phi) \sin n\phi d\phi, \quad n=1,2,3,\dots$$

$$b_n = \frac{r_0^n}{\pi} \int_0^{2\pi} f(\phi) \cos n\phi d\phi, \quad n=1,2,3,\dots$$

Then, the solution becomes

$$\begin{aligned} T(r, \phi) &= \frac{1}{2\pi} \int_0^{2\pi} f(\phi) d\phi + \frac{1}{\pi} \sum_{n=1}^{\infty} \left(\frac{r_0}{r}\right)^n \left\{ \sin n\phi \int_0^{2\pi} f(\phi') \sin n\phi' d\phi' \right. \\ &\quad \left. + \cos n\phi \int_0^{2\pi} f(\phi') \cos n\phi' d\phi' \right\} \end{aligned}$$

or

$$T(r, \phi) = \frac{1}{\pi} \int_0^{2\pi} \left\{ \frac{1}{2} + \sum_{n=1}^{\infty} \left(\frac{r_0}{r}\right)^n \cos n(\phi - \phi') \right\} f(\phi') d\phi'$$


which can also be written as [See Eq. (5.86)]

$$T(r, \phi) = \frac{1}{2\pi} \int_0^{2\pi} \frac{r^2 - r_0^2}{r^2 - 2rr_0 \cos(\phi - \phi') + r_0^2} f(\phi') d\phi'$$

Note that, as  $r \rightarrow \infty$ ,  $T(r, \phi)$  must approach a constant value given by


$$T(\infty, \phi) = T_\infty = \frac{1}{2\pi} \int_0^{2\pi} f(\phi) d\phi$$

PROB. 5.39: The problem defined by Eqs. (5.62) and (5.63a-d):



$$\theta(r, \phi) = T(r, \phi) - T_0$$

≡



where

$$F(\phi) = \begin{cases} f(\phi) - T_0, & 0 < \phi < \pi \\ -[f(2\pi - \phi) - T_0], & \pi < \phi < 2\pi \end{cases}$$

$$\begin{aligned} \theta(r, \phi) &= \frac{1}{2\pi} \int_0^{2\pi} \frac{r_0^2 - r^2}{r_0^2 - 2rr_0 \cos(\phi - \phi') + r^2} F(\phi') d\phi' \\ &= \frac{1}{2\pi} \int_0^\pi (-)[f(\phi') - T_0] d\phi' - \frac{1}{2\pi} \int_{\pi}^{2\pi} (-)[f(2\pi - \phi') - T_0] d\phi' \\ &\quad \underbrace{\qquad\qquad\qquad}_{\cos(\phi + \phi')} \\ &\quad \frac{1}{2\pi} \int_0^\pi \frac{r_0^2 - r^2}{r_0^2 - 2rr_0 \cos(\phi + \phi' - 2\pi) + r^2} [f(\phi') - T_0] d\phi' \end{aligned}$$


Thus,

$$T(r, \phi) - T_0 = \int_0^\pi [K(r, \phi - \phi') - K(r, \phi + \phi')] [f(\phi') - T_0] d\phi'$$

where

$$K(r, \phi \pm \phi') = \frac{1}{2\pi} \frac{r_0^2 - r^2}{r_0^2 - 2rr_0 \cos(\phi \pm \phi') + r^2}$$

PROB. 5.40:



where

$$F(\phi) = \begin{cases} f(\phi), & 0 < \phi < \pi \\ f(2\pi - \phi), & \pi < \phi < 2\pi \end{cases}$$

Thus,

$$\begin{aligned} T(r, \phi) &= \frac{1}{2\pi} \int_0^{2\pi} \frac{r_0^2 - r^2}{r_0^2 - 2rr_0 \cos(\phi - \phi') + r^2} F(\phi') d\phi' \\ &= \frac{1}{2\pi} \int_0^\pi ( ) f(\phi') d\phi' + \underbrace{\frac{1}{2\pi} \int_\pi^{2\pi} ( ) f(2\pi - \phi') d\phi'}_{\text{brace}} \\ &\quad \downarrow \\ &\quad \frac{1}{2\pi} \int_0^\pi \frac{r_0^2 - r^2}{r_0^2 - 2rr_0 \cos(\phi + \phi' - 2\pi) + r^2} f(\phi') d\phi' \end{aligned}$$

$$\therefore T(r, \phi) = \int_0^\pi [K(r, \phi - \phi') + K(r, \phi + \phi')] f(\phi') d\phi'$$

where

$$K(r, \phi \pm \phi') = \frac{1}{2\pi} \frac{r_0^2 - r^2}{r_0^2 - 2rr_0 \cos(\phi \pm \phi') + r^2}$$

PROB. 5.41:

$$f(\theta) = \begin{cases} \cos\theta, & 0 < \theta < \frac{\pi}{2} \\ 0, & \frac{\pi}{2} < \theta < \pi \end{cases}$$

Let  $x = \cos\theta$ , then

$$F(x) = f(\cos^{-1}x) = \begin{cases} x, & 0 < x < 1 \\ 0, & -1 < x < 0 \end{cases}$$

$$\therefore F(x) = A_0 P_0(x) + A_1 P_1(x) + A_2 P_2(x) + \dots, -1 < x < 1$$

From Eq. (5.135),

$$A_n = \frac{2n+1}{2} \int_{-1}^1 F(x) P_n(x) dx$$

$$n=0: \quad A_0 = \frac{1}{2} \cdot \int_0^1 x \cdot 1 \cdot dx = \frac{1}{4}$$

$$n=1: \quad A_1 = \frac{3}{2} \cdot \int_0^1 x \cdot x \cdot dx = \frac{1}{2}$$

$$n=2: \quad A_2 = \frac{5}{2} \cdot \int_0^1 x \cdot \left\{ \frac{1}{2}(3x^2 - 1) \right\} dx = \frac{5}{16}$$

⋮

$$F(x) = \frac{1}{4} P_0(x) + \frac{1}{2} P_1(x) + \frac{5}{16} P_2(x) + \dots, -1 < x < 1$$

Thus,

$$f(\theta) = \frac{1}{4} P_0(\cos\theta) + \frac{1}{2} P_1(\cos\theta) + \frac{5}{16} P_2(\cos\theta) + \dots, 0 < \theta < \pi$$

where

$$P_0(\cos\theta) = 1, \quad P_1(\cos\theta) = \cos\theta, \quad P_2(\cos\theta) = \frac{1}{2}(3\cos^2\theta - 1), \dots$$

PROB. 5.42: The temperature distribution is given by [See Section 5.4.3, Eq. (5.151)]:

$$T(r, \theta) = \sum_{n=0}^{\infty} K_n r^n P_n(\cos \theta)$$

If the surface temperature is specified as

$$T(r_0, \theta) = f(\theta) = \begin{cases} T_0 \cos \theta, & 0 < \theta < \frac{\pi}{2} \\ 0, & \frac{\pi}{2} < \theta < \pi \end{cases}$$

then

$$f(\theta) = \sum_{n=0}^{\infty} K_n r_0^n P_n(\cos \theta), \quad 0 < \theta < \pi$$

or

$$f(x) = \sum_{n=0}^{\infty} K_n r_0^n P_n(x), \quad -1 < x < 1$$

where  $x = \cos \theta$  and

$$f(x) = \begin{cases} T_0 x, & 0 < x < 1 \\ 0, & -1 < x < 0 \end{cases}$$

Now, the expansion coefficients  $K_n r_0^n$  can be determined from

$$K_n r_0^n = \frac{2n+1}{2} \int_{-1}^1 f(x) P_n(x) dx$$

$$n=0: \quad K_0 = T_0 \frac{1}{2} \int_0^1 x dx = T_0 \frac{1}{4}$$

$$n=1: \quad K_1 r_0 = T_0 \frac{3}{2} \int_0^1 x^2 dx = T_0 \frac{1}{2}$$


$$n=2: \quad K_2 r_0^2 = T_0 \frac{5}{2} \int_0^1 x \left\{ \frac{1}{2} (3x^2 - 1) \right\} dx = T_0 \frac{5}{16}$$

⋮

Thus,

$$\frac{T(r, \theta)}{T_0} = \frac{1}{4} + \frac{1}{2} \left( \frac{r}{r_0} \right) P_1(\cos \theta) + \frac{5}{16} \left( \frac{r}{r_0} \right)^2 P_2(\cos \theta) + \dots$$

PROB. 5.43:



We note that  $T = T(r, \theta)$ . The formulation is then given by

$$\frac{\partial}{\partial r} \left( r^2 \frac{\partial \phi}{\partial r} \right) + \frac{1}{\sin \theta} \frac{\partial}{\partial \theta} \left( \sin \theta \frac{\partial \phi}{\partial \theta} \right) = 0, \quad \phi(r, \theta) = T(r, \theta) - T_0$$

$$\phi(0, \theta) = 0, \quad \frac{\partial \phi(r_0, \theta)}{\partial r} = \frac{q''_w}{k}$$

$$\phi(r, 0) = \text{finite}, \quad \phi(r, \frac{\pi}{2}) = 0$$

$$\text{Let } \phi(r, \theta) = R(r) \cdot \psi(\theta)$$



$$\frac{1}{\sin \theta} \frac{d}{d\theta} \left( \sin \theta \frac{d\psi}{d\theta} \right) + \lambda^2 \psi(\theta) = 0 \quad \frac{d^2 R}{dr^2} + \frac{2}{r} \frac{dR}{dr} - \frac{\lambda^2}{r^2} R(r) = 0$$

$$\downarrow \mu = \cos \theta$$

$$R(0) = 0$$

$$\downarrow \quad B_n = 0$$

$$\frac{d}{d\mu} \left[ (1-\mu^2) \frac{d\psi}{d\mu} \right] + \underbrace{\alpha(\alpha+1)}_{\lambda^2} \psi(\mu) = 0 \quad - \textcircled{1}$$

$$\psi(0) = 0, \quad \psi(1) = \text{finite}$$

$$R_n(r) = A_n r^n + B_n r^{-(n+1)}$$

$$\uparrow$$

In order to satisfy the finiteness condition at  $\mu = 1$ ,  $\alpha = n = 0, 1, 2, \dots$  and the solutions of Eq.(1) are the Legendre polynomials  $P_n(\mu)$ .

On the other hand, to satisfy the condition  $\psi(0)=0$ ,

$$P_n(0) = 0 \Rightarrow n=1, 3, 5, \dots \quad (\text{see Eq. 5.113 in text})$$

Thus,  
↑ Must be odd integer.

$$\psi_n(\mu) = C_n P_n(\mu), \quad n=1, 3, 5, \dots$$

Then, the solution for  $\psi(r, \mu)$  can be constructed as

$$\Phi(r, \mu) = \sum_{n=1, 3, 5, \dots}^{\infty} D_n r^n P_n(\mu), \quad D_n = A_n C_n$$

$$\frac{q''}{k} = \sum_{n=1, 3, 5, \dots}^{\infty} D_n n! \int_0^{(n-1)} P_n(\mu) d\mu, \quad 0 < \mu \leq 1 \quad (\text{Not } -\mu \leq 1) \quad -②$$

Re-visit the orthogonality property of  $P_n(\mu)$ 's over  $-1 \leq \mu \leq 1$ :

$$\begin{aligned} \int_{-1}^1 P_m(\mu) P_n(\mu) d\mu &= \begin{cases} 0, & m \neq n \\ N_n, & m = n \end{cases} & \int_0^1 P_m(\mu) P_n(\mu) d\mu &= \begin{cases} 0, & \text{if } m \text{ and } n \text{ both} \\ & \text{odd (or even),} \\ & \text{and } m \neq n \\ N'_n, & m = n \end{cases} \\ N_n &= \int_{-1}^1 [P_n(\mu)]^2 d\mu = \frac{2}{2n+1} & N'_n &= \int_0^1 [P_n(\mu)]^2 d\mu = \frac{1}{2n+1} \end{aligned}$$

Now, with the use of the orthogonality property over  $0 < \mu \leq 1$ , Eq.(2) yields,

$$D_n n! \int_0^{(n-1)} P_n(\mu) d\mu = \dots = \frac{q''}{k} \left[ P_{n+1}(1) - P_{n+1}(0) - P_{n-1}(1) + P_{n-1}(0) \right]$$

Thus,

$$\Phi(r, \mu) = \frac{q''}{k} \sum_{n=1, 3, 5, \dots}^{\infty} \frac{r^n}{n!} \left[ P_{n+1}(1) - P_{n+1}(0) - P_{n-1}(1) + P_{n-1}(0) \right] P_n(\mu)$$

PROB. 5.44: The formulation in terms of  $\psi(r, \theta) = T(r, \theta) - T_0$ :

$$\frac{\partial}{\partial r} \left( r^2 \frac{\partial \psi}{\partial r} \right) + \frac{1}{\sin \theta} \frac{\partial}{\partial \theta} \left( \sin \theta \frac{\partial \psi}{\partial \theta} \right) = 0$$

$$\psi(r_1, \theta) = 0$$

$$\psi(r_2, \theta) = f(\theta) - T_0$$

Assuming a product solution in the form  $\psi(r, \theta) = R(r) \cdot \phi(\theta)$  yields

$$\psi(r, \theta) = \sum_{n=0}^{\infty} A_n [C_n r^n + B_n r^{-(n+1)}] P_n(\cos \theta)$$

The B.C. at  $r_1$  gives

$$C_n r_1^n + B_n r_1^{-(n+1)} = 0 \Rightarrow B_n = -C_n r_1^{2n+1}$$

Then the solution for  $\psi(r, \theta)$  can be rewritten as

$$\psi(r, \theta) = \sum_{n=0}^{\infty} A_n \left[ \left( \frac{r}{r_1} \right)^n - \left( \frac{r}{r_1} \right)^{-(n+1)} \right] P_n(\cos \theta)$$

The B.C. at  $r_2$  yields

$$f(\theta) - T_0 = \sum_{n=0}^{\infty} A_n \left[ \left( \frac{r_2}{r_1} \right)^n - \left( \frac{r_2}{r_1} \right)^{-(n+1)} \right] P_n(\cos \theta)$$

Thus,

$$A_n \left[ \left( \frac{r_2}{r_1} \right)^n - \left( \frac{r_2}{r_1} \right)^{-(n+1)} \right] = \frac{2n+1}{2} \int_0^\pi [f(\theta) - T_0] P_n(\cos \theta) \sin \theta d\theta$$

Then the solution can be written as

$$T(r, \theta) - T_0 = \frac{1}{2} \sum_{n=0}^{\infty} (2n+1) \frac{\left[ \left( \frac{r}{r_1} \right)^n - \left( \frac{r}{r_1} \right)^{-(n+1)} \right]}{\left[ \left( \frac{r_2}{r_1} \right)^n - \left( \frac{r_2}{r_1} \right)^{-(n+1)} \right]} P_n(\cos \theta) \int_0^\pi [f(\theta') - T_0] P_n(\cos \theta') \sin \theta' d\theta'$$

PROB. 5.45: a) Formulation in terms of

$$\Theta(r, \phi, z) = T(r, \phi, z) - T_1 :$$


$$\frac{\partial^2 \Theta}{\partial r^2} + \frac{1}{r} \frac{\partial \Theta}{\partial r} + \frac{1}{r^2} \frac{\partial^2 \Theta}{\partial \phi^2} + \frac{\partial^2 \Theta}{\partial z^2} = 0$$

$$\Theta(0, \phi, z) = \text{finite}; \quad \Theta(r_0, \phi, z) = f(\phi) - T_1$$

$$\Theta(r, \phi, 0) = 0; \quad \Theta(r, \phi, L) = 0$$

$$\Theta(r, \phi, z) = \Theta(r, \phi + 2\pi, z)$$

$$\frac{\partial \Theta(r, \phi, z)}{\partial \phi} = \frac{\partial \Theta(r, \phi + 2\pi, z)}{\partial \phi}$$



$$\text{Let } \Theta(r, \phi, z) = R(r) \cdot \psi(\phi) \cdot Z(z)$$

$$\frac{d^2 Z}{dz^2} + \lambda^2 Z = 0$$

$$Z(0) = 0$$

$$Z(L) = 0$$

↓

$$Z_n(z) = A_n \sin \lambda_n z$$

$$\lambda_n = \frac{n\pi}{L}, \quad n=1, 2, \dots$$

$$\frac{d^2 \psi}{d\phi^2} + \beta^2 \psi = 0$$

$$\psi(\phi) = \psi(\phi + 2\pi)$$

$$\frac{d \psi(\phi)}{d \phi} = \frac{d \psi(\phi + 2\pi)}{d \phi}$$

↓

$$\psi_m(\phi) = B_m \cos m\phi + C_m \sin m\phi$$

$$m = 0, 1, 2, \dots$$

Thus,

$$\left. \begin{aligned} r^2 \frac{d^2 R_{mn}}{dr^2} + r \frac{dR_{mn}}{dr} - (m^2 + r^2 \lambda_n^2) R_{mn} &= 0 \\ R_{mn}(0) &= \text{finite} \end{aligned} \right\} \quad R_{mn}(r) = D_{mn} I_m(\lambda_n r)$$

$$\therefore \Theta(r, \phi, z) = \sum_{m=0}^{\infty} \sum_{n=1}^{\infty} (a_{mn} \cos m\phi + b_{mn} \sin m\phi) \sin \lambda_n z I_m(\lambda_n r)$$

$$\Rightarrow F(\phi) = f(\phi) - T_1 = \sum_{m=0}^{\infty} \sum_{n=1}^{\infty} (a_{mn} \cos m\phi + b_{mn} \sin m\phi) \sin \lambda_n z I_m(\lambda_n r_0)$$

$$= \sum_{n=1}^{\infty} \left[ \sum_{m=0}^{\infty} I_m(\lambda_n r_0) (a_{mn} \cos m\phi + b_{mn} \sin m\phi) \right] \sin \lambda_n z$$

$$\therefore \sum_{m=0}^{\infty} I_m(\lambda_n r_0) (a_{mn} \cos m\phi + b_{mn} \sin m\phi) = \frac{2}{L} \int_0^L F(\phi) \sin \lambda_n z dz$$

$$= \frac{2}{n\pi} F(\phi) [1 - (-1)^n]$$

$$\frac{2}{n\pi} F(\phi) [1 - (-1)^n] = I_0(\lambda_n r_0) a_{0n} + \sum_{m=1}^{\infty} I_m(\lambda_n r_0) (a_{mn} \cos m\phi + b_{mn} \sin m\phi)$$

$$I_0(\lambda_n r_0) a_{0n} = \frac{1}{2\pi} \int_0^{2\pi} \frac{2}{n\pi} F(\phi) [1 - (-1)^n] d\phi$$

$$= \frac{1}{n\pi^2} [1 - (-1)^n] \int_0^{2\pi} F(\phi) d\phi$$

$$I_m(\lambda_n r_0) a_{mn} = \frac{1}{\pi} \int_0^{2\pi} \frac{2}{n\pi} F(\phi) [1 - (-1)^n] \cos m\phi d\phi$$

$$= \frac{2}{n\pi^2} [1 - (-1)^n] \int_0^{2\pi} F(\phi) \cos m\phi d\phi$$

$$I_m(\lambda_n r_0) b_{mn} = \frac{1}{\pi} \int_0^{2\pi} \frac{2}{n\pi} F(\phi) [1 - (-1)^n] \sin m\phi d\phi$$

$$= \frac{2}{\pi r^2} [1 - (-1)^n] \int_0^{2\pi} F(\phi) \sin m\phi d\phi$$

Thus,

$$\begin{aligned} \theta(r, \phi, z) &= T(r, \phi, z) - T_1 \\ &= \frac{1}{\pi^2} \int_0^{2\pi} F(\phi) d\phi \cdot \sum_{n=1}^{\infty} \frac{[1 - (-1)^n]}{n} \frac{I_0(\frac{n\pi}{L}r)}{I_0(\frac{n\pi}{L}r_0)} \sin \frac{n\pi}{L} z \\ &\quad + \frac{2}{\pi^2} \sum_{m=1}^{\infty} \sum_{n=1}^{\infty} \frac{[1 - (-1)^n]}{n} \frac{I_m(\frac{n\pi}{L}r)}{I_m(\frac{n\pi}{L}r_0)} \sin \frac{n\pi}{L} z \\ &\quad \cdot \left\{ \cos m\phi \int_0^{2\pi} F(\phi') \cos m\phi' d\phi' \right. \\ &\quad \left. + \sin m\phi \int_0^{2\pi} F(\phi') \sin m\phi' d\phi' \right\} \end{aligned}$$

Or,

$$\begin{aligned} T(r, \phi, z) - T_1 &= \frac{2}{\pi^2} \left\{ \frac{1}{2} \int_0^{2\pi} [f(\phi') - T_1] d\phi' \sum_{n=1}^{\infty} \frac{[1 - (-1)^n]}{n} \frac{I_0(\frac{n\pi}{L}r)}{I_0(\frac{n\pi}{L}r_0)} \sin \frac{n\pi}{L} z \right. \\ &\quad \left. + \sum_{n=1}^{\infty} \frac{[1 - (-1)^n]}{n} \sin \frac{n\pi}{L} z \sum_{m=1}^{\infty} \frac{I_m(\frac{n\pi}{L}r)}{I_m(\frac{n\pi}{L}r_0)} \int_0^{2\pi} [f(\phi') - T_1] \cos m(\phi - \phi') d\phi' \right\} \end{aligned}$$


b) Since

$$\int_0^{2\pi} \cos m(\phi - \phi') d\phi' = 0, \quad m = 1, 2, \dots$$

when  $f(\phi) = T_0 = \text{Const.}$ , the solution reduces to

$$T(r, z) - T_1 = (T_0 - T_1) \frac{2}{\pi} \sum_{n=1}^{\infty} \frac{[1 - (-1)^n]}{n} \frac{I_0(\frac{n\pi}{L}r)}{I_0(\frac{n\pi}{L}r_0)} \sin \frac{n\pi}{L} z$$

PROB. 5.46:



a) Energy balance on the system shown gives:

$$\frac{\partial q_x}{\partial x} \Delta x + \frac{\partial q_y}{\partial y} \Delta y + 2h \Delta x \Delta y (T - T_\infty) = 0$$

$$\frac{\partial}{\partial x} (-k \Delta y \delta \frac{\partial T}{\partial x}) \Delta x + \frac{\partial}{\partial y} (-k \Delta x \delta \frac{\partial T}{\partial y}) \Delta y + 2h \Delta x \Delta y (T - T_\infty) = 0$$

Assume  $k = \text{const.}$

---


$$\therefore \frac{\partial^2 T}{\partial x^2} + \frac{\partial^2 T}{\partial y^2} - m^2 (T - T_\infty) = 0, \quad m^2 = \frac{2h}{k\delta}$$


---

b) Formulation in terms of  $\theta(x, y) = T(x, y) - T_\infty$  with  $\theta_0 = T_0 - T_\infty$ :

$$\frac{\partial^2 \theta}{\partial x^2} + \frac{\partial^2 \theta}{\partial y^2} - m^2 \theta = 0$$

$$\theta(0, y) = \theta_0; \quad \theta(a, y) = \theta_0$$

$$\theta(x, 0) = \theta_0; \quad \theta(x, a) = \theta_0$$

Consider the following auxiliary problem:

$$\left\{ \begin{array}{l} \frac{\partial^2 \theta_1}{\partial x^2} + \frac{\partial^2 \theta_1}{\partial y^2} - m^2 \theta_1 = 0 \\ \theta_1(0, y) = 0; \quad \theta_1(a, y) = 0 \\ \theta_1(x, 0) = 0; \quad \theta_1(x, a) = \theta_0 \end{array} \right\}$$

Let  $\theta_1(x, y) = X(x) \cdot Y(y)$

$$\left\{ \begin{array}{l} \frac{d^2 X}{dx^2} + \lambda^2 X = 0 \\ X(0) = 0 \\ X(a) = 0 \end{array} \right. \quad \left. \begin{array}{l} X_n(x) = A_n \sin \lambda_n x \\ \lambda_n = \frac{n\pi}{a}, \quad n=1, 2, \dots \end{array} \right| \quad \left. \begin{array}{l} \frac{d^2 Y_n}{dy^2} - (m^2 + \lambda_n^2) Y_n = 0 \\ Y_n(0) = 0 \\ Y_n(y) = B_n \sinh \sqrt{\lambda_n^2 + m^2} y \end{array} \right|$$

$$\therefore \theta_1(x, y) = \sum_{n=1}^{\infty} a_n \sin \lambda_n x \sinh \sqrt{\lambda_n^2 + m^2} y$$

The B.C. at  $y=0$   
gives  $\Rightarrow \theta_0 = \sum_{n=1}^{\infty} [a_n \sinh \sqrt{\lambda_n^2 + m^2} a] \sin \lambda_n x$

$$\therefore a_n \sinh \sqrt{\lambda_n^2 + m^2} a = \frac{2\theta_0}{a} \int_0^a \sin \lambda_n x \, dx = \frac{2\theta_0}{n\pi} [1 - (-1)^n]$$



Thus,

$$\theta_1(x, y) = \frac{2\theta_0}{\pi} \sum_{n=1}^{\infty} \frac{[1 - (-1)^n]}{n} \frac{\sin \lambda_n x \sinh \sqrt{\lambda_n^2 + m^2} y}{\sinh \sqrt{\lambda_n^2 + m^2} a}$$

Now, by the method of superposition we obtain

$$\begin{aligned} \frac{\theta(x, y)}{\theta_0} &= \frac{T(x, y) - T_\infty}{T_0 - T_\infty} = \frac{2}{\pi} \sum_{n=1}^{\infty} \frac{[1 - (-1)^n]}{n \sinh \sqrt{\lambda_n^2 + m^2} a} \left\{ \sin \lambda_n x \left[ \sinh \sqrt{\lambda_n^2 + m^2} y \right. \right. \\ &\quad \left. \left. + \sinh \sqrt{\lambda_n^2 + m^2} (a-y) \right] + \sin \lambda_n y \left[ \sinh \sqrt{\lambda_n^2 + m^2} x + \sinh \sqrt{\lambda_n^2 + m^2} (a-x) \right] \right\} \end{aligned}$$

PROB. 5.47:



a) Energy (thermal) balance on the system shown gives:

$$q_x + q_y + \dot{q} \Delta x \Delta y \delta = q_x + \frac{\partial q_x}{\partial x} \Delta x + q_y + \frac{\partial q_y}{\partial y} \Delta y + 2h \Delta x \Delta y (T - T_\infty)$$

$$q_x = -k \Delta y \delta \frac{\partial T}{\partial x} \quad \text{and} \quad q_y = -k \Delta x \delta \frac{\partial T}{\partial y}$$



$$\dot{q} \Delta x \Delta y \delta = \frac{\partial}{\partial x} \left( -k \Delta y \delta \frac{\partial T}{\partial x} \right) \Delta x + \frac{\partial}{\partial y} \left( -k \Delta x \delta \frac{\partial T}{\partial y} \right) \Delta y + 2h \Delta x \Delta y (T - T_0)$$



Assume  $k = \text{const.}$

---


$$\frac{\partial^2 T}{\partial x^2} + \frac{\partial^2 T}{\partial y^2} - m^2 (T - T_0) + \frac{\dot{q}}{k} = 0, \quad m^2 = \frac{2h}{k\delta}$$


---

b) Formulation in terms of

$$\frac{\theta(x,y) = T(x,y) - T_{\infty} - \frac{q}{km^2}}{\left\{ \begin{array}{l} \frac{\partial^2 \theta}{\partial x^2} + \frac{\partial^2 \theta}{\partial y^2} - m^2 \theta = 0 \\ \theta(0,y) = \theta_0; \quad \theta(a,y) = \theta_0 \\ \theta(x,0) = \theta_0; \quad \theta(x,a) = \theta_0 \end{array} \right\}}$$

where  $\theta_0 = T_0 - T_{\infty} - \frac{q}{km^2}$

Consider the following auxiliary problem:

$$\frac{\partial^2 \theta_1}{\partial x^2} + \frac{\partial^2 \theta_1}{\partial y^2} - m^2 \theta_1 = 0$$

$$\begin{aligned} \theta_1(0,y) &= 0; \quad \theta_1(a,y) = 0 \\ \theta_1(x,0) &= 0; \quad \theta_1(x,a) = \theta_0 \end{aligned}$$

The solution to the auxiliary problem can be obtained as  
(See Prob. 5.46) :

$$\theta_1(x,y) = \frac{2\theta_0}{\pi} \sum_{n=1}^{\infty} \frac{[1-(-1)^n]}{n} \frac{\sin \lambda_n x \sinh \sqrt{m^2 + \lambda_n^2} y}{\sinh \sqrt{m^2 + \lambda_n^2} a}$$

Now, the method of superposition gives

$$\begin{aligned} \frac{T(x,y) - T_{\infty} - \frac{q}{km^2}}{T_0 - T_{\infty} - \frac{q}{km^2}} &= \frac{2}{\pi} \sum_{n=1}^{\infty} \frac{[1-(-1)^n]}{n \sinh \sqrt{m^2 + \lambda_n^2} a} \left\{ \sin \lambda_n x \left[ \sinh \sqrt{m^2 + \lambda_n^2} y \right. \right. \\ &\quad \left. \left. + \sinh \sqrt{m^2 + \lambda_n^2} (a-y) \right] + \sin \lambda_n y \left[ \sinh \sqrt{m^2 + \lambda_n^2} x + \sinh \sqrt{m^2 + \lambda_n^2} (a-x) \right] \right\} \end{aligned}$$

PROB. 5.48: Formulation in terms of  $\theta(r,z) = T(r,z) - T_0$ :

$$\frac{\partial^2 \theta}{\partial r^2} + \frac{1}{r} \frac{\partial \theta}{\partial r} + \frac{\partial^2 \theta}{\partial z^2} + \frac{q_0}{k} e^{-\beta z} = 0$$

$$\theta(0,z) = \text{finite}; \quad \theta(r_0, z) = 0$$

$$\theta(r, 0) = 0; \quad \theta(r, \infty) = 0$$


Let

$$\theta(r,z) = \psi(r,z) + e^{-\beta z} \phi(r)$$

$$\frac{\partial^2 \psi}{\partial r^2} + \frac{1}{r} \frac{\partial \psi}{\partial r} + \frac{\partial^2 \psi}{\partial z^2} = 0$$

$$\psi(0,z) = \text{finite}; \quad \psi(r_0, z) = 0$$

$$\psi(r, 0) = -\phi(r); \quad \psi(r, \infty) = 0$$



$$\frac{d^2 \phi}{dr^2} + \frac{1}{r} \frac{d\phi}{dr} + \beta^2 \phi(r) = -\frac{q_0}{k}$$

$$\phi(0) = \text{finite}$$

$$\phi(r_0) = 0$$

$$\Downarrow \text{Let } \psi(r,z) = R(r) \cdot Z(z)$$

$$\psi(r,z) = \sum_{n=1}^{\infty} A_n e^{-\lambda_n z} J_0(\lambda_n r)$$

where  $\lambda_n$ 's are the positive roots of

$$J_0(\lambda_n r_0) = 0, \quad n=1,2,3,\dots$$

$$\Downarrow$$

$$A_n = \frac{1}{N_n} \int_0^{r_0} \{-\phi(r)\} J_0(\lambda_n r) r dr$$

$$= \dots = \frac{1}{N_n} \frac{q_0}{k} \frac{r_0}{\lambda_n} \frac{J_1(\lambda_n r_0)}{\beta^2 - \lambda_n^2}$$

$$\phi(r) = B J_0(\beta r) + C Y_0(\beta r) - \frac{q_0}{k \beta^2}$$

$$\phi(0) = \text{finite} \Rightarrow C = 0$$

$$\phi(r_0) = 0 \Rightarrow B = \frac{q_0}{k \beta^2 J_0(\beta r_0)}$$

$$\phi(r) = \frac{q_0}{k \beta} \left[ \frac{J_0(\beta r)}{J_0(\beta r_0)} - 1 \right]$$

$$\therefore \Theta(r, z) = T(r, z) - T_0 = \frac{q_0}{k\beta^2} \left[ \frac{J_0(\beta r)}{J_0(\beta r_0)} - 1 \right] e^{-\beta z} + \frac{2q_0}{r_0 k} \sum_{n=1}^{\infty} \frac{\bar{e}^{\lambda_n z}}{\lambda_n (\beta^2 - \lambda_n^2)} \frac{J_0(\lambda_n r)}{J_1(\lambda_n r_0)}$$

$$T(r, z) - T_0 = \frac{q_0}{k} \left\{ \left[ \frac{J_0(\beta r)}{J_0(\beta r_0)} - 1 \right] \frac{\bar{e}^{\beta z}}{\beta^2} + \frac{2}{r_0} \sum_{n=1}^{\infty} \frac{\bar{e}^{\lambda_n z}}{\lambda_n (\beta^2 - \lambda_n^2)} \frac{J_0(\lambda_n r)}{J_1(\lambda_n r_0)} \right\}$$

Note that the above solution does not converge if  $\beta = \lambda_n$ ,  $n=1, 2, 3, \dots$

Solution if  $\beta = \lambda_n$ ,  $n=1, 2, 3, \dots$

First do the following expansion:

$$\phi(r) = \sum_{n=1}^{\infty} B_n J_0(\lambda_n r), \quad 0 \leq r \leq r_0$$

$$\therefore B_n = \frac{1}{N_n} \int_0^{r_0} \phi(r) J_0(\lambda_n r) r dr = -A_n$$

Thus, the solution can be written as

$$T(r, z) - T_0 = \frac{2q_0}{kr_0} \sum_{n=1}^{\infty} \frac{\bar{e}^{\lambda_n z} - \bar{e}^{\beta z}}{\lambda_n (\beta^2 - \lambda_n^2)} \cdot \frac{J_0(\lambda_n r)}{J_1(\lambda_n r_0)}$$

Now, take the following limit,

$$\lim_{\beta \rightarrow \lambda_n} \frac{\bar{e}^{\lambda_n z} - \bar{e}^{\beta z}}{\lambda_n (\beta^2 - \lambda_n^2)} = \lim_{\beta \rightarrow \lambda_n} \frac{z \bar{e}^{\beta z}}{2\lambda_n \beta} = \frac{z \bar{e}^{\lambda_n z}}{2\lambda_n^2}$$

Then, the solution becomes

$$T(r, z) - T_0 = \frac{qz}{kr_0} \sum_{n=1}^{\infty} \frac{\bar{e}^{\lambda_n z}}{\lambda_n^2} \frac{J_0(\lambda_n r)}{J_1(\lambda_n r_0)}, \quad \beta = \lambda_n, n=1, 2, 3, \dots$$

PROB. 5.49: Formulation in terms of  $\phi(r, \phi) = T(r, \phi) - T_f$ :

$$\frac{\partial^2 \phi}{\partial r^2} + \frac{1}{r} \frac{\partial \phi}{\partial r} + \frac{1}{r^2} \frac{\partial^2 \phi}{\partial \phi^2} = 0$$

$$k \left. \frac{\partial \phi}{\partial r} \right|_{r=r_i} = h \phi(r_i, \phi); \quad k \left. \frac{\partial \phi}{\partial r} \right|_{r=r_o} = Q''_w(\phi) = \begin{cases} Q''_w, & 0 < \phi < \pi \\ 0, & \pi < \phi < 2\pi \end{cases}$$

$$\phi(r, \phi) = \phi(r, \phi + 2\pi); \quad \frac{\partial \phi(r, \phi)}{\partial \phi} = \frac{\partial \phi(r, \phi + 2\pi)}{\partial \phi}$$

Seeking a product solution in the form  $\phi(r, \phi) = R(r) \cdot \psi(\phi)$ ,

$$\lambda_n = n = 0, 1, 2, \dots$$

$$\psi_n(\phi) = \begin{cases} A_0 + B_0 \phi, & n=0 \\ A_n \sin n\phi + B_n \cos n\phi, & n=1, 2, 3, \dots \end{cases}$$

$$R_n(r) = \begin{cases} C_0 \ln r + D_0, & n=0 \\ C_n r^n + D_n r^{-n}, & n=1, 2, 3, \dots \end{cases}$$

The general solution may now be constructed as

$$\begin{aligned} \phi(r, \phi) &= (C_0 \ln r + D_0)(A_0 + B_0 \phi) \\ &\quad + \sum_{n=1}^{\infty} (C_n r^n + D_n r^{-n})(A_n \sin n\phi + B_n \cos n\phi) \end{aligned}$$

Note that  $\phi(r, \phi)$  must be single valued. Thus,  $B_0 \equiv 0$ , and the general solution can be re-arranged as

$$\begin{aligned} \phi(r, \phi) &= (a_0 + b_0 \ln r) \\ &\quad + \sum_{n=1}^{\infty} [(a_n r^n + b_n r^{-n}) \sin n\phi + (c_n r^n + d_n r^{-n}) \cos n\phi] \end{aligned}$$

B.C. at  $r=r_i$  gives,

$$\begin{aligned} \frac{k}{h} \left\{ \frac{b_0}{r_i} + \sum_{n=1}^{\infty} [n a_n r_i^{(n-1)} - n b_n r_i^{-(n+1)}] \sin n\phi + [n c_n r_i^{(n-1)} - n d_n r_i^{-(n+1)}] \cos n\phi \right\} \\ = (a_0 + b_0 \ln r_i) + \sum_{n=1}^{\infty} [(a_n r_i^n + b_n \bar{r}_i^n) \sin \phi + (c_n r_i^n + d_n \bar{r}_i^n) \cos \phi] \end{aligned}$$

Thus,

$$\frac{k}{h} \frac{b_0}{r_i} = a_0 + b_0 \ln r_i \quad \dots \quad (1)$$

$$(r_i^n - n \frac{k}{h} r_i^{(n-1)}) a_n + (\bar{r}_i^n + n \frac{k}{h} \bar{r}_i^{(n+1)}) b_n = 0 \quad \dots \quad (2)$$

$$(r_i^n - n \frac{k}{h} r_i^{(n-1)}) c_n + (\bar{r}_i^n + n \frac{k}{h} \bar{r}_i^{(n+1)}) d_n = 0 \quad \dots \quad (3)$$

B.C. at  $r=r_o$  gives,

$$k \left\{ \frac{b_0}{r_o} + \sum_{n=1}^{\infty} [n a_n r_o^{(n-1)} - n b_n r_o^{-(n+1)}] \sin n\phi + [n c_n r_o^{(n-1)} - n d_n r_o^{-(n+1)}] \cos n\phi \right\} = Q''_w(\phi)$$

From this we obtain:

$$\frac{k b_0}{r_o} = \frac{1}{2\pi} \int_0^{2\pi} Q''_w(\phi) d\phi = \frac{Q''_w}{2\pi} \int_0^\pi d\phi = \frac{Q''_w}{2} \quad \dots \quad (4)$$

$$k [n a_n r_o^{(n-1)} - n b_n r_o^{-(n+1)}] = \frac{1}{\pi} \int_0^{2\pi} Q''_w(\phi) \sin n\phi d\phi = \dots = \frac{Q''_w}{\pi} \frac{[1 - (-1)^n]}{n} \quad (5)$$

$$k [n c_n r_o^{(n-1)} - n d_n r_o^{-(n+1)}] = \frac{1}{\pi} \int_0^{2\pi} Q''_w(\phi) \cos n\phi d\phi = \dots = 0 \quad (6)$$

Equations (1) & (4) give,

$$b_0 = \frac{Q''_w r_o}{2k} \quad \text{and} \quad a_0 = \frac{Q''_w r_o}{2k} \left( \ln r_i - \frac{k}{h r_i} \right)^{-1}$$

Equations (2), (3), (5) & (6) yield, for each  $n=1, 2, 3, \dots$ , a set of algebraic set of equation, the solution of which yields

$a_n, b_n, c_n$  and  $d_n$ ,  $n=1, 2, 3, \dots$

# CHAPTER 6

## UNSTEADY HEAT CONDUCTION

### SOLUTIONS WITH SEPARATION OF VARIABLES

---

PROB. 6.1: At any instant  $t$ :

$$\dot{q}V + hA[T_{\infty} - T(t)] = C\rho V \frac{dT}{dt}$$

$$\text{Let } \theta(t) = T(t) - T_{\infty}$$

$$\frac{d\theta}{dt} + \frac{hA}{\rho V C} \theta(t) = \frac{\dot{q}}{\rho C} = \frac{\dot{q}_0}{\rho C} e^{-\beta t}$$

$$T(0) = T_{\infty} \Rightarrow \theta(0) = 0$$

$$\therefore \theta(t) = C_1 e^{mt} + \theta_p, \quad m = \frac{hA}{\rho V C}$$

$$\text{Particular solution: } \theta_p = C_2 e^{\beta t} \rightarrow -C_2 \beta e^{\beta t} + m C_2 e^{\beta t} = \frac{\dot{q}_0}{\rho C} e^{\beta t}$$

$$\theta(t) = C_1 e^{mt} + \frac{\dot{q}_0}{\rho C(m-\beta)} e^{\beta t} \rightarrow C_2 = \frac{\dot{q}_0}{\rho C(m-\beta)}$$

$$\theta(0) = 0 \Rightarrow C_1 + \frac{\dot{q}_0}{\rho C(m-\beta)} = 0 \rightarrow C_1 = -\frac{\dot{q}_0}{\rho C(m-\beta)}$$

$$\theta(t) = T(t) - T_{\infty} = \frac{\dot{q}_0}{\rho C(m-\beta)} [e^{\beta t} - e^{mt}], \quad \beta \neq m$$


$$\theta(t) = T(t) - T_{\infty} = \frac{\dot{q}_0}{\rho C} t e^{\beta t}, \quad \beta = m \text{ (why?)}$$

The maximum temperature will be reached when

$$\frac{dT}{dt} = 0 \Rightarrow t = \begin{cases} \frac{1}{m-\beta} \ln \frac{m}{\beta}, & \beta \neq m \\ \frac{1}{\beta}, & \beta = m \end{cases}$$

and it will be


$$T_{\max} - T_{\infty} = \begin{cases} \frac{\dot{q}_0}{\rho C(m-\beta)} \left[ \left(\frac{m}{\beta}\right)^{\frac{m}{m-\beta}} - \left(\frac{m}{\beta}\right)^{-\frac{m}{m-\beta}} \right], & \beta \neq m \\ \frac{\dot{q}_0}{\rho C} \frac{1}{\beta e^{\beta}}, & \beta = m \end{cases}$$



PROB. 6.2:

a) Energy balance:

$$\left\{ \begin{array}{l} h A_s (T_f - T) = \rho c A_s \delta \frac{dT}{dt} \\ \frac{dT}{dt} + \frac{h}{\rho c \delta} (T - T_i - At) = 0 \\ T(0) = T_i \end{array} \right\}$$



$$\left[ \begin{array}{l} \frac{d\theta}{d\tau} + \theta = \tau, \quad \theta = \frac{T - T_i}{mA}, \quad \tau = \frac{t}{m} \quad \& \quad m = \frac{\rho c \delta}{h} \\ \theta(0) = 0 \end{array} \right]$$

$$\begin{aligned} \frac{d}{d\tau}(e^\tau \theta) &= e^\tau \tau \quad \rightarrow e^\tau \theta(\tau) - \theta(0) = \int_0^\tau \tau e^\tau d\tau \\ &= \tau e^\tau \Big|_0^\tau - \int_0^\tau e^\tau d\tau \\ &= \tau e^\tau - e^\tau + 1 \\ \therefore \theta(\tau) &= \frac{\tau - 1 + e^{-\tau}}{e^\tau} \end{aligned}$$

b) For Large times as  $t \rightarrow \infty$ :

$$\theta(\tau) = \tau - 1$$

$$\therefore \frac{T - T_i}{mA} = \frac{t}{m} - 1$$

PROB. 6.3: The solution for  $T = T(x, t)$  is given, from Eq. (6.23), by

$$T(x, t) = T_1 - (T_1 - T_2) \frac{x}{L} + \sum_{n=1}^{\infty} A_n e^{-\left(\frac{n\pi}{L}\right)^2 \alpha t} \sin \frac{n\pi}{L} x$$

with, from Eq. (6.25),

$$\begin{aligned} A_n &= \frac{2}{n\pi} [(-1)^n T_2 - T_1] + \frac{2}{L} \int_0^L T_1 \sin \frac{n\pi}{L} x \, dx \\ &= \frac{2}{n\pi} [(-1)^n T_2 - T_1] + \frac{2}{n\pi} [1 - (-1)^n] T_1 \end{aligned}$$

Thus,


$$\begin{aligned} T(x, t) &= T_1 - (T_1 - T_2) \frac{x}{L} \\ &+ \frac{2}{\pi} \sum_{n=1}^{\infty} \frac{[1 - (-1)^n] T_1 + (-1)^n T_2 - T_1}{n} e^{-\left(\frac{n\pi}{L}\right)^2 \alpha t} \sin \frac{n\pi}{L} x \end{aligned}$$

PROB. 6.4: Formulation of the problem:

$$\frac{\partial^2 T}{\partial x^2} = \frac{1}{\alpha} \frac{\partial T}{\partial t}, \quad \alpha = \frac{k}{\rho c}$$

$$T(x, 0) = T_{\infty}$$

$$T(0, t) = T_f; \quad -k \left. \frac{\partial T}{\partial x} \right|_{x=L} = h [T(L, t) - T_{\infty}]$$



$$\text{Let } T(x, t) = T_t(x, t) + T_s(x)$$

$$\frac{\partial^2 T_t}{\partial x^2} = \frac{1}{\alpha} \frac{\partial T_t}{\partial t}$$

$$\frac{\partial^2 T_s}{\partial x^2} = 0$$

$$T_t(x, 0) = T_{\infty} - T_s(x)$$

$$T_s(0) = T_f$$

$$T_t(0, t) = 0; \quad -k \left. \frac{\partial T_t}{\partial x} \right|_{x=L} = h T_t(L, t)$$

$$-k \frac{d T_s(L)}{d x} = h [T_s(L) - T_{\infty}]$$

$$\therefore T_s(x) = T_f + Ax, \quad A = -\frac{T_f - T_\infty}{(1 + \frac{1}{Bi})} \frac{1}{L}, \quad Bi = \frac{hL}{k}$$

Now, Let  $T_f(x,t) = X(x) \cdot \Gamma(t)$ .

$$\frac{d^2X}{dx^2} + \lambda^2 X = 0 \quad \leftarrow \qquad \rightarrow \quad \frac{d\Gamma_n}{dt} + \alpha \lambda_n^2 \Gamma_n = 0$$

$$X(0) = 0$$

$$h X(L) + k \frac{dX(L)}{dx} = 0$$

$$\Gamma_n(t) = B_n e^{-\alpha \lambda_n^2 t}$$



$$X_n(x) = A_n \sin \lambda_n x$$

where  $\lambda_n$ 's are the positive roots of  $\tan \lambda L = -\frac{\lambda L}{Bi}$ ,  $Bi = \frac{hL}{k}$ .

$$\therefore T_t(x,t) = \sum_{n=1}^{\infty} a_n e^{-\alpha \lambda_n^2 t} \sin \lambda_n x$$



$$T_\infty - T_s(x) = \sum_{n=1}^{\infty} a_n \sin \lambda_n x$$

From Table 4.1.

$$\therefore a_n = \frac{1}{N_n} \int_0^L [T_\infty - T_s(x)] \sin \lambda_n x dx, \quad N_n = \frac{1}{2\lambda_n} [\lambda_n L - \sin \lambda_n L \cos \lambda_n L]$$


$$= \dots = \frac{T_\infty - T_f}{N_n \lambda_n} = \frac{2(T_\infty - T_f)}{[\lambda_n L - \sin \lambda_n L \cos \lambda_n L]}$$

⋮

$\frac{T(x,t) - T_f}{T_\infty - T_f} = \frac{Bi}{1+Bi} \frac{x}{L} + 2 \sum_{n=1}^{\infty} \frac{e^{-\alpha \lambda_n^2 t}}{[\lambda_n L - \sin \lambda_n L \cos \lambda_n L]} \sin \lambda_n x$
--

PROB. 6.5:

a) Formulation in terms of  $\theta(x,t) = T(x,t) - T_w$ :



$$\left\{ \begin{array}{l} \frac{\partial^2 \theta}{\partial x^2} = \frac{1}{\alpha} \frac{\partial \theta}{\partial t} \\ \theta(x,0) = T_i - T_w \\ \frac{\partial \theta(0,t)}{\partial x} = 0; \quad \theta(L,t) = 0 \end{array} \right\}$$

Let  $\theta(x,t) = X(x) \cdot \Gamma(t)$

$$\left\{ \begin{array}{l} \frac{d^2 X}{dx^2} + \lambda^2 X = 0 \\ \frac{dX(0)}{dx} = 0 \\ X(L) = 0 \end{array} \right.$$

$$X_n(x) = A_n \cos \lambda_n x$$

where  $\lambda_n$ 's are zeros of

$$\cos \lambda L = 0$$

$$\lambda_n = \frac{(2n-1)\pi}{2L}, \quad n=1, 2, 3, \dots$$

$$\frac{d\Gamma_n}{dt} + \alpha \lambda_n^2 \Gamma_n = 0$$

↓

$$\Gamma_n(t) = B_n e^{-\alpha \lambda_n^2 t}$$

$$\therefore \theta(x,t) = \sum_{n=1}^{\infty} a_n e^{-\alpha \lambda_n^2 t} \cos \lambda_n x$$

$$\Rightarrow T_i - T_w = \sum_{n=1}^{\infty} a_n \cos \lambda_n x$$

$$\begin{aligned} a_n &= \frac{2}{L} (T_i - T_w) \int_0^L \cos \lambda_n x \, dx = \frac{2(T_i - T_w)}{L} \frac{\sin \lambda_n x}{\lambda_n} \Big|_0^L \\ &= \frac{4(T_i - T_w)(-1)^{n+1}}{(2n-1)\pi} \end{aligned}$$

$$\frac{\theta(x,t)}{T_i - T_w} = \frac{T(x,t) - T_w}{T_i - T_w} = \frac{4}{\pi} \sum_{n=1}^{\infty} \frac{(-1)^{n+1}}{2n-1} e^{-\alpha \lambda_n^2 t} \cos \lambda_n x$$

b)  $T_m / g / L A = \int_0^L T / g / A dx \Rightarrow T_m(t) = \frac{1}{L} \int_0^L T(x, t) dx$

$$\boxed{\frac{T_m(t) - T_w}{T_i - T_w} = \frac{8}{\pi^2} \sum_{n=1}^{\infty} \frac{-\alpha \lambda_n^2 t}{(2n-1)^2}}$$

$$\Downarrow \quad \Leftarrow \frac{T_m(t) - T_w}{T_i - T_w} = \frac{1}{L} \int_0^L \theta(x, t) dx$$


c)  $q(t) = -k A \left. \frac{\partial T}{\partial x} \right|_{x=L} \Rightarrow$

$$\boxed{q(t) = 2kA \frac{T_i - T_w}{L} \sum_{n=1}^{\infty} e^{-\alpha \lambda_n^2 t}}$$

d)  $Q(t) = \int_0^t q(t) dt \quad \downarrow$

$$\boxed{\frac{Q(t)}{Q_i} = \frac{8}{\pi^2} \sum_{n=1}^{\infty} \frac{1 - e^{-\alpha \lambda_n^2 t}}{(2n-1)^2}, \quad Q_i = \text{sc AL}(T_i - T_w)}$$

PROB. 6.6:



a) Formulation of the problem in terms of

$$\theta(x, t) = T(x, t) - T_w :$$

$$\frac{\partial^2 \theta}{\partial x^2} = \frac{1}{\alpha} \frac{\partial \theta}{\partial t}$$

$$\theta(x, 0) = \theta_i$$

$$\boxed{\theta_L' = T_L - T_w}$$

$$\boxed{\theta(0, t) = 0; \theta(L, t) = \theta_L'}$$

$$\text{Let } \theta(x, t) = \theta_t(x, t) + \theta_s(x)$$

$$\frac{\partial^2 \theta_t}{\partial x^2} = \frac{1}{\alpha} \frac{\partial \theta_t}{\partial t} \quad \leftarrow \quad \rightarrow \quad \frac{d^2 \theta_s}{dx^2} = 0$$

$$\theta_t(x, 0) = \theta_i - \theta_s(x)$$

$$\theta_s(0) = 0; \theta_s(L) = \theta_L'$$

$$\theta_t(0, t) = 0; \theta_t(L, t) = 0$$

$$\therefore \boxed{\theta_s(x) = \theta_i \frac{x}{L}}$$

$$\therefore \theta_t(x,t) = \sum_{n=1}^{\infty} A_n e^{-\alpha \lambda_n^2 t} \sin \lambda_n x ; \quad \lambda_n = \frac{n\pi}{L}, \quad n=1,2,3,\dots$$



$$\theta_i \left[ 1 - \frac{x}{L} \right] = \sum_{n=1}^{\infty} A_n \sin \lambda_n x \Rightarrow A_n = \frac{2\theta_i}{L} \int_0^L \left[ 1 - \frac{x}{L} \right] \sin \lambda_n x dx = \frac{2\theta_i}{n\pi}$$

$$\therefore \boxed{\frac{\theta(x,t)}{\theta_i} = \frac{x}{L} + \frac{2}{\pi} \sum_{n=1}^{\infty} \frac{1}{n} e^{-\alpha \lambda_n^2 t} \sin \lambda_n x}$$

b)

$$\cancel{\int_0^L \theta(x,t) dx} = \int_0^L T_m dx \Rightarrow T_m = \frac{1}{L} \int_0^L T dx$$

$$\therefore \boxed{\frac{T_m - T_w}{T_i - T_w} = \frac{1}{2} + \frac{L}{\pi^2} \sum_{n=1}^{\infty} \frac{1}{n^2} [1 - (-1)^n] e^{-\alpha \lambda_n^2 t}}$$

c)

$$q(0,t) = -k \left. \frac{\partial T}{\partial x} \right|_{x=0} = -\frac{k\theta_i}{L} \left[ 1 + 2 \sum_{n=1}^{\infty} e^{-\alpha \lambda_n^2 t} \right]$$

$$q(L,t) = -k \left. \frac{\partial T}{\partial x} \right|_{x=L} = -\frac{k\theta_i}{L} \left[ 1 + 2 \sum_{n=1}^{\infty} (-1)^n e^{-\alpha \lambda_n^2 t} \right]$$

PROB. 6.7: Formulation in terms of  $\theta(r,t) = T(r,t) - T_w$ :

$$\frac{\partial^2 \theta}{\partial r^2} + \frac{1}{r} \frac{\partial \theta}{\partial r} = \frac{1}{\alpha} \frac{\partial \theta}{\partial t}, \quad \alpha = \frac{k}{\rho c}$$

$$\theta(r,0) = T_i - T_w$$

$$\theta(0,t) = \text{finite}; \quad \theta(r_0, t) = 0$$

$$\text{Let } \theta(r,t) = R(r) \cdot \Gamma(t)$$



$$r^2 \frac{d^2 R}{dr^2} + r \frac{dR}{dr} + \lambda^2 r^2 R = 0 \quad | \quad \frac{d \Gamma_n}{dt} + \alpha \lambda_n^2 \Gamma_n = 0$$

$$R(0) = \text{finite}$$

$$R(r_0) = 0$$

$$\Gamma_n(t) = B_n e^{-\alpha \lambda_n^2 t}$$



$$R_n(r) = A_n J_0(\lambda_n r), \text{ where } \lambda_n \text{'s are zeros of } J_0(\lambda r_0) = 0.$$



$$\Theta(r, t) = \sum_{n=1}^{\infty} a_n e^{-\alpha \lambda_n^2 t} J_0(\lambda_n r)$$

$$\therefore T_i - T_w = \sum_{n=1}^{\infty} a_n J_0(\lambda_n r)$$

$$\Rightarrow a_n = \frac{1}{N_n} \int_0^{r_0} (T_i - T_w) J_0(\lambda_n r) r dr = \frac{T_i - T_w}{N_n} \frac{r_0}{\lambda_n} J_1(\lambda_n r_0); N_n = \frac{r_0^2}{2} J_1^2(\lambda_n r_0)$$

$$\therefore \boxed{\frac{\Theta(r, t)}{T_i - T_w} = \frac{T(r, t) - T_w}{T_i - T_w} = \frac{2}{r_0} \sum_{n=1}^{\infty} \frac{J_0(\lambda_n r)}{\lambda_n J_1(\lambda_n r_0)} e^{-\alpha \lambda_n^2 t}}$$

b)

$$\pi r_0^2 g_C T_m = \int_0^{r_0} g_C T 2\pi r dr \Rightarrow T_m = \frac{2}{r_0^2} \int_0^{r_0} T r dr$$

$$\Rightarrow \boxed{\frac{T_m - T_w}{T_i - T_w} = \frac{4}{r_0^2} \sum_{n=1}^{\infty} \frac{e^{-\alpha \lambda_n^2 t}}{\lambda_n^2}}$$

c)

$$q' = -k 2\pi r_0 \left( \frac{\partial T}{\partial r} \right)_{r=r_0} = 4\pi k (T_i - T_w) \sum_{n=1}^{\infty} e^{-\alpha \lambda_n^2 t}$$

d)

$$Q = \int_0^t q' dt \Rightarrow \frac{Q}{Q_i} = \frac{4}{r_0^2} \sum_{n=1}^{\infty} \frac{1 - e^{-\alpha \lambda_n^2 t}}{\lambda_n^2}, Q_i = \pi r_0^2 g_C (T_i - T_\infty)$$


PROB. 6.11:

$$2L = 1 \text{ in.} \Rightarrow L = 0.5 \text{ in}$$

$$\alpha = 8.8 \times 10^{-8} \text{ m}^2/\text{s}$$

Since the surface temperature is specified,

$$Bi \rightarrow \infty \quad \text{or} \quad \frac{1}{Bi} = 0$$



a)

$$\frac{T_c - 140}{20 - 140} = \frac{130 - 140}{20 - 140} = 0.083 \quad \Rightarrow \text{from Fig. 6.9} \quad \frac{\alpha t}{L^2} = 1.1$$

$$\begin{aligned} \therefore t &= 1.1 \frac{L^2}{\alpha} = 1.1 \times \frac{(0.5 \times 2.54 \times 10^{-2})^2}{8.8 \times 10^{-8}} \\ &= 2016 \text{ sec} = \underline{33 \text{ min } 36 \text{ sec}} \end{aligned}$$

b)

$$\frac{T_p - 140}{T_c - 140} \approx 0.72 \quad \leftarrow \text{From Fig. 6.10 with } \frac{x}{L} = 0.5$$

$$\begin{aligned} \therefore T_p &= 140 + 0.72 (130 - 140) = 140 - 7.2 \\ &= \underline{132.8^\circ C} \end{aligned}$$

c)

$$\frac{T_p - 140}{T_c - 140} = \frac{130 - 140}{T_c - 140} = 0.72 \quad \Rightarrow T_c = 140 - \frac{10}{0.72} = 126.1^\circ C$$

$$\frac{T_c - 140}{20 - 140} = \frac{13.9}{120} = 0.116 \quad \Rightarrow \text{from Fig. 6.9} \quad \frac{\alpha t}{L^2} = 1$$

$$\therefore t = \frac{L^2}{\alpha} = 1833 \text{ sec} = \underline{30 \text{ min } 32 \text{ sec}}$$

PROB. 6.14: Since  $T_0 = T_s$  at all times, the copper sphere is a lumped heat capacity system. Thus,

$$\frac{T - T_{\infty}}{T_i - T_{\infty}} = \exp\left(-\frac{hA}{gCV} t\right) \rightarrow h = -\frac{gCV}{At} \ln \frac{T - T_{\infty}}{T_i - T_{\infty}}$$

At  $t = 5 \text{ min}$ ,  $T = 50^\circ\text{C}$ .

$$\therefore h = -\frac{(8990)(381)(\frac{4}{3}\pi r_0^3)}{(4\pi r_0^2)(300)} \ln \frac{50-20}{100-20}$$

$$= 112 \text{ W/(m}^2\text{.K)}$$

At  $t = 10 \text{ min}$ ,  $T = 31.2^\circ\text{C}$ .

$$\therefore h = -\frac{(\quad)(\quad)(\quad)}{(\quad)(600)} \ln \frac{31.2-20}{100-20}$$

$$= 112.25 \text{ W/(m}^2\text{.K)}$$

$$\underline{h \approx 112.12 \text{ W/(m}^2\text{.K)}}$$

Now, consider the rubber sphere : The same  $h$  applies. (Why?)

$$Bi = \frac{h r_0}{k} = \frac{(112.12)(0.03)}{0.43} = 7.8 \leftarrow \text{Distributed system}$$

$$Fo = \frac{\alpha t}{r_0^2} = \frac{(0.43)}{(1070)(1630)} \times \frac{(30 \times 60)}{(0.03)^2} = 0.493$$


$$\left. \begin{array}{l} Fo = 0.493 \\ 1/Bi = 0.128 \end{array} \right\} \xrightarrow{\text{Fig. 6.16}} \frac{T_c - T_{\infty}}{T_i - T_{\infty}} = 0.045$$

$$\therefore T_c = 0.045(100-20) + 20 = \underline{23.6^\circ\text{C}}$$

$$\text{Fig. 6.17} \implies \frac{T_s - T_{\infty}}{T_c - T_{\infty}} = 0.25$$

$$\therefore T_s = 0.25(23.6 - 20) + 20 = \underline{20.9^\circ\text{C}}$$

PROB. 6.16: Consider a control volume of infinitesimal dimensions  $\Delta x$ ,  $\Delta y$  and  $\Delta z$  as shown.



where

$$q_{x1x} = -k \frac{\partial T}{\partial x} \Delta y \Delta z + T c g V \Delta y \Delta z \stackrel{\text{Negligible}}{\approx} T c g V \Delta y \Delta z$$

$$q_{y1y} = -k \Delta x \Delta z \frac{\partial T}{\partial y}$$

$$q_{z1z} = -k \Delta x \Delta y \frac{\partial T}{\partial z}$$

Energy balance on the c.v. gives:

$$\begin{aligned} q_{x1x} + q_{y1y} + q_{z1z} &= q_{x1x+\Delta x} + q_{y1y+\Delta y} + q_{z1z+\Delta z} \\ &= q_{x1x} + \frac{\partial q_{x1x}}{\partial x} \Delta x + q_{y1y} + \frac{\partial q_{y1y}}{\partial y} \Delta y + q_{z1z} + \frac{\partial q_{z1z}}{\partial z} \Delta z \\ &\Downarrow \\ \frac{\partial^2 T}{\partial y^2} + \frac{\partial^2 T}{\partial z^2} &= \frac{V}{\alpha} \frac{\partial T}{\partial x} = \frac{V}{\alpha} \frac{\partial T}{\partial t} \frac{\partial}{\partial x} = \frac{1}{\alpha} \frac{\partial T}{\partial t} \end{aligned}$$

Note that  $T(x, y, z) \equiv T(y, z, t)$  with  $x = Vt$ .

Thus,

$$\underbrace{\left\{ \frac{T(y, z, t) - T_{\infty}}{T_i - T_{\infty}} \right\}_{bar}}_{\Phi(y, z, t)} = \underbrace{\left\{ \frac{T(y, t) - T_{\infty}}{T_i - T_{\infty}} \right\}_{slab}}_{\Phi_1(y, t)} \times \underbrace{\left\{ \frac{T(z, t) - T_{\infty}}{T_i - T_{\infty}} \right\}_{slab}}_{\Phi_2(z, t)}$$

$$\underbrace{\frac{150-25}{500-25}}_{\phi(0,0,t)} = 0.263 = \underbrace{\left[ \frac{T(0,t)-25}{500-25} \right]_{\text{slab}}}_{\phi_1(0,t)} \times \underbrace{\left[ \frac{T(0,t)-25}{500-25} \right]_{\text{slab}}}_{\phi_2(0,t)}^{2b}$$

Also

$$\alpha = \frac{k}{\rho c} = \frac{230}{2707 \times 896} = 9.48 \times 10^{-5} \text{ m}^2/\text{s}$$

$$\frac{k}{h a} = \frac{230}{5000 \times \underbrace{2.5 \times 10^{-2}}_a} = 1.84 \quad \text{and} \quad \frac{k}{h b} = \frac{3.68}{\underbrace{1.25 \times 10^{-2}}_b} = 3.68$$

Construct the following iteration table using Fig. 6.9:

$t$ (sec)	$\frac{\alpha t}{a^2}$	$\phi_1(0,t)$	$\frac{\alpha t}{b^2}$	$\phi_2(0,t)$	$\phi_1 \cdot \phi_2$
10	1.52	0.52	6.07	0.24	0.1248
5	0.76	0.8	3.04	0.41	0.328
7	1.06	0.7	4.25	0.37	0.259 $\cong 0.263$

$$\therefore t \cong 7 \text{ sec} \Rightarrow L \cong 0.5 \times 7 = 3.5 \text{ m}$$

b) The maximum surface temperature that can be reached after the bars emerge from the cooling tank would be the mean temperature at  $L = 3.5 \text{ m}$  ( $t = 7 \text{ sec}$ ). Since the cross-section is rectangular,

$$\bar{T}(t) = \frac{1}{ab} \int_0^a \int_0^b T(y, z, t) dy dz \rightarrow \underbrace{\frac{\bar{T}(t) - T_{\infty}}{T_i - T_{\infty}}}_{\bar{\Phi}(t)} = \frac{1}{ab} \int_0^a \int_0^b \phi dy dz$$

$$\therefore \bar{\Phi}(t) = \underbrace{\frac{1}{a} \int_0^a \phi_1(y, t) dy}_{\bar{\Phi}_1(t)} \cdot \underbrace{\frac{1}{b} \int_0^b \phi_2(z, t) dz}_{\bar{\Phi}_2(t)} = \bar{\Phi}_1(t) \cdot \bar{\Phi}_2(t)$$

Now construct the following "numerical integration" table by making use of Fig. 6.10:

$\frac{y}{a}$ or $\frac{z}{b}$	$\phi_1(y, t)/\phi_1(0, t)$ *	$\phi_2(z, t)/\phi_2(0, t)$ **	$\phi_1(y, t)$	$\phi_2(z, t)$
0.0	1.0	1.0	0.7	0.37
0.2	0.98	0.99	0.686	0.366
0.4	0.96	0.97	0.672	0.359
0.6	0.92	0.95	0.644	0.352
0.8	0.84	0.93	0.588	0.344
1.0	0.78	0.88	0.546	0.326

Means:

$$\bar{\phi}_1 \approx \frac{1}{5} \left[ \frac{0.7+0.686}{2} + \frac{0.686+0.672}{2} + \dots + \frac{0.588+0.546}{2} \right] = 0.643$$

$$\bar{\phi}_2 \approx \frac{1}{5} \left[ \frac{0.37+0.366}{2} + \frac{0.366+0.359}{2} + \dots + \frac{0.344+0.326}{2} \right] = 0.354$$

$$\therefore \bar{\phi} = 0.643 \times 0.354 = 0.228$$


$$\Rightarrow T_m = 0.228(T_i - T_\infty) + T_\infty = 0.228(500 - 25) + 25 \approx \underline{\underline{133^\circ C}}$$

\* Read from Fig. 6.10 with  $\frac{k}{ha} = 1.84$ . Also note that  $\phi_1(0, t) = 0.7$  from Part (a).

\*\* Read from Fig. 6.10 with  $\frac{k}{hb} = 3.68$ . Also note that  $\phi_2(0, t) = 0.37$  from Part (a)

PROB. 6.19:

Formulation of the problem:



$$\frac{\partial^2 T}{\partial z^2} = \frac{1}{\alpha} \frac{\partial T}{\partial t}$$

$$T(z, 0) = T_1 + \frac{T_2 - T_1}{H} z$$

$$\frac{\partial T(0, t)}{\partial z} = 0; T(H, t) = T_2$$

Note that the temperature distribution will not depend on  $r$ . (why?)

$$\therefore T(z, t) - T_2 = \sum_{n=1}^{\infty} A_n e^{-\alpha \lambda_n^2 t} \cos \lambda_n z$$

where  $\lambda_n$ 's are the positive roots of  $\cos \lambda_n H = 0$ .

$$\therefore \lambda_n = \frac{2n-1}{H} \frac{\pi}{2}, n = 1, 2, 3, \dots$$

Applying the initial condition yields

$$(T_1 - T_2) \left(1 - \frac{z}{H}\right) = \sum_{n=1}^{\infty} A_n \cos \lambda_n z$$

$$\Rightarrow A_n = \frac{2}{H} (T_1 - T_2) \int_0^H \left(1 - \frac{z}{H}\right) \cos \lambda_n z dz = \frac{2(T_1 - T_2)}{H^2 \lambda_n^2}$$

$$\therefore \boxed{\frac{T(z, t) - T_2}{T_1 - T_2} = \frac{2}{H^2} \sum_{n=1}^{\infty} \frac{1}{\lambda_n^2} e^{-\alpha \lambda_n^2 t} \cos \lambda_n z}$$

PROB. 6.20: Formulation of the problem:

$$\left\{ \begin{array}{l} \frac{\partial^2 T}{\partial x^2} = \frac{1}{\alpha} \frac{\partial T}{\partial t} \\ T(x, 0) = T_i \\ \left. \frac{\partial T}{\partial x} \right|_{x=0} = 0 ; \quad k \left. \frac{\partial T}{\partial x} \right|_{x=L} = \frac{q''_w}{k} \end{array} \right\}$$

$$\text{Let } T(x, t) = \psi(x, t) + \phi(x) + \Gamma(t)$$

$\frac{\partial^2 \psi}{\partial x^2} = \frac{1}{\alpha} \frac{\partial \psi}{\partial t}$   
 $\psi(x, 0) = -\phi(x)$   
 $\left. \frac{\partial \psi(0, t)}{\partial x} \right|_0 = 0 ; \quad \left. \frac{\partial \psi(L, t)}{\partial x} \right|_L = 0$

$\downarrow$   
 $\downarrow$   
 $\downarrow$   
 $\downarrow$

$\frac{d^2 \phi}{dx^2} = \frac{1}{\alpha} \frac{d \phi}{dt} = \text{const} = C$   
 $\Gamma(0) = T_i$   
 $\left. \frac{d \phi(0)}{dx} \right|_0 = 0 ; \quad \left. \frac{d \phi(L)}{dx} \right|_L = \frac{q''_w}{k}$

$$\frac{d \Gamma}{dt} = \alpha C \rightarrow \Gamma(t) = \alpha C t + D \quad \& \quad \Gamma(0) = T_i \rightarrow D = T_i$$

$$\therefore \Gamma(t) = \alpha C t + T_i$$

$$\frac{d^2 \phi}{dx^2} = C \rightarrow \phi(x) = \frac{C x^2}{2} + A x + B$$

$$\left. \frac{d \phi(0)}{dx} \right|_0 = 0 \rightarrow A = 0$$

$$\left. \frac{d \phi(L)}{dx} \right|_L = \frac{q''_w}{k} \rightarrow C = \frac{q''_w}{k L}$$

Thus,

$$\phi(x) = \frac{q''_w}{2 k L} x^2 + B$$

$$\Gamma(t) = \frac{\alpha q''_w}{k L} t + T_i$$

On the other hand, let  $\psi(x,t) = X(x) \cdot \Sigma(t)$ .

$$\left\{ \begin{array}{l} \frac{d^2X}{dx^2} + \lambda^2 X = 0 \\ \frac{dX(0)}{dx} = 0 \\ \frac{dX(L)}{dx} = 0 \end{array} \right\} \quad \begin{array}{l} X_n(x) = a_n \cos \lambda_n x \\ \text{where} \\ \lambda_n = \frac{n\pi}{L}, n=0,1,2,\dots \end{array} \quad \begin{array}{l} \frac{d\Sigma}{dt} + \alpha \lambda_n^2 \Sigma_n = 0 \\ \downarrow \\ \Sigma_n(t) = b_n e^{-\alpha \lambda_n^2 t} \end{array}$$

$$\therefore \psi(x,t) = C_0 + \sum_{n=1}^{\infty} C_n e^{-\alpha \lambda_n^2 t} \cos \lambda_n x$$



$$-\phi(x) = -\frac{q''_w}{2kL} x^2 - B = C_0 + \sum_{n=1}^{\infty} C_n \cos \lambda_n x$$



$$C_0 + B = \frac{1}{N_0} \int_0^L \left( -\frac{q''_w}{2kL} x^2 \right) dx, \quad N_0 = L$$

$$= -\frac{q''_w L}{6k}$$


$$C_n = \frac{1}{N_n} \int_0^L \left( -\frac{q''_w}{2kL} x^2 \right) \cos \lambda_n x dx, \quad N_n = \frac{L}{2}, n=1,2,\dots$$

$$= \dots = -\frac{2q''_w L}{k} \frac{(-1)^n}{(n\pi)^2}$$

⋮

$$T(x,t) - T_i = \frac{q''_w L}{k} \left[ \frac{\alpha t}{L^2} + \frac{1}{2} \left( \frac{x}{L} \right)^2 - \frac{1}{6} - \frac{2}{\pi L^2} \sum_{n=1}^{\infty} \frac{(-1)^n}{n^2} e^{-\alpha \lambda_n^2 t} \cos \lambda_n x \right]$$

PROB. 6.21:



Formulation of the problem:

$$\frac{\partial^2 T}{\partial r^2} + \frac{1}{r} \frac{\partial T}{\partial r} = \frac{1}{\alpha} \frac{\partial T}{\partial t}$$

$$T(r, 0) = T_i$$

$$\left. \frac{\partial T}{\partial r} \right|_{r=0} = 0; \quad k \left( \frac{\partial T}{\partial r} \right)_{r=r_0} = q''_w$$

Based on the physics of the problem, let

$$T(r, t) = \psi(r, t) + \phi(r) + \Omega(t)$$

which yields

$$\frac{\partial^2 \psi}{\partial r^2} + \frac{1}{r} \frac{\partial \psi}{\partial r} = \frac{1}{\alpha} \frac{\partial \psi}{\partial t} \quad ; \quad \frac{d^2 \phi}{dr^2} + \frac{1}{r} \frac{d\phi}{dr} = \frac{1}{\alpha} \frac{d\Omega}{dt} = \text{const} = C_1$$

$$\psi(r, 0) = -\phi(r)$$

$$\Omega(0) = T_i$$

$$\left. \frac{\partial \psi(r_0, t)}{\partial r} \right|_0 = 0; \quad \left. \frac{\partial \psi(r_0, t)}{\partial r} \right|_0 = 0 \quad ; \quad \left. \frac{d\phi(r)}{dr} \right|_0 = 0; \quad \left. \frac{d\phi(r_0)}{dr} \right|_0 = \frac{q''_w}{k}$$

$$\frac{d\Omega}{dt} = \alpha C_1 \rightarrow \Omega(t) = \alpha C_1 t + C_2 \quad \& \quad \Omega(0) = T_i \rightarrow C_2 = T_i$$

$$\Omega(t) = \alpha C_1 t + T_i$$

$$\frac{d^2 \phi}{dr^2} + \frac{1}{r} \frac{d\phi}{dr} = C_1 \rightarrow \phi(r) = C_1 \frac{r^2}{4} + C_3 \ln r + C_4$$

$$\left. \frac{d\phi(r)}{dr} \right|_0 = 0 \rightarrow C_3 = 0$$

$$\left. \frac{d\phi(r_0)}{dr} \right|_0 = \frac{q''_w}{k} \rightarrow C_1 = \frac{2q''_w}{r_0 k}$$

Thus,

$$\phi(r) = \frac{q''_w}{2r_0 k} r^2 + C_4$$

$$\Omega(t) = \frac{2\alpha q''_w}{r_0 k} t + T_i$$

$$\left\{ \begin{array}{l} \frac{\partial^2 \psi}{\partial r^2} + \frac{1}{r} \frac{\partial \psi}{\partial r} = \frac{1}{\alpha} \frac{\partial \psi}{\partial t} \\ \psi(r, 0) = -\frac{q_w''}{2r_0 k} r^2 - C_4 \\ \frac{\partial \psi(0, t)}{\partial r} = 0, \quad \frac{\partial \psi(r_0, t)}{\partial r} = 0 \end{array} \right\} \text{Let } \bar{\psi}(r, t) = \psi(r, t) + C_4 = R(r) \Gamma(t)$$

Thus,

$$\left\{ \begin{array}{l} \frac{d^2 R}{dr^2} + \frac{1}{r} \frac{dR}{dr} + \lambda^2 R = 0 \\ \frac{dR(0)}{dr} = 0, \quad \frac{dR(r_0)}{dr} = 0 \end{array} \right\} \quad R_n(r) = A_n J_0(\lambda_n r)$$

where  $\lambda_n$ 's are the positive roots of  $J_1(\lambda_n r_0) = 0, n = 0, 1, 2, \dots$

Note that  $\lambda_0 = 0$  is an eigenvalue.

$$\frac{d \Gamma_n}{dt} + \alpha \lambda_n^2 \Gamma_n = 0 \rightarrow \Gamma_n(t) = B_n e^{-\alpha \lambda_n^2 t}$$

$$\therefore \bar{\psi}(r, t) = C_0 + \sum_{n=1}^{\infty} C_n e^{-\alpha \lambda_n^2 t} J_0(\lambda_n r)$$

$$\Rightarrow -\frac{q_w''}{2r_0 k} r^2 = C_0 + \sum_{n=1}^{\infty} C_n J_0(\lambda_n r)$$

$$C_0 = -\frac{2}{r_0^2} \int_0^{r_0} \frac{q_w''}{2r_0 k} r^2 \cdot r dr = -\frac{q_w'' r_0}{4k}$$

$$C_n = -\frac{1}{N_n} \int_0^{r_0} \frac{q_w''}{2r_0 k} r^2 J_0(\lambda_n r) r dr = -\frac{2q_w''}{r_0 k \lambda_n^2 J_0(\lambda_n r_0)}, \quad N_n = \frac{r_0^2}{2} J_0^2(\lambda_n r_0)$$

$$\therefore T(r, t) = \psi(r, t) + \phi(r) + \varphi(t)$$

$$= -\frac{q_w'' r_0}{4k} - \frac{2q_w''}{r_0 k} \sum_{n=1}^{\infty} e^{-\alpha \lambda_n^2 t} \frac{J_0(\lambda_n r)}{\lambda_n^2 J_0(\lambda_n r_0)} - C_4 \\ + \frac{q_w''}{2r_0 k} r^2 + C_4 + \frac{2\alpha q_w''}{r_0 k} t + T_i$$

$$T(r, t) - T_i = \frac{q_w''}{k r_0} \left[ \frac{r^2}{2} + 2\alpha t - \frac{r_0^2}{4} - 2 \sum_{n=1}^{\infty} \frac{J_0(\lambda_n r)}{\lambda_n^2 J_0(\lambda_n r_0)} e^{-\alpha \lambda_n^2 t} \right]$$

PROB. 6.22: Formulation in terms of  $\theta(r,t) = T(r,t) - T_i$ :

$$\frac{\partial^2 \theta}{\partial r^2} + \frac{2}{r} \frac{\partial \theta}{\partial r} = \frac{1}{\alpha} \frac{\partial \theta}{\partial t}$$

$$\theta(r,0) = 0$$

$$\theta(0,t) = \text{finite}; \quad k \left. \frac{\partial \theta}{\partial r} \right|_{r=r_0} = q''_w$$

||

$$\underline{U(r,t) = r \cdot \theta(r,t)}$$



$$\frac{\partial^2 U}{\partial r^2} = \frac{1}{\alpha} \frac{\partial U}{\partial t}$$

$$U(r_0) = 0$$

$$U(0,t) = 0; \quad \left. \frac{\partial U}{\partial r} \right|_{r=r_0} = \frac{U(r_0,t)}{r_0} = r_0 \frac{q''_w}{k}$$

||

$$\underline{U(r,t) = \psi(r,t) + \phi(r) + r \Omega(t)}$$



$$\frac{\partial^2 \psi}{\partial r^2} = \frac{1}{\alpha} \frac{\partial \psi}{\partial t}$$

$$\left| \frac{1}{r} \frac{d^2 \phi}{dr^2} = \frac{1}{\alpha} \frac{d \Omega}{dt} = \text{const.} = C \right.$$

$$\psi(r_0) = -\phi(r)$$

$$\Omega(0) = 0$$

$$\psi(0,t) = 0; \quad \left. \frac{\partial \psi}{\partial r} \right|_{r=r_0} = \frac{\partial \psi(r_0,t)}{r_0} = 0$$

$$\left| \phi(0) = 0; \quad \frac{d \phi(r_0)}{dr} - \frac{\phi(r_0)}{r_0} = r_0 \frac{q''_w}{k} \right.$$

$$\Omega(t) = \alpha c t + A' \rightarrow \Omega(t) = \alpha c t$$

$$\phi(r) = C \frac{r^3}{6} + Br + B' \rightarrow \left( \frac{C}{2} r_0^2 + B' \right) - \left( \frac{Cr_0^2}{6} + B' \right) = r_0 \frac{q''_w}{k}$$

$$C = \frac{3q''_w}{kr_0}$$

$$\Omega(t) = \frac{3\alpha q''_{tw}}{k r_0} t$$

$$\phi(r) = \frac{3q''_{tw}}{kr_0} \frac{r^3}{6} + Br = \frac{q''_{tw} r^3}{2kr_0} + Br$$

$$\frac{\partial^2 \psi}{\partial r^2} = \frac{1}{\alpha} \frac{\partial \psi}{\partial t}$$

$$\psi(r, 0) = -\frac{q''_{tw} r^3}{2kr_0} - Br$$

$$\psi(0, t) = 0; \quad \left. \frac{\partial \psi}{\partial r} \right|_{r=r_0} - \frac{\partial \psi(r_0, t)}{r_0} = 0$$

$\Downarrow$   $\bar{\psi}(r, t) = \psi(r, t) + Br$

$$\frac{\partial^2 \bar{\psi}}{\partial r^2} = \frac{1}{\alpha} \frac{\partial \bar{\psi}}{\partial t}$$

$$\bar{\psi}(r, 0) = -\frac{q''_{tw} r^3}{2kr_0}$$

$$\bar{\psi}(0, t) = 0; \quad \left. \frac{\partial \bar{\psi}}{\partial r} \right|_{r=r_0} - \frac{\bar{\psi}(r_0, t)}{r_0} = 0$$

$\Downarrow$   
 $\bar{\psi}(r, t) = R(r) \Gamma(t)$

$\Downarrow$

$$\left\{ \begin{array}{l} \frac{d^2 R}{dr^2} + \lambda^2 R = 0 \\ R(0) = 0 \\ \frac{dR(r_0)}{dr} - \frac{R(r_0)}{r_0} = 0 \end{array} \right\}$$

$$\frac{d\Gamma_n}{dt} + \alpha \lambda_n^2 \Gamma_n = 0$$

$\downarrow$   
 $\Gamma_n(t) = C_n e^{-\alpha \lambda_n^2 t}$

Note that  $\lambda_0 = 0$  is an eigenvalue and the corresponding eigenfunction is (see Prob. 4.7)

$$R_0(r) = A_0 r .$$

When  $\lambda_n \neq 0$ ,

$$R_n(r) = A_n \sin \lambda_n r$$

where

$$\tan \lambda_n r_0 = \lambda_n r_0 , \quad \lambda_n = 1, 2, 3, \dots$$

$$\therefore \frac{\bar{\Psi}(r,t) = A_0 r + \sum_{n=1}^{\infty} A_n e^{-\alpha \lambda_n^2 t} \sin \lambda_n r}{A_0 c_0 \quad A_n c_n}$$

$$\bar{\Psi}(r_0) = -\frac{q_w'' r_0^3}{2kr_0} \Rightarrow -\frac{q_w'' r_0^3}{2kr_0} = A_0 r_0 + \sum_{n=1}^{\infty} A_n \sin \lambda_n r_0$$

$$A_0 = \frac{1}{N_0} \int_0^{r_0} \left( -\frac{q_w'' r^3}{2kr_0} \right) r dr , \quad N_0 = \int_0^{r_0} r^2 dr = \frac{r_0^3}{3}$$

$$= \dots = -\frac{3}{10} \frac{q_w'' r_0}{k}$$

$$A_n = \frac{1}{N_n} \int_0^{r_0} \left( -\frac{q_w'' r^3}{2kr_0} \right) \sin \lambda_n r dr , \quad N_n = \int_0^{r_0} \sin^2 \lambda_n r dr = \dots = \frac{r_0}{2} \sin^2 \lambda_n r_0$$

$$= \dots = -2 \frac{q_w''}{k} \frac{1}{\lambda_n^2 \sin \lambda_n r_0}$$

$$\vdots$$

$$\therefore T(r,t) - T_i = \frac{T(r,t)}{r} = \frac{q_w''}{kr_0} \left[ \frac{5r^2 - 3r_0^2}{10} + 3\alpha t - 2 \frac{r_0}{r} \sum_{n=1}^{\infty} \frac{\sin \lambda_n r}{\lambda_n^2 \sin \lambda_n r_0} e^{-\alpha \lambda_n^2 t} \right]$$

PROB. 6.23: Assume constant thermophysical properties ( $k, \rho, c$ ).

Formulation:

$$\frac{\partial^2 T}{\partial r^2} + \frac{2}{r} \frac{\partial T}{\partial r} = \frac{1}{\alpha} \frac{\partial T}{\partial t}$$

$$T(r, 0) = T_i$$

$$T(r_1, t) = 0 ; -k \left. \frac{\partial T}{\partial r} \right|_{r=r_2} = h T(r_2, t)$$

$$\text{Let } \underline{T(\eta, t) = r \cdot T(r, t) ; \eta = r - r_1} \\ \downarrow$$

$$\frac{\partial^2 U}{\partial \eta^2} = \frac{1}{\alpha} \frac{\partial U}{\partial t}$$

$$U(\eta, 0) = (\eta + r_1) T_i$$

$$U(0, t) = 0 ; \left[ k \frac{\partial U}{\partial \eta} + \left( h - \frac{k}{r_2} \right) U(\eta, t) \right]_{\eta=r_2-r_1} = 0$$

$$\therefore U(\eta, t) = \sum_{n=1}^{\infty} A_n e^{-\alpha \lambda_n^2 t} \sin \lambda_n \eta , \quad \text{where } \lambda_n \text{'s are the positive roots of}$$

$$\Rightarrow A_n = \frac{1}{N_n} \int_0^{r_2-r_1} (r_1 + \eta) T_i \cdot \sin \lambda_n \eta d\eta \quad \cot \lambda (r_2 - r_1) = -\frac{Bi-1}{\lambda r_2}$$

$$= \frac{T_i}{N_n} \left[ \frac{Bi}{\lambda_n^2} \sin \lambda_n (r_2 - r_1) + \frac{r_1}{\lambda_n} \right] \quad \text{with } Bi = \frac{h r_2}{k}$$

where

$$N_n = \frac{1}{2\lambda_n} \left[ \lambda_n (r_2 - r_1) - \sin \lambda_n (r_2 - r_1) \cos \lambda_n (r_2 - r_1) \right]$$

$$\therefore \boxed{T(r, t) = \frac{2T_i}{r} \sum_{n=1}^{\infty} \frac{[Bi \sin \lambda_n (r_2 - r_1) + r_1 \lambda_n] e^{-\alpha \lambda_n^2 t} \sin \lambda_n (r - r_1)}{\lambda_n [\lambda_n (r_2 - r_1) - \sin \lambda_n (r_2 - r_1) \cos \lambda_n (r_2 - r_1)]}}$$


PROB. 6.24: Formulation of the problem

in terms of  $\theta(x,t) = T(x,t) - T_w$ :

$$\frac{\partial^2 \theta}{\partial x^2} + \frac{\dot{q}}{k} = \frac{1}{\alpha} \frac{\partial \theta}{\partial t}$$

$$\theta(x,0) = T_i(x) - T_w \triangleq \theta_i(x)$$

$$\theta(0,t) = \theta(L,t) = 0$$



Let  $\theta(x,t) = \psi(x,t) + \phi(x)$

$$\left\{ \begin{array}{l} \frac{\partial^2 \psi}{\partial x^2} = \frac{1}{\alpha} \frac{\partial \psi}{\partial t} \\ \psi(x,0) = \theta_i(x) - \phi(x) \\ \psi(0,t) = \psi(L,t) = 0 \end{array} \right\}$$

$$\left\{ \begin{array}{l} \frac{d^2 \phi}{dx^2} + \frac{\dot{q}}{k} = 0 \\ \phi(0) = \phi(L) = 0 \end{array} \right\}$$

$$\psi(x,t) = \sum_{n=1}^{\infty} A_n e^{-\alpha \lambda_n^2 t} \sin \lambda_n x, \quad \lambda_n = \frac{n\pi}{L}$$

$$\phi(x) = \frac{\dot{q} L^2}{2k} \left[ \frac{x}{L} - \left( \frac{x}{L} \right)^2 \right]$$

$$\theta_i(x) - \phi(x) = \sum_{n=1}^{\infty} A_n \sin \lambda_n x \rightarrow A_n = \frac{2}{L} \int_0^L [\theta_i(x) - \phi(x)] \sin \lambda_n x dx$$

$$\therefore A_n = \frac{2}{L} \left\{ \int_0^L T_i(x) \sin \lambda_n x dx - \frac{1 - (-1)^n}{\lambda_n} \left[ T_\infty - \frac{\dot{q}}{k} \frac{1}{\lambda_n^2} \right] \right\}$$

Thus, the solution becomes

$$T(x,t) - T_w = \frac{\dot{q} L^2}{2k} \left[ \frac{x}{L} - \left( \frac{x}{L} \right)^2 \right] + \frac{2}{L} \sum_{n=1}^{\infty} e^{-\alpha \lambda_n^2 t} \sin \lambda_n x$$

$$\cdot \left\{ \int_0^L T_i(x') \sin \lambda_n x' dx' - \frac{1 - (-1)^n}{\lambda_n} \left[ T_\infty + \frac{\dot{q}}{k} \frac{1}{\lambda_n^2} \right] \right\}$$

PROB. 6.26: Formulation of the problem:

$$\left\{ \begin{array}{l} \frac{\partial^2 T}{\partial x^2} + \frac{\dot{q}(x,t)}{k} = \frac{1}{\alpha} \frac{\partial T}{\partial t} \\ T(x,0) = T_i(x) \\ T(0,t) = T_w(t); \quad T(L,t) = T_w(t) \end{array} \right\}$$

$$\text{Let } \theta(x,t) = T(x,t) - T_w(t)$$

$$\downarrow$$

$$\left\{ \begin{array}{l} \frac{\partial^2 \theta}{\partial x^2} + \frac{\dot{q}(x,t)}{k} = \frac{1}{\alpha} \left[ \frac{\partial \theta}{\partial t} + \frac{dT_w}{dt} \right] \\ \theta(x,0) = T_i(x) - T_w(0) \\ \theta(0,t) = 0; \quad \theta(L,t) = 0 \end{array} \right\}$$

Let

$$\theta(x,t) = \sum_{n=1}^{\infty} A_n(t) \sin \frac{n\pi}{L} x \quad \leftarrow \text{satisfies the B.C.'s.}$$

and

$$\frac{\dot{q}(x,t)}{k} = \sum_{n=1}^{\infty} B_n(t) \sin \frac{n\pi}{L} x$$

$$\therefore B_n(t) = \frac{2}{L} \int_0^L \frac{\dot{q}(x,t)}{k} \sin \frac{n\pi}{L} x dx$$

$$\frac{dT_w}{dt} = \sum_{n=1}^{\infty} C_n(t) \sin \frac{n\pi}{L} x$$

$$\therefore C_n(t) = \frac{2}{L} \int_0^L \frac{dT_w}{dt} \sin \frac{n\pi}{L} x dx = \frac{2}{n\pi} [1 - (-1)^n] \frac{dT_w}{dt}$$

Substituting the above expansions into the DE yields

$$\sum_{n=1}^{\infty} \left[ -\lambda_n^2 A_n(t) + B_n(t) \right] \sin \lambda_n x = \frac{1}{\alpha} \sum_{n=1}^{\infty} \left[ \frac{dA_n}{dt} + C_n(t) \right] \sin \lambda_n x$$

↓ where  $\lambda_n = \frac{n\pi}{L}$ .

$$\frac{dA_n}{dt} + \alpha \lambda_n^2 A_n(t) = \alpha B_n(t) - C_n(t)$$

$$\frac{d}{dt} \left[ e^{\alpha \lambda_n^2 t} A_n(t) \right] = e^{\alpha \lambda_n^2 t} [\alpha B_n(t) - C_n(t)]$$

$$\therefore A_n(t) = e^{-\alpha \lambda_n^2 t} A_n(0) + e^{-\alpha \lambda_n^2 t} \int_0^t e^{\alpha \lambda_n^2 t'} [\alpha B_n(t') - C_n(t')] dt'$$

Now, apply the I.C. to get

$$T_i(x) - T_w(0) = \sum_{n=1}^{\infty} A_n(0) \sin \lambda_n x$$

$$\therefore A_n(0) = \frac{2}{L} \int_0^L [T_i(x) - T_w(0)] \sin \lambda_n x dx$$


$$= \frac{2}{L} \int_0^L T_i(x) \sin \lambda_n x dx - \frac{2}{n\pi} [1 - (-1)^n] T_w(0)$$

⋮

$$\boxed{\Theta(x,t) = T(x,t) - T_w(t) = \frac{2}{L} \sum_{n=1}^{\infty} e^{-\alpha \lambda_n^2 t} \sin \lambda_n x \left\{ \int_0^L T_i(x) \sin \lambda_n x dx' \right. \\ \left. - \frac{[1 - (-1)^n]}{\lambda_n} \left[ T_w(0) + \int_0^t e^{\alpha \lambda_n^2 t'} \frac{dT_w}{dt'} dt' \right] \right. \\ \left. + \frac{1}{\beta C} \int_0^t e^{\alpha \lambda_n^2 t'} \int_0^L \dot{q}(x',t) \sin \lambda_n x' dx' dt' \right\}}$$

PROB. 6.27: Formulation in terms of  $\theta(r,t) = T(r,t) - T_i$ :

$$\left\{ \begin{array}{l} \frac{\partial^2 \theta}{\partial r^2} + \frac{1}{r} \frac{\partial \theta}{\partial r} + \frac{q}{k} = \frac{1}{\alpha} \frac{\partial \theta}{\partial t} \\ \theta(r,0) = 0 \\ \theta(0,t) = \text{finite}; \quad \theta(r_0, t) = 0 \end{array} \right.$$



Let  $\theta(r,t) = \theta_t(r,t) + \theta_s(r)$

$$\begin{array}{ll} \frac{\partial^2 \theta_t}{\partial r^2} + \frac{1}{r} \frac{\partial \theta_t}{\partial r} = \frac{1}{\alpha} \frac{\partial \theta_t}{\partial t} & | \\ \theta_t(r,0) = -\theta_s(r) & | \\ \theta_t(0,t) = \text{finite}; \quad \theta_t(r_0, t) = 0 & | \\ \downarrow & | \\ \theta_t(r,t) = \sum_{n=1}^{\infty} a_n e^{-\alpha \lambda_n^2 t} J_0(\lambda_n r) & | \end{array} \quad \begin{array}{ll} \frac{d^2 \theta_s}{dr^2} + \frac{1}{r} \frac{d \theta_s}{dr} + \frac{q}{k} = 0 & | \\ \theta_s(0) \neq \infty & | \\ \theta_s(r_0) = 0 & | \\ \downarrow & | \\ \theta_s(r) = \frac{q r_0^2}{4k} \left[ 1 - \left( \frac{r}{r_0} \right)^2 \right] & | \end{array}$$

where  $\lambda_n$ 's are the positive roots of  $J_0(\lambda_n r_0) = 0$ .

$$\begin{aligned} \therefore -\dot{\theta}_s(r) &= -\frac{\dot{q} r_0^2}{4k} \left[ 1 - \left( \frac{r}{r_0} \right)^2 \right] = \sum_{n=1}^{\infty} a_n \dot{J}_0(\lambda_n r) \\ \downarrow \\ a_n &= -\frac{1}{N_n} \frac{\dot{q} r_0^2}{4k} \int_0^{r_0} \left[ 1 - \left( \frac{r}{r_0} \right)^2 \right] \dot{J}_0(\lambda_n r) r dr, \quad N_n = \frac{r_0^2}{2} \dot{J}_1^2(\lambda_n r_0) \\ &= \dots = -\frac{2 \dot{q}}{k r_0 \lambda_n^3 \dot{J}_1(\lambda_n r_0)} \end{aligned}$$

$$\therefore T(r,t) - T_i = -\frac{2 \dot{q}}{k r_0} \sum_{n=1}^{\infty} e^{-\alpha \lambda_n^2 t} \frac{J_0(\lambda_n r)}{\lambda_n^3 \dot{J}_1(\lambda_n r_0)} + \frac{\dot{q} r_0^2}{4k} \left[ 1 - \left( \frac{r}{r_0} \right)^2 \right]$$

PROB. 6.28: Formulation in terms of  $\theta(r,t) = T(r,t) - T_i$ :

$$\frac{\partial^2 \theta}{\partial r^2} + \frac{2}{r} \frac{\partial \theta}{\partial r} + \frac{\dot{q}}{k} = \frac{1}{\alpha} \frac{\partial \theta}{\partial t}$$

$$\theta(r,0) = 0$$

$$\theta(0,t) = \text{finite}; \quad \theta(r_0, t) = 0$$

↓      Let  $\theta(r,t) = \theta_t(r,t) + \theta_s(r)$

$$\frac{\partial^2 \theta_t}{\partial r^2} + \frac{2}{r} \frac{\partial \theta_t}{\partial r} = \frac{1}{\alpha} \frac{\partial \theta_t}{\partial t} \quad \frac{d^2 \theta_s}{dr^2} + \frac{2}{r} \frac{d \theta_s}{dr} + \frac{\dot{q}}{k} = 0$$

$$\theta_t(r,0) = -\theta_s(r)$$

$$\theta_s(0) = \text{finite}$$

$$\theta_t(0,t) = \text{finite}; \quad \theta_t(r_0, t) = 0$$

$$\theta_s(r_0) = 0$$

$$\downarrow$$

$$U(r,t) = r \theta_t(r,t)$$

$$\downarrow$$

$$\frac{\partial^2 U}{\partial r^2} = \frac{1}{\alpha} \frac{\partial U}{\partial t}$$

$$U(r,0) = -r \theta_s(r)$$

$$U(0,t) = 0; \quad U(r_0, t) = 0$$

$$\downarrow$$

$$\boxed{\theta_s(r) = \frac{\dot{q} r_0^2}{6 k} \left[ 1 - \left( \frac{r}{r_0} \right)^2 \right]}$$

$$U(r,t) = \sum_{n=1}^{\infty} A_n e^{-\alpha \lambda_n^2 t} \sin \lambda_n r, \quad \lambda_n = \frac{n\pi}{r_0}, \quad n=1, 2, \dots$$

$$\therefore -\frac{\dot{q}r_0^2}{6k} \left[ r - \frac{r^3}{r_0^2} \right] = \sum_{n=1}^{\infty} A_n \sin \lambda_n r$$



$$A_n = \frac{2}{r_0} \int_0^{r_0} \left( -\frac{\dot{q}r_0^2}{6k} \right) \left( r - \frac{r^3}{r_0^2} \right) \sin \lambda_n r dr \\ = \dots = (-1)^n \frac{2 \dot{q}r_0^3}{k(n\pi)^3}$$

$$\therefore U(r,t) = 2 \frac{\dot{q}r_0^3}{k\pi^3} \sum_{n=1}^{\infty} \frac{(-1)^n}{n^3} e^{-\alpha \lambda_n^2 t} \sin \lambda_n r$$



$$T(r,t) - T_i = \frac{\dot{q}r_0^2}{6k} \left[ 1 - \left( \frac{r}{r_0} \right)^2 \right] + \frac{2 \dot{q}r_0^3}{k\pi^3 r} \sum_{n=1}^{\infty} \frac{(-1)^n}{n^3} e^{-\alpha \lambda_n^2 t} \sin \lambda_n r$$

$$\frac{T(r,t) - T_i}{\dot{q}r_0^2 / 6k} = 1 - \left( \frac{r}{r_0} \right)^2 + \frac{12}{\pi^3} \frac{r_0}{r} \sum_{n=1}^{\infty} \frac{(-1)^n}{n^3} e^{-\alpha \lambda_n^2 t} \sin \lambda_n r$$

The temperature distribution as  $t \rightarrow \infty$ :

$$T_s(r,t) - T_i = \frac{\dot{q}r_0^2}{6k} \left[ 1 - \left( \frac{r}{r_0} \right)^2 \right]$$

PROB. 6.29: Formulation in terms of  $\theta(x,t) = T(x,t) - T_i$  (see Prob. 2.9):

$$\left\{ \begin{array}{l} \frac{\partial^2 \theta}{\partial x^2} + \frac{\dot{q}_o e^{-\beta t}}{k} = \frac{1}{\alpha} \frac{\partial \theta}{\partial t} \\ \theta(x,0) = 0 \\ \frac{\partial \theta}{\partial x} \Big|_{x=0} = 0; \quad \theta(L,t) = 0 \end{array} \right\}$$

Let  $\theta(x,t) = e^{-\beta t} \phi(x,t)$

↓

$$\left\{ \begin{array}{l} \frac{\partial^2 \phi}{\partial x^2} + \frac{\dot{q}_o}{k} = \frac{1}{\alpha} \left[ -\beta \phi + \frac{\partial \phi}{\partial t} \right] \\ \phi(x,0) = 0 \\ \frac{\partial \phi}{\partial x} \Big|_{x=0} = 0; \quad \phi(L,t) = 0 \end{array} \right\}$$

Let  $\phi(x,t) = \psi(x,t) + \Omega(x)$

↓

$$\frac{\partial^2 \psi}{\partial x^2} = \frac{1}{\alpha} \left[ \frac{\partial \psi}{\partial t} - \beta \psi \right] \quad \frac{d^2 \Omega}{dx^2} + \frac{\beta}{\alpha} \Omega = - \frac{\dot{q}_o}{k}$$

$$\psi(x,0) = -\Omega(x)$$

$$\frac{d\Omega(0)}{dx} = 0$$

$$\frac{\partial \psi}{\partial x} \Big|_{x=0} = 0; \quad \psi(L,t) = 0$$

$$\Omega(L) = 0$$

↓

$$\Omega(x) = \frac{\dot{q}_o \alpha}{k \beta} \left[ \frac{\cos \sqrt{\frac{\beta}{\alpha}} x}{\cos \sqrt{\frac{\beta}{\alpha}} L} - 1 \right]$$

Let  $\psi(x,t) = X(x)\Gamma(t)$



$$\left\{ \begin{array}{l} \frac{d^2X}{dx^2} + \lambda^2 X = 0 \\ \frac{dX(0)}{dx} = 0 \\ X(L) = 0 \end{array} \right\} \quad \left| \begin{array}{l} X_n(x) = A_n \cos \lambda_n x \\ \lambda_n = \frac{2n-1}{L} \frac{\pi}{2}, n=1,2,\dots \end{array} \right. \quad \left| \begin{array}{l} \frac{d\Gamma_n}{dt} + (\alpha \lambda_n^2 - \beta) \Gamma_n = 0 \\ \Gamma_n(t) = B_n e^{-(\alpha \lambda_n^2 - \beta)t} \end{array} \right.$$

$$\therefore \psi(x,t) = \sum_{n=1}^{\infty} a_n e^{-(\alpha \lambda_n^2 - \beta)t} \cos \lambda_n x$$

$$\Rightarrow -\mathcal{Q}(x) = \frac{\dot{q}_0 \alpha}{k\beta} \left[ 1 - \frac{\cos \sqrt{\frac{\beta}{\alpha}} x}{\cos \sqrt{\frac{\beta}{\alpha}} L} \right] = \sum_{n=1}^{\infty} a_n \cos \lambda_n x$$

$$a_n = -\frac{2}{L} \int_0^L \mathcal{Q}(x) \cos \lambda_n x dx = -\frac{2}{L} \frac{\dot{q}_0 \alpha}{k} \frac{(-1)^n}{\lambda_n(\beta - \alpha \lambda_n^2)}, n=1,2,\dots$$

$$\therefore \phi(x,t) = \mathcal{Q}(x) - \frac{2}{L} \frac{\dot{q}_0 \alpha}{k} \sum_{n=1}^{\infty} \frac{(-1)^n}{\lambda_n(\beta - \alpha \lambda_n^2)} e^{-(\alpha \lambda_n^2 - \beta)t} \cos \lambda_n x$$

$$\theta(x,t) = T(x,t) - T_i = e^{-\beta t} \phi(x,t)$$

$$\therefore \boxed{\theta(x,t) = \frac{\dot{q}_0 \alpha}{k\beta} \left[ \frac{\cos \sqrt{\frac{\beta}{\alpha}} x}{\cos \sqrt{\frac{\beta}{\alpha}} L} - 1 \right] e^{-\beta t} - \frac{2}{L} \frac{\dot{q}_0 \alpha}{k} \sum_{n=1}^{\infty} (-1)^n \frac{e^{-\alpha \lambda_n^2 t} \cos \lambda_n x}{\lambda_n(\beta - \alpha \lambda_n^2)}}$$


PROB. 6.30: Formulation in terms of

$$\theta(x,t) = T(x,t) - T_i :$$

$$\frac{\partial^2 \theta}{\partial x^2} + \frac{\dot{q}_o}{k} [1 + \beta \theta] = \frac{1}{\alpha} \frac{\partial \theta}{\partial t}$$

$$\theta(x,0) = 0$$

$$\left. \frac{\partial \theta}{\partial x} \right|_{x=0} = 0, \quad \theta(L,t) = 0$$



$$\text{Let } \theta(x,t) = \psi(x,t) + \phi(x)$$

$$\frac{\partial^2 \psi}{\partial x^2} + \frac{\dot{q}_o \beta}{k} \psi = \frac{1}{\alpha} \frac{\partial \psi}{\partial t}$$

$$\psi(x,0) = -\phi(x)$$

$$\left. \frac{\partial \psi}{\partial x} \right|_{x=0} = 0, \quad \psi(L,t) = 0$$

$$\frac{\partial^2 \phi}{\partial x^2} + \frac{\dot{q}_o \beta}{k} \phi = -\frac{\dot{q}_o}{k}$$

$$\left. \frac{d\phi(x)}{dx} \right|_{x=0} = 0$$

$$\phi(L) = 0$$

$$\text{Let } \psi(x,t) = X(x) \cdot \Gamma(t)$$

$$\left\{ \begin{array}{l} \frac{d^2 X}{dx^2} + \lambda^2 X = 0 \\ \left. \frac{dX(x)}{dx} \right|_{x=0} = 0 \\ X(L) = 0 \end{array} \right\} \Rightarrow \begin{aligned} X_n(x) &= A_n \cos \lambda_n x \\ \lambda_n &= \frac{(2n-1)\pi}{2L}, \quad n=1, 2, 3, \dots \end{aligned}$$

and

$$\frac{d\Gamma_n}{dt} + \alpha \left( \lambda_n^2 - \frac{\dot{q}_o \beta}{k} \right) \Gamma_n = 0 \rightarrow \Gamma_n(t) = B_n e^{-\alpha \left( \lambda_n^2 - \frac{\dot{q}_o \beta}{k} \right) t}$$

Thus,

$$\psi(x,t) = \sum_{n=1}^{\infty} a_n \cos \lambda_n x e^{-\alpha (\lambda_n^2 - \frac{\dot{q}_0 \beta}{k}) t}$$

$$\therefore -\phi(x) = \sum_{n=1}^{\infty} a_n \cos \lambda_n x, \quad a_n = \frac{2}{L} \int_0^L \{-\phi(x)\} \cos \lambda_n x dx$$

Note that  $\phi(x) = -\frac{1}{\beta} - \frac{k}{\dot{q}_0 \beta} \frac{d^2 \phi}{dx^2}$ . Then,

$$\begin{aligned} \int_0^L \phi(x) \cos \lambda_n x dx &= - \int_0^L \left[ \frac{1}{\beta} + \frac{k}{\dot{q}_0 \beta} \frac{d^2 \phi}{dx^2} \right] \cos \lambda_n x dx \\ &= \frac{(-1)^n}{\beta \lambda_n} + \frac{k \lambda_n^2}{\dot{q}_0 \beta} \int_0^L \phi(x) \cos \lambda_n x dx \end{aligned}$$

$$\therefore \int_0^L \phi(x) \cos \lambda_n x dx = \frac{\dot{q}_0}{k \lambda_n} \frac{(-1)^n}{[\dot{q}_0 \beta / k - \lambda_n^2]}$$

Thus,

$$a_n = \frac{2}{L} \frac{\dot{q}_0}{k \lambda_n} \frac{(-1)^{n+1}}{[\dot{q}_0 \beta / k - \lambda_n^2]}$$

and

$$\boxed{\psi(x,t) = \frac{2\dot{q}_0}{KL} \sum_{n=1}^{\infty} \frac{(-1)^{n+1}}{\lambda_n [\dot{q}_0 \beta / k - \lambda_n^2]} \cos \lambda_n x \cdot e^{-\alpha (\lambda_n^2 - \frac{\dot{q}_0 \beta}{k}) t}}$$

This holds for either sign of the product  $\dot{q}_0 \beta$ .

$$\therefore \theta(x,t) = \psi(x,t) + \phi(x)$$

### Solution for $\phi(x)$

Case I:  $\dot{q}_0\beta < 0$ .

$$\text{Let } r^2 = -\frac{\dot{q}_0\beta}{k}. \text{ Then, } \phi(x) = \frac{1}{\beta} \left[ \frac{\cosh rx}{\cosh rL} - 1 \right].$$

Thus,

$$\begin{aligned} \theta(x,t) &= T(x,t) - T_i = \frac{1}{\beta} \left[ \frac{\cosh rx}{\cosh rL} - 1 \right] \\ &\quad + \frac{2\dot{q}_0}{kL} \sum_{n=1}^{\infty} \frac{(-1)^n}{\lambda_n[r^2 + \lambda_n^2]} \cos \lambda_n x \cdot \exp[-\alpha(\lambda_n^2 + r^2)t] \end{aligned}$$

Since the exponential terms in this result all tend to zero as  $t \rightarrow \infty$ , the steady-state solution is given by  $\phi(x)$  for all values of  $\dot{q}_0\beta < 0$ .

Case II:  $\dot{q}_0\beta > 0$ . Let  $p^2 = \frac{\dot{q}_0\beta}{k}$ . Then,

$$\phi(x) = \frac{1}{\beta} \left[ \frac{\cos px}{\cos pL} - 1 \right].$$

Thus,

$$\begin{aligned} \theta(x,t) &= T(x,t) - T_i = \frac{1}{\beta} \left[ \frac{\cos px}{\cos pL} - 1 \right] \\ &\quad + \frac{2\dot{q}_0}{kL} \sum_{n=1}^{\infty} \frac{(-1)^n}{\lambda_n[p^2 - \lambda_n^2]} \cos \lambda_n x \cdot \exp[-\alpha(\lambda_n^2 - p^2)t] \end{aligned}$$

In this case, the exponential terms vanish as  $t \rightarrow \infty$  if  $p^2 < \lambda_1^2$  or  $\frac{\dot{q}_0\beta}{k} < (\frac{\pi}{2L})^2$

and, if this inequality holds, the steady-state solution is given by  $\phi(x)$ . If  $p^2 > \lambda_1^2$ , no steady-state solution exists, because heat would be generated at a rate too great for its removal to be possible.


PROB. 6.33: Formulation of the problem:

$$\frac{\partial^2 T}{\partial x^2} = \frac{1}{\alpha} \frac{\partial T}{\partial t}, \quad \alpha = \frac{k}{\rho c}$$

$$T(x,0) = \begin{cases} T_1, & 0 < x < a \\ T_2, & a < x < a+b \end{cases}$$

$$\frac{\partial T(0,t)}{\partial x} = 0; \quad \frac{\partial T(a+b,t)}{\partial x} = 0$$

$$\text{Let } T(x,t) = X(x) \cdot \Gamma(t)$$



$$\left\{ \begin{array}{l} \frac{d^2 X}{dx^2} + \lambda^2 X = 0 \\ \frac{dX(0)}{dx} = 0 \\ \frac{dX(a+b)}{dx} = 0 \end{array} \right\} \Rightarrow \begin{array}{l} X_n(x) = A_n \cos \lambda_n x \\ \text{where } \lambda_n \text{ are zeros of } \sin \lambda_n (a+b) = 0 \\ \therefore \lambda_n = \frac{n\pi}{a+b}, n = 0, 1, 2, \dots \end{array} \quad \left| \begin{array}{l} \frac{d\Gamma_n}{dt} + \alpha \lambda_n^2 \Gamma_n = 0 \\ \downarrow \\ \Gamma_n(t) = B_n e^{-\alpha \lambda_n^2 t} \end{array} \right.$$

$$\therefore T(x,t) = \sum_{n=0}^{\infty} a_n e^{-\alpha \lambda_n^2 t} \cos \lambda_n x = a_0 + \sum_{n=1}^{\infty} a_n e^{-\alpha \lambda_n^2 t} \cos \lambda_n x$$

$$\Rightarrow T(x,0) = a_0 + \sum_{n=1}^{\infty} a_n \cos \lambda_n x$$

Thus,

$$a_0 = \frac{1}{a+b} \int_0^{a+b} T(x,0) dx = \frac{1}{a+b} \left\{ T_1 \int_0^a dx + T_2 \int_a^{a+b} dx \right\} = \frac{a T_1 + b T_2}{a+b}$$

$$a_n = \frac{2}{a+b} \int_0^{a+b} T(x,0) \cos \lambda_n x dx = \dots = \frac{2}{a+b} \frac{(T_1 - T_2) \sin \lambda_n a}{\lambda_n}$$

$$\boxed{\therefore T(x,t) = \frac{a T_1 + b T_2}{a+b} + \frac{2(T_1 - T_2)}{\pi} \sum_{n=1}^{\infty} \frac{\sin \lambda_n a \cos \lambda_n x}{n} e^{-\alpha \lambda_n^2 t}}$$

As  $t \rightarrow \infty$ ,  $T_s = \frac{a T_1 + b T_2}{a+b}$  ← Steady-state temperature

PROB. 6.34: The formulation in terms of  $\theta(x,t) = T(x,t) - (T_i + At)$ :

$$\left\{ \begin{array}{l} \frac{\partial^2 \theta}{\partial x^2} - \frac{A}{\alpha} = \frac{1}{\alpha} \frac{\partial \theta}{\partial t} \\ \theta(x,0) = 0 \\ \left. \frac{\partial \theta}{\partial x} \right|_{x=0} = 0; \quad \theta(L,t) = 0 \end{array} \right\}$$

Let  $\theta(x,t) = \psi(x,t) + \phi(x)$

$$\frac{\partial^2 \psi}{\partial x^2} = \frac{1}{\alpha} \frac{\partial \psi}{\partial t}$$

$$\psi(x,0) = -\phi(x)$$

$$\left. \frac{\partial \psi(x,t)}{\partial x} \right|_{x=0} = 0; \quad \psi(L,t) = 0$$

$$\Downarrow$$

$$\psi(x,t) = \sum_{n=1}^{\infty} a_n e^{-\alpha \lambda_n^2 t} \cos \lambda_n x$$

$$\frac{\partial^2 \phi}{\partial x^2} - \frac{A}{\alpha} = 0$$

$$\frac{d\phi(0)}{dx} = 0$$

$$\phi(L) = 0$$

$$\Downarrow$$

$$\phi(x) = \frac{AL^2}{2\alpha} \left[ 1 - \left( \frac{x}{L} \right)^2 \right]$$

where  $\lambda_n = \frac{2n-1}{L} \frac{\pi}{2}$  (See Table 4.1)

$$\therefore -\phi(x) = \frac{AL^2}{2\alpha} \left[ 1 - \left( \frac{x}{L} \right)^2 \right] = \sum_{n=1}^{\infty} a_n \cos \lambda_n x$$

$$\Downarrow$$

$$a_n = \frac{2}{L} \int_0^L \{-\phi(x)\} \cos \lambda_n x \, dx = \dots = \frac{2A}{\alpha L} \frac{(-1)^{n+1}}{\lambda_n^3}$$

$$\therefore \theta(x,t) = -\frac{2A}{\alpha L} \sum_{n=1}^{\infty} \frac{(-1)^n}{\lambda_n^3} e^{-\alpha \lambda_n^2 t} \cos \lambda_n x - \frac{AL^2}{2\alpha} \left[ 1 - \left( \frac{x}{L} \right)^2 \right]$$

or

---


$$T(x,t) - T_i = A \left\{ t - \frac{L^2}{2\alpha} \left[ 1 - \left( \frac{x}{L} \right)^2 \right] - \frac{2}{\alpha L} \sum_{n=1}^{\infty} \frac{(-1)^n}{\lambda_n^3} e^{-\alpha \lambda_n^2 t} \cos \lambda_n x \right\}$$


---

PROB. 6.35: Formulation of the problem:

$$\left\{ \begin{array}{l} \frac{\partial^2 T}{\partial r^2} + \frac{1}{r} \frac{\partial T}{\partial r} = \frac{1}{\alpha} \frac{\partial T}{\partial t} \\ T(r, 0) = T_i \\ T(0, t) = \text{finite}; \quad T(r_0, t) = T_i + At \end{array} \right\}$$

$$\text{Let } \theta(r, t) = T(r, t) - (T_i + At)$$



$$\frac{\partial^2 \theta}{\partial r^2} + \frac{1}{r} \frac{\partial \theta}{\partial r} = \frac{1}{\alpha} \left[ \frac{\partial \theta}{\partial t} + A \right]$$

$$\theta(r, 0) = 0$$

$$\theta(0, t) = \text{finite}; \quad \theta(r_0, t) = 0$$

The solution for  $\theta(r, t)$  may now be written as (see Prob. 6.27):

$$\theta(r, t) = \frac{2A}{\alpha r_0} \sum_{n=1}^{\infty} e^{-\alpha \lambda_n^2 t} \frac{J_0(\lambda_n r)}{\lambda_n^3 J_1(\lambda_n r_0)} - \frac{Ar_0^2}{4\alpha} \left[ 1 - \left( \frac{r}{r_0} \right)^2 \right]$$

Thus,

$$T(r, t) - T_i = \frac{Ar_0^2}{4\alpha} \left\{ 4 \frac{\alpha t}{r_0^2} - 1 + \left( \frac{r}{r_0} \right)^2 + \frac{8}{r_0^3} \sum_{n=1}^{\infty} e^{-\alpha \lambda_n^2 t} \frac{J_0(\lambda_n r)}{\lambda_n^3 J_1(\lambda_n r_0)} \right\}$$

where  $\lambda_n$ 's are the positive roots of  $J_0(\lambda r_0) = 0$ .

PROB. 6.37:

Formulation  $\Rightarrow \left\{ \begin{array}{l} \frac{\partial^2 T}{\partial s^2} = \frac{1}{\alpha} \frac{\partial T}{\partial t} \\ T(s, 0) = T_i(s) \\ T(s, t) = T(s+2L, t) \\ \frac{\partial T(s, t)}{\partial s} = \frac{\partial T(s+2L, t)}{\partial s} \end{array} \right\}$

Let  $T(s, t) = X(s) \cdot \Gamma(t)$

$$\left\{ \begin{array}{l} \frac{d^2 X}{ds^2} + \lambda X = 0 \\ X(s) = X(s+2L) \\ \frac{dX(s)}{ds} = \frac{dX(s+2L)}{ds} \end{array} \right\} \Rightarrow \lambda_n = \frac{n\pi}{L}, n = 0, 1, 2, \dots$$

$$X_n(s) = A_n \sin \lambda_n s + B_n \cos \lambda_n s$$

$$\frac{d\Gamma_n}{dt} - \alpha \lambda_n^2 \Gamma_n = 0 \rightarrow \Gamma_n(t) = C_n e^{-\alpha \lambda_n^2 t}$$

$$\therefore T(s, t) = b_0 + \sum_{n=1}^{\infty} e^{-\alpha \lambda_n^2 t} [A_n \sin \lambda_n s + B_n \cos \lambda_n s] \quad (1)$$

$$T(s, 0) = T_i(s) \Rightarrow T_i(s) = b_0 + \sum_{n=1}^{\infty} [a_n \sin \lambda_n s + b_n \cos \lambda_n s]$$

↑  
Complete Fourier series

$$\therefore b_0 = \frac{1}{2L} \int_0^{2L} T_i(s) ds, \quad a_n = \frac{1}{L} \int_0^{2L} T_i(s) \sin \lambda_n s ds, \quad b_n = \frac{1}{L} \int_0^{2L} T_i(s) \cos \lambda_n s ds$$

Substitution of  $b_0, a_n$  &  $b_n$  into Eq (1) gives  $T(s, t), t > 0$ .

Also, note that the steady-state temp.  $T(s) = \lim_{t \rightarrow \infty} T(s, t) = b_0$ .

PROB. 6.38: Let  $\theta(x, y, t) = T(x, y, t) - T_i$ .


Then,

$$\frac{\partial^2 \theta}{\partial x^2} + \frac{\partial^2 \theta}{\partial y^2} = \frac{1}{\alpha} \frac{\partial \theta}{\partial t}$$

$$\theta(x, y, 0) = 0$$

$$\theta(0, y, t) = 0; \quad \theta(L, y, t) = T_{\infty} - T_i$$

$$\frac{\partial \theta(x, 0, t)}{\partial y} = 0; \quad \theta(x, b, t) = T_{\infty} - T_i$$



$$\text{Let } \theta(x, y, t) = \theta_s(x, y) + \theta_t(x, y, t)$$

$$\frac{\partial^2 \theta_s}{\partial x^2} + \frac{\partial^2 \theta_s}{\partial y^2} = 0$$

$$\theta_s(0, y) = 0; \quad \theta_s(L, y) = T_{\infty} - T_i$$

$$\frac{\partial \theta_s(x, 0)}{\partial y} = 0; \quad \theta_s(x, b) = T_{\infty} - T_i$$

$$\frac{\partial^2 \theta_t}{\partial x^2} + \frac{\partial^2 \theta_t}{\partial y^2} = \frac{1}{\alpha} \frac{\partial \theta_t}{\partial t}$$

$$\theta_t(x, y, 0) = -\theta_s(x, y)$$

$$\theta_t(0, y, t) = 0; \quad \theta_t(L, y, t) = 0$$

$$\frac{\partial \theta_t(x, 0, t)}{\partial y} = 0; \quad \theta_t(x, b, t) = 0$$

$$\theta_s(x, y) - (T_{\infty} - T_i) = \sum_{n=1}^{\infty} A_n \sinh \frac{2n-1}{b} \frac{\pi}{2} (L-x) \cos \frac{2n-1}{b} \frac{\pi}{2} y$$

$$\Rightarrow -(T_{\infty} - T_i) = \sum_{n=1}^{\infty} A_n \sinh \frac{2n-1}{b} \frac{\pi}{2} L \cdot \cos \frac{2n-1}{b} \frac{\pi}{2} y$$

$$\begin{aligned} \therefore A_n \sin \frac{2n-1}{b} \frac{\pi}{2} L &= -\frac{2}{b} (T_{\infty} - T_i) \int_0^b \cos \frac{2n-1}{b} \frac{\pi}{2} y \, dy \\ &= \dots = \frac{4}{\pi} \frac{(-1)^n}{(2n-1)} (T_{\infty} - T_i) \end{aligned}$$

$$\therefore \boxed{\theta_s(x, y) = (T_{\infty} - T_i) \left[ 1 + \frac{4}{\pi} \sum_{n=1}^{\infty} \frac{(-1)^n}{(2n-1)} \frac{\sinh \frac{(2n-1)\pi}{b} \frac{\pi}{2} (L-x) \cos \frac{2n-1}{b} \frac{\pi}{2} y}{\sinh \frac{2n-1}{b} \frac{\pi}{2} L} \right]}$$

On the other hand,

$$\theta_t(x, y, t) = \sum_{m=1}^{\infty} \sum_{k=1}^{\infty} B_{mk} \sin \frac{m\pi}{L} x \cdot \cos \frac{2k-1}{b} \frac{\pi}{2} y \cdot \exp \left\{ - \left[ \left( \frac{m\pi}{L} \right)^2 + \left( \frac{2k-1}{b} \frac{\pi}{2} \right)^2 \right] t \right\}$$

$$\Rightarrow -\theta_s(x, y) = \sum_{m=1}^{\infty} \sum_{k=1}^{\infty} B_{mk} \sin \frac{m\pi}{L} x \cdot \cos \frac{(2k-1)\pi}{2b} y$$

$$\therefore B_{mk} = - \frac{4}{Lb} \int_0^L \int_0^b \theta_s(x, y) \sin \frac{m\pi}{L} x \cos \frac{(2k-1)\pi}{2b} y dx dy$$

$$-\frac{Lb}{4} B_{mk} = (T_{\infty} - T_i) \int_0^L \int_0^b \sin \frac{m\pi}{L} x \cos \frac{(2k-1)\pi}{2b} y dx dy$$

$$+ \sum_{n=1}^{\infty} A_n \int_0^L \sinh \frac{(2n-1)\pi}{2b} (L-x) \sin \frac{m\pi}{L} x dx \underbrace{\int_0^b \cos \frac{(2n-1)\pi}{2b} y \cos \frac{(2k-1)\pi}{2b} y dy}_{\delta_{nk} \frac{b}{2}}$$

⋮

$$B_{mk} = - \frac{8(T_{\infty} - T_i)}{m(2k-1)\pi^2} (-1)^k \left[ 1 - (-1)^m \right] - \frac{8(-1)^k}{\pi L(2k-1)} \frac{\int_0^L \sinh \frac{(2k-1)\pi}{2b} (L-x) \sin \frac{m\pi}{L} x dx}{\sinh \frac{2k-1}{b} \frac{\pi}{2} L}$$

Thus,

$$\frac{T(x, y, t) - T_i}{T_{\infty} - T_i} = 1 + \frac{4}{\pi} \sum_{n=1}^{\infty} \frac{(-1)^n}{(2n-1)} \frac{\sinh \frac{(2n-1)\pi}{2b} (L-x) \cos \frac{(2n-1)\pi}{2b} y}{\sinh \frac{(2n-1)\pi}{2b} L}$$

$$- \frac{8}{\pi} \sum_{m=1}^{\infty} \sum_{k=1}^{\infty} \frac{(-1)^k}{(2k-1)} \left\{ \frac{[1 - (-1)^m]}{m\pi} + \frac{1}{L} \frac{\int_0^L \sinh \frac{(2k-1)\pi}{2b} (L-x') \sin \frac{m\pi}{L} x' dx'}{\sin \frac{2k-1}{b} \frac{\pi}{2} L} \right\}$$

$$\cdot \sin \frac{m\pi}{L} x \cos \frac{(2k-1)\pi}{2b} y \exp \left\{ - \left[ \left( \frac{m\pi}{L} \right)^2 + \left( \frac{2k-1}{b} \frac{\pi}{2} \right)^2 \right] t \right\}$$

PROB. 6.39: Formulation in terms of  $\theta(r, \phi, t) = T(r, \phi, t) - T_w$ :

$$\frac{\partial^2 \theta}{\partial r^2} + \frac{1}{r} \frac{\partial \theta}{\partial r} + \frac{1}{r^2} \frac{\partial^2 \theta}{\partial \phi^2} = \frac{1}{\alpha} \frac{\partial \theta}{\partial t}$$

$$\theta(r, \phi, 0) = F(r, \phi) - T_w$$

$$\theta(0, \phi, t) = \text{finite}; \quad \theta(r_0, \phi, t) = 0$$

$$\theta(r, 0, t) = 0; \quad \theta(r, \pi, t) = 0$$

$$\text{Let } \theta(r, \phi, t) = R(r) \cdot \Phi(\phi) \cdot \Gamma(t)$$

Then,

$$\frac{d^2 \Phi}{d\phi^2} + \lambda^2 \Phi = 0$$

$$\Phi(0) = 0 \quad \Rightarrow \quad \Phi_n(\phi) = A_n \sin n\phi, \quad n=1, 2, 3, \dots$$

$$\Phi(\pi) = 0$$

and

$$r^2 \frac{d^2 R}{dr^2} + r \frac{dR}{dr} + (\beta^2 r^2 - n^2) R = 0$$

$$R(0) = \text{finite} \quad \Rightarrow \quad R_{mn}(r) = B_{mn} J_n(\beta_{mn} r)$$

$$R(r_0) = 0$$

where  $\beta_{mn}$  are the zeros of

finally,

$$\frac{d \Gamma_{mn}}{dt} + \alpha \beta_{mn}^2 \Gamma_{mn} = 0$$

$$J_n(\beta r_0) = 0$$



$$\beta_{mn}, \quad m=1, 2, \dots$$

$$\Gamma_{mn} = C_{mn} e^{-\alpha \beta_{mn}^2 t}$$

Thus,

$$\theta(r, \phi, t) = \sum_{m=1}^{\infty} \sum_{n=1}^{\infty} a_{mn} J_n(\beta_{mn} r) \sin(n\phi) e^{-\alpha \beta_{mn}^2 t}$$



$$F(r, \phi) - T_w = \sum_{m=1}^{\infty} \sum_{n=1}^{\infty} a_{mn} J_n(\beta_{mn} r) \sin(n\phi)$$



$$a_{mn} = \frac{2}{\pi} \frac{1}{N_{mn}} \int_0^{r_0} \int_0^{\pi} [F(r, \phi) - T_w] \sin(n\phi) J_n(\beta_{mn} r) r dr d\phi$$

where, from Table 4.2,

$$N_{nm} = \frac{r_0^2}{2} J_{n+1}^2(\beta_{mn} r_0)$$

Then the solution can be written as

$$T(r, \phi, t) - T_w = \frac{4}{\pi r_0^2} \sum_{m=1}^{\infty} \sum_{n=1}^{\infty} \left\{ \frac{J_n(\beta_{mn} r)}{J_{n+1}^2(\beta_{mn} r_0)} \sin(n\phi) e^{-\alpha \beta_{mn}^2 t} \right.$$

$$\left. \cdot \int_0^{r_0} \int_0^{\pi} [F(r, \phi) - T_w] \sin(n\phi) J_n(\beta_{mn} r) r dr d\phi \right\}$$

PROB. 6.40: The formulation in terms of  $\Theta(r, z, t) = T(r, z, t) - T_i$ :

$$\left\{ \begin{array}{l} \frac{\partial^2 \Theta}{\partial r^2} + \frac{1}{r} \frac{\partial \Theta}{\partial r} + \frac{\partial^2 \Theta}{\partial z^2} = \frac{1}{\alpha} \frac{\partial \Theta}{\partial t} \\ \Theta(r, z, 0) = 0 \\ \Theta(0, z, t) = \text{finite}; \quad k \frac{\partial \Theta}{\partial r} \Big|_{r=r_0} = q_w'' \\ \Theta(r, 0, t) = 0; \quad \Theta(r, H, t) = 0 \end{array} \right\}$$

$$\Theta(r, z, t) = \Theta_t(r, z, t) + \Theta_s(r, z)$$

$$\left\{ \begin{array}{ll} \frac{\partial^2 \Theta_t}{\partial r^2} + \frac{1}{r} \frac{\partial \Theta_t}{\partial r} + \frac{\partial^2 \Theta_t}{\partial z^2} = \frac{1}{\alpha} \frac{\partial \Theta_t}{\partial t} & | \quad \frac{\partial^2 \Theta_s}{\partial r^2} + \frac{1}{r} \frac{\partial \Theta_s}{\partial r} + \frac{\partial^2 \Theta_s}{\partial z^2} = 0 \\ \Theta_t(r, z, 0) = -\Theta_s(r, z) & | \quad \Theta_s(0, z) = \text{finite} \\ \Theta_t(0, z, t) = \text{finite}; \quad \frac{\partial \Theta_t}{\partial r} \Big|_{r=r_0} = 0 & | \quad \frac{\partial \Theta_s}{\partial r} \Big|_{r=r_0} = \frac{q_w''}{k} \\ \Theta_t(r, 0, t) = 0; \quad \Theta_t(r, H, t) = 0 & | \quad \Theta_s(r, 0) = 0; \quad \Theta_s(r, H) = 0 \end{array} \right.$$

Solution for  $\Theta_s(r, z)$ :

$$\Theta_s(r, z) = \sum_{n=1}^{\infty} A_n I_0\left(\frac{n\pi}{H} r\right) \sin \frac{n\pi}{H} z$$

$\Downarrow$

$$\frac{\partial \Theta_s}{\partial r} \Big|_{r=r_0} = \frac{q_w''}{k} = \frac{\pi}{H} \sum_{n=1}^{\infty} A_n n I_1\left(\frac{n\pi}{H} r_0\right) \sin \frac{n\pi}{H} z$$

Thus,

$$A_n \frac{n\pi}{H} I_1\left(\frac{n\pi}{H} r_0\right) = \frac{2}{H} \left(\frac{q_w''}{k}\right) \int_0^H \sin \frac{n\pi}{H} z dz \Rightarrow A_n = \frac{q_w''}{k} \frac{2H}{(n\pi)^2} \frac{[1 - (-1)^n]}{I_1\left(\frac{n\pi}{H} r_0\right)}$$

$$\therefore \Theta_s(r, z) = \frac{q_w''}{k} \frac{2H}{\pi^2} \sum_{n=1}^{\infty} \frac{[1 - (-1)^n]}{n^2} \frac{J_0(\frac{n\pi r}{H})}{J_1(\frac{n\pi r_0}{H})} \sin \frac{n\pi}{H} z$$

Solution for  $\Theta_t(r, z, t)$ :

$$\Theta_t(r, z, t) = \sum_{m=1}^{\infty} \sum_{n=0}^{\infty} B_{mn} e^{-(\lambda_n^2 + \beta_m^2)t} \cdot J_0(\lambda_n r) \cdot \sin \beta_m z, \quad \beta_m = \frac{m\pi}{H}, m=1, 2, 3, \dots$$

and  $\lambda_n$ 's are positive roots of (see Table 4.2)

$$\frac{d J_0(\lambda r_0)}{dr} = 0 \Rightarrow -\lambda J_1(\lambda r_0) = 0$$

$\lambda_0 = 0$

$J_1(\lambda_n r_0) = 0 \Rightarrow \lambda_n, n=1, 2, 3, \dots$

Thus,

$$-\Theta_s(r, z) = \sum_{m=1}^{\infty} \sum_{n=0}^{\infty} B_{mn} J_0(\lambda_n r) \sin \beta_m z$$

Or,

$$-\Theta_s(r, z) = \sum_{m=1}^{\infty} B_{m0} \sin \beta_m z + \sum_{m=1}^{\infty} \sum_{n=1}^{\infty} B_{mn} J_0(\lambda_n r) \sin \beta_m z$$

Then,

$$\underbrace{- \int_0^{r_0} \int_0^H \Theta_s(r, z) \sin \beta_m z r dr dz}_{-\frac{q_w'' r_0}{k} \frac{[1 - (-1)^m]}{\beta^3}} = B_{m0} \underbrace{\int_0^{r_0} r dr}_{r_0^2/2} \underbrace{\int_0^H \sin^2 \beta_m z dz}_{H/2}$$

$$\therefore B_{m0} = -\frac{4}{H r_0} \frac{q_w''}{k} \frac{[1 - (-1)^m]}{\beta^3}$$

and

$$\begin{aligned}
 & - \underbrace{\int_0^{r_0} \int_0^H \Theta_s(r, z) \sin \beta_m z J_0(\lambda_n r) r dr dz}_{\vdots} = B_{mn} \underbrace{\int_0^{r_0} J_0^2(\lambda_n r) r dr}_{\frac{r_0^2}{2} J_0^2(\lambda_n r_0)} \underbrace{\int_0^H \sin^2 \beta_m z dz}_{\frac{H}{2}} \\
 & - \frac{q''_w r_0}{k} J_0(\lambda_n r_0) \frac{[1 - (-1)^m]}{\beta_m [\lambda_n^2 + \beta_m^2]}
 \end{aligned}$$

$$\therefore B_{mn} = - \frac{4}{H r_0} \frac{q''_w}{k} \frac{1}{J_0(\lambda_n r_0)} \frac{[1 - (-1)^m]}{\beta_m [\lambda_n^2 + \beta_m^2]}$$

Note that  $B_{mn}|_{n=0} = B_{mo}$ . Thus,

---


$$\Theta_t(r, z, t) = - \frac{4}{H r_0} \frac{q''_w}{k} \sum_{m=1}^{\infty} \sum_{n=0}^{\infty} \frac{[1 - (-1)^m]}{\beta_m [\lambda_n^2 + \beta_m^2]} e^{-(\lambda_n^2 + \beta_m^2)t} \frac{J_0(\lambda_n r)}{J_0(\lambda_n r_0)} \sin \beta_m z$$


---

Hence, the solution is given by

$$\begin{aligned}
 T(r, z, t) - T_i &= \frac{2H}{\pi^2} \frac{q''_w}{k} \sum_{n=1}^{\infty} \frac{[1 - (-1)^n]}{n^2} \frac{I_0(\frac{n\pi}{H} r)}{I_1(\frac{n\pi}{H} r_0)} \sin \frac{n\pi}{H} z \\
 &\quad - \frac{4}{H r_0} \frac{q''_w}{k} \sum_{m=1}^{\infty} \sum_{n=0}^{\infty} \frac{[1 - (-1)^m]}{\beta_m [\lambda_n^2 + \beta_m^2]} e^{-(\lambda_n^2 + \beta_m^2)t} \frac{J_0(\lambda_n r)}{J_0(\lambda_n r_0)} \sin \beta_m z
 \end{aligned}$$

PROB. 6.42: The formulation in terms of  $\Theta(x, y, t) = T(x, y, t) - T_i$ :

$$\left\{ \begin{array}{l} \frac{\partial^2 \Theta}{\partial x^2} + \frac{\partial^2 \Theta}{\partial y^2} + \frac{\dot{q}}{k} = \frac{1}{\alpha} \frac{\partial \Theta}{\partial t} \\ \Theta(x, y, 0) = 0 \\ \Theta(0, y, t) = \Theta(a, y, t) = \Theta(x, 0, t) = \Theta(x, b, t) = 0 \end{array} \right\}$$

$$\text{Let } \Theta(x, y, t) = \Theta_t(x, y, t) + \Theta_s(x, y)$$

$$\left\{ \begin{array}{l} \frac{\partial^2 \Theta_t}{\partial x^2} + \frac{\partial^2 \Theta_t}{\partial y^2} = \frac{1}{\alpha} \frac{\partial \Theta_t}{\partial t} \\ \Theta_t(x, y, 0) = -\Theta_s(x, y) \\ \Theta_t(0, y, t) = \Theta_t(a, y, t) = \Theta(x, 0, t) = \Theta(x, b, t) = 0 \end{array} \right. \quad \left. \begin{array}{l} \frac{\partial^2 \Theta_s}{\partial x^2} + \frac{\partial^2 \Theta_s}{\partial y^2} + \frac{\dot{q}}{k} = 0 \\ \Theta_s(0, y) = \Theta_s(a, y) = \Theta_s(x, 0) = \Theta_s(x, b) = 0 \end{array} \right.$$

Solution for  $\Theta_s(x, y)$ : Let  $\Theta_s(x, y) = \psi(x, y) + \phi(x)$

$$\left\{ \begin{array}{l} \frac{\partial^2 \psi}{\partial x^2} + \frac{\partial^2 \psi}{\partial y^2} = 0 \\ \psi(0, y) = \psi(a, y) = 0 \\ \psi(x, 0) = \psi(x, b) = -\phi(x) \end{array} \right. \quad \left. \begin{array}{l} \frac{d^2 \phi}{dx^2} + \frac{\dot{q}}{k} = 0 \\ \phi(0) = \phi(a) = 0 \\ \therefore \phi(x) = \frac{\dot{q}a^2}{2k} \left( \frac{x}{a} \right) \left( 1 - \frac{x}{a} \right) \end{array} \right.$$

Let  $\psi(x, y) = \psi_1(x, y) + \psi_2(x, y)$

$$\left\{ \begin{array}{l} \frac{\partial^2 \psi_1}{\partial x^2} + \frac{\partial^2 \psi_1}{\partial y^2} = 0 \\ \psi_1(0, y) = \psi_1(a, y) = 0 \\ \psi_1(x, 0) = 0; \psi_1(x, b) = -\phi(x) \end{array} \right. \quad \left. \begin{array}{l} \frac{\partial^2 \psi_2}{\partial x^2} + \frac{\partial^2 \psi_2}{\partial y^2} = 0 \\ \psi_2(0, y) = \psi_2(a, y) = 0 \\ \psi_2(x, 0) = -\phi(x); \psi_2(x, b) = 0 \end{array} \right.$$

Solution for  $\psi_1(x, y)$ : See Eq. (5.17) in text.

$$\psi_1(x, y) = \frac{2}{a} \sum_{n=1}^{\infty} \frac{\sin \frac{n\pi}{a} x \sinh \frac{n\pi}{a} y}{\sinh \frac{n\pi}{a} b} \int_0^a \{-\phi(x')\} \sin \frac{n\pi}{a} x' dx'$$

Here,

$$\begin{aligned} \int_0^a \phi(x) \sin \frac{n\pi}{a} x dx &= -\phi(x) \left[ \frac{\cos \frac{n\pi}{a} x}{(n\pi/a)} \right]_0^a + \int_0^a \frac{\cos \frac{n\pi}{a} x}{(n\pi/a)} \frac{d\phi}{dx} dx \\ &= \frac{d\phi}{dx} \left[ \frac{\sin \frac{n\pi}{a} x}{(n\pi/a)^2} \right]_0^a - \left( \frac{a}{n\pi} \right)^2 \int_0^a \sin \frac{n\pi}{a} x \frac{d^2\phi}{dx^2} dx \\ &= -\left( \frac{a}{n\pi} \right)^2 \int_0^a \sin \frac{n\pi}{a} x \left\{ -\frac{q}{k} \right\} dx \\ &= -\left( \frac{a}{n\pi} \right)^2 \frac{q}{k} \left[ \frac{\cos \frac{n\pi}{a} x}{n\pi/a} \right]_0^a = \left( \frac{a}{n\pi} \right)^3 \frac{q}{k} [1 - (-1)^n] \end{aligned}$$

$$\therefore \psi_1(x, y) = -\frac{2a^2 q}{k\pi^3} \sum_{n=1}^{\infty} \frac{[1 - (-1)^n]}{n^3} \frac{\sin \frac{n\pi}{a} x \sinh \frac{n\pi}{a} y}{\sinh \frac{n\pi}{a} b}$$

Then, the solution for  $\psi_2(x, y)$  can be written as

$$\psi_2(x, y) = \frac{2a^2 q}{k\pi^3} \sum_{n=1}^{\infty} \frac{[1 - (-1)^n]}{n^3} \frac{\sin \frac{n\pi}{a} x \sinh \frac{n\pi}{a} (b-y)}{\sinh \frac{n\pi}{a} b}$$

Thus,

---


$$\theta_s(x, y) = \frac{q a^2}{2k} \left( \frac{x}{a} \right) \left( 1 - \frac{x}{a} \right)$$

$$- \frac{2a^2 q}{k\pi^3} \sum_{n=1}^{\infty} \frac{[1 - (-1)^n]}{n^3} \frac{\sin \frac{n\pi}{a} x}{\sinh \frac{n\pi}{a} b} \left[ \sinh \frac{n\pi}{a} y + \sinh \frac{n\pi}{a} (b-y) \right]$$


---

Solution for  $\theta_t(x, y, t)$ :

$$\theta_t(x, y, t) = \sum_{m=1}^{\infty} \sum_{n=1}^{\infty} A_{mn} \sin \frac{n\pi}{a} x \cdot \sin \frac{m\pi}{b} y \cdot e^{-[(\frac{n\pi}{a})^2 + (\frac{m\pi}{b})^2] \alpha t}$$



$$-\theta_s(x, y) = \sum_{m=1}^{\infty} \sum_{n=1}^{\infty} A_{mn} \sin \frac{n\pi}{a} x \cdot \sin \frac{m\pi}{b} y$$

$$\therefore - \int_0^a \int_0^b \theta_s(x, y) \sin \frac{n\pi}{a} x \cdot \sin \frac{m\pi}{b} y dx dy = A_{mn} \underbrace{\int_0^a \sin^2 \frac{n\pi}{a} x dx}_{\frac{a}{2}} \cdot \underbrace{\int_0^b \sin^2 \frac{m\pi}{b} y dy}_{\frac{b}{2}}$$

$$A_{mn} = - \frac{4}{ab} \int_0^a \int_0^b \theta_s(x, y) \sin \frac{n\pi}{a} x \cdot \sin \frac{m\pi}{b} y dx dy$$

⋮

$$A_{mn} = - \frac{\dot{q}}{k} \frac{1}{(\frac{n\pi}{a})^2 + (\frac{m\pi}{b})^2}$$

Thus, the final solution is

$$\begin{aligned} T(x, y, t) - T_i &= \frac{\dot{q} a^2}{2k} \left( \frac{x}{a} \right) \left( 1 - \frac{x}{a} \right) \\ &\quad - \frac{2a^2 \dot{q}}{k\pi^3} \sum_{n=1}^{\infty} \frac{[1 - (-1)^n]}{n^3} \frac{\sin \frac{n\pi}{a} x}{\sin \frac{n\pi}{a} b} \left[ \sinh \frac{n\pi}{a} y + \sinh \frac{n\pi}{a} (b-a) \right] \\ &\quad - \frac{\dot{q}}{k} \sum_{n=1}^{\infty} \sum_{m=1}^{\infty} \frac{\sin \frac{n\pi}{a} x \cdot \sin \frac{m\pi}{b} y}{(\frac{n\pi}{a})^2 + (\frac{m\pi}{b})^2} e^{-[(\frac{n\pi}{a})^2 + (\frac{m\pi}{b})^2] \alpha t} \end{aligned}$$

PROB. 6.43: Formulation in terms of  $\theta(r, z, t) = T(r, z, t) - T_i$ :

$$\frac{\partial^2 \theta}{\partial r^2} + \frac{1}{r} \frac{\partial \theta}{\partial r} + \frac{\partial^2 \theta}{\partial z^2} + \frac{\dot{q}}{k} = \frac{1}{\alpha} \frac{\partial \theta}{\partial t}$$

$$\theta(r, z, 0) = 0$$

$$\theta(0, z, t) = \text{finite}; \quad \theta(r_0, z, t) = 0$$

$$\theta(r, 0, t) = 0; \quad \theta(r, H, t) = 0$$

$$\text{Let } \theta(r, z, t) = \theta_t(r, z, t) + \theta_s(r, z)$$

$$\frac{\partial^2 \theta_t}{\partial r^2} + \frac{1}{r} \frac{\partial \theta_t}{\partial r} + \frac{\partial^2 \theta_t}{\partial z^2} = \frac{1}{\alpha} \frac{\partial \theta_t}{\partial t}$$

$$\theta_t(r, z, 0) = -\theta_s(r, z)$$

$$\theta_t(0, z, t) = \text{finite}; \quad \theta_t(r_0, z, t) = 0$$

$$\theta_t(r, 0, t) = 0; \quad \theta_t(r, H, t) = 0$$

$$\frac{\partial^2 \theta_s}{\partial r^2} + \frac{1}{r} \frac{\partial \theta_s}{\partial r} + \frac{\partial^2 \theta_s}{\partial z^2} + \frac{\dot{q}}{k} = 0$$

$$\theta_s(0, z) = \text{finite}; \quad \theta_s(r_0, z) = 0$$

$$\theta_s(r, 0) = 0; \quad \theta_s(r, H) = 0$$

From Prob. 5.21:

$$\theta_s(r, z) = \frac{\dot{q}H^2}{2k} \left[ \left(\frac{z}{H}\right) - \left(\frac{z}{H}\right)^2 \right] - \frac{2\dot{q}H^2}{\pi^3 k} \sum_{n=1}^{\infty} \frac{[1 - (-1)^n]}{n^3} \frac{J_0\left(\frac{n\pi r}{H}\right)}{J_0\left(\frac{n\pi r_0}{H}\right)} \sin \frac{n\pi}{H} z$$

On the other hand,

$$\theta_t(r, z, t) = \sum_{m=1}^{\infty} \sum_{n=1}^{\infty} A_{mn} e^{-\alpha[\lambda_n^2 + \beta_m^2]t} J_0(\lambda_n r) \sin \beta_m z$$

where

$$\beta_m = \frac{m\pi}{H}, \quad m = 1, 2, 3, \dots$$

and  $\lambda_n$ 's are the positive roots of  $J_0(\lambda_n r_0) = 0$ .

Thus,

$$-\Theta_s(r, z) = \sum_{m=1}^{\infty} \sum_{n=1}^{\infty} A_{mn} J_0(\lambda_n r) \sin \beta_m z$$

$$\therefore -\underbrace{\int_0^H \int_0^{r_0} \Theta_s(r, z) \sin \beta_m z J_0(\lambda_n r) r dr dz}_{\vdots} = A_{mn} \underbrace{\int_0^{r_0} J_0^2(\lambda_n r) r dr}_{\frac{r_0^2}{2} J_1^2(\lambda_n r_0)} \cdot \underbrace{\int_0^H \sin^2 \beta_m z dz}_{\frac{H}{2}}$$

$$\frac{\dot{q} r_0}{k} \frac{[1 - (-1)^m] J_1(\lambda_n r_0)}{\beta_m \lambda_n (\lambda_n^2 - \beta_m^2)}$$

$$\therefore A_{mn} = \frac{4}{r_0 H} \frac{\dot{q}}{k} \frac{[1 - (-1)^m]}{\beta_m \lambda_n (\lambda_n^2 - \beta_m^2) J_1(\lambda_n r_0)}$$

Now, the solution can be written as

$$\begin{aligned} T(r, z, t) - T_i &= \frac{4}{r_0 H} \frac{\dot{q}}{k} \sum_{m=1}^{\infty} \sum_{n=1}^{\infty} \frac{[1 - (-1)^m]}{\beta_m \lambda_n (\lambda_n^2 - \beta_m^2)} \frac{J_0(\lambda_n r)}{J_1(\lambda_n r_0)} \sin \beta_m z e^{-\alpha(\lambda_n^2 + \beta_m^2)t} \\ &\quad + \frac{\dot{q} H^2}{2k} \left\{ \frac{z}{H} - \left(\frac{z}{H}\right)^2 - \frac{4}{\pi^3} \sum_{n=1}^{\infty} \frac{[1 - (-1)^n]}{n^3} \frac{I_0\left(\frac{n\pi}{H} r\right)}{I_0\left(\frac{n\pi}{H} r_0\right)} \sin \frac{n\pi}{H} z \right\} \end{aligned}$$

# CHAPTER 7

## SOLUTIONS WITH INTEGRAL TRANSFORMS

---

PROB. 7.1: From Prob. 4.6:

$$f(x) = \sum_{n=1}^{\infty} \left\{ \frac{1}{N_n} \int_0^L f(x') \cos \lambda_n x' dx' \right\} \cos \lambda_n x \quad \text{with } N_n = \frac{1}{2\lambda_n} [\lambda_n L + \sin \lambda_n L \cos \lambda_n L]$$

and  $\lambda_n$ 's are the positive roots of  $\lambda \tan \lambda L = \frac{\alpha}{\beta}$ . Now, rewrite the expansion in two parts:

$$\begin{aligned} \bar{f}_n &= \int_0^L f(x') K_n(x') dx' && \leftarrow \text{Transform} \\ f(x) &= \sum_{n=1}^{\infty} \bar{f}_n \cdot K_n(x) && \leftarrow \text{Inversion} \end{aligned} \quad \begin{array}{l} K_n(x) = \frac{\cos \lambda_n x}{\sqrt{N_n}} \\ \uparrow \\ \text{Kernel} \end{array}$$

PROB. 7.2: From Prob. 4.16;

$$f(r) = \sum_{n=1}^{\infty} \left\{ \frac{1}{N_n} \int_a^b f(r') \phi_n(r') r' dr' \right\} \phi_n(r)$$

where

$$\phi_n(r) = J_0(\lambda_n r) Y_0(\lambda_n b) - J_0(\lambda_n b) Y_0(\lambda_n r) \quad \& \quad N_n = \frac{2}{\pi^2 \lambda_n^2} \left[ 1 - \frac{J_0^2(\lambda_n b)}{J_0^2(\lambda_n a)} \right]$$

and  $\lambda_n$ 's are the positive roots of

$$J_0(\lambda a) Y_0(\lambda b) - J_0(\lambda b) Y_0(\lambda a) = 0$$

Now, rewrite the expansion in two parts:

$$\begin{aligned} \bar{f}_n &= \int_a^b f(r') K_n(r') r' dr' && \leftarrow \text{Transform} \\ f(r) &= \sum_{n=1}^{\infty} \bar{f}_n \cdot K_n(r) && \leftarrow \text{Inversion} \end{aligned} \quad \begin{array}{l} K_n(r) = \frac{\phi_n(r)}{\sqrt{N_n}} \\ \uparrow \\ \text{Kernel} \end{array}$$

PROB. 7.3: Re-formulating the problem given by Eqs. (7.16) in terms of  $\Theta(x,t) = T(x,t) - T_{\infty}(t)$  we get :

$$\left\{ \begin{array}{l} \frac{\partial^2 \Theta}{\partial x^2} + F(x,t) = \frac{1}{\alpha} \frac{\partial \Theta}{\partial x} \\ \Theta(x,0) = \Theta_i(x) \\ \Theta(0,t) = \Theta(L,t) = 0 \end{array} \right\} \text{ where } \left\{ \begin{array}{l} F(x,t) = \frac{q(x,t)}{k} - \frac{1}{\alpha} \frac{dT_{\infty}}{dt} \\ \Theta_i(x) = T_i(x) - T_{\infty}(0) \end{array} \right.$$

Solution to this problem by the integral transform method (see Sec. 7.3) is given by

$$\Theta(x,t) = \frac{2}{L} \sum_{n=1}^{\infty} e^{-\alpha \lambda_n^2 t} \sin \lambda_n x \times \left\{ \int_0^L \Theta_i(x') \sin \lambda_n x' dx' \right. \\ \left. + \alpha \int_0^t e^{\alpha \lambda_n^2 t'} \int_0^L F(x',t') \sin \lambda_n x' dx' dt' \right\}, \quad \lambda_n = \frac{n\pi}{L}$$

Substituting the definitions of  $F(x,t)$  and  $\Theta_i(x)$  into this solution we obtain:

$$\Theta(x,t) = T(x,t) - T_{\infty}(t) = \frac{2}{L} \sum_{n=1}^{\infty} e^{-\alpha \lambda_n^2 t} \sin \lambda_n x \times \left\{ \int_0^L T_i(x') \sin \lambda_n x' dx' \right. \\ \left. - \frac{1}{\lambda_n} [1 - (-1)^n] \left[ T_{\infty}(0) + \int_0^t e^{\alpha \lambda_n^2 t'} \frac{dT_{\infty}}{dt'} dt' \right] \right. \\ \left. + \frac{1}{\rho C} \int_0^t e^{\alpha \lambda_n^2 t} \int_0^L q(x',t') \sin \lambda_n x' dx' dt' \right\}$$

This is the same result as given by Eq. (7.33).

PROB. 7.9:

- Because of symmetry, consider only one half of the rod.
- Assume  $k = \text{Constant}$ .

Formulation in terms of  $\theta(r, z) = T(r, z) - T_\infty$ :

$$\left. \begin{array}{l} \frac{1}{r} \frac{d}{dr} \left( r \frac{\partial \theta}{\partial r} \right) + \frac{\partial^2 \theta}{\partial r^2} + \frac{\partial^2 \theta}{\partial z^2} + \frac{\dot{q}(z)}{k} = 0 \\ \theta(0, z) = \text{finite}; \quad -k \frac{\partial \theta}{\partial r} \Big|_{r=r_0} = h \theta(r_0, z) \\ \frac{\partial \theta}{\partial z} \Big|_{z=0} = \frac{\partial \theta}{\partial z} \Big|_{z=L} = 0 \end{array} \right\} \text{where } \dot{q}(z) = \begin{cases} \dot{q}, & 0 < z < \frac{L}{2} \\ 0, & \frac{L}{2} < z < L \end{cases}$$

Define  $\bar{\theta}_n(r) = \int_0^L \theta(r, z) K_n(z) dz \leftarrow \text{Transform}$

with

$$\theta(r, z) = \sum_{n=0}^{\infty} \bar{\theta}_n(r) K_n(z) \leftarrow \text{Inversion}$$

where  $K_n(z)$  are the normalized eigenfunctions of

$$\left. \begin{array}{l} \frac{d^2 Z}{dz^2} + \lambda^2 Z = 0 \\ \frac{dZ(0)}{dz} = 0 \\ \frac{dZ(L)}{dz} = 0 \end{array} \right\} \text{From Table 7.1, } K_n(z) = \frac{\cos \lambda_n z}{\sqrt{N_n}}, \quad \lambda_n = \frac{n\pi}{L}, \quad n=0, 1, 2, \dots$$

$$N_n = \begin{cases} L, & n=0 \\ \frac{L}{2}, & n=1, 2, 3, \dots \end{cases}$$

Take the transform of the heat conduction equation:

$$\underbrace{\int_0^L \frac{1}{r} \frac{d}{dr} \left( r \frac{\partial \theta}{\partial r} \right) K_n(z) dz}_{\frac{1}{r} \frac{d}{dr} \left( r \frac{d\bar{\theta}_n}{dr} \right)} + \underbrace{\int_0^L \frac{\partial^2 \theta}{\partial z^2} K_n(z) dz}_{I_n} + \underbrace{\int_0^L \frac{\dot{q}(z)}{k} K_n(z) dz}_{J_n} = 0$$

$$I_n = \int_0^L \frac{\partial^2 \theta}{\partial z^2} K_n(z) dz = \left. \frac{\partial \theta}{\partial z} K_n(z) \right|_{z=0}^{z=L} - \int_0^L \frac{\partial \theta}{\partial z} \frac{dK_n}{dz} dz$$

$$= -\theta(r, z) \left. \frac{dK_n}{dz} \right|_{z=0}^{z=L} + \int_0^L \underbrace{\frac{d^2 K_n}{dz^2}}_{-\lambda_n^2 K_n(z)} dz = -\lambda_n^2 \bar{\theta}_n(r)$$

$$J_n = \int_0^L \frac{\dot{q}(z)}{k} K_n(z) dz = \frac{\dot{q}}{k} \int_0^{L/2} K_n(z) dz$$

$$\vdots$$

$$= \begin{cases} \frac{\dot{q}}{2k} \sqrt{L}, & n=0 \\ \frac{\dot{q}}{k} \sqrt{\frac{2}{L}} \frac{1}{\lambda_n} \sin \lambda_n \frac{L}{2}, & n=1, 2, 3, \dots \end{cases}$$

Thus,

$$\frac{1}{r} \frac{d}{dr} \left( r \frac{d\bar{\theta}_n}{dr} \right) - \lambda_n^2 \bar{\theta}_n(r) = -J_n$$

with

$$\bar{\theta}_n(0) = \text{finite}; \quad -k \frac{d\bar{\theta}_n}{dr} \Big|_{r=r_0} = h \bar{\theta}_n(r_0)$$

When  $n=0$ ,

$$\bar{\theta}_0(r) = J_0 \frac{r_0^2}{4} \left[ 1 + \frac{2k}{hr_0} - \left( \frac{r}{r_0} \right)^2 \right]$$

When  $n=1, 2, 3, \dots$ ,

$$\bar{\theta}_n(r) = \frac{J_n}{\lambda_n^2} \left[ 1 - \frac{h I_0(\lambda_n r)}{k \lambda_n I_1(\lambda_n r_0) + h I_0(\lambda_n r_0)} \right]$$

Thus, inverting  $\bar{\theta}_n(r)$ ,

$$\theta(r, z) = \frac{J_n r_0^2}{4} \left[ 1 + \frac{2k}{hr_0} - \left( \frac{r}{r_0} \right)^2 \right] \frac{1}{\sqrt{L}} + \sum_{n=1}^{\infty} \frac{J_n}{\lambda_n^2} \left[ 1 - \frac{h I_0(\lambda_n r)}{k \lambda_n I_1(\lambda_n r_0) + h I_0(\lambda_n r_0)} \right] \frac{\cos \lambda_n z}{\sqrt{L/2}}$$

Or

$$\theta(r, z) = T(r, z) - T_0 = \frac{\dot{q} r_0^2}{8k} \left[ 1 + \frac{2k}{hr_0} - \left( \frac{r}{r_0} \right)^2 \right] + \frac{2\dot{q}}{kL} \sum_{n=1}^{\infty} \frac{\sin \lambda_n \frac{L}{2}}{\lambda_n^3} \left[ 1 - \frac{h I_0(\lambda_n r)}{k \lambda_n I_1(\lambda_n r_0) + h I_0(\lambda_n r_0)} \right] \cos \lambda_n z$$

## Alternative Solution

Define  $\bar{\bar{\theta}}_{mn} = \int_0^{r_0} \bar{\theta}_n(r) K_0(\beta_m r) r dr \quad \leftarrow \text{Transform}$

With  $\bar{\theta}_n(r) = \sum_{m=1}^{\infty} \bar{\bar{\theta}}_{mn} K_0(\beta_m r) \quad \leftarrow \text{Inversion}$

where  $K_0(\beta_m r)$  are normalized eigenfunctions of

$$r^2 \frac{d^2 R}{dr^2} + r \frac{dR}{dr} + \beta^2 r^2 R = 0$$

$R(0) = \text{finite}$

$$h R(r_0) + k \frac{R(r_0)}{dr} = 0$$

From Table 7.3,

$$K_0(\beta_m r) = \frac{\sqrt{2}}{r_0} \frac{1}{\left[1 + \left(\frac{h}{k\beta_m}\right)^2\right]^{1/2}} \frac{J_0(\beta_m r)}{J_0(\beta_m r_0)}$$

where  $\beta_m$ 's are the positive roots of

$$h J_0(\beta r_0) + k \frac{d J_0(\beta r_0)}{dr} = 0 \Rightarrow \frac{J_0(\beta r_0)}{J_1(\beta r_0)} = \frac{Bk}{h}, \quad m=1, 2, 3, \dots$$

Thus,  $K_0(\beta_m r)$  can also be written as

$$K_0(\beta_m r) = \frac{\sqrt{2}}{r_0} \frac{J_0(\beta_m r)}{\left[J_0^2(\beta_m r_0) + J_1^2(\beta_m r_0)\right]^{1/2}}$$

Now, obtain the second transform of  $\bar{\theta}_n(r)$  problem:

$$\underbrace{\int_0^{r_0} \frac{1}{r} \frac{d}{dr} \left( r \frac{d \bar{\theta}_n}{dr} \right) K_0(\beta_m r) r dr - \lambda_n^2 \bar{\bar{\theta}}_{mn}}_{-\beta_m^2 \bar{\bar{\theta}}_{mn}} = -J_n \int_0^{r_0} K_0(\beta_m r) r dr$$

↓ Integrating twice by parts

$$-\beta_m^2 \bar{\bar{\theta}}_{mn}$$

Furthermore,

$$\int_0^{r_0} K_0(\beta_m r) r dr = \frac{\sqrt{2}}{r_0} \frac{1}{[J_0^2(\beta_m r_0) + J_1^2(\beta_m r_0)]^{1/2}} \underbrace{\int_0^{r_0} J_0(\beta_m r) r dr}_{\frac{r_0}{\beta_m} J_1(\beta_m r_0)}$$

$$= \frac{\sqrt{2}}{\beta_m [1 + (\frac{\beta_m k}{h})^2]^{1/2}}$$

Thus,

$$(\lambda_n^2 + \beta_m^2) \bar{\bar{\theta}}_{mn} = \frac{\sqrt{2} J_n}{\beta_m [1 + (\frac{\beta_m k}{h})^2]^{1/2}}$$

$$\bar{\bar{\theta}}_{mn} = \frac{\sqrt{2} J_n}{\beta_m (\lambda_n^2 + \beta_m^2) [1 + (\frac{\beta_m k}{h})^2]^{1/2}}$$

Then,

$$\begin{aligned} \theta(r, z) &= \sum_{n=0}^{\infty} \left\{ \sum_{m=1}^{\infty} \bar{\bar{\theta}}_{mn} K_0(\beta_m r) \right\} K_n(z) \\ &= \sum_{m=1}^{\infty} \left\{ \sum_{n=0}^{\infty} \bar{\bar{\theta}}_{mn} K_n(z) \right\} K_0(\beta_m r) \\ &\vdots \end{aligned}$$

$$\begin{aligned} \theta(r, z) &= \frac{2}{r_0} \frac{q}{k} \sum_{m=1}^{\infty} \frac{1}{\beta_m [1 + (\frac{\beta_m k}{h})^2]^{1/2}} \left\{ \frac{1}{2\beta_m^2} \right. \\ &\quad \left. + \frac{2}{L} \sum_{n=1}^{\infty} \frac{\sin \lambda_n \frac{L}{2} \cos \lambda_n z}{\lambda_n (\lambda_n^2 + \beta_m^2)} \right\} \frac{J_0(\beta_m r)}{[J_0^2(\beta_m r_0) + J_1^2(\beta_m r_0)]^{1/2}} \end{aligned}$$

PROB. 7.11: Formulation in terms of  $\Theta(x,y) = T(x,y) - T_0$ :

$$\frac{\partial^2 \Theta}{\partial x^2} + \frac{\partial^2 \Theta}{\partial y^2} - m(\Theta - \Theta_\infty) = 0, \quad \Theta_\infty = T_\infty - T_0$$

$$\Theta(0,y) = \Theta(a,y) = 0$$

$$\Theta(x,0) = \Theta(x,b) = 0$$

In the interval  $(0,a)$ , define a finite Fourier transform of  $\Theta(x,y)$  with respect to  $x$  as:

$$\bar{\Theta}(\lambda_n, y) = \int_0^a \Theta(x, y) K(\lambda_n, x) dx$$

with the inversion

$$\Theta(x, y) = \sum_{n=1}^{\infty} \bar{\Theta}(\lambda_n, y) K(\lambda_n, x)$$

with

$$(\text{from Table 7.1}) \quad K(\lambda_n, x) = \sqrt{\frac{2}{a}} \sin \lambda_n x, \quad \lambda_n = \frac{n\pi}{a}, \quad n=1, 2, 3, \dots$$

Now, obtain the transform of the differential equation:

$$\underbrace{\int_0^a \frac{\partial^2 \Theta}{\partial x^2} K(\lambda_n, x) dx}_{= \dots = -\lambda_n^2 \bar{\Theta}(\lambda_n, y)} + \frac{d^2 \bar{\Theta}}{dy^2} - m^2 \bar{\Theta}(\lambda_n, y) + m^2 \Theta_\infty \underbrace{\int_0^a K(\lambda_n, x) dx}_{\sqrt{\frac{2}{a}} \frac{1-(-1)^n}{\lambda_n}} = 0$$

Thus,

$$\frac{d^2 \bar{\Theta}}{dy^2} - (\lambda_n^2 + m^2) \bar{\Theta}(\lambda_n, y) = -m^2 \Theta_\infty \sqrt{\frac{2}{a}} \frac{1-(-1)^n}{\lambda_n} \quad \text{--- (1)}$$

$$\therefore \bar{\Theta}(\lambda_n, y) = A_n \sinh \sqrt{\lambda_n^2 + m^2} y + B_n \cosh \sqrt{\lambda_n^2 + m^2} y + \underbrace{m^2 \Theta_\infty \sqrt{\frac{2}{a}} \frac{1-(-1)^n}{\lambda_n (\lambda_n^2 + m^2)}}_{\text{Particular solution}} \bar{\Theta}_p$$

$$\theta(x, 0) = 0 \Rightarrow \bar{\theta}(\lambda_n, 0) = 0 \Rightarrow B_n = 0$$

$$\theta(x, b) = 0 \Rightarrow \bar{\theta}(\lambda_n, b) = 0 \Rightarrow A_n = -\frac{\bar{\theta}_p}{\sinh \sqrt{\lambda_n^2 + m^2} b}$$

Thus,

$$\bar{\theta}(\lambda_n, y) = \bar{\theta}_p \left\{ 1 - \frac{\sinh \sqrt{\lambda_n^2 + m^2} y}{\sinh \sqrt{\lambda_n^2 + m^2} b} \right\}$$

and

$$\theta(x, y) = \sum_{n=1}^{\infty} \bar{\theta}_p \left\{ 1 - \frac{\sinh \sqrt{\lambda_n^2 + m^2} y}{\sinh \sqrt{\lambda_n^2 + m^2} b} \right\} K(\lambda_n x)$$



$$\frac{\theta(x, y)}{\theta_{\infty}} = \frac{T(x, y) - T_0}{T_{\infty} - T_0} = \frac{2m^2}{a} \sum_{n=1}^{\infty} \frac{1 - (-1)^n}{\lambda_n^2 (\lambda_n^2 + m^2)} \sin \lambda_n x \left\{ 1 - \frac{\sinh \sqrt{\lambda_n^2 + m^2} y}{\sinh \sqrt{\lambda_n^2 + m^2} b} \right\}$$

### Alternative approach

Define  $\bar{\theta}(\lambda_n, \beta_k) = \int_0^b \bar{\theta}(\lambda_n, y) K(\beta_k y) dy$

with

$$\bar{\theta}(\lambda_n, y) = \sum_{k=1}^{\infty} \bar{\theta}(\lambda_n, \beta_k) K(\beta_k y)$$

and

$$K(\beta_k y) = \sqrt{\frac{2}{b}} \sin \beta_k y, \beta_k = \frac{k\pi}{b}, k=1, 2, 3, \dots$$

Now, one more transform of Eq. ① in the y-direction gives

$$\int_0^b \frac{d\bar{\theta}}{dy^2} K(\beta_k y) dy - (\lambda_n^2 + m^2) \bar{\theta}(\lambda_n, \beta_k) = -m^2 \theta_\infty \sqrt{\frac{2}{ab}} \frac{1-(-1)^n}{\lambda_n} \int_0^b K(\beta_k y) dy$$

$$= \dots = -\beta_k^2 \bar{\theta}(\lambda_n, \beta_k)$$

$$\sqrt{\frac{2}{b}} \frac{1-(-1)^k}{\beta_k}$$

Thus,

$$(\lambda_n^2 + \beta_k^2 + m^2) \bar{\theta}(\lambda_n, \beta_k) = m^2 \theta_\infty \sqrt{\frac{4}{ab}} \frac{[1-(-1)^n][1-(-1)^k]}{\lambda_n \beta_k}$$

$$\bar{\theta}(\lambda_n, \beta_k) = m^2 \theta_\infty \frac{2}{\sqrt{ab}} \frac{[1-(-1)^n][1-(-1)^k]}{\lambda_n \beta_k (\lambda_n^2 + \beta_k^2 + m^2)}$$



$$\boxed{\frac{\theta(x,y)}{\theta_\infty} = \frac{T(x,y) - T_0}{T_\infty - T_0} = m^2 \frac{4}{ab} \sum_{n=1}^{\infty} \sum_{k=1}^{\infty} \frac{[1-(-1)^n][1-(-1)^k]}{\lambda_n \beta_k (\lambda_n^2 + \beta_k^2 + m^2)} \sin \lambda_n x \cdot \sin \beta_k y}$$


PROB. 7.13: Formulation:

$$\frac{\partial^2 T}{\partial x^2} + \frac{\dot{q}(x,t)}{k} = \frac{1}{\alpha} \frac{\partial T}{\partial t}$$

$$T(x,0) = T_i(x)$$

$$k \frac{\partial T}{\partial x} \Big|_{x=0} = h_1 [T(0,t) - T_\infty(t)]$$

$$-k \frac{\partial T}{\partial x} \Big|_{x=L} = h_2 [T(L,t) - T_\infty(t)]$$



Define,

with  $\bar{T}_n(t) = \int_0^L T(x,t) K_n(x) dx \leftarrow \text{Transform}$

$$T(x,t) = \sum_{n=1}^{\infty} \bar{T}_n(t) K_n(x) \leftarrow \text{Inversion}$$

where, from Table 7.1,

$$K_n(x) = \sqrt{2} \frac{\lambda_n \cos \lambda_n x - H_1 \sin \lambda_n x}{\left\{ (\lambda_n^2 + H_1^2) \left[ L + \frac{H_2}{\lambda_n^2 + H_1^2} \right] - H_1 \right\}^{1/2}} \quad \text{with } H_1 = -\frac{h_1}{k} \text{ & } H_2 = \frac{h_2}{k}$$

Where  $\lambda_n$  are the positive roots of

$$\tan \lambda L = \frac{\lambda (H_2 - H_1)}{\lambda^2 + H_1 H_2} \Rightarrow \lambda_n, n=1, 2, 3, \dots$$

Now, the transform of the D.E.:

$$\int_0^L \frac{\partial^2 T}{\partial x^2} K_n(x) dx + \frac{1}{k} \int_0^L \dot{q}(x,t) K_n(x) dx = \frac{1}{\alpha} \int_0^L \frac{\partial T}{\partial t} K_n(x) dx$$

which yields

$$\bar{T}_n(t) = \bar{e}^{-\alpha \lambda_n^2 t} \left[ \bar{T}_i(\lambda_n) + \alpha \int_0^t F_n(t') e^{\alpha \lambda_n^2 t'} dt' \right]$$

where

$$F_n(t) = \left[ \frac{h_2}{k} K_n(L) + \frac{h_1}{k} K_n(0) \right] T_\infty(t) + \frac{1}{k} \bar{q}_n(t)$$

$$\bar{T}_i(\lambda_n) = \int_0^L T_i(x) K_n(x) dx \quad \& \quad \bar{q}_n(t) = \int_0^L \dot{q}(x,t) K_n(x) dx$$

Then, inversion results in

$$\therefore T(x,t) = \sum_{n=1}^{\infty} \bar{e}^{-\alpha \lambda_n^2 t} \left\{ \bar{T}_i(\lambda_n) + \alpha \int_0^t F_n(t') e^{\alpha \lambda_n^2 t'} dt' \right\}$$

PROB. 7.15: Formulation in terms of  $\theta(x,t) = T(x,t) - T_i$ :

$$\frac{\partial^2 \theta}{\partial x^2} + \frac{\dot{q}_o e^{-\beta t}}{k} = \frac{1}{\alpha} \frac{\partial \theta}{\partial t}$$

$$\theta(x,0) = 0$$

$$\left. \frac{\partial \theta}{\partial x} \right|_{x=0} = 0; \quad \theta(L,t) = 0$$

Define,

$$\text{Transform} \rightarrow \bar{\Theta}_n(t) = \int_0^L \theta(x,t) K_n(x) dx$$

$$\text{Inversion} \rightarrow \theta(x,t) = \sum_{n=1}^{\infty} \bar{\Theta}_n(t) K_n(x)$$

with the kernel  $K_n(x) = \sqrt{\frac{2}{L}} \cos \lambda_n x, \quad \lambda_n = \frac{(2n-1)\pi}{L}, n=1,2,3,\dots$

From Table 7.1 (Note that B.C. at  $x=0$  is 2<sup>nd</sup> kind  
and at  $x=L$  is 1<sup>st</sup> kind)

Transform of the D.E.:

$$\underbrace{\int_0^L \frac{\partial^2 \theta}{\partial x^2} K_n(x) dx}_{-\lambda_n^2 \bar{\Theta}_n(t)} + \frac{\dot{q}_o e^{-\beta t}}{k} \underbrace{\int_0^L K_n(x) dx}_{\sqrt{\frac{2}{L}}} = \frac{1}{\alpha} \underbrace{\int_0^L \frac{\partial \theta}{\partial t} K_n(x) dx}_{\frac{d \bar{\Theta}_n(t)}{dt}}$$

↓ Integrating by parts twice  
 $- \lambda_n^2 \bar{\Theta}_n(t)$

$$\sqrt{\frac{2}{L}} \frac{(-1)^{n+1}}{\lambda_n} \frac{d \bar{\Theta}_n(t)}{dt}$$

$$\Rightarrow \frac{d \bar{\Theta}_n}{dt} + \alpha \lambda_n^2 \bar{\Theta}_n = (-1)^{n+1} \sqrt{\frac{2}{L}} \frac{\dot{q}_o \alpha e^{-\beta t}}{k \lambda_n}$$

with the I.C. :  $\bar{\Theta}_n(0) = 0$

Thus,

$$\bar{\theta}_n(t) = (-1)^n \sqrt{\frac{2}{L}} \frac{\alpha \dot{q}_0}{k \lambda_n} \frac{e^{-\beta t} - e^{\alpha \lambda_n^2 t}}{\beta - \alpha \lambda_n^2}$$

Note that if  $\beta = \alpha \lambda_n^2$ , then

$$\frac{e^{-\beta t} - e^{\alpha \lambda_n^2 t}}{\beta - \alpha \lambda_n^2} = -t e^{-\beta t}$$

Inverting  $\bar{\theta}_n(t)$  we obtain

---


$$\theta(x, t) = \frac{2}{L} \frac{\alpha \dot{q}_0}{k} \sum_{n=1}^{\infty} (-1)^n \frac{e^{-\beta t} - e^{\alpha \lambda_n^2 t}}{\lambda_n(\beta - \alpha \lambda_n^2)} \cos \lambda_n x$$


---

It can further be shown that (when  $\beta \neq \alpha \lambda_n^2$ )

$$\frac{\cos \sqrt{\frac{\beta}{\alpha}} x}{\cos \sqrt{\frac{\beta}{\alpha}} L} - 1 = \frac{2\beta}{L} \sum_{n=1}^{\infty} (-1)^n \frac{\cos \lambda_n x}{\lambda_n(\beta - \alpha \lambda_n^2)}$$

Therefore, the solution can be rewritten as (when  $\beta \neq \alpha \lambda_n^2$ )

---


$$\theta(x, t) = \frac{\alpha \dot{q}_0}{k \beta} \left[ \frac{\cos \sqrt{\beta/\alpha} x}{\cos \sqrt{\beta/\alpha} L} - 1 \right] e^{-\beta t} - \frac{2}{L} \frac{\alpha \dot{q}_0}{k} \sum_{n=1}^{\infty} (-1)^n \frac{e^{-\alpha \lambda_n^2 t} \cos \lambda_n x}{\lambda_n(\beta - \alpha \lambda_n^2)}$$


---

PROB. 7.16: The formulation of the problem in terms of  $\theta(x,t) = T(x,t) - T_i$ :

$$\frac{\partial^2 \theta}{\partial x^2} + \frac{\dot{q}_o}{k} [1 + \beta \theta] = \frac{1}{\alpha} \frac{\partial \theta}{\partial t}$$

$$\theta(x,0) = 0$$

$$\left. \frac{\partial \theta}{\partial x} \right|_{x=0} = 0; \quad \theta(L,t) = 0$$

Define

$$\bar{\Theta}_n(t) = \int_0^L \theta(x,t) K_n(x) dx \leftarrow \text{Transform}$$

with

$$\theta(x,t) = \sum_{n=1}^{\infty} \bar{\Theta}_n(t) \cdot K_n(x) \leftarrow \text{Inversion}$$

and (from Table 7.):

$$\text{Kernel} \rightarrow K_n(x) = \sqrt{\frac{2}{L}} \cos \lambda_n x, \quad \lambda_n = \frac{(2n-1)\pi}{2L}, n=1,2,3,\dots$$

Now, obtain the transform of heat conduction equation:

$$\underbrace{\int_0^L \frac{\partial^2 \theta}{\partial x^2} K_n(x) dx}_{-\lambda_n^2 \bar{\Theta}_n(t)} + \underbrace{\frac{\dot{q}_o}{k} \int_0^L K_n(x) dx}_{-\frac{(-1)^n}{\lambda_n} \sqrt{\frac{2}{L}}} + \frac{\beta \dot{q}_o}{k} \bar{\Theta}_n(t) = \frac{1}{\alpha} \frac{d \bar{\Theta}_n}{dt}$$

Thus,

$$\frac{1}{\alpha} \frac{d \bar{\Theta}_n}{dt} + \left( \lambda_n^2 - \frac{\beta \dot{q}_o}{k} \right) \bar{\Theta}_n(t) = -\sqrt{\frac{2}{L}} \frac{(-1)^n}{\lambda_n}$$

$$\bar{\Theta}_n(t) = A_n \exp \left\{ -\alpha \left( \lambda_n^2 - \frac{\beta \dot{q}_o}{k} \right) t \right\} + \bar{\Theta}_p$$

where

$$\bar{\Theta}_p = \sqrt{\frac{2}{L}} \frac{\dot{q}_o}{\lambda_n k} \frac{(-1)^n}{(-\lambda_n^2 + \dot{q}_o \beta/k)} \leftarrow \text{Particular solution}$$

$$\theta(x, 0) = 0 \Rightarrow \bar{\theta}_n(0) = 0 \Rightarrow A_n = -\bar{\theta}_p$$

$$\therefore \bar{\theta}_n(t) = \bar{\theta}_p \left[ 1 - \exp \left\{ -\alpha \left( \lambda_n^2 - \frac{\beta \dot{q}_0}{k} \right) t \right\} \right]$$

Now, inversion gives

$$\begin{aligned} \theta(x, t) &= \sum_{n=1}^{\infty} \bar{\theta}_p \left[ 1 - \exp \left\{ -\alpha \left( \lambda_n^2 - \frac{\beta \dot{q}_0}{k} \right) t \right\} \right] \cdot K_n(x) \\ &\downarrow \\ \theta(x, t) &= \frac{2\dot{q}_0}{kL} \sum_{n=1}^{\infty} \frac{(-1)^n}{\lambda_n [\dot{q}_0 \beta / k - \lambda_n^2]} \left\{ 1 - \exp \left[ -\alpha \left( \lambda_n^2 - \frac{\beta \dot{q}_0}{k} \right) t \right] \right\} \cos \lambda_n x \end{aligned}$$

Note that this solution holds for either sign of the product  $\beta \dot{q}_0$ .

### Steady-state temperatures:

Case I:  $\beta \dot{q}_0 < 0$

Since the exponential terms in this result all tend to zero as  $t \rightarrow \infty$ , the steady-state solution is given by

$$\theta_s(x) = \frac{2\dot{q}_0}{kL} \sum_{n=1}^{\infty} \frac{(-1)^n}{\lambda_n [\dot{q}_0 \beta / k - \lambda_n^2]} = \dots = \frac{1}{\beta} \left[ \frac{\cosh rx}{\cosh rL} - 1 \right], r = \sqrt{\frac{-\beta \dot{q}_0}{k}}$$

Case II:  $\beta \dot{q}_0 > 0$

In this case, the exponential terms vanish as  $t \rightarrow \infty$ , if


$$P^2 = \frac{\beta \dot{q}_0}{k} < \lambda_1^2 \quad \text{or} \quad \frac{\dot{q}_0 \beta}{k} < \left( \frac{\pi}{2L} \right)^2$$

and

$$\theta_s(x) = \frac{2\dot{q}_0}{kL} \sum_{n=1}^{\infty} \frac{(-1)^n}{\lambda_n [\beta \dot{q}_0 / k - \lambda_n^2]} = \dots = \frac{1}{\beta} \left[ \frac{\cos px}{\cos pL} - 1 \right]$$

If  $P^2 > \lambda_1^2$ , no steady-state solution exists, because heat would be generated at a rate too great for its removal to be possible.

PROB. 7.23: Assume constant thermophysical properties ( $k, g, c$ ). Then the formulation of the problem is



$$\frac{\partial^2 T}{\partial r^2} + \frac{1}{r} \frac{\partial T}{\partial r} + \frac{1}{r^2} \frac{\partial^2 T}{\partial \phi^2} = \frac{1}{\alpha} \frac{\partial T}{\partial t}$$

$$T(r, \phi, 0) = T_i(r, \phi)$$

$$T(0, \phi, t) = \text{finite}; T(b, \phi, t) = T_3(\phi, t)$$

$$T(r, 0, t) = T_1(r, t); T(r, \frac{\pi}{2}, t) = T_2(r, t)$$

Define, in the interval  $(0, \frac{\pi}{2})$ , a transform of  $T(r, \phi, t)$  with respect to  $\phi$  as

$$\bar{T}(r, n, t) = \int_0^{\pi/2} T(r, \phi, t) K(n, \phi) d\phi$$

with

$$T(r, \phi, t) = \sum_{n=1}^{\infty} \bar{T}(r, n, t) \cdot K(n, \phi) \quad n=1, 2, 3, \dots$$

Now, obtain the transform of the heat conduction equation w.r.t  $\phi$ :

$$\frac{\partial^2 \bar{T}}{\partial r^2} + \frac{1}{r} \frac{\partial \bar{T}}{\partial r} + \frac{1}{r^2} \underbrace{\int_0^{\pi/2} \frac{\partial^2 T}{\partial \phi^2} K(n, \phi) d\phi}_{(-1)^n T_2(r, t) \frac{4n}{\sqrt{\pi}}} = \frac{1}{\alpha} \frac{\partial \bar{T}}{\partial t}$$

$$(-1)^n T_2(r, t) \frac{4n}{\sqrt{\pi}} + T_1(r, t) \frac{4n}{\sqrt{\pi}} - 4n^2 \bar{T}(r, n, \phi)$$

Thus,

$$\frac{\partial^2 \bar{T}}{\partial r^2} + \frac{1}{r} \frac{\partial \bar{T}}{\partial r} + \frac{1}{r^2} \frac{4n}{\sqrt{\pi}} [T_1(r, t) + (-1)^n T_2(r, t)] - \frac{4n^2}{r^2} \bar{T}(r, n, t) = \frac{1}{\alpha} \frac{\partial \bar{T}}{\partial t}$$

Now, define a Hankel transform as

$$\bar{\bar{T}}(\lambda_m, n, t) = \int_0^b \bar{T}(r, n, t) K_n(\lambda_m, r) r dr$$

$$\bar{T}(r, n, t) = \sum_{m=1}^{\infty} \bar{\bar{T}}(\lambda_m, n, t) K_n(\lambda_m, r)$$

where the kernel  $K_n(\lambda_m, r)$  are the normalized eigenfunctions of the following eigenvalue problem:

$$r^2 \frac{d^2 R}{dr^2} + r \frac{dR}{dr} + (\lambda^2 r^2 - 4n^2) R(r) = 0$$

$$R(0) = \text{finite}$$

$$R(b) = 0$$

Then, from Table 7.3,

$$K_n(\lambda_m, r) = \frac{\sqrt{2}}{b} \frac{J_{2n}(\lambda_m r)}{J_{2n+1}(\lambda_m b)}$$

and the eigenvalues  $\lambda_m$  are the positive zeros of

$$J_{2n}(\lambda b) = 0$$

Now, a second transform gives

$$\int_0^b \left\{ \frac{\partial^2 \bar{T}}{\partial r^2} + \frac{1}{r} \frac{\partial \bar{T}}{\partial r} \right\} K_n(\lambda_m, r) f dr + \frac{4n}{\sqrt{\pi}} \int_0^b \frac{T_1(r, t) + (-1)^n T_2(r, t)}{r^2} K_n(\lambda_m, r) f dr$$

$$\underbrace{\frac{1}{r} \frac{\partial}{\partial r} \left( r \frac{\partial \bar{T}}{\partial r} \right)}_{\vdots} + \underbrace{-4n^2 \int_0^b \frac{\bar{T}(r, n, t)}{r^2} K_n(\lambda_m, r) f dr}_{\xrightarrow{\text{def}}} = \frac{1}{\alpha} \frac{d \bar{T}}{dt}$$

$$-b \bar{T}_3(n, t) \underbrace{\frac{d K_n(\lambda_m, b)}{dr}}_{-\frac{\sqrt{2}}{b} \lambda_m} - \lambda_m^2 \bar{T}(\lambda_m, n, t) + \underbrace{4n^2 \int_0^b \frac{\bar{T}(r, n, t)}{r} K_n(\lambda_m, r) f dr}_{F_n(t)}$$

Thus, we have

$$\frac{1}{\alpha} \frac{d \bar{T}}{dt} + \lambda_m^2 \bar{T}(\lambda_m, n, t) = F_n(t)$$

where

$$F_n(t) = \sqrt{2} \lambda_m \bar{T}_3(n, t) + \frac{4n}{\pi} \int_0^b \frac{T_1(r, t) + (-1)^n T_2(r, t)}{r} K_n(\lambda_m, r) dr$$

Now, the solution of Eq. ① can be found as

$$\int_0^t e^{\alpha \lambda_m^2 t'} \left[ \frac{1}{\alpha} \frac{d \bar{T}}{dt} + \lambda_m^2 \bar{T}(\lambda_m, n, t') \right] dt' = \int_0^t e^{\alpha \lambda_m^2 t'} F_n^m(t') dt'$$

$$\frac{1}{\alpha} \frac{d}{dt} \left[ e^{\alpha \lambda_m^2 t'} \bar{T}(\lambda_m, n, t') \right]$$

Thus,

$$\bar{T}(\lambda_m, n, t) = \bar{T}_i(\lambda_m, n, 0) + \alpha \int_0^t e^{\alpha \lambda_m^2 t'} F_n^m(t') dt'$$

Therefore, the two inversions yield

$$T(r, \phi, t) = \sum_{n=1}^{\infty} \sum_{m=1}^{\infty} e^{\alpha \lambda_m^2 t} \left[ \bar{T}_i(\lambda_m, n) + \alpha \int_0^t e^{\alpha \lambda_m^2 t'} F_n^m(t') dt' \right] K_n(\lambda_m r) K(n, \phi)$$

Or,

$$\boxed{\bar{T}(r, \phi, t) = \frac{2}{b\sqrt{\pi}} \sum_{n=1}^{\infty} \sum_{m=1}^{\infty} e^{-\alpha \lambda_m^2 t} \sin 2n\phi \cdot \frac{J_{2n}(\lambda_m r)}{J_{2n+1}(\lambda_m b)} \\ \times \left[ \bar{T}_i(\lambda_m, n) + \alpha \int_0^t e^{\alpha \lambda_m^2 t'} F_n^m(t') dt' \right]}$$

where

$$\bar{T}_i(\lambda_m, n) = \frac{2\sqrt{2}}{b\sqrt{\pi} J_{2n+1}(\lambda_m b)} \int_0^b \int_0^{\pi/2} T_i(r, \phi) \sin 2n\phi J_{2n}(\lambda_m r) r dr d\phi$$

and

$$F_n^m(t) = \sqrt{2} \lambda_m \bar{T}_3(n, t) + \frac{4n}{\pi} \int_0^b \frac{T_1(r, t) + (-1)^n T_2(r, t)}{r} K_n(\lambda_m r) dr$$

with

$$\bar{T}_3(n, t) = \frac{2}{\sqrt{\pi}} \int_0^{\pi/2} T_3(\phi, t) \sin 2n\phi d\phi \quad \frac{\sqrt{2}}{b} \frac{J_{2n}(\lambda_m b)}{J_{2n+1}(\lambda_m b)}$$

PROB. 7.24: If the surface temperatures are independent of time (in Prob. 7.23), then

$$F_n^m = \sqrt{2} \lambda_m \bar{T}_3(n) + \frac{4\sqrt{2}n}{b\pi} \int_0^b \frac{T_1(r) + (-1)^n T_2(r)}{r J_{2n+1}(\lambda_m b)} J_{2n}(\lambda_m r) dr$$

where

$$\bar{T}_3(n) = \frac{2}{\sqrt{\pi}} \int_0^{\pi/2} T_3(\phi) \sin 2n\phi d\phi$$

Also,

$$\int_0^t e^{\alpha \lambda_m^2 t'} F_n^m dt' = F_n^m \int_0^t e^{\alpha \lambda_m^2 t'} dt' = F_n^m \frac{e^{\alpha \lambda_m^2 t} - 1}{\alpha \lambda_m^2}$$

and

$$e^{-\alpha \lambda_m^2 t} \int_0^t e^{\alpha \lambda_m^2 t'} F_n^m dt' = F_n^m \underbrace{\frac{1 - e^{-\alpha \lambda_m^2 t}}{\alpha \lambda_m^2}}$$

As  $t \rightarrow \infty$ ,  $\frac{1}{\alpha \lambda_m^2}$

Therefore, from the solution of Prob. 7.23, as  $t \rightarrow \infty$

$$T(r, \phi) = \frac{2}{b\sqrt{\pi}} \sum_{n=1}^{\infty} \sum_{m=1}^{\infty} F_n^m \frac{\sin 2n\phi \cdot J_{2n}(\lambda_m r)}{\lambda_m^2 J_{2n+1}(\lambda_m b)}$$

# CHAPTER 8

## SOLUTIONS WITH LAPLACE TRANSFORMS

---

PROB. B.1:

$$\begin{aligned}
 \mathcal{L}\{\sin\omega t\} &= \int_0^\infty e^{-pt} \sin\omega t dt = -e^{-pt} \frac{\cos\omega t}{\omega} \Big|_0^\infty - \frac{p}{\omega} \int_0^\infty \cos\omega t e^{-pt} dt \\
 &= \frac{1}{\omega} - \frac{p}{\omega} \left[ e^{-pt} \frac{\sin\omega t}{\omega} \Big|_0^\infty + \underbrace{\frac{p}{\omega} \int_0^\infty \sin\omega t e^{-pt} dt}_{\mathcal{L}\{\sin\omega t\}} \right] \\
 &= \frac{1}{\omega} - \frac{p^2}{\omega^2} \mathcal{L}\{\sin\omega t\}
 \end{aligned}$$

$$\therefore \mathcal{L}\{\sin\omega t\} = \frac{\omega}{p^2+\omega^2}, \quad p>0$$

Also, from above

$$\mathcal{L}\{\sin\omega t\} = \frac{1}{\omega} - \frac{p}{\omega} \mathcal{L}\{\cos\omega t\}$$

$$\therefore \mathcal{L}\{\cos\omega t\} = \frac{\omega}{p} \left[ \frac{1}{\omega} - \mathcal{L}\{\sin\omega t\} \right] = \frac{\omega}{p} \left[ \frac{1}{\omega} - \frac{\omega}{p^2+\omega^2} \right] = \frac{p}{p^2+\omega^2}, \quad p>0$$


---

PROB. B.2:

$$\mathcal{L}\left\{\frac{df}{dt}\right\} = \int_0^\infty e^{-pt} \frac{df}{dt} dt = e^{-pt} f(t) \Big|_0^\infty + p \underbrace{\int_0^\infty e^{-pt} f(t) dt}_{\bar{f}(p)} = -f(0) + p \bar{f}(p), \quad p>0$$


---

PROB. B.3:

$$\mathcal{L}\left\{\int_0^t f(t') dt'\right\} = \mathcal{L}\{\bar{F}(t)\}, \quad \bar{F}(t) = \int_0^t f(t') dt'$$

Using the result of Prob. B.2:

$$\bar{F}(p) = \mathcal{L}\left\{\int_0^t f(t') dt'\right\} = \frac{1}{p} \left[ \mathcal{L}\left\{\frac{df}{dt}\right\} + f(0) \right] = \frac{1}{p} \bar{f}(p)$$

PROB. B.4:

$$\bar{g}(p) \cdot \bar{h}(p) = \int_0^\infty e^{-pt} g(t) dt \cdot \int_0^\infty e^{-pt} h(t) dt \\ = \int_0^\infty \int_0^\infty e^{-p(u+v)} g(u) h(v) du dv$$

$$\begin{array}{ll} \leftarrow & \rightarrow \\ = \int_0^\infty g(u) \left\{ \int_0^\infty e^{-p(u+v)} h(v) dv \right\} du & = \int_0^\infty h(v) \left\{ \int_0^\infty e^{-p(u+v)} g(u) du \right\} dv \\ u+v=t \rightarrow dv=dt & u+v=t \rightarrow du=dt \\ = \int_0^\infty g(u) \left\{ \int_u^\infty e^{-pt} h(t-u) dt \right\} du & = \int_0^\infty h(v) \left\{ \int_v^\infty e^{-pt} g(t-v) dt \right\} dv \\ = \int_0^\infty \left\{ \int_u^\infty e^{-pt} g(u) h(t-u) dt \right\} du & = \int_0^\infty \left\{ \int_v^\infty e^{-pt} h(v) g(t-v) dt \right\} dv \end{array}$$

Change the order of integration

$$\begin{array}{ll} = \int_0^\infty \left\{ \int_0^t e^{-pt} g(u) h(t-u) du \right\} dt & = \int_0^\infty \left\{ \int_0^t e^{-pt} h(v) g(t-v) dv \right\} dt \\ = \int_0^\infty e^{-pt} \left\{ \int_0^t g(u) h(t-u) du \right\} dt & = \int_0^\infty e^{-pt} \left\{ \int_0^t h(v) g(t-v) dv \right\} dt \\ = L \left\{ \int_0^t g(u) h(t-u) du \right\} & = L \left\{ \int_0^t h(v) g(t-v) dv \right\} \end{array}$$

$$\therefore L^{-1} \{ \bar{g}(p) \cdot \bar{h}(p) \} = \int_0^t g(t-v) h(v) dv = \int_0^t h(t-u) g(u) du$$

PROB. 8.5:

$$\mathcal{L}^{-1}\left\{\frac{\cosh mx}{p \cosh mL}\right\} = \mathcal{L}^{-1}\left\{\frac{P(p)}{Q(p)}\right\} = \sum_{n=1}^{\infty} \frac{P(p_n)}{\frac{dQ(p_n)}{dp}} e^{p_n t}, \quad m = \sqrt{\frac{p}{\alpha}}$$

$$P(p) = \cosh mx = \sum_{n=0}^{\infty} \frac{x^{2n}}{(2n)!} \left(\frac{p}{\alpha}\right)^n$$

$$Q(p) = p \cosh mL = \sum_{n=0}^{\infty} \frac{L^{2n}}{(2n)!} \frac{p^{n+1}}{\alpha^n}$$

where  $p_n$ 's are the zeros of

$$p \cosh mL = 0 \quad \begin{array}{l} P_0 = 0 \\ \nearrow \\ \end{array} \quad \begin{array}{l} \rightarrow \\ \searrow \\ \cosh mL = 0 \rightarrow \cos imL = 0 \end{array}$$

$$\therefore i\sqrt{\frac{P_n}{\alpha}} L = \frac{(2n-1)\pi}{2}$$

$$p_n = -\frac{(2n-1)^2 \pi^2 \alpha}{4L^2}, \quad n = 1, 2, \dots$$

$$P(0) = 1, \quad P(p_n) = \cosh \left[ -i \frac{(2n-1)\pi}{2L} x \right] = \cos \frac{(2n-1)\pi x}{2L}$$

$$\frac{dQ}{dp} = \cosh mL + \frac{L}{2} \sqrt{\frac{p}{\alpha}} \sinh mL$$

$$\begin{aligned} \frac{dQ(0)}{dp} &= 1, \quad \frac{dQ(p_n)}{dp} = \cosh \left[ -i \frac{(2n-1)\pi}{2} \right] - \frac{L}{2} i \frac{(2n-1)\pi}{2L} \sinh \left[ -i \frac{(2n-1)\pi}{2} \right] \\ &= -\frac{(2n-1)\pi}{4} \sin \frac{(2n-1)\pi}{2} = -\frac{(2n-1)\pi}{4} (-1)^{n+1} \end{aligned}$$

$$\therefore \mathcal{L}^{-1}\left\{\frac{\cosh mx}{p \cosh mL}\right\} = 1 + \frac{4}{\pi} \sum_{n=1}^{\infty} \frac{(-1)^n}{2n-1} e^{-\frac{(2n-1)^2 \pi^2 x}{4L^2} t} \cdot \cos \frac{(2n-1)\pi x}{2L}$$

PROB. B.6: Formulation of the problem:

$$\frac{\partial^2 T}{\partial x^2} = \frac{1}{\alpha} \frac{\partial T}{\partial t}$$

$$T(x,0) = T_i$$

$$-k \frac{\partial T}{\partial x} \Big|_{x=0} = h [T_\infty - T(0,t)]; \lim_{x \rightarrow \infty} T(x,t) = T_\infty$$

The transformed problem:

$$\frac{d^2 \bar{T}}{dx^2} - \frac{P}{\alpha} \bar{T}(x,p) = -\frac{1}{\alpha} T_i$$

$$\therefore \bar{T}(x,p) = A e^{mx} + B \bar{e}^{mx} + \frac{T_i}{P}$$

$$\lim_{x \rightarrow \infty} \bar{T}(x,p) = \frac{T_i}{P} \rightarrow A = 0$$

$$-k \frac{d \bar{T}}{dx} \Big|_{x=0} = h \left[ \frac{1}{P} T_\infty - \bar{T}(0,p) \right] \rightarrow k m B = h \left[ \frac{1}{P} T_\infty - B - \frac{T_i}{P} \right]$$

$$B = \frac{h(T_\infty - T_i)}{P(km + h)}$$

$$\therefore \bar{T}(x,p) = \frac{T_i}{P} + \frac{h(T_\infty - T_i)}{P(km + h)} \bar{e}^{mx}$$

The inversion gives

$$T(x,t) = T_i + (T_\infty - T_i) \frac{h}{k} \mathcal{L}^{-1} \left\{ \frac{\bar{e}^{mx}}{P(m + \frac{h}{k})} \right\}$$

$$= T_i + (T_\infty - T_i) \left\{ \operatorname{erfc} \left[ \frac{x}{2\sqrt{\alpha t}} \right] \right.$$

$$\left. - e^{-hx/k + \alpha h^2 t / k^2} \operatorname{erfc} \left[ \frac{x}{2\sqrt{\alpha t}} + \frac{h}{k} \sqrt{\alpha t} \right] \right\}$$

PROB. B.7: Formulation:

$$\frac{\partial^2 T}{\partial x^2} = \frac{1}{\alpha} \frac{\partial T}{\partial t}$$

$$T(x,0) = T_i$$

$$-k \frac{\partial T(x,0)}{\partial x} = q''_{tw}; \quad \lim_{x \rightarrow \infty} T(x,t) = T_i$$

The transformed problem:

$$\frac{d^2 \bar{T}}{dx^2} - \frac{P}{\alpha} \bar{T}(x,p) = -\frac{T_i}{\alpha} \rightarrow \bar{T}(x,p) = A e^{mx} + B e^{-mx} + \frac{T_i}{P}$$

$$m = \sqrt{\frac{P}{\alpha}}$$

$$\left. \bar{T}(x,p) \right|_{x \rightarrow \infty} = \frac{T_i}{P} \rightarrow A = 0$$

$$-k \frac{d\bar{T}(0,p)}{dx} = \frac{q''_{tw}}{P} \rightarrow k_m B = \frac{q''_{tw}}{P}$$

$$B = \frac{q''_{tw}}{k_m P} = \frac{q''_{tw}}{k_p} \sqrt{\frac{\alpha}{P}}$$

$$\therefore \bar{T}(x,p) = \frac{T_i}{P} + \frac{q''_{tw}}{k} \sqrt{\frac{\alpha}{P}} \frac{e^{-mx}}{P}$$

The inversion gives


$$T(x,t) = T_i + \frac{q''_{tw}}{k} \mathcal{L}^{-1} \left\{ \frac{e^{-\sqrt{\frac{P}{\alpha}} x}}{P \sqrt{\alpha/P}} \right\}$$

↓ Appendix D, Item 28

$$\therefore T(x,t) - T_i = \frac{q''_{tw}}{k} \left[ 2 \left( \frac{\alpha t}{\pi} \right)^{1/2} e^{-x^2/4\alpha t} - x \operatorname{erfc} \left( \frac{x}{2\sqrt{\alpha t}} \right) \right]$$

PROB. 8.8:

Formulation:



$$\frac{\partial^2 T}{\partial x^2} = \frac{1}{\alpha} \frac{\partial T}{\partial t}$$

$$T(x,0) = T_i$$

$$T(0,t) = F(t); \quad T(L,t) = T_i$$

The transformed problem:

$$\frac{d^2 \bar{T}}{dx^2} - \frac{P}{\alpha} \bar{T}(x,p) = - \frac{T_i}{\alpha}$$

$$\bar{T}(x,p) = A \sinh mx + B \cosh mx + \frac{T_i}{P}$$

$$m = \sqrt{P/\alpha}$$

$$\bar{T}(0,p) = \bar{F}(p) = B + \frac{T_i}{P} \rightarrow B = \bar{F}(p) - \frac{T_i}{P}$$

$$\bar{T}(L,p) = \frac{T_i}{P} = A \sinh mL + B \cosh mL + \frac{T_i}{P} \rightarrow A = -B \frac{\cosh mL}{\sinh mL}$$

$$\therefore \bar{T}(x,p) = B \left[ -\frac{\cosh mL}{\sinh mL} \sinh mx + \cosh mx \right] + \frac{T_i}{P}$$

$$= \left[ \bar{F}(p) - \frac{T_i}{P} \right] \frac{\sinh [m(L-x)]}{\sinh mL} + \frac{T_i}{P}$$

$$T(x,t) = \mathcal{L}^{-1} \left\{ \bar{F}(p) \frac{\sinh [m(L-x)]}{\sinh mL} \right\} - T_i \mathcal{L}^{-1} \left\{ \frac{\sinh [m(L-x)]}{P \sinh mL} \right\} + T_i$$

$$T(x,t) - T_i = \int_{t'=0}^t F(t-t') \mathcal{L}^{-1} \left\{ \frac{\sinh [m(L-x)]}{\sinh mL} \right\} dt'$$

$$- T_i \left[ \frac{L-x}{L} + \frac{2}{\pi} \sum_{n=1}^{\infty} \frac{(-1)^n}{n} e^{-\alpha n^2 \pi^2 t / L^2} \sin \frac{n \pi (L-x)}{L} \right]$$

$$T(x,t) - T_i = \frac{2\pi\alpha}{L^2} \int_{t'=0}^t F(t-t') \left[ \sum_{n=1}^{\infty} (-1)^n e^{-\alpha n^2 \pi^2 t'/L^2} \sin \frac{n\pi(L-x)}{L} \right] dt'$$

$$- T_i \left[ \frac{L-x}{L} + \frac{2}{\pi} \sum_{n=1}^{\infty} \frac{(-1)^n}{n} e^{-\alpha n^2 \pi^2 t/L^2} \sin \frac{n\pi(L-x)}{L} \right]$$

Since  $\sin \frac{n\pi(L-x)}{L} = -(-1)^n \sin \frac{n\pi}{L} x$

$$T(x,t) - T_i = \frac{2\pi\alpha}{L^2} \int_{t'=0}^t F(t-t') \left[ \sum_{n=1}^{\infty} n e^{-\alpha n^2 \pi^2 t'/L^2} \sin \frac{n\pi}{L} x \right] dt'$$

$$- T_i \left[ 1 - \frac{x}{L} - \frac{2}{\pi} \sum_{n=1}^{\infty} \frac{1}{n} e^{-\alpha n^2 \pi^2 t/L^2} \sin \frac{n\pi}{L} x \right]$$

In particular, if  $F(t) = T_w = \text{const.}$

$$T(x,t) - T_i = \frac{2}{\pi} T_w \sum_{n=1}^{\infty} \frac{1}{n} \left[ 1 - e^{-\alpha n^2 \pi^2 t/L^2} \right] \sin \frac{n\pi}{L} x$$

$$- T_i \left[ 1 - \frac{x}{L} - \frac{2}{\pi} \sum_{n=1}^{\infty} \frac{1}{n} e^{-\alpha n^2 \pi^2 t/L^2} \sin \frac{n\pi}{L} x \right]$$


or

$$\frac{T(x,t) - T_i}{T_w - T_i} = 1 - \frac{x}{L} - \frac{2}{\pi} \sum_{n=1}^{\infty} \frac{1}{n} e^{-\alpha n^2 \pi^2 t/L^2} \sin \frac{n\pi}{L} x$$

where we have used

$$1 - \frac{x}{L} = \frac{2}{\pi} \sum_{n=1}^{\infty} \frac{1}{n} \sin \frac{n\pi}{L} x$$

PROB. B.9: (a)



$$\frac{\partial^2 T^{(1)}}{\partial x^2} = \frac{1}{\alpha_1} \frac{\partial T^{(1)}}{\partial t} ; \quad \frac{\partial^2 T^{(2)}}{\partial x^2} = \frac{1}{\alpha_2} \frac{\partial T^{(2)}}{\partial t}$$

$$T^{(1)}(x,0) = T_1 ; \quad T^{(2)}(x,0) = T_2$$

$$\lim_{x \rightarrow -\infty} T^{(1)}(x,t) = T_1 ; \quad \lim_{x \rightarrow \infty} T^{(2)}(x,t) = T_2$$

$$\rightarrow T^{(1)}(0,t) = T^{(2)}(0,t)$$

$$k_1 \left. \frac{\partial T^{(1)}}{\partial x} \right|_{x=0} = k_2 \left. \frac{\partial T^{(2)}}{\partial x} \right|_{x=0}$$

Assuming perfect contact  
at the interface

The formulation of the problem in terms of

$$\theta_1(x,t) = T^{(1)}(x,t) - T_1 \text{ and } \theta_2(x,t) = T^{(2)}(x,t) - T_2 :$$

$$\frac{\partial^2 \theta_1}{\partial x^2} = \frac{1}{\alpha_1} \frac{\partial \theta_1}{\partial t} ; \quad \frac{\partial^2 \theta_2}{\partial x^2} = \frac{1}{\alpha_2} \frac{\partial \theta_2}{\partial t}$$

$$\theta_1(x,0) = 0 ; \quad \theta_2(x,0) = 0$$

$$\lim_{x \rightarrow -\infty} \theta_1(x,t) = 0 ; \quad \lim_{x \rightarrow \infty} \theta_2(x,t) = 0$$

$$\theta_1(0,t) + T_1 = \theta_2(0,t) + T_2$$

$$k_1 \left. \frac{\partial \theta_1}{\partial x} \right|_{x=0} = k_2 \left. \frac{\partial \theta_2}{\partial x} \right|_{x=0}$$

Let

$$\bar{\theta}_1(x,p) = \int_0^\infty e^{-pt} \theta_1(x,t) dt \text{ and } \bar{\theta}_2(x,p) = \int_0^\infty e^{-pt} \theta_2(x,t) dt$$

Then, the transformed differential equations and boundary conditions become

$$\frac{d^2\bar{\theta}_1}{dx^2} - \frac{P}{\alpha_1} \bar{\theta}_1(x, P) = 0 \quad ; \quad \frac{d^2\bar{\theta}_2}{dx^2} - \frac{P}{\alpha_2} \bar{\theta}_2(x, P) = 0$$

$$\lim_{x \rightarrow -\infty} \bar{\theta}_1(x, P) = 0 \quad ; \quad \lim_{x \rightarrow \infty} \bar{\theta}_2(x, P) = 0$$

$$\bar{\theta}_1(0, P) + \frac{T_1}{P} = \bar{\theta}_2(0, P) + \frac{T_2}{P}$$

$$k_1 \left. \frac{d\bar{\theta}_1}{dx} \right|_{x=0} = k_2 \left. \frac{d\bar{\theta}_2}{dx} \right|_{x=0}$$

The solutions of the differential equations:

$$\bar{\theta}_1(x, P) = A_1 e^{m_1 x} + A_2 \bar{e}^{-m_1 x}, \quad m_1^2 = \frac{P}{\alpha_1}$$

$$\bar{\theta}_2(x, P) = B_1 e^{m_2 x} + B_2 \bar{e}^{-m_2 x}, \quad m_2^2 = \frac{P}{\alpha_2}$$

$$\lim_{x \rightarrow -\infty} \bar{\theta}_1 = 0 \Rightarrow A_2 = 0 \quad \text{and} \quad \lim_{x \rightarrow \infty} \bar{\theta}_2 = 0 \Rightarrow B_1 = 0$$

The boundary conditions at  $x=0$  give

$$\left. \begin{array}{l} A_1 + \frac{T_1}{P} = B_2 + \frac{T_2}{P} \\ k_1 m_1 A_1 = -k_2 m_2 B_2 \end{array} \right\} \quad \begin{array}{l} A_1 = -\frac{T_1 - T_2}{P} \frac{1}{1+m} \\ B_2 = \frac{T_1 - T_2}{P} \frac{m}{1+m} \end{array}$$

where

$$m = \frac{k_1}{k_2} \sqrt{\frac{\alpha_2}{\alpha_1}} = \sqrt{\frac{k_1 s_1 c_1}{k_2 s_2 c_2}}$$

Thus,

$$\bar{\theta}_1(x, P) = -\frac{T_1 - T_2}{P} \frac{1}{1+m} e^{\sqrt{\frac{P}{\alpha_1}} x} \quad \text{and} \quad \bar{\theta}_2(x, P) = \frac{T_1 - T_2}{P} \frac{m}{1+m} \bar{e}^{-\sqrt{\frac{P}{\alpha_2}} x}$$

The inversions give

$$\theta_1(x,t) = \frac{T_2 - T_1}{1+m} \operatorname{erfc}\left(-\frac{x}{2\sqrt{\alpha_1 t}}\right), \quad x < 0$$

$$\theta_2(x,t) = m \frac{T_1 - T_2}{1+m} \operatorname{erfc}\left(\frac{x}{2\sqrt{\alpha_2 t}}\right), \quad x > 0$$

Thus,

$$T^{(1)}(x,t) = T_1 + \frac{T_2 - T_1}{1+m} \operatorname{erfc}\left(\frac{|x|}{2\sqrt{\alpha_1 t}}\right), \quad x < 0$$

$$T^{(2)}(x,t) = T_2 + m \frac{T_1 - T_2}{1+m} \operatorname{erfc}\left(\frac{x}{2\sqrt{\alpha_2 t}}\right), \quad x > 0$$

or

$$T^{(1)}(x,t) = \frac{m T_1 + T_2}{1+m} \operatorname{erfc}\left(\frac{|x|}{2\sqrt{\alpha_1 t}}\right), \quad x < 0$$

$$T^{(2)}(x,t) = \frac{m T_1 + T_2}{1+m} \operatorname{erfc}\left(\frac{x}{2\sqrt{\alpha_2 t}}\right), \quad x > 0$$

$$\therefore T(x,t) = \frac{m T_1 + T_2}{1+m} \operatorname{erfc}\left(\frac{|x|}{2\sqrt{\alpha t}}\right), \quad -\infty < x < \infty$$

where  $\alpha = \begin{cases} \alpha_1 & \text{when } x < 0 \\ \alpha_2 & \text{when } x > 0 \end{cases}$

(b) The interface temperature :

$$T(0,t) = T^{(1)}(0,t) = T^{(2)}(0,t) = \frac{m T_1 + T_2}{1+m}, \quad t \geq 0$$

↑  
Constant

PROB. 8.10: Formulation of the problem (see Prob. 6.1):

$$\frac{d\theta}{dt} + m \theta(t) = \frac{\dot{q}_0}{\rho C} e^{-\beta t}, \quad m = \frac{hA}{\rho V c}$$

$$\theta(0) = 0$$

The transformed problem:

$$\bar{\theta}(p) = \frac{\dot{q}_0}{\rho C} \frac{1}{p+\beta} \frac{1}{p+m} = \begin{cases} \frac{\dot{q}_0}{\rho C} \frac{1}{m-\beta} \left[ \frac{1}{p+\beta} - \frac{1}{p+m} \right], & \beta \neq m \\ \frac{\dot{q}_0}{\rho C} \frac{1}{(p+\beta)^2}, & \beta = m \end{cases}$$

The inversion gives:

$$\theta(t) = T(t) - T_{\infty} = \begin{cases} \frac{\dot{q}_0}{\rho C} \frac{1}{m-\beta} [\bar{e}^{\beta t} - \bar{e}^{mt}], & \beta \neq m \\ \frac{\dot{q}_0}{\rho C} t \bar{e}^{-\beta t}, & \beta = m \end{cases}$$

The maximum temperature will be reached when

$$\frac{dT}{dt} = 0 \Rightarrow t = \begin{cases} \frac{1}{m-\beta} \ln \frac{m}{\beta}, & \beta \neq m \\ \frac{1}{\beta}, & \beta = m \end{cases}$$

and it will be

$$T_{\max} - T_{\infty} = \begin{cases} \frac{\dot{q}_0}{\rho C} \frac{1}{m-\beta} \left[ \left(\frac{m}{\beta}\right)^{-\frac{1}{m-\beta}} - \left(\frac{m}{\beta}\right)^{-\frac{m}{m-\beta}} \right], & \beta \neq m \\ \frac{\dot{q}_0}{\rho C} \frac{1}{\beta}, & \beta = m \end{cases}$$

PROB. 8.12: Formulation of the problem in terms of  $\theta(x,t) = T(x,t) - T_i$ :

$$\left\{ \begin{array}{l} \frac{\partial^2 \theta}{\partial x^2} = \frac{1}{\alpha} \frac{\partial \theta}{\partial t} \\ \theta(x,0) = 0 \\ \frac{\partial \theta}{\partial x} \Big|_{x=0} = 0; \quad k \frac{\partial \theta}{\partial x} \Big|_{x=L} = q''_w \end{array} \right\}$$

$$\text{Let } \bar{\theta}(x,p) = \sum \{ \theta(x,t) \} = \int_0^\infty e^{pt} \theta(x,t) dt$$

Then, the transformed differential equation and the boundary conditions become

$$\frac{d^2 \bar{\theta}}{dx^2} - m^2 \bar{\theta}(x,p) = 0, \quad m^2 = \frac{p}{\alpha}$$

$$\frac{d \bar{\theta}}{dx} \Big|_{x=0} = 0; \quad \frac{d \bar{\theta}}{dx} \Big|_{x=L} = \frac{q''_w}{kp}$$



$$\bar{\theta}(x,p) = \frac{q''_w}{kpm} \frac{\cosh mx}{\sinh mL}$$

$$\frac{\cosh mx}{\sinh mL} = \frac{e^{mx} + e^{-mx}}{e^{mL} - e^{-mL}} = e^{-mL} \frac{e^{mx} + e^{-mx}}{1 - e^{2mL}} = \left[ e^{-m(L-x)} + e^{-m(L+x)} \right] \frac{1}{1 - e^{2mL}}$$

$$= \left[ e^{-m(L-x)} + e^{-m(L+x)} \right] \sum_{k=0}^{\infty} e^{-2kmL}$$

$$= \sum_{k=0}^{\infty} \left\{ \exp[-m(nL-x)] + \exp[-m(nL+x)] \right\}$$

where  $n = 2k+1$


$$\therefore \bar{\theta}(x,p) = \frac{q''_{tw}}{kp^m} \sum_{k=0}^{\infty} \left\{ \exp[-m(nL-x)] + \exp[-m(nL+x)] \right\}$$

From Appendix D,

$$\frac{e^{-x\sqrt{p/\alpha}}}{p\sqrt{p/\alpha}} = 2 \left( \frac{\alpha t}{\pi} \right)^{1/2} e^{-x^2/4\alpha t} \times \operatorname{erfc} \left[ \frac{x}{2(\alpha t)^{1/2}} \right]$$

Thus, the inversion gives

$$\begin{aligned} \theta(x,t) &= \frac{q''_{tw}}{k} \sum_{k=0}^{\infty} \left\{ 2 \left( \frac{\alpha t}{\pi} \right)^{1/2} e^{-(nL-x)^2/4\alpha t} - (nL-x) \operatorname{erfc} \left[ \frac{nL-x}{2(\alpha t)^{1/2}} \right] \right. \\ &\quad \left. + 2 \left( \frac{\alpha t}{\pi} \right)^{1/2} e^{-(nL+x)^2/4\alpha t} - (nL+x) \operatorname{erfc} \left[ \frac{nL+x}{2(\alpha t)^{1/2}} \right] \right\} \end{aligned}$$



$$\theta(x,t) = \frac{2q''_{tw}\sqrt{\alpha t}}{k} \sum_{k=0}^{\infty} \left\{ i \operatorname{erfc} \frac{(2k+1)L-x}{2\sqrt{\alpha t}} + i \operatorname{erfc} \frac{(2k+1)L+x}{2\sqrt{\alpha t}} \right\}$$

where

$$i \operatorname{erfc}(x) = \frac{1}{\sqrt{\pi t}} e^{-x^2} - x \operatorname{erfc}(x)$$

PROB. 8.14: Formulation in terms of  $\theta(r,t) = T(r,t) - T_i$ :

$$\frac{\partial^2 \theta}{\partial r^2} + \frac{2}{r} \frac{\partial \theta}{\partial r} = \frac{1}{\alpha} \frac{\partial \theta}{\partial t}$$

$$\theta(r,0) = 0$$

$$\theta(0,t) = \text{finite}; \quad k \left. \frac{\partial \theta}{\partial r} \right|_{r=r_0} = q''_{tw}$$

The transformed problem:

$$\frac{d^2 \bar{\theta}}{dr^2} + \frac{2}{r} \frac{d\bar{\theta}}{dr} = \frac{1}{\alpha} \left[ p \bar{\theta}(r,p) - \theta(r,0) \right] \quad \text{where } \bar{\theta}(r,p) = \int_0^\infty e^{-pt} \theta(r,t) dt$$

$$\Rightarrow \frac{d^2(r\bar{\theta})}{dr^2} - m^2(r\bar{\theta}) = 0, \quad m^2 = \frac{p}{\alpha}$$

$$\therefore \bar{\theta}(r,p) = \frac{1}{r} [C_1 \sinh mr + C_2 \cosh mr]$$

$$\bar{\theta}(0,p) = \text{finite} \Rightarrow C_2 \equiv 0$$

$$\left. \frac{d\bar{\theta}}{dr} \right|_{r=r_0} = \frac{1}{p} \frac{q''_{tw}}{k} \Rightarrow C_1 = \frac{q''_{tw} r_0^2}{k} \frac{1}{p} \frac{1}{m r_0 \cosh m r_0 - \sinh m r_0}$$

$$\therefore \bar{\theta}(r,p) = \frac{q''_{tw} r_0}{k} \left( \frac{r_0}{r} \right) \frac{1}{p} \frac{\sinh mr}{m r_0 \cosh m r_0 - \sinh m r_0}$$

Thus,

$$\theta(r,t) = \frac{q''_{tw} r_0}{k} \left( \frac{r_0}{r} \right) \mathcal{L}^{-1} \left\{ \frac{1}{p} \frac{\sinh mr}{m r_0 \cosh m r_0 - \sinh m r_0} \right\}$$

Note that

$$\mathcal{L}^{-1} \left\{ \frac{1}{p} \frac{\sinh mr}{m r_0 \cosh m r_0 - \sinh m r_0} \right\} = \int_0^t \mathcal{L}^{-1} \left\{ \frac{\sinh mr}{m r_0 \cosh m r_0 - \sinh m r_0} \right\} dt$$

On the other hand,

$$\mathcal{L}^{-1} \left\{ \frac{\sinh mr}{mr_0 \cosh mr_0 - \sinh mr_0} \right\} = \mathcal{L}^{-1} \left\{ \frac{P(p)}{Q(p)} \right\} = \sum_{n=0}^{\infty} \frac{P(p_n)}{dQ(p_n)} e^{p_n t}$$

where  $p_n$ 's are the zeros of

Heaviside's expansion theorem

$$mr_0 \cosh mr_0 - \sinh mr_0 = 0 \quad \text{or} \quad mr_0 \cos i m r_0 + i \sin i m r_0 = 0, i = \sqrt{-1}$$

or

$$\tan imr_0 = imr_0 \quad \text{or} \quad \tan \lambda r_0 = \lambda r_0 \quad \text{where } \lambda^2 = -m^2 = -\frac{p}{\alpha}$$



$$\lambda_n^2 \left( = -\frac{p_n}{\alpha} \right), \quad n=0, 1, 2, \dots$$

For  $n=0$ ,

$$\lambda_0 = 0 \Rightarrow \underbrace{p_0}_{m=0} = 0 \Rightarrow P(0) = 0$$

$$\frac{dQ}{dp} = \frac{r_0}{2\sqrt{\alpha p}} \left\{ \cosh mr_0 + mr_0 \sinh mr_0 - \cosh mr_0 \right\} = \frac{r_0^2}{2\alpha} \sinh mr_0$$

$$\frac{dQ(0)}{dp} = 0$$

that the first term in Heaviside's expansion is  $\frac{0}{0}$ . Thus, this term can be calculated as

$$\lim_{p_n \rightarrow 0} \frac{\sinh mr}{\frac{r_0^2}{2\alpha} \sinh mr_0} = \frac{2\alpha}{r_0^2} \lim_{p \rightarrow 0} \frac{r \cosh mr \cdot \frac{dm}{dp}}{r_0 \cosh mr_0 \cdot \frac{dm}{dp}} = \frac{2\alpha}{r_0^2} \left( \frac{r}{r_0} \right)$$

$$\text{For } n \geq 1, \quad p_n = -\alpha \lambda_n^2 \rightarrow m_n = -i \lambda_n$$

where  $\lambda_n$ 's are non-zero zeros of  $\tan \lambda r_0 = \lambda r_0$

$$P(p_n) = \sinh m_n r = \sinh(-i \lambda_n r) = -i \sin \lambda_n r$$

$$\frac{dQ}{dp} = \frac{r_0^2}{2\alpha} \sinh m_n r_0 = \frac{r_0}{2\alpha} \sinh (-i\lambda_n) r_0 = -\frac{r_0}{2\alpha} i \sin \lambda_n r_0$$

Therefore,

$$\mathcal{L}^{-1} \left\{ \frac{\sinh mr}{mr_0 \cosh mr_0 - \sinh mr_0} \right\} = \frac{2\alpha}{r_0^2} \left( \frac{r}{r_0} \right) + \underbrace{\sum_{n=1}^{\infty} \frac{2\alpha}{r_0^2} \frac{\sin \lambda_n r}{\sin \lambda_n r_0} e^{-\alpha \lambda_n^2 t}}_{\text{and}}$$

and

$$\mathcal{L}^{-1} \left\{ \frac{1}{p} \frac{\sinh mr}{mr_0 \cosh mr_0 - \sinh mr_0} \right\} = \int_0^t \left\{ \dots \right\} dt$$

$$= \frac{2\alpha r}{r_0^3} t + \frac{2}{r_0^2} \sum_{n=1}^{\infty} \frac{\sin \lambda_n r}{\lambda_n^2 \sin \lambda_n r_0} (1 - e^{-\alpha \lambda_n^2 t})$$

Now, the solution can be written as

$$\theta(r,t) = T(r,t) - T_i = \frac{q''_w}{kr_0} \left\{ 2\alpha t + 2 \frac{r_0}{r} \sum_{n=1}^{\infty} \frac{\sin \lambda_n r}{\lambda_n^2 \sin \lambda_n r_0} (1 - e^{-\alpha \lambda_n^2 t}) \right\}$$

It can be shown that

$$2 \frac{r_0}{r} \sum_{n=1}^{\infty} \frac{\sin \lambda_n r}{\lambda_n^2 \sin \lambda_n r_0} = \frac{r_0^2}{2} \left[ \left( \frac{r}{r_0} \right)^2 - \frac{3}{5} \right] = \frac{5r^2 - 3r_0^2}{10}$$

Then, the solution can be rewritten as

$$T(r,t) - T_i = \frac{q''_w}{kr_0} \left\{ \frac{5r^2 - 3r_0^2}{10} + 2\alpha t - 2 \frac{r_0}{r} \sum_{n=1}^{\infty} \frac{\sin \lambda_n r}{\lambda_n^2 \sin \lambda_n r_0} e^{-\alpha \lambda_n^2 t} \right\}$$

PROB.B.16: formulation in terms of  $\theta(r,t) = T(r,t) - T_i$ :

$$\frac{\partial^2 \theta}{\partial r^2} + \frac{1}{r} \frac{\partial \theta}{\partial r} + \frac{\dot{q}}{k} = \frac{1}{\alpha} \frac{\partial \theta}{\partial t}$$

$$\theta(r,0) = 0$$

$$\theta(0,t) = \text{finite}; \quad \theta(r_0, t) = 0$$

The transformed problem:

$$\frac{d^2 \bar{\theta}}{dr^2} + \frac{1}{r} \frac{d\bar{\theta}}{dr} + \frac{\dot{q}}{kp} = \frac{1}{\alpha} [p \bar{\theta}(r,p) - \bar{\theta}(r_0)]$$

$$\bar{\theta}(0,p) = \text{finite}; \quad \bar{\theta}(r_0, p) = 0$$



$$\bar{\theta}(r,p) = \frac{\dot{q}\alpha}{kp^2} \left[ 1 - \frac{J_0(mr)}{J_0(mr_0)} \right], \quad m = \sqrt{\frac{p}{\alpha}}$$

$$\therefore \theta(r,t) = \frac{\dot{q}\alpha}{k} \left[ t - \int_0^t \left\{ 1 - 2 \sum_{k=1}^{\infty} \frac{J_0(\lambda_k \frac{r}{r_0})}{\lambda_k J_1(\lambda_k)} e^{-\frac{\alpha \lambda_k^2}{r_0^2} t'} \right\} dt' \right]$$

where  $\lambda_k$ 's are the zeros of  $J_0(z)$ .

Thus,

$$\theta(r,t) = -2 \frac{\dot{q}r_0^2}{k} \left\{ \underbrace{\sum_{n=1}^{\infty} \frac{J_0(\lambda_n \frac{r}{r_0})}{\lambda_n^3 J_1(\lambda_n)} e^{-\frac{\alpha \lambda_n^2}{r_0^2} t}}_{-\sum_{n=1}^{\infty} \frac{J_0(\lambda_n \frac{r}{r_0})}{\lambda_n^3 J_1(\lambda_n)}} \right\} + \frac{1}{8} \left[ 1 - \left( \frac{r}{r_0} \right)^2 \right]$$

or

$$T(r,t) - T_i = -2 \frac{\dot{q}r_0^2}{k} \sum_{n=1}^{\infty} \frac{J_0(\lambda_n \frac{r}{r_0})}{\lambda_n^3 J_1(\lambda_n)} e^{-\frac{\alpha \lambda_n^2}{r_0^2} t} + \frac{\dot{q}r_0^2}{4k} \left[ 1 - \left( \frac{r}{r_0} \right)^2 \right]$$

PROB. 8.17: The formulation in terms of  $\theta(r,t) = T(r,t) - T_i$ :

$$\frac{\partial^2 \theta}{\partial r^2} + \frac{2}{r} \frac{\partial \theta}{\partial r} + \frac{\dot{q}}{k} = \frac{1}{\alpha} \frac{\partial \theta}{\partial t}$$

$$\theta(r,0) = 0$$

$$\theta(0,t) = \text{finite}; \quad \theta(r_0, t) = 0$$

The transformed problem:

$$\frac{d^2 \bar{\theta}}{dr^2} + \frac{2}{r} \frac{d\bar{\theta}}{dr} + \frac{\dot{q}}{kp} = \frac{1}{\alpha} [p \bar{\theta}(r,p) - \theta(r,0)]$$

where

$$\bar{\theta}(r,p) = \int_0^\infty e^{-pt} \theta(r,t) dt$$

$$\frac{d^2 \bar{\theta}}{dr^2} + \frac{2}{r} \frac{d\bar{\theta}}{dr} - m^2 \bar{\theta}(r,p) = -\frac{\dot{q}}{kp}, \quad m^2 = \frac{p}{\alpha}$$

or

$$\frac{d^2}{dr^2}(r\bar{\theta}) - m^2(r\bar{\theta}) = -\frac{\dot{q}}{kp}r$$

$$\therefore \bar{\theta}(r,p) = \frac{1}{r} \left\{ C_1 \sinh mr + C_2 \cosh mr \right\} + \frac{\dot{q}\alpha}{kp^2}$$

$$\bar{\theta}(0,p) = \text{finite} \Rightarrow C_2 = 0$$

$$\bar{\theta}(r_0, p) = 0 \Rightarrow C_1 = -\frac{\dot{q}\alpha r_0}{kp^2 \sinh mr_0}$$

$$\bar{\theta}(r,p) = \frac{\dot{q}\alpha}{kp^2} \left[ 1 - \frac{r_0}{r} \frac{\sinh mr}{\sinh mr_0} \right]$$

$$\therefore \theta(r,t) = \frac{\dot{q}\alpha}{k} \left[ t - \frac{r_0}{r} \text{erf} \left\{ \frac{\sinh mr}{\sqrt{p^2 \sinh mr_0}} \right\} \right]$$

$$\begin{aligned}
 \frac{\sinh mr}{\sinh mr_0} &= \frac{e^{mr} - e^{-mr}}{e^{mr_0} - e^{-mr_0}} = \left[ \bar{e}^{m(r_0-r)} - \bar{e}^{m(r_0+tr)} \right] \frac{1}{1 - \bar{e}^{2mr_0}} \\
 &= \left[ \bar{e}^{m(r_0-r)} - \bar{e}^{-m(r_0+tr)} \right] \sum_{k=0}^{\infty} \bar{e}^{2mkf_0} \\
 &= \sum_{k=0}^{\infty} \left[ \bar{e}^{-m(nr_0-r)} - \bar{e}^{-m(nr_0+r)} \right], \quad n = 2k+1
 \end{aligned}$$

Thus,

$$\mathcal{L}^{-1} \left\{ \frac{\sinh mr}{p^2 \sinh mr_0} \right\} = \sum_{k=0}^{\infty} \left[ \mathcal{L}^{-1} \left\{ \frac{\bar{e}^{-m(nr_0-r)}}{p^2} \right\} - \mathcal{L}^{-1} \left\{ \frac{\bar{e}^{-m(nr_0+r)}}{p^2} \right\} \right]$$

From Appendix D, item 29:

$$\begin{aligned}
 \mathcal{L}^{-1} \left\{ \frac{\bar{e}^{-x\sqrt{p/a}}}{p^2} \right\} &= t + \frac{x^2}{2a} \operatorname{erfc} \left[ \frac{x}{2\sqrt{at}} \right] - x \left( \frac{t}{\pi a} \right)^{1/2} \bar{e}^{-x^2/4at} \\
 &= t - x \sqrt{\frac{t}{a}} i \operatorname{erfc} \left[ \frac{x}{2\sqrt{at}} \right]
 \end{aligned}$$

$$\text{where } i \operatorname{erfc}(x) = \frac{1}{\sqrt{\pi}} \bar{e}^{-x^2} - x \operatorname{erfc}(x)$$

Thus,

$$\mathcal{L}^{-1} \left\{ \frac{\sinh mr}{p^2 \sinh mr_0} \right\} = \sqrt{\frac{t}{a}} \sum_{k=0}^{\infty} \left\{ (nr_0+r) i \operatorname{erfc} \left[ \frac{nr_0+r}{2\sqrt{at}} \right] - (nr_0-r) i \operatorname{erfc} \left[ \frac{nr_0-r}{2\sqrt{at}} \right] \right\}$$

Then,

$$\boxed{
 \begin{aligned}
 \frac{T(r,t) - T_i}{q \alpha / k} &= t - \frac{r_0}{r} \sqrt{\frac{t}{a}} \sum_{k=0}^{\infty} \left\{ [(2k+1)r_0+r] i \operatorname{erfc} \left[ \frac{(2k+1)r_0+r}{2\sqrt{at}} \right] \right. \\
 &\quad \left. - [(2k+1)r_0-r] i \operatorname{erfc} \left[ \frac{(2k+1)r_0-r}{2\sqrt{at}} \right] \right\}
 \end{aligned}$$

PROB. 8.19: Refer to Section 8.10 in text.

$$\begin{aligned}
 \frac{\bar{\theta}}{\theta_w} &= \frac{r_0}{r} \frac{\sinh mr}{p \sinh mr_0} = \frac{r_0}{rp} \frac{e^{mr} - e^{-mr}}{e^{mr_0} - e^{-mr_0}} \\
 &= \frac{r_0}{rp} \frac{e^{-m(r_0-r)} - e^{-m(r_0+r)}}{1 - e^{-2mr_0}} \\
 &= \frac{r_0}{r} \frac{1}{p} \left[ e^{-m(r_0-r)} - e^{-m(r_0+r)} \right] \sum_{k=0}^{\infty} e^{-2mkf_0} \\
 &= \frac{r_0}{r} \frac{1}{p} \sum_{k=0}^{\infty} \left\{ e^{-m(nr_0-r)} - e^{-m(nr_0+r)} \right\}
 \end{aligned}$$

$$n = 2k+1$$

$$\therefore \frac{\theta(r,t)}{\theta_w} = \frac{r_0}{r} \sum_{k=0}^{\infty} \left[ f^{-1} \left\{ \frac{e^{-m(nr_0-r)}}{p} \right\} - f^{-1} \left\{ \frac{e^{-m(nr_0+r)}}{p} \right\} \right]$$

$$\boxed{\frac{T(r,t) - T_i}{T_w - T_i} = \frac{r_0}{r} \sum_{k=0}^{\infty} \left\{ \operatorname{erfc} \left[ \frac{(2k+1)r_0 - r}{2\sqrt{kt}} \right] - \operatorname{erfc} \left[ \frac{(2k+1)r_0 + r}{2\sqrt{kt}} \right] \right\}}$$

PROB. 8.20:

$$\left\{ \begin{array}{l} \frac{\partial^2 T}{\partial x^2} = \frac{1}{\alpha} \frac{\partial T}{\partial t} \\ T(x, 0) = T_i \\ T(0, t) = T_0 + (\Delta T)_0 \sin \omega t \\ T(x, t) \Big|_{x \rightarrow \infty} = T_i \end{array} \right\} \xrightarrow{\theta(x, t) = \frac{T(x, t) - T_0}{(\Delta T)_0}} \left\{ \begin{array}{l} \frac{\partial^2 \theta}{\partial x^2} = \frac{1}{\alpha} \frac{\partial \theta}{\partial t} \\ \theta(x, 0) = \frac{T_i - T_0}{(\Delta T)_0} = \theta_i \\ \theta(0, t) = \sin \omega t \\ \theta(x, t) \Big|_{x \rightarrow \infty} = \theta_i \end{array} \right\}$$

$$\frac{d^2 \bar{\theta}}{dx^2} = \frac{1}{\alpha} \{ p \bar{\theta}(x, p) - \theta_i \}, \quad \bar{\theta}(x, p) = \int_0^\infty e^{pt} \theta(x, t) dt$$

$$\bar{\theta}(x, p) = A e^{mx} + B \bar{e}^{-mx} + \frac{\theta_i}{p}, \quad m = \sqrt{\frac{p}{\alpha}}$$

$$\bar{\theta}(x, p) \Big|_{x \rightarrow \infty} \rightarrow \frac{\theta_i}{p} \Rightarrow A \equiv 0$$

$$\bar{\theta}(0, p) = \frac{\omega}{p^2 + \omega^2} = B + \frac{\theta_i}{p} \Rightarrow B = \frac{\omega}{p^2 + \omega^2} - \frac{\theta_i}{p}$$

$$\therefore \bar{\theta}(x, p) = \frac{\theta_i}{p} (1 - \bar{e}^{-mx}) + \frac{\omega}{p^2 + \omega^2} \bar{e}^{-mx}$$

↓

$$\theta(x, t) = \theta_i \left[ 1 - \mathcal{L}^{-1} \left\{ \frac{\bar{e}^{-x \sqrt{p/\alpha}}}{p} \right\} \right] + \omega \mathcal{L}^{-1} \left\{ \frac{\bar{e}^{-x \sqrt{p/\alpha}}}{p^2 + \omega^2} \right\}$$

$$= \theta_i \left[ 1 - \operatorname{erfc} \left( \frac{x}{2\sqrt{\alpha t}} \right) \right] + \underbrace{\int_0^t \sin \omega t' \frac{x e^{-\frac{x^2}{4\alpha(t-t')}}}{2\sqrt{\pi\alpha(t-t')^3}} dt'}_{\operatorname{erf} \left( \frac{x}{2\sqrt{\alpha t}} \right)}$$

$$\text{Let } \frac{x}{2\sqrt{\alpha(t-t')}} = \mu \Rightarrow t-t' = \frac{x^2}{4\alpha\mu^2} \quad \& \quad dt' = \frac{4\sqrt{\alpha}(t-t')^{3/2}}{x} d\mu$$

Thus,

$$\int_0^t \sin \omega t' \frac{x e^{-\frac{x^2}{4\alpha}(t-t')}}{2\sqrt{\pi\alpha(t-t')^3}} dt' = \frac{2}{\pi} \int_x^\infty \sin \left\{ \omega \left[ t - \frac{x^2}{4\alpha\mu^2} \right] \right\} e^{-\mu^2} d\mu$$

Also, by a known definite integral,\*

$$\frac{2}{\sqrt{\pi}} \int_0^\infty \sin \left\{ \omega \left[ t - \frac{x^2}{4\alpha\mu^2} \right] \right\} e^{-\mu^2} d\mu = e^{-x\sqrt{\frac{\omega}{2\alpha}}} \sin \left\{ \omega t - x \left( \frac{\omega}{2\alpha} \right)^{1/2} \right\}$$

it follows that

$$\begin{aligned} \frac{2}{\sqrt{\pi}} \int_x^\infty \sin \left\{ \omega \left[ t - \frac{x^2}{4\alpha\mu^2} \right] \right\} e^{-\mu^2} d\mu &= e^{-x\sqrt{\frac{\omega}{2\alpha}}} \sin \left\{ \omega t - x \left( \frac{\omega}{2\alpha} \right)^{1/2} \right\} \\ &\quad - \frac{2}{\sqrt{\pi}} \int_0^{2\sqrt{\alpha t}} \sin \left\{ \omega \left[ t - \frac{x^2}{4\alpha\mu^2} \right] \right\} e^{-\mu^2} d\mu \end{aligned}$$

Then, the solution can be written as

$$\begin{aligned} \theta(x,t) &= \frac{T(x,t) - T_0}{(\Delta T)_0} = e^{-x\sqrt{\frac{\omega}{2\alpha}}} \sin \left\{ \omega t - x \left( \frac{\omega}{2\alpha} \right)^{1/2} \right\} \\ &\quad + \frac{T_i - T_0}{(\Delta T)_0} \operatorname{erf} \left( \frac{x}{2\sqrt{\alpha t}} \right) - \frac{2}{\sqrt{\pi}} \int_0^{2\sqrt{\alpha t}} \sin \left\{ \omega \left[ t - \frac{x^2}{4\alpha\mu^2} \right] \right\} e^{-\mu^2} d\mu \end{aligned}$$

Note that, as  $t \rightarrow \infty$ , the last two terms die away, leaving the first term which is a steady oscillation of  $\frac{2\pi}{\omega}$ .

\* See Ref.[3], Carslaw and Jaeger, p. 65.

PROB. 8.25: The formulation in terms of  $\theta(x,t) = T(x,t) - T_i$ :

$$\frac{\partial^2 \theta}{\partial x^2} = \frac{1}{\alpha} \frac{\partial \theta}{\partial t}$$

$$\theta(x_0) = 0$$

$$\frac{\partial \theta}{\partial x} \Big|_{x=0} = 0; \quad \theta(L,t) = At$$

The transformed problem:

$$\frac{d^2 \bar{\theta}}{dx^2} - \frac{P}{\alpha} \bar{\theta}(x,p) = 0, \quad \bar{\theta}(x,p) = \int_0^\infty e^{-pt} \theta(x,t) dt$$

$$\frac{d\bar{\theta}}{dx} \Big|_{x=0} = 0; \quad \theta(L,p) = \frac{A}{p^2}$$



$$\bar{\theta}(x,p) = \frac{A}{p^2} \frac{\cosh mx}{\cosh mL}, \quad m^2 = \frac{P}{\alpha}$$

$$\therefore \theta(x,t) = A \int_0^t \mathcal{L}^{-1} \left\{ \frac{\cosh mx}{p \cosh mL} \right\} dt$$

Transform No. 40,  
Appendix D

$$= A \int_0^t \left\{ 1 + \frac{2}{L} \sum_{n=1}^{\infty} \frac{(-1)^n}{\lambda_n} e^{-\alpha \lambda_n^2 t} \cos \lambda_n x \right\} dt, \quad \lambda_n = \frac{(2n-1)\pi}{2L}$$

$$\therefore \theta(x,t) = T(x,t) - T_i = A \left\{ t + \frac{2}{\alpha L} \sum_{n=1}^{\infty} \frac{(-1)^n}{\lambda_n^3} \left( 1 - e^{-\alpha \lambda_n^2 t} \right) \cos \lambda_n x \right\}$$

or

---


$$T(x,t) - T_i = A \left\{ t - \frac{L^2}{2\alpha} \left[ 1 - \left( \frac{x}{L} \right)^2 \right] - \frac{2}{\alpha L} \sum_{n=1}^{\infty} \frac{(-1)^n}{\lambda_n^3} e^{-\alpha \lambda_n^2 t} \cos \lambda_n x \right\}$$


---

Here, we used the relation,

$$1 - \left( \frac{x}{L} \right)^2 = \frac{4}{L^3} \sum_{n=1}^{\infty} \frac{(-1)^{n+1}}{\lambda_n^3} \cos \lambda_n x, \quad 0 \leq x \leq L$$

Note that the above solution does not converge rapidly for small values of  $t$ . An alternative solution that converges rapidly for small values of  $t$  can be obtained as follows:

$$\frac{\cosh mx}{\cosh mL} = \sum_{k=0}^{\infty} (-1)^k \left\{ e^{-m(nL-x)} + e^{-m(nL+x)} \right\}, \quad n=2k+1$$

Thus,

$$\begin{aligned}\theta(x,t) &= \mathcal{L}^{-1} \left\{ \frac{A}{p^2} \frac{\cosh mx}{\cosh mL} \right\} \\ &= A \sum_{k=0}^{\infty} (-1)^k \mathcal{L}^{-1} \left\{ \frac{e^{-m(nL-x)}}{p^2} + \frac{e^{-m(nL+x)}}{p^2} \right\}\end{aligned}$$

Transform No. 29, Appendix D:

$$\mathcal{L}^{-1} \left\{ \frac{e^{-x\sqrt{p/a}}}{p^2} \right\} = \left( t + \frac{x^2}{2a} \right) \operatorname{erfc} \left[ \frac{x}{2\sqrt{at}} \right] - x \left( \frac{t}{\pi a} \right)^{1/2} e^{-x^2/4at}$$

Then, it follows that

---


$$\begin{aligned}\theta(x,t) &= T(x,t) - T_i = A \sum_{k=0}^{\infty} (-1)^k \left\{ \left[ t + \frac{(nL-x)^2}{2\alpha} \right] \operatorname{erf} \left[ \frac{nL-x}{2\sqrt{\alpha t}} \right] \right. \\ &\quad + \left[ t + \frac{(nL+x)^2}{2\alpha} \right] \operatorname{erfc} \left[ \frac{nL+x}{2\sqrt{\alpha t}} \right] \\ &\quad - (nL-x) \left( \frac{t}{\pi \alpha} \right)^{1/2} \exp \left[ - \frac{(nL-x)^2}{4\alpha t} \right] \\ &\quad \left. - (nL+x) \left( \frac{t}{\pi \alpha} \right)^{1/2} \exp \left[ - \frac{(nL+x)^2}{4\alpha t} \right] \right\}\end{aligned}$$


---

PROB. 8.26: The formulation in terms of  $\Theta(r,t) = T(r,t) - T_i$ :

$$\frac{\partial^2 \Theta}{\partial r^2} + \frac{1}{r} \frac{\partial \Theta}{\partial r} = \frac{1}{\alpha} \frac{\partial \Theta}{\partial t}$$

$$\Theta(r,0) = 0$$

$$\Theta(0,t) = \text{finite}; \quad \Theta(r_0, t) = At$$

The transformed problem:

$$\frac{d^2 \bar{\Theta}}{dr^2} + \frac{1}{r} \frac{d\bar{\Theta}}{dr} - \frac{P}{\alpha} \bar{\Theta}(r,p) = 0, \quad \bar{\Theta}(r,p) = \int_0^\infty e^{-pt} \Theta(r,t) dt$$

$$\bar{\Theta}(0,p) = \text{finite}; \quad \bar{\Theta}(r_0, p) = \frac{A}{p^2}$$



$$\bar{\Theta}(r,p) = \frac{A}{p^2} \frac{I_0(mr)}{I_0(mr_0)}, \quad m^2 = \frac{P}{\alpha}$$

$$\therefore \Theta(r,t) = A \int_0^t \left\{ 1 - 2 \sum_{n=1}^{\infty} \frac{J_0(\lambda_n \frac{r}{r_0})}{\lambda_n J_1(\lambda_n)} e^{-\alpha \frac{\lambda_n^2}{r_0^2} t'} \right\} dt'$$

where  $\lambda_n$  are the zeros of  $J_0(z)$ . (See Prob. 8.16)

Thus,

$$\Theta(r,t) = A \left\{ t + \frac{2r_0^2}{\alpha} \sum_{n=1}^{\infty} \frac{J_0(\lambda_n \frac{r}{r_0})}{\lambda_n^3 J_1(\lambda_n)} e^{-\frac{\alpha \lambda_n^2}{r_0^2} t} - \underbrace{\frac{2r_0^2}{\alpha} \sum_{n=1}^{\infty} \frac{J_0(\lambda_n \frac{r}{r_0})}{\lambda_n J_1(\lambda_n)}}_{\frac{1}{8} \left[ 1 - \left( \frac{r}{r_0} \right)^2 \right]} \right\}$$

or

$$\boxed{\Theta(r,t) = \frac{Ar_0^2}{4\alpha} \left\{ 4 \frac{\alpha t}{r_0^2} - \left[ 1 - \left( \frac{r}{r_0} \right)^2 \right] + 8 \sum_{n=1}^{\infty} \frac{J_0(\lambda_n \frac{r}{r_0})}{\lambda_n^3 J_1(\lambda_n)} \exp\left(-\frac{\alpha \lambda_n^2}{r_0^2} t\right) \right\}}$$

PROB. 8.27: The formulation in terms of  $\theta(r,t) = T(r,t) - T_i$ :

$$\frac{\partial^2 \theta}{\partial r^2} + \frac{2}{r} \frac{\partial \theta}{\partial r} = \frac{1}{\alpha} \frac{\partial \theta}{\partial t}$$

$$\theta(r,0) = 0$$

$$\theta(0,t) = \text{finite}; \quad \theta(r_0, t) = At$$

The transformed problem:

$$\boxed{\frac{d^2 \bar{\theta}}{dr^2} + \frac{2}{r} \frac{d\bar{\theta}}{dr} - \frac{P}{\alpha} \bar{\theta}(r,p) = 0, \quad \bar{\theta}(r,p) = \int_0^\infty e^{-pt} \theta(r,t) dt}$$

$$\bar{\theta}(0,p) = \text{finite}; \quad \bar{\theta}(r_0, p) = \frac{A}{p^2}$$

$$\rightarrow \frac{d^2}{dr^2} (r\bar{\theta}) - m^2 (r\bar{\theta}) = 0, \quad m^2 = \frac{P}{\alpha}$$

$$\therefore \bar{\theta}(r,p) = \frac{1}{r} \left\{ C_1 \cosh mr + C_2 \sinh mr \right\}$$

$$\bar{\theta}(0,p) = \text{finite} \Rightarrow C_1 = 0$$

$$\bar{\theta}(r_0, p) = \frac{A}{p^2} \Rightarrow \frac{1}{r_0} C_2 \sinh mr_0 = \frac{A}{p^2} \Rightarrow C_2 = \frac{A r_0}{p^2 \sinh mr_0}$$

$$\bar{\theta}(r,p) = A \frac{r_0}{r} \frac{\sinh mr}{p^2 \sinh mr_0} = A \frac{r_0}{r} \frac{1}{p^2} \frac{e^{mr} - \bar{e}^{mr}}{e^{mr_0} - \bar{e}^{mr_0}}$$

$$= A \frac{r_0}{r} \frac{1}{p^2} \left[ \bar{e}^{m(r_0-r)} - \bar{e}^{m(r_0+r)} \right] \sum_{k=0}^{\infty} e^{-2mk r_0}$$

$$= A \frac{r_0}{r} \frac{1}{p^2} \sum_{k=0}^{\infty} \left\{ \bar{e}^{-m(n r_0 - r)} - \bar{e}^{-m(n r_0 + r)} \right\}, \quad n = 2k+1$$

$$\theta(r,t) = A \frac{r_0}{r} \sum_{k=0}^{\infty} \left[ \mathcal{L}^{-1} \left\{ \frac{\bar{e}^{-m(nr_0-r)}}{p^2} \right\} - \mathcal{L}^{-1} \left\{ \frac{\bar{e}^{-m(nr_0+r)}}{p^2} \right\} \right]$$

From Appendix D, Item 29:

$$\begin{aligned} \mathcal{L}^{-1} \left\{ \frac{\bar{e}^{-x\sqrt{p/a}}}{p^2} \right\} &= \left( t + \frac{x^2}{2a} \right) \operatorname{erfc} \left[ \frac{x}{2\sqrt{at}} \right] - x \left( \frac{t}{\pi a} \right)^{1/2} e^{-x^2/4at} \\ &= t \operatorname{erfc} \left[ \frac{x}{2\sqrt{at}} \right] - x \sqrt{\frac{t}{a}} i \operatorname{erfc} \left[ \frac{x}{2\sqrt{at}} \right] \end{aligned}$$

where

$$i \operatorname{erfc}(x) = \frac{1}{\sqrt{\pi}} e^{-x^2} - x \operatorname{erfc}(x)$$

Then,

$$\mathcal{L}^{-1} \left\{ \frac{\bar{e}^{-m(nr_0+r)}}{p^2} \right\} = t \operatorname{erfc} \left[ \frac{(nr_0+r)}{2\sqrt{at}} \right] - (nr_0+r) \sqrt{\frac{t}{a}} i \operatorname{erfc} \left[ \frac{nr_0+r}{2\sqrt{at}} \right]$$

$$n = 2k+1$$

Thus,

$$\theta(r,t) = A \frac{r_0 t}{r} \sum_{k=0}^{\infty} \left\{ \operatorname{erfc} \left[ \frac{(2k+1)r_0 - r}{2\sqrt{at}} \right] + \operatorname{erfc} \left[ \frac{(2k+1)r_0 + r}{2\sqrt{at}} \right] \right\}$$

$$+ A \frac{r_0}{r} \sqrt{\frac{t}{a}} \sum_{k=0}^{\infty} \left\{ [(2k+1)r_0 + r] i \operatorname{erfc} \left[ \frac{(2k+1)r_0 + r}{2\sqrt{at}} \right] \right.$$

$$\left. - [(2k+1)r_0 - r] i \operatorname{erfc} \left[ \frac{(2k+1)r_0 - r}{2\sqrt{at}} \right] \right\}$$





**CRC Press**

Taylor & Francis Group  
an **informa** business

[www.taylorandfrancisgroup.com](http://www.taylorandfrancisgroup.com)

6000 Broken Sound Parkway, NW  
Suite 300, Boca Raton, FL 33487  
270 Madison Avenue  
New York, NY 10016  
2 Park Square, Milton Park  
Abingdon, Oxon OX14 4RN, UK

K14686

9 781466 507845