Data Science Analysis - Quiz 3

Pushkal Mishra EE20BTECH11042

April 28, 2023

Importing Libraries

```
[1]: import numpy as np from scipy.stats import chi2 import matplotlib.pyplot as plt
```

Question 1

```
[2]: def calculate_score(H0_exp1, H0_exp2, H0_exp1_err, H0_exp2_err):

# Significance from p-value is also known as Z-score

z_score = np.abs(H0_exp1 - H0_exp2) / np.sqrt(H0_exp1_err ** 2 + H0_exp2_err

→** 2)

return z_score
```

```
[3]: H0_exp1, H0_exp2 = 67, 71
H0_exp1_err, H0_exp2_err = 2.3, 1.3

z_score = calculate_score(H0_exp1, H0_exp2, H0_exp1_err, H0_exp2_err)

print(f"The Z-score between measurements: {z_score:.5f} sigmas.")
```

The Z-score between measurements: 1.51402 sigmas.

Question 2

```
[4]: free_params = 2
confidence_interval = 0.85

# This formula is from the slides of Second and Third week on page 17
del_chi2 = chi2(free_params).ppf(confidence_interval)

print(f"The value for constant del Chi2 for an {int(confidence_interval * 100)}%

→confidence interval: {del_chi2:.5f}")
```

The value for constant del Chi2 for an 85% confidence interval: 3.79424

Question 3

```
[5]: def calculate_del_BIC(num_data, chi2_vals, free_params):
         A_chi2, B_chi2 = chi2_vals
         A_free, B_free = free_params
         # Note that in this question, the reduced Chi2 value is given
         # To find out Chi2 value, we need to multiply it with number of free
      → parameters (given in slides)
         BIC_A = (A_free * np.log(num_data)) - (2 * np.log(A_free * A_chi2))
         BIC_B = (B_free * np.log(num_data)) - (2 * np.log(B_free * B_chi2))
         return BIC_A, BIC_B, (BIC_A - BIC_B)
[6]: num_data_points = 10
     A_{red_chi2}, A_{free} = 1.3, 6
     B_{red_chi2}, B_{free} = 0.7, 8
     BIC_A, BIC_B, del_BIC = calculate_del_BIC(num_data_points, [A_red_chi2,_
     →B_red_chi2], [A_free, B_free])
     print(f"BIC of Model A: {BIC_A:.5f}")
     print(f"BIC of Model B: {BIC_B:.5f}\n")
     print(f"The difference in BIC of two models are: {del_BIC:.5f}")
    BIC of Model A: 9.70726
    BIC of Model B: 14.97515
```

The difference in BIC of two models are: -5.26788

Question 4

```
[7]: mean = 10
    number = 1000
    random_variables = np.random.poisson(mean, number)
    print(random_variables)

fig = plt.figure(figsize = (12, 6))
    plt.hist(random_variables, bins = 20, label = "Poisson Random Numbers")
    plt.title("Histogram plot for Poisson distributed random variables", size = 15)
    plt.xlabel("Value", size = 15)
    plt.ylabel("Count", size = 15)
    plt.legend()
    plt.show()
```

[6 6 11 10 8 11 9 5 9 10 9 12 9 14 16 11 5 8 5 8 11 6 6 12 6 14 10 9 7 14 10 7 14 6 7 7 2 7 11 7 9 7 9 10 16 8 7 9 14 9 4 13 8 11 15 17 9 15 11 4 14 12 14 5 6 10 14 8 9 12 7 13

