

CS 105: Department Introductory Course on Discrete Structures

Instructor : S. Akshay

Aug 28, 2023

Lecture 09 – Basic Mathematical Structures
Uncountable Sets and relations

Countable and countably infinite sets

Definition

- ▶ Set C is called **countably infinite**, if there is a bijection from set C to \mathbb{N} .
- ▶ A set is **countable** if it is finite or countably infinite.

Countable and countably infinite sets

Definition

- ▶ Set C is called **countably infinite**, if there is a bijection from set C to \mathbb{N} .
- ▶ A set is **countable** if it is finite or countably infinite.

Properties

- ▶ Union of countable sets is countable (e.g., $\mathbb{Z} \cup \{\sqrt{2}\}$)
- ▶ Cartesian product of countable sets is countable (e.g., $\mathbb{Z} \times \mathbb{N}$)

Countable and countably infinite sets

Definition

- ▶ Set C is called **countably infinite**, if there is a bijection from set C to \mathbb{N} .
- ▶ A set is **countable** if it is finite or countably infinite.

Properties

- ▶ Union of countable sets is countable (e.g., $\mathbb{Z} \cup \{\sqrt{2}\}$)
- ▶ Cartesian product of countable sets is countable (e.g., $\mathbb{Z} \times \mathbb{N}$)

Qn. Show that set of primes P is countable.

▶ Proof 1

1. In Lecture 1, we showed that set of primes is infinite.

Countable and countably infinite sets

Definition

- ▶ Set C is called **countably infinite**, if there is a bijection from set C to \mathbb{N} .
- ▶ A set is **countable** if it is finite or countably infinite.

Properties

- ▶ Union of countable sets is countable (e.g., $\mathbb{Z} \cup \{\sqrt{2}\}$)
- ▶ Cartesian product of countable sets is countable (e.g., $\mathbb{Z} \times \mathbb{N}$)

Qn. Show that set of primes P is countable.

▶ Proof 1

1. In Lecture 1, we showed that set of primes is infinite.
2. So there is surjection from P to \mathbb{N} .

Countable and countably infinite sets

Definition

- ▶ Set C is called **countably infinite**, if there is a bijection from set C to \mathbb{N} .
- ▶ A set is **countable** if it is finite or countably infinite.

Properties

- ▶ Union of countable sets is countable (e.g., $\mathbb{Z} \cup \{\sqrt{2}\}$)
- ▶ Cartesian product of countable sets is countable (e.g., $\mathbb{Z} \times \mathbb{N}$)

Qn. Show that set of primes P is countable.

▶ Proof 1

1. In Lecture 1, we showed that set of primes is infinite.
2. So there is surjection from P to \mathbb{N} .
3. Now, $P \subseteq \mathbb{N}$. $Id : P \rightarrow \mathbb{N}$ is an injection.

Countable and countably infinite sets

Definition

- ▶ Set C is called **countably infinite**, if there is a bijection from set C to \mathbb{N} .
- ▶ A set is **countable** if it is finite or countably infinite.

Properties

- ▶ Union of countable sets is countable (e.g., $\mathbb{Z} \cup \{\sqrt{2}\}$)
- ▶ Cartesian product of countable sets is countable (e.g., $\mathbb{Z} \times \mathbb{N}$)

Qn. Show that set of primes P is countable.

▶ Proof 1

1. In Lecture 1, we showed that set of primes is infinite.
2. So there is surjection from P to \mathbb{N} .
3. Now, $P \subseteq \mathbb{N}$. $Id : P \rightarrow \mathbb{N}$ is an injection. Conclude by Schroder-Bernstein Theorem.

Countable and countably infinite sets

Definition

- ▶ Set C is called **countably infinite**, if there is a bijection from set C to \mathbb{N} .
- ▶ A set is **countable** if it is finite or countably infinite.

Properties

- ▶ Union of countable sets is countable (e.g., $\mathbb{Z} \cup \{\sqrt{2}\}$)
- ▶ Cartesian product of countable sets is countable (e.g., $\mathbb{Z} \times \mathbb{N}$)

Qn. Show that set of primes P is countable.

▶ **Proof 1**

1. In Lecture 1, we showed that set of primes is infinite.
2. So there is surjection from P to \mathbb{N} .
3. Now, $P \subseteq \mathbb{N}$. $Id : P \rightarrow \mathbb{N}$ is an injection. Conclude by Schroder-Bernstein Theorem.

- ▶ **Proof 2** Show $f : P \rightarrow \mathbb{N}$ by f maps i^{th} prime to i is a bijection

Countable sets and functions

Are the following sets countable?

- ▶ the set of all integers \mathbb{Z}
- ▶ $\mathbb{N} \times \mathbb{N}$
- ▶ $\mathbb{N} \times \mathbb{N} \times \mathbb{N}$
- ▶ the set of rationals \mathbb{Q}
- ▶ the set of all (finite and infinite) subsets of \mathbb{N}
- ▶ the set of all real numbers \mathbb{R}

Comparing \mathbb{N} and set of all subsets of \mathbb{N}

Theorem (Cantor, 1891)

There is no bijection between \mathbb{N} and the set of all subsets of \mathbb{N} .

Comparing \mathbb{N} and set of all subsets of \mathbb{N}

Theorem (Cantor, 1891)

There is no bijection between \mathbb{N} and the set of all subsets of \mathbb{N} .

Comparing \mathbb{N} and set of all subsets of \mathbb{N}

Theorem (Cantor, 1891)

There is no bijection between \mathbb{N} and the set of all subsets of \mathbb{N} .

Comparing \mathbb{N} and set of all subsets of \mathbb{N}

Theorem (Cantor, 1891)

There is no bijection between \mathbb{N} and the set of all subsets of \mathbb{N} .

Proof by contradiction: Suppose there is such a bijection, say f . This would imply that each $i \in \mathbb{N}$ maps to some set $f(i) \subseteq \mathbb{N}$.

	0	1	2	3	...
$f(0)$	✓	×	×	×	...
$f(1)$	✓	×	✓	✓	...
$f(2)$	×	×	×	×	...
$f(3)$	×	✓	×	✓	...

Comparing \mathbb{N} and set of all subsets of \mathbb{N}

Theorem (Cantor, 1891)

There is no bijection between \mathbb{N} and the set of all subsets of \mathbb{N} .

Proof by contradiction: Suppose there is such a bijection, say f . This would imply that each $i \in \mathbb{N}$ maps to some set $f(i) \subseteq \mathbb{N}$.

	0	1	2	3	...
$f(0)$	✓ ×	×	×	×	...
$f(1)$	✓	× ✓	✓	✓	...
$f(2)$	×	×	× ✓	×	...
$f(3)$	×	✓	×	✓ ×	...

- Consider the set $S \subseteq \mathbb{N}$ obtained by switching the diagonal elements, i.e., $S = \{i \in \mathbb{N} \mid i \notin f(i)\}$.

Comparing \mathbb{N} and set of all subsets of \mathbb{N}

Theorem (Cantor, 1891)

There is no bijection between \mathbb{N} and the set of all subsets of \mathbb{N} .

Proof by contradiction: Suppose there is such a bijection, say f . This would imply that each $i \in \mathbb{N}$ maps to some set $f(i) \subseteq \mathbb{N}$.

	0	1	2	3	...
$f(0)$	✓ ×	×	×	×	...
$f(1)$	✓	✗✓	✓	✓	...
$f(2)$	×	×	✗✓	×	...
$f(3)$	×	✓	×	✓ ×	...

- ▶ Consider the set $S \subseteq \mathbb{N}$ obtained by switching the diagonal elements, i.e., $S = \{i \in \mathbb{N} \mid i \notin f(i)\}$.
- ▶ As f is bij, $\exists j \in \mathbb{N}, f(j) = S$.

Comparing \mathbb{N} and set of all subsets of \mathbb{N}

Theorem (Cantor, 1891)

There is no bijection between \mathbb{N} and the set of all subsets of \mathbb{N} .

Proof by contradiction: Suppose there is such a bijection, say f . This would imply that each $i \in \mathbb{N}$ maps to some set $f(i) \subseteq \mathbb{N}$.

	0	1	2	3	...
$f(0)$	✓ ×	×	×	×	...
$f(1)$	✓	× ✓	✓	✓	...
$f(2)$	×	×	× ✓	×	...
$f(3)$	×	✓	×	✓ ×	...

- ▶ Consider the set $S \subseteq \mathbb{N}$ obtained by switching the diagonal elements, i.e., $S = \{i \in \mathbb{N} \mid i \notin f(i)\}$.
- ▶ As f is bij, $\exists j \in \mathbb{N}, f(j) = S$.
- ▶ S and $f(j)$ differ at position j , for any j .

Comparing \mathbb{N} and set of all subsets of \mathbb{N}

Theorem (Cantor, 1891)

There is no bijection between \mathbb{N} and the set of all subsets of \mathbb{N} .

Proof by contradiction: Suppose there is such a bijection, say f . This would imply that each $i \in \mathbb{N}$ maps to some set $f(i) \subseteq \mathbb{N}$.

	0	1	2	3	...
$f(0)$	✓✗	×	×	×	...
$f(1)$	✓	✗✓	✓	✓	...
$f(2)$	×	×	✗✓	×	...
$f(3)$	×	✓	×	✓✗	...

- ▶ Consider the set $S \subseteq \mathbb{N}$ obtained by switching the diagonal elements, i.e., $S = \{i \in \mathbb{N} \mid i \notin f(i)\}$.
- ▶ As f is bij, $\exists j \in \mathbb{N}, f(j) = S$.
- ▶ S and $f(j)$ differ at position j , for any j .
- ▶ Thus, $S \neq f(j)$ for all $j \in \mathbb{N}$, which is a contradiction! \square

Cantor's diagonalization

Does this proof look familiar??

Cantor's diagonalization

Does this proof look familiar??




Figure: Cantor and Russell

Cantor's diagonalization

Does this proof look familiar??




Figure: Cantor and Russell

- ▶ $S = \{i \in \mathbb{N} \mid i \notin f(i)\}$ is like the one from Russell's paradox.

Cantor's diagonalization

Does this proof look familiar??




Figure: Cantor and Russell

- ▶ $S = \{i \in \mathbb{N} \mid i \notin f(i)\}$ is like the one from Russell's paradox.
- ▶ If $\exists j \in \mathbb{N}$ such that $f(j) = S$, then we have a contradiction.
 - ▶ If $j \in S$, then $j \notin f(j) = S$.
 - ▶ If $j \notin S$, then $j \notin f(j)$, which implies $j \in S$.

Cantor's diagonalization

Does this proof look familiar??




Figure: Cantor and Russell

In fact, using diagonalization Cantor showed that...

- ▶ There cannot be a bijection between **any** set and its power set (i.e., its set of subsets). **(H.W)**
- ▶ So there is an infinite hierarchy of “larger” infinities...

Cantor's diagonalization

Does this proof look familiar??




Figure: Cantor and Russell

In fact, using diagonalization Cantor showed that...

- ▶ There cannot be a bijection between **any** set and its power set (i.e., its set of subsets). **(H.W)**
- ▶ So there is an infinite hierarchy of “larger” infinities...
- ▶ There is no bijection from \mathbb{R} to \mathbb{N} **(H.W)**. Moreover, there is a bijection from \mathbb{R} to set of subsets of \mathbb{N} .

One infinity is “strictly” bigger than another!

Theorem (Cantor, 1891)

There is no bijection between \mathbb{N} and the set of all subsets of \mathbb{N} .

One infinity is “strictly” bigger than another!

Theorem (Cantor, 1891)

There is no bijection between \mathbb{N} and the set of all subsets of \mathbb{N} .

- ▶ But, there is a surjection from set of all subsets of \mathbb{N} to \mathbb{N} .
- ▶ Thus, the “size” of $\mathcal{P}(\mathbb{N})$ is strictly greater than \mathbb{N} !

One infinity is “strictly” bigger than another!

Theorem (Cantor, 1891)

There is no bijection between \mathbb{N} and the set of all subsets of \mathbb{N} .

- ▶ But, there is a surjection from set of all subsets of \mathbb{N} to \mathbb{N} .
- ▶ Thus, the “size” of $\mathcal{P}(\mathbb{N})$ is strictly greater than \mathbb{N} !

Can there be some set whose “size” is in between the two?

One infinity is “strictly” bigger than another!

Theorem (Cantor, 1891)

There is no bijection between \mathbb{N} and the set of all subsets of \mathbb{N} .

- ▶ But, there is a surjection from set of all subsets of \mathbb{N} to \mathbb{N} .
- ▶ Thus, the “size” of $\mathcal{P}(\mathbb{N})$ is strictly greater than \mathbb{N} !

Can there be some set whose “size” is in between the two?

Cantor’s Continuum hypothesis

There is no set whose “cardinality” is strictly between \mathbb{N} and $\mathcal{P}(\mathbb{N})$ (i.e., between naturals and reals).




Figure: 1st of Hilbert’s 23 problems for the 20th century in 1900.

What did the world think about these proofs (in 1890s?)



(a) Kronecker



(b) Poincaré



(c) Theologians

- ▶ **Kronecker:** Only constructive proofs are proofs! “Scientific Charlatan”, “Corruptor of youth”!
- ▶ **Poincaré:** Set theory is a “disease” from which mathematics will be cured.
- ▶ **Christian Theologians:** God=Uniqueness of an absolute infinity. So, what is all this different infinities...?!

What did the world think about these proofs (in 1890s?)



(a) Kronecker



(b) Poincaré



(c) Theologians

- ▶ **Kronecker:** Only constructive proofs are proofs! “Scientific Charlatan”, “Corruptor of youth”!
- ▶ **Poincaré:** Set theory is a “disease” from which mathematics will be cured.
- ▶ **Christian Theologians:** God=Uniqueness of an absolute infinity. So, what is all this different infinities...?!
- ▶ **Hilbert:** No one can expel us from the paradise that Cantor has created for us.

Summary and moving on...

- ▶ Finite and infinite sets.
- ▶ Using functions to compare sets: focus on bijections.
- ▶ Countable, countably infinite and uncountable sets.
- ▶ Cantor's diagonalization argument (A new powerful proof technique!).

Summary and moving on...

- ▶ Finite and infinite sets.
- ▶ Using functions to compare sets: focus on bijections.
- ▶ Countable, countably infinite and uncountable sets.
- ▶ Cantor's diagonalization argument (A new powerful proof technique!).

Next: Basic Mathematical Structures – Relations

Relations

Definition: Function

Let A, B be two sets. A **function** f from A to B is a subset R of $A \times B$ such that

- (i) $\forall a \in A, \exists b \in B$ such that $(a, b) \in R$, and
- (ii) if $(a, b) \in R$ and $(a, c) \in R$, then $b = c$.

- Now, suppose A is the set of all Btech students and B is the set of all courses. Clearly, we can assign to each student the set of courses he/she is taking. Is this a function?

Relations

Definition: Function

Let A, B be two sets. A **function** f from A to B is a subset R of $A \times B$ such that

- (i) $\forall a \in A, \exists b \in B$ such that $(a, b) \in R$, and
- (ii) if $(a, b) \in R$ and $(a, c) \in R$, then $b = c$.

- ▶ Now, suppose A is the set of all Btech students and B is the set of all courses. Clearly, we can assign to each student the set of courses he/she is taking. Is this a function?
- ▶ By removing the two extra assumptions in the defn, we get:

Definition: Relation

- ▶ A **relation** R from A to B is a subset of $A \times B$. If $(a, b) \in R$, we also write this as $a R b$.
- ▶ Thus, a relation is a way to relate the elements of two (not necessarily different) sets.

Examples and representations of relations

We write $R(A, B)$ for a relation from A to B and just $R(A)$ if $A = B$. Also if A is clear from context, we just write R .

Examples of relations

- ▶ All functions are relations.
- ▶ $R_1(\mathbb{Z}) = \{(a, b) \mid a, b \in \mathbb{Z}, a - b \text{ is even}\}.$
- ▶ $R_2(\mathbb{Z}) = \{(a, b) \mid a, b \in \mathbb{Z}, a \leq b\}.$
- ▶ Let S be a set, $R_3(\mathcal{P}(S)) = \{(A, B) \mid A, B \subseteq S, A \subseteq B\}.$
- ▶ Relational databases are practical examples.

Examples and representations of relations

We write $R(A, B)$ for a relation from A to B and just $R(A)$ if $A = B$. Also if A is clear from context, we just write R .

Examples of relations

- ▶ All functions are relations.
- ▶ $R_1(\mathbb{Z}) = \{(a, b) \mid a, b \in \mathbb{Z}, a - b \text{ is even} \}$.
- ▶ $R_2(\mathbb{Z}) = \{(a, b) \mid a, b \in \mathbb{Z}, a \leq b\}$.
- ▶ Let S be a set, $R_3(\mathcal{P}(S)) = \{(A, B) \mid A, B \subseteq S, A \subseteq B\}$.
- ▶ Relational databases are practical examples.

Representations of a relation from A to B .

- ▶ As a set of **ordered pairs of elements**, i.e., subset of $A \times B$.
- ▶ As a **directed graph**.
- ▶ As a **(database) table**.

Use of relations

Practical application in relational databases: IMDB, university records, etc.

Use of relations

Practical application in relational databases: IMDB, university records, etc.

But, why study relations in this course?

Use of relations

Practical application in relational databases: IMDB, university records, etc.

But, why study relations in this course?

- ▶ Functions were special kinds of relations that were useful to compare sets.

Use of relations

Practical application in relational databases: IMDB, university records, etc.

But, why study relations in this course?

- ▶ Functions were special kinds of relations that were useful to compare sets.
- ▶ Are there other **special** relations? What are they useful for?

Use of relations

Practical application in relational databases: IMDB, university records, etc.

But, why study relations in this course?

- ▶ Functions were special kinds of relations that were useful to compare sets.
- ▶ Are there other **special** relations? What are they useful for?
 - ▶ Equivalence relations
 - ▶ Partial orders

Partitions of a set – grouping “like” elements

Examples

- ▶ Natural numbers are partitioned into even and odd.
- ▶ This class is partitioned into sets of students from **same** hostel.

How do you define a partition?

Partitions of a set – grouping “like” elements

Examples

- ▶ Natural numbers are partitioned into even and odd.
- ▶ This class is partitioned into sets of students from **same** hostel.

How do you define a partition?

Definition

A partition of a set S is a

Partitions of a set – grouping “like” elements

Examples

- ▶ Natural numbers are partitioned into even and odd.
- ▶ This class is partitioned into sets of students from **same** hostel.

How do you define a partition?

Definition

A partition of a set S is a **set P of its subsets** such that

Partitions of a set – grouping “like” elements

Examples

- ▶ Natural numbers are partitioned into even and odd.
- ▶ This class is partitioned into sets of students from **same** hostel.

How do you define a partition?

Definition

A partition of a set S is a **set P of its subsets** such that

- ▶ if $S' \in P$, then $S' \neq \emptyset$.
- ▶ $\bigcup_{S' \in P} S' = S$: its union covers entire set S .
- ▶ If $S_1, S_2 \in P$, then $S_1 \cap S_2 = \emptyset$: sets are disjoint.

Partitions of a set – grouping “like” elements

Examples

- ▶ Natural numbers are partitioned into even and odd.
- ▶ This class is partitioned into sets of students from **same** hostel.

How do you define a partition?

Definition

A partition of a set S is a **set P of its subsets** such that

- ▶ if $S' \in P$, then $S' \neq \emptyset$.
- ▶ $\bigcup_{S' \in P} S' = S$: its union covers entire set S .
- ▶ If $S_1, S_2 \in P$, then $S_1 \cap S_2 = \emptyset$: sets are disjoint.

Can you think of two trivial partitions that any set must have?

Interpreting partitions as relations

Thus, a partition divides S into subsets that each contain (only) elements that share some common property. e.g.,

Interpreting partitions as relations

Thus, a partition divides S into subsets that each contain (only) elements that share some common property. e.g.,

- ▶ Evenness or oddness (formally, remainder modulo 2!).
- ▶ Same hostel...

Interpreting partitions as relations

Thus, a partition divides S into subsets that each contain (only) elements that share some common property. e.g.,

- ▶ Evenness or oddness (formally, remainder modulo 2!).
- ▶ Same hostel...
- ▶ But, this sounds like relation, right? Which one?

Interpreting partitions as relations

Thus, a partition divides S into subsets that each contain (only) elements that share some common property. e.g.,

- ▶ Evenness or oddness (formally, remainder modulo 2!).
- ▶ Same hostel...
- ▶ But, this sounds like relation, right? Which one?

Relation generated by a partition

- ▶ Clearly all elements in a set of the partition are related by the “sameness” or “likeness” property.
- ▶ So can we define this as a relation?

Interpreting partitions as relations

Thus, a partition divides S into subsets that each contain (only) elements that share some common property. e.g.,

- ▶ Evenness or oddness (formally, remainder modulo 2!).
- ▶ Same hostel...
- ▶ But, this sounds like relation, right? Which one?

Relation generated by a partition

- ▶ Clearly all elements in a set of the partition are related by the “sameness” or “likeness” property.
- ▶ So can we define this as a relation? aRb if a is “like” b .

Interpreting partitions as relations

Thus, a partition divides S into subsets that each contain (only) elements that share some common property. e.g.,

- ▶ Evenness or oddness (formally, remainder modulo 2!).
- ▶ Same hostel...
- ▶ But, this sounds like relation, right? Which one?

Relation generated by a partition

- ▶ Clearly all elements in a set of the partition are related by the “sameness” or “likeness” property.
- ▶ So can we define this as a relation? aRb if a is “like” b .
- ▶ Formally, we define $R(S)$ by aRb if a and b belong to the same set in the partition of S .

Interpreting partitions as relations

Thus, a partition divides S into subsets that each contain (only) elements that share some common property. e.g.,

- ▶ Evenness or oddness (formally, remainder modulo 2!).
- ▶ Same hostel...
- ▶ But, this sounds like relation, right? Which one?

Relation generated by a partition

- ▶ Clearly all elements in a set of the partition are related by the “sameness” or “likeness” property.
- ▶ So can we define this as a relation? aRb if a is “like” b .
- ▶ Formally, we define $R(S)$ by aRb if a and b belong to the same set in the partition of S .

What properties does this relation have?