

CS 105: Department Introductory Course on Discrete Structures

Instructor : S. Akshay

Aug 29, 2023

Lecture 10 – Basic Mathematical Structures
Equivalence relations and partitions

Recap: Proofs and Structures

Chapter 1: Proofs

1. Propositions, predicates
2. Types of proofs, axioms
3. Mathematical Induction, Well-ordering principle
4. Strong Induction

Recap: Proofs and Structures

Chapter 1: Proofs

1. Propositions, predicates
2. Types of proofs, axioms
3. Mathematical Induction, Well-ordering principle
4. Strong Induction

Chapter 2: Sets and Functions

1. Finite and infinite sets.
2. Using functions to compare sets: focus on bijections.
3. Countable, countably infinite and uncountable sets.
4. Cantor's diagonalization (New/powerful proof technique!).

Recap: Proofs and Structures

Chapter 1: Proofs

1. Propositions, predicates
2. Types of proofs, axioms
3. Mathematical Induction, Well-ordering principle
4. Strong Induction

Chapter 2: Sets and Functions

1. Finite and infinite sets.
2. Using functions to compare sets: focus on bijections.
3. Countable, countably infinite and uncountable sets.
4. Cantor's diagonalization (New/powerful proof technique!).

Chapter 3: Relations

Relations

Definition: Relation

- ▶ A **relation** R from A to B is a subset of $A \times B$. If $(a, b) \in R$, we also write this as $a R b$.

We write $R(A, B)$ for a relation from A to B and just $R(A)$ if $A = B$. Also if A is clear from context, we just write R .

Relations

Definition: Relation

- ▶ A **relation** R from A to B is a subset of $A \times B$. If $(a, b) \in R$, we also write this as $a R b$.

We write $R(A, B)$ for a relation from A to B and just $R(A)$ if $A = B$. Also if A is clear from context, we just write R .

Examples of relations

- ▶ All functions are relations.
- ▶ $R_1(\mathbb{Z}) = \{(a, b) \mid a, b \in \mathbb{Z}, a - b \text{ is even} \}$.
- ▶ $R_2(\mathbb{Z}) = \{(a, b) \mid a, b \in \mathbb{Z}, a \leq b\}$.
- ▶ Let S be a set, $R_3(\mathcal{P}(S)) = \{(A, B) \mid A, B \subseteq S, A \subseteq B\}$.

Relations

Definition: Relation

- ▶ A **relation** R from A to B is a subset of $A \times B$. If $(a, b) \in R$, we also write this as $a R b$.

We write $R(A, B)$ for a relation from A to B and just $R(A)$ if $A = B$. Also if A is clear from context, we just write R .

Examples of relations

- ▶ All functions are relations.
- ▶ $R_1(\mathbb{Z}) = \{(a, b) \mid a, b \in \mathbb{Z}, a - b \text{ is even} \}$.
- ▶ $R_2(\mathbb{Z}) = \{(a, b) \mid a, b \in \mathbb{Z}, a \leq b\}$.
- ▶ Let S be a set, $R_3(\mathcal{P}(S)) = \{(A, B) \mid A, B \subseteq S, A \subseteq B\}$.

Representations of a relation from A to B .

As a set of **ordered pairs of elements**, i.e., subset of $A \times B$; As a **directed graph**; As a **(database) table**.

Partitions of a set – grouping “like” elements

Definition

A partition of a set S is a

Partitions of a set – grouping “like” elements

Definition

A partition of a set S is a **set P of its subsets** such that

Partitions of a set – grouping “like” elements

Definition

A partition of a set S is a **set P of its subsets** such that

- ▶ if $S' \in P$, then $S' \neq \emptyset$.
- ▶ $\bigcup_{S' \in P} S' = S$: its union covers entire set S .
- ▶ If $S_1, S_2 \in P$, then $S_1 \cap S_2 = \emptyset$: sets are disjoint.

Examples

- ▶ Natural numbers are partitioned into even and odd.
- ▶ Class is partitioned into sets of students from **same** hostel.

Interpreting partitions as relations

Thus, a partition divides S into subsets that each contain (only) elements that share some common property. e.g.,

Interpreting partitions as relations

Thus, a partition divides S into subsets that each contain (only) elements that share some common property. e.g.,

- ▶ Evenness or oddness (formally, remainder modulo 2!).
- ▶ Same hostel...

Interpreting partitions as relations

Thus, a partition divides S into subsets that each contain (only) elements that share some common property. e.g.,

- ▶ Evenness or oddness (formally, remainder modulo 2!).
- ▶ Same hostel...
- ▶ But, this sounds like relation, right? Which one?

Interpreting partitions as relations

Thus, a partition divides S into subsets that each contain (only) elements that share some common property. e.g.,

- ▶ Evenness or oddness (formally, remainder modulo 2!).
- ▶ Same hostel...
- ▶ But, this sounds like relation, right? Which one?

Every Partition gives rise to a Relation

- ▶ All elements in a set of the partition are related by the “sameness” or “likeness” property.
- ▶ We can define this as a relation!

Interpreting partitions as relations

Thus, a partition divides S into subsets that each contain (only) elements that share some common property. e.g.,

- ▶ Evenness or oddness (formally, remainder modulo 2!).
- ▶ Same hostel...
- ▶ But, this sounds like relation, right? Which one?

Every Partition gives rise to a Relation

- ▶ All elements in a set of the partition are related by the “sameness” or “likeness” property.
- ▶ We can define this as a relation! aRb if a is “like” b .

Interpreting partitions as relations

Thus, a partition divides S into subsets that each contain (only) elements that share some common property. e.g.,

- ▶ Evenness or oddness (formally, remainder modulo 2!).
- ▶ Same hostel...
- ▶ But, this sounds like relation, right? Which one?

Every Partition gives rise to a Relation

- ▶ All elements in a set of the partition are related by the “sameness” or “likeness” property.
- ▶ We can define this as a relation! aRb if a is “like” b .
- ▶ Formally, we define $R(S)$ by aRb if a and b belong to the same set in the partition of S .

Interpreting partitions as relations

Thus, a partition divides S into subsets that each contain (only) elements that share some common property. e.g.,

- ▶ Evenness or oddness (formally, remainder modulo 2!).
- ▶ Same hostel...
- ▶ But, this sounds like relation, right? Which one?

Every Partition gives rise to a Relation

- ▶ All elements in a set of the partition are related by the “sameness” or “likeness” property.
- ▶ We can define this as a relation! aRb if a is “like” b .
- ▶ Formally, we define $R(S)$ by aRb if a and b belong to the same set in the partition of S .

What properties does this relation have?

Properties of a relation generated by a partition

1. All elements must be related to themselves

Properties of a relation generated by a partition

1. All elements must be related to themselves
 - ▶ A relation $R(S)$ is called **reflexive** if for all $a \in S$, aRa .

Properties of a relation generated by a partition

1. All elements must be related to themselves
 - ▶ A relation $R(S)$ is called **reflexive** if for all $a \in S$, aRa .
2. If a is “like” b , then b must be “like” a .

Properties of a relation generated by a partition

1. All elements must be related to themselves
 - ▶ A relation $R(S)$ is called **reflexive** if for all $a \in S$, aRa .
2. If a is “like” b , then b must be “like” a .
 - ▶ A relation $R(S)$ is called **symmetric** if for all $a, b \in S$, we have aRb implies bRa .

Properties of a relation generated by a partition

1. All elements must be related to themselves
 - ▶ A relation $R(S)$ is called **reflexive** if for all $a \in S$, aRa .
2. If a is “like” b , then b must be “like” a .
 - ▶ A relation $R(S)$ is called **symmetric** if for all $a, b \in S$, we have aRb implies bRa .
3. If a is “like” b and b is “like” c , then a must be “like” c .

Properties of a relation generated by a partition

1. All elements must be related to themselves
 - ▶ A relation $R(S)$ is called **reflexive** if for all $a \in S$, aRa .
2. If a is “like” b , then b must be “like” a .
 - ▶ A relation $R(S)$ is called **symmetric** if for all $a, b \in S$, we have aRb implies bRa .
3. If a is “like” b and b is “like” c , then a must be “like” c .
 - ▶ A relation $R(S)$ called **transitive** if for all $a, b, c \in S$, we have aRb and bRc implies aRc .

Properties of a relation generated by a partition

1. All elements must be related to themselves
 - ▶ A relation $R(S)$ is called **reflexive** if for all $a \in S$, aRa .
2. If a is “like” b , then b must be “like” a .
 - ▶ A relation $R(S)$ is called **symmetric** if for all $a, b \in S$, we have aRb implies bRa .
3. If a is “like” b and b is “like” c , then a must be “like” c .
 - ▶ A relation $R(S)$ called **transitive** if for all $a, b, c \in S$, we have aRb and bRc implies aRc .

Any other defining properties?

Properties of a relation generated by a partition

1. All elements must be related to themselves
 - ▶ A relation $R(S)$ is called **reflexive** if for all $a \in S$, aRa .
2. If a is “like” b , then b must be “like” a .
 - ▶ A relation $R(S)$ is called **symmetric** if for all $a, b \in S$, we have aRb implies bRa .
3. If a is “like” b and b is “like” c , then a must be “like” c .
 - ▶ A relation $R(S)$ called **transitive** if for all $a, b, c \in S$, we have aRb and bRc implies aRc .

Any other defining properties?

Definition

A relation which satisfies all these three properties is called an **equivalence relation**.

Properties of a relation generated by a partition

1. All elements must be related to themselves
 - ▶ A relation $R(S)$ is called **reflexive** if for all $a \in S$, aRa .
2. If a is “like” b , then b must be “like” a .
 - ▶ A relation $R(S)$ is called **symmetric** if for all $a, b \in S$, we have aRb implies bRa .
3. If a is “like” b and b is “like” c , then a must be “like” c .
 - ▶ A relation $R(S)$ called **transitive** if for all $a, b, c \in S$, we have aRb and bRc implies aRc .

Any other defining properties?

Definition

A relation which satisfies all these three properties is called an **equivalence relation**.

Thus, from any **partition**, we get an **equivalence relation**.

Properties of a relation generated by a partition

1. All elements must be related to themselves
 - ▶ A relation $R(S)$ is called **reflexive** if for all $a \in S$, aRa .
2. If a is “like” b , then b must be “like” a .
 - ▶ A relation $R(S)$ is called **symmetric** if for all $a, b \in S$, we have aRb implies bRa .
3. If a is “like” b and b is “like” c , then a must be “like” c .
 - ▶ A relation $R(S)$ called **transitive** if for all $a, b, c \in S$, we have aRb and bRc implies aRc .

Any other defining properties?

Definition

A relation which satisfies all these three properties is called an **equivalence relation**.

Thus, from any **partition**, we get an **equivalence relation**. Is the converse true?

Examples

- ▶ **Reflexive:** $\forall a \in S, aRa$.
- ▶ **Symmetric:** $\forall a, b \in S, aRb$ implies bRa .
- ▶ **Transitive:** $\forall a, b, c \in S, aRb, bRc$ implies aRc .
- ▶ **Equivalence:** Reflexive, Symmetric and Transitive.

Examples

- ▶ **Reflexive:** $\forall a \in S, aRa$.
- ▶ **Symmetric:** $\forall a, b \in S, aRb$ implies bRa .
- ▶ **Transitive:** $\forall a, b, c \in S, aRb, bRc$ implies aRc .
- ▶ **Equivalence:** Reflexive, Symmetric and Transitive.

Relation	Refl.	Sym.	Trans.	Equiv.
aR_4b if students a and b take same set of courses	✓	✓	✓	✓
aR_5b if student a takes course b				
$\{(a, b) \mid a, b \in \mathbb{Z}, (a - b) \bmod 2 = 0\}$				
$\{(a, b) \mid a, b \in \mathbb{Z}, a \leq b\}$				
$\{(a, b) \mid a, b \in \mathbb{Z}, a < b\}$				
$\{(a, b) \mid a, b \in \mathbb{Z}, a \mid b\}$				
$\{(a, b) \mid a, b \in \mathbb{R}, a - b < 1\}$				
$\{((a, b), (c, d)) \mid (a, b), (c, d) \in \mathbb{Z} \times (\mathbb{Z} \setminus \{0\}), (ad = bc)\}$				

Partitions of a set – grouping “like” elements

Definition

A partition of a set S is a set P of its subsets such that

- ▶ if $S' \in P$, then $S' \neq \emptyset$.
- ▶ $\bigcup_{S' \in P} S' = S$: its union covers entire set S .
- ▶ If $S_1, S_2 \in P$, then $S_1 \cap S_2 = \emptyset$: sets are disjoint.

Example: natural numbers partitioned into even and odd...

Theorem

Every partition of set S gives rise to a **canonical** equivalence relation R on S , namely,

- ▶ aRb if a and b belong to the same set in the partition of S .

Partitions of a set – grouping “like” elements

Definition

A partition of a set S is a set P of its subsets such that

- ▶ if $S' \in P$, then $S' \neq \emptyset$.
- ▶ $\bigcup_{S' \in P} S' = S$: its union covers entire set S .
- ▶ If $S_1, S_2 \in P$, then $S_1 \cap S_2 = \emptyset$: sets are disjoint.

Example: natural numbers partitioned into even and odd...

Theorem

Every partition of set S gives rise to a **canonical** equivalence relation R on S , namely,

- ▶ **aRb if a and b belong to the same set in the partition of S .**

Is the converse true? Can we generate a partition from every equivalence relation?

Equivalence classes

Definition

- ▶ Let R be an equivalence relation on set S , and let $a \in S$.
- ▶ Then the **equivalence class** of a , denoted $[a]$, is the set of all elements related to it, i.e., $[a] = \{b \in S \mid (a, b) \in R\}$.

Equivalence classes

Definition

- ▶ Let R be an equivalence relation on set S , and let $a \in S$.
- ▶ Then the **equivalence class** of a , denoted $[a]$, is the set of all elements related to it, i.e., $[a] = \{b \in S \mid (a, b) \in R\}$.

In $R = \{(a, b) \in \mathbb{Z} \times \mathbb{Z} \mid (a - b) \bmod 5 = 0\}$, what are $[0]$, $[1]$?

Equivalence classes

Definition

- ▶ Let R be an equivalence relation on set S , and let $a \in S$.
- ▶ Then the **equivalence class** of a , denoted $[a]$, is the set of all elements related to it, i.e., $[a] = \{b \in S \mid (a, b) \in R\}$.

In $R = \{(a, b) \in \mathbb{Z} \times \mathbb{Z} \mid (a - b) \bmod 5 = 0\}$, what are $[0]$, $[1]$?

Lemma

Let R be an equivalence relation on S . Let $a, b \in S$. Then, the following statements are equivalent:

1. aRb
2. $[a] = [b]$
3. $[a] \cap [b] \neq \emptyset$.

Equivalence classes

Definition

- ▶ Let R be an equivalence relation on set S , and let $a \in S$.
- ▶ Then the **equivalence class** of a , denoted $[a]$, is the set of all elements related to it, i.e., $[a] = \{b \in S \mid (a, b) \in R\}$.

In $R = \{(a, b) \in \mathbb{Z} \times \mathbb{Z} \mid (a - b) \bmod 5 = 0\}$, what are $[0]$, $[1]$?

Lemma

Let R be an equivalence relation on S . Let $a, b \in S$. Then, the following statements are equivalent:

1. aRb
2. $[a] = [b]$
3. $[a] \cap [b] \neq \emptyset$.

Proof Sketch: (1) to (2) symm and trans, (2) to (3) refl, (3) to (1) symm and trans. (H.W.: Redo the proof formally.)

From equivalence relations to partitions

Theorem

1. Let R be an equivalence relation on S . Then, the equivalence classes of R form a partition of S .

From equivalence relations to partitions

Theorem

1. Let R be an equivalence relation on S . Then, the equivalence classes of R form a partition of S .
2. Conversely, given a partition P of S , there is an equivalence relation R whose equivalence classes are exactly the sets of P .

From equivalence relations to partitions

Theorem

1. Let R be an equivalence relation on S . Then, the equivalence classes of R form a partition of S .
2. Conversely, given a partition P of S , there is an equivalence relation R whose equivalence classes are exactly the sets of P .

Proof sketch of (1): Union, non-emptiness follows from reflexivity. The rest (pairwise disjointness) follows from the previous lemma.

(H.W.): Write the formal proofs of (1) and (2).

More “applications” of equivalence relations

Defining new objects using equivalence relations

Consider

$$R = \{((a, b), (c, d)) \mid (a, b), (c, d) \in \mathbb{Z} \times (\mathbb{Z} \setminus \{0\}), (ad = bc)\}.$$

More “applications” of equivalence relations

Defining new objects using equivalence relations

Consider

$$R = \{((a, b), (c, d)) \mid (a, b), (c, d) \in \mathbb{Z} \times (\mathbb{Z} \setminus \{0\}), (ad = bc)\}.$$

- ▶ Then the equivalence classes of R define the rational numbers.
- ▶ e.g., $\left[\frac{1}{2}\right] = \left[\frac{2}{4}\right]$ are two names for the same rational number.
- ▶ Indeed, when we write $\frac{p}{q}$ we implicitly mean $\left[\frac{p}{q}\right]$.

More “applications” of equivalence relations

Defining new objects using equivalence relations

Consider

$$R = \{((a, b), (c, d)) \mid (a, b), (c, d) \in \mathbb{Z} \times (\mathbb{Z} \setminus \{0\}), (ad = bc)\}.$$

- ▶ Then the equivalence classes of R define the rational numbers.
- ▶ e.g., $\left[\frac{1}{2}\right] = \left[\frac{2}{4}\right]$ are two names for the same rational number.
- ▶ Indeed, when we write $\frac{p}{q}$ we implicitly mean $\left[\frac{p}{q}\right]$.
- ▶ With this definition, why are addition and multiplication “well-defined”?

More “applications” of equivalence relations

Defining new objects using equivalence relations

Consider

$$R = \{((a, b), (c, d)) \mid (a, b), (c, d) \in \mathbb{Z} \times (\mathbb{Z} \setminus \{0\}), (ad = bc)\}.$$

- ▶ Then the equivalence classes of R define the rational numbers.
- ▶ e.g., $\left[\frac{1}{2}\right] = \left[\frac{2}{4}\right]$ are two names for the same rational number.
- ▶ Indeed, when we write $\frac{p}{q}$ we implicitly mean $\left[\frac{p}{q}\right]$.
- ▶ With this definition, why are addition and multiplication “well-defined”?

Can we define **integers** and **real numbers** starting from naturals by using equivalence classes?