

# CS 105: Department Introductory Course on Discrete Structures

Instructor : S. Akshay

Aug 31, 2023

Lecture 11 – Basic Mathematical Structures  
Equivalence relations and partially ordered sets

# Recap: Proofs and Structures

## Chapter 1: Proofs

1. Propositions, predicates
2. Types of proofs, axioms
3. Mathematical Induction, Well-ordering principle
4. Strong Induction

# Recap: Proofs and Structures

## Chapter 1: Proofs

1. Propositions, predicates
2. Types of proofs, axioms
3. Mathematical Induction, Well-ordering principle
4. Strong Induction

## Chapter 2: Sets and Functions

1. Finite and infinite sets.
2. Using functions to compare sets: focus on bijections.
3. Countable, countably infinite and uncountable sets.
4. Cantor's diagonalization (New/powerful proof technique!).

# Recap: Proofs and Structures

## Chapter 1: Proofs

1. Propositions, predicates
2. Types of proofs, axioms
3. Mathematical Induction, Well-ordering principle
4. Strong Induction

## Chapter 2: Sets and Functions

1. Finite and infinite sets.
2. Using functions to compare sets: focus on bijections.
3. Countable, countably infinite and uncountable sets.
4. Cantor's diagonalization (New/powerful proof technique!).

## Chapter 3: Relations

1. Equivalence Relations
2. Partial Orders

# Examples

- ▶ **Reflexive:**  $\forall a \in S, aRa$ .
- ▶ **Symmetric:**  $\forall a, b \in S, aRb$  implies  $bRa$ .
- ▶ **Transitive:**  $\forall a, b, c \in S, aRb, bRc$  implies  $aRc$ .
- ▶ **Equivalence:** Reflexive, Symmetric and Transitive.

# Examples

- ▶ **Reflexive:**  $\forall a \in S, aRa$ .
- ▶ **Symmetric:**  $\forall a, b \in S, aRb$  implies  $bRa$ .
- ▶ **Transitive:**  $\forall a, b, c \in S, aRb, bRc$  implies  $aRc$ .
- ▶ **Equivalence:** Reflexive, Symmetric and Transitive.

Relation	Refl.	Sym.	Trans.	Equiv.
$aR_4b$ if students $a$ and $b$ take same set of courses	✓	✓	✓	✓
$aR_5b$ if student $a$ takes course $b$				
$\{(a, b) \mid a, b \in \mathbb{Z}, (a - b) \bmod 2 = 0\}$				
$\{(a, b) \mid a, b \in \mathbb{Z}, a \leq b\}$				
$\{(a, b) \mid a, b \in \mathbb{Z}, a < b\}$				
$\{(a, b) \mid a, b \in \mathbb{Z}, a \mid b\}$				
$\{(a, b) \mid a, b \in \mathbb{R},  a - b  < 1\}$				
$\{((a, b), (c, d)) \mid (a, b), (c, d) \in \mathbb{Z} \times (\mathbb{Z} \setminus \{0\}), (ad = bc)\}$				

# Equivalence classes

## Definition

- ▶ Let  $R$  be an equivalence relation on set  $S$ , and let  $a \in S$ .
- ▶ Then the **equivalence class** of  $a$ , denoted  $[a]$ , is the set of all elements related to it, i.e.,  $[a] = \{b \in S \mid (a, b) \in R\}$ .

# Equivalence classes

## Definition

- ▶ Let  $R$  be an equivalence relation on set  $S$ , and let  $a \in S$ .
- ▶ Then the **equivalence class** of  $a$ , denoted  $[a]$ , is the set of all elements related to it, i.e.,  $[a] = \{b \in S \mid (a, b) \in R\}$ .

In  $R = \{(a, b) \in \mathbb{Z} \times \mathbb{Z} \mid (a - b) \bmod 5 = 0\}$ , what are  $[0]$ ,  $[1]$ ?



# Equivalence classes

## Definition

- ▶ Let  $R$  be an equivalence relation on set  $S$ , and let  $a \in S$ .
- ▶ Then the **equivalence class** of  $a$ , denoted  $[a]$ , is the set of all elements related to it, i.e.,  $[a] = \{b \in S \mid (a, b) \in R\}$ .

In  $R = \{(a, b) \in \mathbb{Z} \times \mathbb{Z} \mid (a - b) \bmod 5 = 0\}$ , what are  $[0]$ ,  $[1]$ ?

## Lemma

Let  $R$  be an equivalence relation on  $S$ . Let  $a, b \in S$ . Then, the following statements are equivalent:

1.  $aRb$
2.  $[a] = [b]$
3.  $[a] \cap [b] \neq \emptyset$ .

# Equivalence classes

## Definition

- ▶ Let  $R$  be an equivalence relation on set  $S$ , and let  $a \in S$ .
- ▶ Then the **equivalence class** of  $a$ , denoted  $[a]$ , is the set of all elements related to it, i.e.,  $[a] = \{b \in S \mid (a, b) \in R\}$ .

In  $R = \{(a, b) \in \mathbb{Z} \times \mathbb{Z} \mid (a - b) \bmod 5 = 0\}$ , what are  $[0]$ ,  $[1]$ ?

## Lemma

Let  $R$  be an equivalence relation on  $S$ . Let  $a, b \in S$ . Then, the following statements are equivalent:

1.  $aRb$
2.  $[a] = [b]$
3.  $[a] \cap [b] \neq \emptyset$ .

Proof Sketch: (1) to (2) symm and trans, (2) to (3) refl, (3) to (1) symm and trans. (H.W.: Redo the proof formally.)

# From equivalence relations to partitions

## Theorem

1. Let  $R$  be an equivalence relation on  $S$ . Then, the equivalence classes of  $R$  form a partition of  $S$ .

# From equivalence relations to partitions

## Theorem

1. Let  $R$  be an equivalence relation on  $S$ . Then, the equivalence classes of  $R$  form a partition of  $S$ .
2. Conversely, given a partition  $P$  of  $S$ , there is an equivalence relation  $R$  whose equivalence classes are exactly the sets of  $P$ .

# From equivalence relations to partitions

## Theorem

1. Let  $R$  be an equivalence relation on  $S$ . Then, the equivalence classes of  $R$  form a partition of  $S$ .
2. Conversely, given a partition  $P$  of  $S$ , there is an equivalence relation  $R$  whose equivalence classes are exactly the sets of  $P$ .

Proof sketch of (1): Union, non-emptiness follows from reflexivity. The rest (pairwise disjointness) follows from the previous lemma.

(H.W.): Write the formal proofs of (1) and (2).

## More “applications” of equivalence relations

### Defining new objects using equivalence relations

Consider

$$R = \{((a, b), (c, d)) \mid (a, b), (c, d) \in \mathbb{Z} \times (\mathbb{Z} \setminus \{0\}), (ad = bc)\}.$$

# More “applications” of equivalence relations

## Defining new objects using equivalence relations

Consider

$$R = \{((a, b), (c, d)) \mid (a, b), (c, d) \in \mathbb{Z} \times (\mathbb{Z} \setminus \{0\}), (ad = bc)\}.$$

- ▶ Then the equivalence classes of  $R$  define the rational numbers.
- ▶ e.g.,  $\left[\frac{1}{2}\right] = \left[\frac{2}{4}\right]$  are two names for the same rational number.
- ▶ Indeed, when we write  $\frac{p}{q}$  we implicitly mean  $\left[\frac{p}{q}\right]$ .

# More “applications” of equivalence relations

## Defining new objects using equivalence relations

Consider

$$R = \{((a, b), (c, d)) \mid (a, b), (c, d) \in \mathbb{Z} \times (\mathbb{Z} \setminus \{0\}), (ad = bc)\}.$$

- ▶ Then the equivalence classes of  $R$  define the rational numbers.
- ▶ e.g.,  $\left[\frac{1}{2}\right] = \left[\frac{2}{4}\right]$  are two names for the same rational number.
- ▶ Indeed, when we write  $\frac{p}{q}$  we implicitly mean  $\left[\frac{p}{q}\right]$ .
- ▶ With this definition, why are addition and multiplication “well-defined”?



# More “applications” of equivalence relations

## Defining new objects using equivalence relations

Consider

$$R = \{((a, b), (c, d)) \mid (a, b), (c, d) \in \mathbb{Z} \times (\mathbb{Z} \setminus \{0\}), (ad = bc)\}.$$

- ▶ Then the equivalence classes of  $R$  define the rational numbers.
- ▶ e.g.,  $\left[\frac{1}{2}\right] = \left[\frac{2}{4}\right]$  are two names for the same rational number.
- ▶ Indeed, when we write  $\frac{p}{q}$  we implicitly mean  $\left[\frac{p}{q}\right]$ .
- ▶ With this definition, why are addition and multiplication “well-defined”?

Can we define **integers** and **real numbers** starting from naturals by using equivalence classes?

## Moving on another special relation: Partial Orders

Consider  $\{(a, b) \mid a, b \in \mathbb{Z}, a \leq b\}$ .

## Moving on another special relation: Partial Orders

Consider  $\{(a, b) \mid a, b \in \mathbb{Z}, a \leq b\}$ . This is reflexive, transitive but not symmetric. In fact, it is quite different!

## Partial Orders

Consider  $\{(a, b) \mid a, b \in \mathbb{Z}, a \leq b\}$ . This is reflexive, transitive but not symmetric. In fact, it is quite different!

### Anti-symmetric

A relation  $R$  on  $S$  is **anti-symmetric** if for all  $a, b \in S$   $aRb$  and  $bRa$  implies  $a = b$ .

# Partial Orders

Consider  $\{(a, b) \mid a, b \in \mathbb{Z}, a \leq b\}$ . This is reflexive, transitive but not symmetric. In fact, it is quite different!

## Anti-symmetric

A relation  $R$  on  $S$  is **anti-symmetric** if for all  $a, b \in S$   $aRb$  and  $bRa$  implies  $a = b$ .

Examples:

- ▶  $R_1(\mathbb{Z}) = \{(a, b) \mid a, b \in \mathbb{Z}, a \leq b\}$ .
- ▶  $R_2(\mathcal{P}(S)) = \{(A, B) \mid A, B \in \mathcal{P}(S), A \subseteq B\}$ .

# Partial Orders

Consider  $\{(a, b) \mid a, b \in \mathbb{Z}, a \leq b\}$ . This is reflexive, transitive but not symmetric. In fact, it is quite different!

## Anti-symmetric

A relation  $R$  on  $S$  is **anti-symmetric** if for all  $a, b \in S$   $aRb$  and  $bRa$  implies  $a = b$ .

Examples:

- ▶  $R_1(\mathbb{Z}) = \{(a, b) \mid a, b \in \mathbb{Z}, a \leq b\}$ .
- ▶  $R_2(\mathcal{P}(S)) = \{(A, B) \mid A, B \in \mathcal{P}(S), A \subseteq B\}$ .

## Definition

A **partial order** is a relation which is **reflexive**, **transitive** and **anti-symmetric**.

## Partial orders and equivalences relations

- ▶ **Reflexive:**  $\forall a \in S, aRa$ .
- ▶ **Symmetric:**  $\forall a, b \in S, aRb$  implies  $bRa$ .
- ▶ **Anti-symmetric:**  $\forall a, b \in S, aRb, bRa$  implies  $a = b$ .
- ▶ **Transitive:**  $\forall a, b, c \in S, aRb, bRc$  implies  $aRc$ .

## Partial orders and equivalences relations

- ▶ **Reflexive:**  $\forall a \in S, aRa$ .
- ▶ **Symmetric:**  $\forall a, b \in S, aRb$  implies  $bRa$ .
- ▶ **Anti-symmetric:**  $\forall a, b \in S, aRb, bRa$  implies  $a = b$ .
- ▶ **Transitive:**  $\forall a, b, c \in S, aRb, bRc$  implies  $aRc$ .

	Reflexive	Transitive	Symmetric	Anti-symmetric
Equivalence relation	✓	✓	✓	
Partial order	✓	✓		✓



# Partial orders and equivalences relations

- ▶ **Reflexive:**  $\forall a \in S, aRa$ .
- ▶ **Symmetric:**  $\forall a, b \in S, aRb$  implies  $bRa$ .
- ▶ **Anti-symmetric:**  $\forall a, b \in S, aRb, bRa$  implies  $a = b$ .
- ▶ **Transitive:**  $\forall a, b, c \in S, aRb, bRc$  implies  $aRc$ .

	Refl.	Anti-Sym	Trans.	PO
$\{(a, b) \mid a, b \in \mathbb{Z}, a \leq b\}$	✓	✓	✓	✓
$\{(A, B) \mid A, B \in \mathcal{P}(S), A \subseteq B\}$	✓	✓	✓	✓
$\{(a, b) \mid a, b \in \mathbb{Z}, a < b\}$				
$\{(a, b) \mid a, b \in \mathbb{Z}^+, a \mid b\}$				
$\{((a, b), (c, d)) \mid (a, b), (c, d) \in \mathbb{Z}^+ \times \mathbb{Z}^+, a < c \text{ or } (a = c, b \leq d)\}$				

## Partial orders

	Refl.	Anti-Sym	Trans.	PO
$\{(a, b) \mid a, b \in \mathbb{Z}, a \leq b\}$	✓	✓	✓	✓
$\{(A, B) \mid A, B \in \mathcal{P}(S), A \subseteq B\}$	✓	✓	✓	✓
$\{(a, b) \mid a, b \in \mathbb{Z}, a < b\}$				
$\{(a, b) \mid a, b \in \mathbb{Z}^+, a \mid b\}$				
$\{((a, b), (c, d)) \mid (a, b), (c, d) \in \mathbb{Z}^+ \times \mathbb{Z}^+, a < c \text{ or } (a = c, b \leq d)\}$				

- We use  $\preceq$  to denote partial orders and write  $a \preceq b$  instead of  $aRb$ .

## Partial orders

	Refl.	Anti-Sym	Trans.	PO
$\{(a, b) \mid a, b \in \mathbb{Z}, a \leq b\}$	✓	✓	✓	✓
$\{(A, B) \mid A, B \in \mathcal{P}(S), A \subseteq B\}$	✓	✓	✓	✓
$\{(a, b) \mid a, b \in \mathbb{Z}, a < b\}$				
$\{(a, b) \mid a, b \in \mathbb{Z}^+, a \mid b\}$				
$\{((a, b), (c, d)) \mid (a, b), (c, d) \in \mathbb{Z}^+ \times \mathbb{Z}^+, a < c \text{ or } (a = c, b \leq d)\}$				

- ▶ We use  $\preceq$  to denote partial orders and write  $a \preceq b$  instead of  $aRb$ .
- ▶ Why is it called “partial” order?

# Partial orders

	Refl.	Anti-Sym	Trans.	PO
$\{(a, b) \mid a, b \in \mathbb{Z}, a \leq b\}$	✓	✓	✓	✓
$\{(A, B) \mid A, B \in \mathcal{P}(S), A \subseteq B\}$	✓	✓	✓	✓
$\{(a, b) \mid a, b \in \mathbb{Z}, a < b\}$				
$\{(a, b) \mid a, b \in \mathbb{Z}^+, a \mid b\}$				
$\{((a, b), (c, d)) \mid (a, b), (c, d) \in \mathbb{Z}^+ \times \mathbb{Z}^+, a < c \text{ or } (a = c, b \leq d)\}$				

- ▶ We use  $\preceq$  to denote partial orders and write  $a \preceq b$  instead of  $aRb$ .
- ▶ Why is it called “partial” order? Because, not all pairs of elements are “comparable” (i.e., related by  $\preceq$ ). It is important

# Partial orders

	Refl.	Anti-Sym	Trans.	PO
$\{(a, b) \mid a, b \in \mathbb{Z}, a \leq b\}$	✓	✓	✓	✓
$\{(A, B) \mid A, B \in \mathcal{P}(S), A \subseteq B\}$	✓	✓	✓	✓
$\{(a, b) \mid a, b \in \mathbb{Z}, a < b\}$				
$\{(a, b) \mid a, b \in \mathbb{Z}^+, a \mid b\}$				
$\{((a, b), (c, d)) \mid (a, b), (c, d) \in \mathbb{Z}^+ \times \mathbb{Z}^+, a < c \text{ or } (a = c, b \leq d)\}$				

- ▶ We use  $\preceq$  to denote partial orders and write  $a \preceq b$  instead of  $aRb$ .
- ▶ Why is it called “partial” order? Because, not all pairs of elements are “comparable” (i.e., related by  $\preceq$ ).
- ▶ A total order is a partial order  $\preceq$  on  $S$  in which every pair of elements is comparable **an important lemma**
  - ▶ i.e.,  $\forall a, b \in S$ , either  $a \preceq b$  or  $b \preceq a$ .

# Partial orders

	Refl.	Anti-Sym	Trans.	PO	TO
$\{(a, b) \mid a, b \in \mathbb{Z}, a \leq b\}$	✓	✓	✓	✓	✓
$\{(A, B) \mid A, B \in \mathcal{P}(S), A \subseteq B\}$	✓	✓	✓	✓	×
$\{(a, b) \mid a, b \in \mathbb{Z}, a < b\}$					
$\{(a, b) \mid a, b \in \mathbb{Z}^+, a \mid b\}$					
$\{((a, b), (c, d)) \mid (a, b), (c, d) \in \mathbb{Z}^+ \times \mathbb{Z}^+, a < c \text{ or } (a = c, b \leq d)\}$					

- ▶ We use  $\preceq$  to denote partial orders and write  $a \preceq b$  instead of  $aRb$ .
- ▶ Why is it called “partial” order? Because, not all pairs of elements are “comparable” (i.e., related by  $\preceq$ ).
- ▶ A total order is a partial order  $\preceq$  on  $S$  in which every pair of elements is comparable
  - ▶ i.e.,  $\forall a, b \in S$ , either  $a \preceq b$  or  $b \preceq a$ .

## Partial orders

	Refl.	Anti-Sym	Trans.	PO	TO
$\{(a, b) \mid a, b \in \mathbb{Z}, a \leq b\}$	✓	✓	✓	✓	✓
$\{(A, B) \mid A, B \in \mathcal{P}(S), A \subseteq B\}$	✓	✓	✓	✓	×
$\{(a, b) \mid a, b \in \mathbb{Z}, a < b\}$					
$\{(a, b) \mid a, b \in \mathbb{Z}^+, a \mid b\}$					
$\{((a, b), (c, d)) \mid (a, b), (c, d) \in \mathbb{Z}^+ \times \mathbb{Z}^+, a < c \text{ or } (a = c, b \leq d)\}$					

- ▶ We use  $\preceq$  to denote partial orders and write  $a \preceq b$  instead of  $aRb$ .
- ▶ Why is it called “partial” order? Because, not all pairs of elements are “comparable” (i.e., related by  $\preceq$ ).
- ▶ A total order is a partial order  $\preceq$  on  $S$  in which every pair of elements is comparable
- ▶ Qn: Can a relation be symmetric and anti-symmetric?

# Partial orders

	Refl.	Anti-Sym	Trans.	PO	TO
$\{(a, b) \mid a, b \in \mathbb{Z}, a \leq b\}$	✓	✓	✓	✓	✓
$\{(A, B) \mid A, B \in \mathcal{P}(S), A \subseteq B\}$	✓	✓	✓	✓	×
$\{(a, b) \mid a, b \in \mathbb{Z}, a < b\}$					
$\{(a, b) \mid a, b \in \mathbb{Z}^+, a \mid b\}$					
$\{((a, b), (c, d)) \mid (a, b), (c, d) \in \mathbb{Z}^+ \times \mathbb{Z}^+, a < c \text{ or } (a = c, b \leq d)\}$					

- ▶ We use  $\preceq$  to denote partial orders and write  $a \preceq b$  instead of  $aRb$ .
- ▶ Why is it called “partial” order? Because, not all pairs of elements are “comparable” (i.e., related by  $\preceq$ ).
- ▶ A total order is a partial order  $\preceq$  on  $S$  in which every pair of elements is comparable
- ▶ Qn: Can a relation be symmetric and anti-symmetric?
- ▶ Qn: Can a relation be neither symmetric nor anti-symmetric?



# Partially ordered sets (Posets)

## Definition

A set  $S$  together with a partial order  $\preceq$  on  $S$ , is called a **partially-ordered set** or **poset**, denoted  $(S, \preceq)$ .

# Partially ordered sets (Posets)

## Definition

A set  $S$  together with a partial order  $\preceq$  on  $S$ , is called a **partially-ordered set** or **poset**, denoted  $(S, \preceq)$ .



## Examples

- ▶  $(\mathbb{Z}, \leq)$ : integers with the usual less than or equal to relation.
- ▶  $(\mathcal{P}(S), \subseteq)$ : powerset of any set with the subset relation.
- ▶  $(\mathbb{Z}^+, |)$ : positive integers with divisibility relation.

# Graphical representation of relations: posets

Recall: any relation on a set can be represented as a **graph** with

- ▶ nodes as elements of the set and
- ▶ directed edges between them indicating the ordered pairs that are related.




- ▶ Did these come from posets?
- ▶ Do graphs defined by posets have any “special” properties?

## Graphical representation of relations: posets

- ▶ Let  $S = \{1, 2, 3\}$ . Recall the poset  $(\mathcal{P}(S), \subseteq)$ .
- ▶ How does the graph of  $(\mathcal{P}(S), \subseteq)$  look like?

## Graphical representation of relations: posets

- ▶ Let  $S = \{1, 2, 3\}$ . Recall the poset  $(\mathcal{P}(S), \subseteq)$ .
- ▶ How does the graph of  $(\mathcal{P}(S), \subseteq)$  look like?



## Graphical representation of relations: posets

- ▶ Let  $S = \{1, 2, 3\}$ . Recall the poset  $(\mathcal{P}(S), \subseteq)$ .
- ▶ How does the graph of  $(\mathcal{P}(S), \subseteq)$  look like?




Figure: Graph of a poset and its **Hasse diagram**

# Graphical representation of relations: posets

- Let  $S = \{1, 2, 3\}$ . Recall the poset  $(\mathcal{P}(S), \subseteq)$ .




Figure: Graph of a poset and its **Hasse diagram**

- What is “special” about these graphs?

# Graphical representation of relations: posets

- Let  $S = \{1, 2, 3\}$ . Recall the poset  $(\mathcal{P}(S), \subseteq)$ .




Figure: Graph of a poset and its **Hasse diagram**

- What is “special” about these graphs?
- **Graphs of posets are “acyclic” (except for self-loops).** yes it is indeed
- Starting from a node and following the directed edges (except self-loops), one can't come back to the same node.



# Graphical representation of relations: posets

- Let  $S = \{1, 2, 3\}$ . Recall the poset  $(\mathcal{P}(S), \subseteq)$ .




Figure: Graph of a poset and its **Hasse diagram**

- What is “special” about these graphs?
- **Graphs of posets are “acyclic” (except for self-loops).**
- Starting from a node and following the directed edges (except self-loops), one can’t come back to the same node.
- Given the Hasse diagram of a poset, its **reflexive transitive closure** gives back the graph of the poset.