

CS 105: DIC on Discrete Structures

Graph theory

Characterizing maximum matchings
via augmenting paths

Lecture 31

Oct 31 2023

Topic 3: Graph theory

Basic concepts

- ▶ Basics: graphs, paths, cycles, walks, trails, ...
- ▶ Cliques and independent sets.
- ▶ Graph representations, isomorphisms and automorphisms.
- ▶ Matchings: perfect, maximal and maximum.

Topic 3: Graph theory

Basic concepts

- ▶ Basics: graphs, paths, cycles, walks, trails, ...
- ▶ Cliques and independent sets.
- ▶ Graph representations, isomorphisms and automorphisms.
- ▶ Matchings: perfect, maximal and maximum.

Characterizations

1. **Eulerian graphs:** Using degrees of vertices.
2. **Bipartite graphs:** Using odd length cycles.
3. **Connected components:** Using cycles.

Topic 3: Graph theory

Basic concepts

- ▶ Basics: graphs, paths, cycles, walks, trails, ...
- ▶ Cliques and independent sets.
- ▶ Graph representations, isomorphisms and automorphisms.
- ▶ Matchings: perfect, maximal and maximum.


Characterizations

1. **Eulerian graphs:** Using degrees of vertices.
2. **Bipartite graphs:** Using odd length cycles.
3. **Connected components:** Using cycles.
4. **Maximum matchings:** Using augmenting paths.

Definitions

- ▶ A **matching** in a graph G is a set of (non-loop) edges with no shared end-points. The vertices incident to edges in a matching are called **matched** or **saturated**. Others are **unsaturated**.
- ▶ A **perfect matching** in a graph is a matching that saturates every vertex.
- ▶ A **maximal matching** in a graph is a matching that cannot be enlarged by adding an edge.
- ▶ A **maximum matching** is a matching of maximum size (# edges) among all matchings in a graph.


Matchings: Pop Quiz



Give an example of the following, if possible:

1. A maximal matching in G which is not a maximum matching.
2. A maximum matching in G . How do you know it is maximum?
No augmenting path.
3. Can there be more than one maximum matching in a graph?
4. A graph which has no perfect matching but has a maximum matching. Is G such a graph? **Odd no of vertices.**

Matchings: Pop Quiz



- ▶ Perfect matching \implies maximum matching \implies maximal matching
- ▶ The reverse directions in the above implications do not hold.

Alternating and Augmenting paths


Definition

- ▶ Given a matching M , an M -alternating path is a path that alternates between edges in M and edges not in M .
- ▶ An M -alternating path whose endpoints are unmatched by M is an M -augmenting path.

Alternating and Augmenting paths

Definition

- ▶ Given a matching M , an M -alternating path is a path that alternates between edges in M and edges not in M .
- ▶ An M -alternating path whose endpoints are unmatched by M is an M -augmenting path.




- ▶ **Ex 1:** Give an example of a matching M in G and
 1. a M -alternating path which is an M -augmenting path and
 2. a M -alternating path which is not an M -augmenting path

Alternating and Augmenting paths

Definition

- ▶ Given a matching M , an M -alternating path is a path that alternates between edges in M and edges not in M .
- ▶ An M -alternating path whose endpoints are unmatched by M is an M -augmenting path.



Characterization of Maximum matchings

- ▶ Clearly, a maximum matching cannot have an M -augmenting path.

Characterization of Maximum matchings

- ▶ Clearly, a maximum matching cannot have an M -augmenting path.
- ▶ In fact, this is the characterization!

Characterization of Maximum matchings

- ▶ Clearly, a maximum matching cannot have an M -augmenting path.
- ▶ In fact, this is the characterization!

Theorem


A matching M in G is a maximum matching iff G has no M -augmenting path.

Characterizing maximum matchings

We need a definition and a lemma.

Definition

If M, M' are matchings in a graph H , the **symmetric difference** $M \triangle M'$ is the set of edges which are either in M or in M' but not both, i.e., $M \triangle M' = (M \setminus M') \cup (M' \setminus M)$. Not in both.




Characterizing maximum matchings

We need a definition and a lemma.

Definition

If M, M' are matchings in a graph H , the **symmetric difference** $M \triangle M'$ is the set of edges which are either in M or in M' but not both, i.e., $M \triangle M' = (M \setminus M') \cup (M' \setminus M)$.




Characterizing maximum matchings

We need a definition and a lemma.

Definition

If M, M' are matchings in a graph H , the **symmetric difference** $M \triangle M'$ is the set of edges which are either in M or in M' but not both, i.e., $M \triangle M' = (M \setminus M') \cup (M' \setminus M)$.




Ex 2: What is the symmetric difference of M (red) and M' (green) in the above graph?

Characterizing maximum matchings

We need a definition and a lemma.


Definition

If M, M' are matchings in a graph H , the **symmetric difference** $M \triangle M'$ is the set of edges which are either in M or in M' but not both, i.e., $M \triangle M' = (M \setminus M') \cup (M' \setminus M)$.



Ex 2: What is the symmetric difference of M (red) and M' (green) in the above graph? Can you generalize this?

Characterizing maximum matchings




Lemma

Every component of the symmetric difference of two matchings is either a path or an even cycle.

Important lemma.

Characterizing maximum matchings




Lemma

Every component of the symmetric difference of two matchings is either a path or an even cycle.

- Let $F = M \Delta M'$. F has at most **2 edges** at each vertex, hence every component is a path or a cycle.

Characterizing maximum matchings




Lemma

Every component of the symmetric difference of two matchings is either a path or an even cycle.

- ▶ Let $F = M \Delta M'$. F has at most 2 edges at each vertex, hence every component is a path or a cycle.
- ▶ Further every path/cycle alternates between edges of $M \setminus M'$ and $M' \setminus M$.

Characterizing maximum matchings



Lemma

Every component of the symmetric difference of two matchings is either a path or an even cycle.

- ▶ Let $F = M \Delta M'$. F has at most 2 edges at each vertex, hence every component is a path or a cycle.
- ▶ Further every path/cycle alternates between edges of $M \setminus M'$ and $M' \setminus M$.
- ▶ Thus, each cycle has even length with equal edges from M and M' . □

Characterizing maximum matchings

Theorem (Berge'57)

A matching M in G is a maximum matching iff G has no M -augmenting path.

Proof:

Characterizing maximum matchings

Theorem (Berge'57)

A matching M in G is a maximum matching iff G has no M -augmenting path.

Proof:

- ▶ One direction is trivial (which one?!).

Characterizing maximum matchings

Theorem (Berge'57)

A matching M in G is a maximum matching iff G has no M -augmenting path.

Proof:

- ▶ One direction is trivial (which one?!).
- ▶ (\Leftarrow) For the other, we will show the contrapositive.

Characterizing maximum matchings

Theorem (Berge'57)

A matching M in G is a maximum matching iff G has no M -augmenting path.

Proof:

- ▶ One direction is trivial (which one?!).
- ▶ (\Leftarrow) For the other, we will show the contrapositive.
- ▶ i.e., if \exists matching M' larger than M , we will construct an M -augmenting path.

Characterizing maximum matchings

Theorem (Berge'57)

A matching M in G is a maximum matching iff G has no M -augmenting path.

Proof:

- ▶ One direction is trivial (which one?!).
- ▶ (\Leftarrow) For the other, we will show the contrapositive.
- ▶ i.e., if \exists matching M' larger than M , we will construct an M -augmenting path.
- ▶ Let $F = M \triangle M'$. By Lemma, F has only paths and even cycles with equal no. of edges from M and M' .

Characterizing maximum matchings

Theorem (Berge'57)

A matching M in G is a maximum matching iff G has no M -augmenting path.

Proof:

- ▶ One direction is trivial (which one?!).
- ▶ (\Leftarrow) For the other, we will show the contrapositive.
- ▶ i.e., if \exists matching M' larger than M , we will construct an M -augmenting path.
- ▶ Let $F = M \Delta M'$. By Lemma, F has only paths and even cycles with equal no. of edges from M and M' .
- ▶ But then since $|M'| > |M|$ it must have a component with more edges in M' than M .

Characterizing maximum matchings

Theorem (Berge'57)

A matching M in G is a maximum matching iff G has no M -augmenting path.

Proof:

- ▶ One direction is trivial (which one?!).
- ▶ (\Leftarrow) For the other, we will show the contrapositive.
- ▶ i.e., if \exists matching M' larger than M , we will construct an M -augmenting path.
- ▶ Let $F = M \triangle M'$. By Lemma, F has only paths and even cycles with equal no. of edges from M and M' .
- ▶ But then since $|M'| > |M|$ it must have a component with more edges in M' than M .
- ▶ This component can only be a path that starts and ends with an edge of M' ; i.e., it is an M -augmenting path in G . □

Perfect matchings in bipartite graphs

- ▶ If there are n women and n men, and each woman is compatible with exactly k men and each man compatible with exactly k women, can they be perfectly matched?

Perfect matchings in bipartite graphs


- ▶ If there are n women and n men, and each woman is compatible with exactly k men and each man compatible with exactly k women, can they be perfectly matched?
- ▶ If there are m jobs and n applicants, when can we find a perfect matching where all m jobs are saturated?

Perfect matchings in bipartite graphs

- ▶ Consider a bipartite graph with X, Y as partitions.
- ▶ If a matching M saturates X , then for every $S \subseteq X$,


Perfect matchings in bipartite graphs

- ▶ Consider a bipartite graph with X, Y as partitions.
- ▶ If a matching M saturates X , then for every $S \subseteq X$, what can we say?




Perfect matchings in bipartite graphs

- ▶ Consider a bipartite graph with X, Y as partitions.
- ▶ If a matching M saturates X , then for every $S \subseteq X$, there must exist at least $|S|$ vertices that have neighbours in S .




Perfect matchings in bipartite graphs

- ▶ Consider a bipartite graph with X, Y as partitions.
- ▶ If a matching M saturates X , then for every $S \subseteq X$, there must exist at least $|S|$ vertices that have neighbours in S .
- ▶ That is, $\forall S \subseteq X, |N(S)| \geq |S|$ (**Hall's Condition**).



Perfect matchings in bipartite graphs

- ▶ Consider a bipartite graph with X, Y as partitions.
- ▶ If a matching M saturates X , then for every $S \subseteq X$, there must exist at least $|S|$ vertices that have neighbours in S .
- ▶ That is, $\forall S \subseteq X, |N(S)| \geq |S|$ (**Hall's Condition**). This is a necessary condition, is it sufficient?



Perfect matchings in bipartite graphs

- ▶ Consider a bipartite graph with X, Y as partitions.
- ▶ If a matching M saturates X , then for every $S \subseteq X$, there must exist at least $|S|$ vertices that have neighbours in S .
- ▶ That is, $\forall S \subseteq X, |N(S)| \geq |S|$ (**Hall's Condition**). This is a necessary condition, is it sufficient?

Theorem (Hall'35)

A bipartite graph G with bipartitions X, Y has a matching that saturates X iff for all $S \subseteq X$, $|N(S)| \geq |S|$.

Characterizing perfect matchings in bipartite graphs

Theorem (Hall'35)

A bipartite graph G with bipartitions X, Y has a matching that saturates X iff for all $S \subseteq X$, $|N(S)| \geq |S|$.

Characterizing perfect matchings in bipartite graphs

Theorem (Hall'35)

A bipartite graph G with bipartitions X, Y has a matching that saturates X iff for all $S \subseteq X$, $|N(S)| \geq |S|$.

Proof: (\implies) is straightforward:

- ▶ Let M be a matching.
- ▶ Then for any $S \subseteq X$, each vertex of S is matched to a distinct vertex in $N(S)$
- ▶ So $|N(S)| \geq |S|$.