

# CS 105: Department Introductory Course on Discrete Structures

Instructor : S. Akshay

Aug 24, 2023

Lecture 08 – Basic Mathematical Structures  
Countable and Uncountable Sets

# Countable and countably infinite sets

## Definition

- ▶ Set  $C$  is called **countably infinite**, if there is a bijection from set  $C$  to  $\mathbb{N}$ .
- ▶ A set is **countable** if it is finite or countably infinite.

Examples: even numbers, number of horses,...

By previous corollary ( $\exists$  surj from any infinite set to  $\mathbb{N}$ )

Countably infinite sets are the “smallest” infinite sets.

## Some questions...

Are the following sets countable?

That is, is there a bijection from these sets to  $\mathbb{N}$ ?

1. the set of all integers  $\mathbb{Z}$
2.  $\mathbb{N} \times \mathbb{N}$
3.  $\mathbb{N} \times \mathbb{N} \times \mathbb{N}$
4. the set of rationals  $\mathbb{Q}$
5. the set of all (finite and infinite) subsets of  $\mathbb{N}$
6. the set of all real numbers  $\mathbb{R}$

## Some questions...

Are the following sets countable?

That is, is there a bijection from these sets to  $\mathbb{N}$ ?

1. the set of all integers  $\mathbb{Z}$
2.  $\mathbb{N} \times \mathbb{N}$
3.  $\mathbb{N} \times \mathbb{N} \times \mathbb{N}$
4. the set of rationals  $\mathbb{Q}$
5. the set of all (finite and infinite) subsets of  $\mathbb{N}$
6. the set of all real numbers  $\mathbb{R}$

To show these it suffices to show that

- ▶ there is an **injection** from these sets to  $\mathbb{N}$

## Some questions...

Are the following sets countable?

That is, is there a bijection from these sets to  $\mathbb{N}$ ?

1. the set of all integers  $\mathbb{Z}$
2.  $\mathbb{N} \times \mathbb{N}$
3.  $\mathbb{N} \times \mathbb{N} \times \mathbb{N}$
4. the set of rationals  $\mathbb{Q}$
5. the set of all (finite and infinite) subsets of  $\mathbb{N}$
6. the set of all real numbers  $\mathbb{R}$

To show these it suffices to show that

- ▶ there is an **injection** from these sets to  $\mathbb{N}$
- ▶ or there is a **surjection** from  $\mathbb{N}$  (or **any countable set**) to these sets.

## Unions of countable sets is countable

Let  $A = \{a_0, \dots\}$  be a countably infinite set and  $B$  be a set. Then, **is  $A \cup B$  countable**, under the following conditions?

1.  $B = \{b_0\}$  is a singleton
2.  $B = \{b_0, \dots, b_n\}$  is a finite set
3.  $B = \{b_0, \dots\}$  is a countably infinite set

## Unions of countable sets is countable

Let  $A = \{a_0, \dots\}$  be a countably infinite set and  $B$  be a set. Then, is  $A \cup B$  countable, under the following conditions?

1.  $B = \{b_0\}$  is a singleton
2.  $B = \{b_0, \dots, b_n\}$  is a finite set
3.  $B = \{b_0, \dots\}$  is a countably infinite set

Can we say  $\{a_0, \dots, b_0, \dots\}$  is a countably infinite set?

## Unions of countable sets is countable

Let  $A = \{a_0, \dots\}$  be a countably infinite set and  $B$  be a set. Then, **is  $A \cup B$  countable**, under the following conditions?

1.  $B = \{b_0\}$  is a singleton
2.  $B = \{b_0, \dots, b_n\}$  is a finite set
3.  $B = \{b_0, \dots\}$  is a countably infinite set

**Can we say**  $\{a_0, \dots, b_0, \dots\}$  is a countably infinite set?

- ▶ But then what is the **position** of  $b_i$  (i.e., natural number corresponding to it)?



## Unions of countable sets is countable

Let  $A = \{a_0, \dots\}$  be a countably infinite set and  $B$  be a set. Then, is  $A \cup B$  countable, under the following conditions?

1.  $B = \{b_0\}$  is a singleton
2.  $B = \{b_0, \dots, b_n\}$  is a finite set
3.  $B = \{b_0, \dots\}$  is a countably infinite set

Can we say  $\{a_0, \dots, b_0, \dots\}$  is a countably infinite set?

- ▶ But then what is the position of  $b_i$  (i.e., natural number corresponding to it)?
- ▶ Rather, choose  $\{a_0, b_0, a_1, b_1, \dots\}$ , then  $b_i$  is at  $(2i + 1)^{th}$  position.

## Unions of countable sets is countable

Let  $A = \{a_0, \dots\}$  be a countably infinite set and  $B$  be a set. Then, is  $A \cup B$  countable, under the following conditions?

1.  $B = \{b_0\}$  is a singleton
2.  $B = \{b_0, \dots, b_n\}$  is a finite set
3.  $B = \{b_0, \dots\}$  is a countably infinite set

Can we say  $\{a_0, \dots, b_0, \dots\}$  is a countably infinite set?

- ▶ But then what is the position of  $b_i$  (i.e., natural number corresponding to it)?
- ▶ Rather, choose  $\{a_0, b_0, a_1, b_1, \dots\}$ , then  $b_i$  is at  $(2i + 1)^{th}$  position.
- ▶ Formally, define a bijection  $f : (A \cup B) \rightarrow \mathbb{N}$  by  $f(a_i) = 2i$  and  $f(b_i) = 2i + 1$

## Unions of countable sets is countable

Let  $A = \{a_0, \dots\}$  be a countably infinite set and  $B$  be a set. Then, is  $A \cup B$  countable, under the following conditions?

1.  $B = \{b_0\}$  is a singleton
2.  $B = \{b_0, \dots, b_n\}$  is a finite set
3.  $B = \{b_0, \dots\}$  is a countably infinite set

Can we say  $\{a_0, \dots, b_0, \dots\}$  is a countably infinite set?

- ▶ But then what is the position of  $b_i$  (i.e., natural number corresponding to it)?
- ▶ Rather, choose  $\{a_0, b_0, a_1, b_1, \dots\}$ , then  $b_i$  is at  $(2i + 1)^{th}$  position.
- ▶ Formally, define a bijection  $f : (A \cup B) \rightarrow \mathbb{N}$  by  $f(a_i) = 2i$  and  $f(b_i) = 2i + 1$
- ▶ Is this correct?

## Products of countable sets are countable

Theorem: The cartesian product of two countably infinite sets is countably infinite

## Products of countable sets are countable

Theorem: The cartesian product of two countably infinite sets is countably infinite

Proof: Let  $A, B$  be countably infinite. Find a way to “number” the elements in  $A \times B = \{(a, b) \mid a \in A, b \in B\}$ .

## Products of countable sets are countable

Theorem: The cartesian product of two countably infinite sets is countably infinite

Proof: Let  $A, B$  be countably infinite. Find a way to “number” the elements in  $A \times B = \{(a, b) \mid a \in A, b \in B\}$ .

- ▶ That is, define a bijection from  $A \times B$  to  $\mathbb{N}$ .

# Products of countable sets are countable

**Theorem:** The cartesian product of two countably infinite sets is countably infinite

**Proof:** Let  $A, B$  be countably infinite. Find a way to “number” the elements in  $A \times B = \{(a, b) \mid a \in A, b \in B\}$ .

- ▶ That is, define a bijection from  $A \times B$  to  $\mathbb{N}$ .

$$f(a_i, b_j) = \left( \sum_{k=1}^{i+j} k \right) + j + 1$$

# Products of countable sets are countable

**Theorem:** The cartesian product of two countably infinite sets is countably infinite

**Proof:** Let  $A, B$  be countably infinite. Find a way to “number” the elements in  $A \times B = \{(a, b) \mid a \in A, b \in B\}$ .

- ▶ That is, define a bijection from  $A \times B$  to  $\mathbb{N}$ .

$$f(a_i, b_j) = \left( \sum_{k=1}^{i+j} k \right) + j + 1$$

## Corollaries

- ▶  $\mathbb{N} \times \mathbb{N}, \mathbb{N} \times \mathbb{N} \times \mathbb{N}, \mathbb{N} \times \mathbb{Z} \times \mathbb{N}$  are countable.



# Products of countable sets are countable

**Theorem:** The cartesian product of two countably infinite sets is countably infinite

**Proof:** Let  $A, B$  be countably infinite. Find a way to “number” the elements in  $A \times B = \{(a, b) \mid a \in A, b \in B\}$ .

- ▶ That is, define a bijection from  $A \times B$  to  $\mathbb{N}$ .

$$f(a_i, b_j) = \left( \sum_{k=1}^{i+j} k \right) + j + 1$$

## Corollaries

- ▶  $\mathbb{N} \times \mathbb{N}$ ,  $\mathbb{N} \times \mathbb{N} \times \mathbb{N}$ ,  $\mathbb{N} \times \mathbb{Z} \times \mathbb{N}$  are countable.
- ▶ The set of (positive) rationals is countable.

# Products of countable sets are countable

**Theorem:** The cartesian product of two countably infinite sets is countably infinite

**Proof:** Let  $A, B$  be countably infinite. Find a way to “number” the elements in  $A \times B = \{(a, b) \mid a \in A, b \in B\}$ .

- ▶ That is, define a bijection from  $A \times B$  to  $\mathbb{N}$ .

$$f(a_i, b_j) = \left( \sum_{k=1}^{i+j} k \right) + j + 1$$

## Corollaries

- ▶  $\mathbb{N} \times \mathbb{N}$ ,  $\mathbb{N} \times \mathbb{N} \times \mathbb{N}$ ,  $\mathbb{N} \times \mathbb{Z} \times \mathbb{N}$  are countable.
- ▶ The set of (positive) rationals is countable.

**Hint:** Show that  $f(a, b) = \begin{cases} a/b & \text{if } b \neq 0 \\ 0 & \text{if } b = 0 \end{cases}$ , is a surjection. How does the result follow?

# Countable sets and functions

Are the following sets countable?

- ▶ the set of all integers  $\mathbb{Z}$
- ▶  $\mathbb{N} \times \mathbb{N}$
- ▶  $\mathbb{N} \times \mathbb{N} \times \mathbb{N}$
- ▶ the set of rationals  $\mathbb{Q}$
- ▶ the set of all (finite and infinite) subsets of  $\mathbb{N}$
- ▶ the set of all real numbers  $\mathbb{R}$

## Comparing $\mathbb{N}$ and set of all subsets of $\mathbb{N}$

Theorem (Cantor, 1891)

There is no bijection between  $\mathbb{N}$  and the set of all subsets of  $\mathbb{N}$ .

# Comparing $\mathbb{N}$ and set of all subsets of $\mathbb{N}$

Theorem (Cantor, 1891)

There is no bijection between  $\mathbb{N}$  and the set of all subsets of  $\mathbb{N}$ .

- ▶ Proving existence just needs one to exhibit a function
- ▶ But how do we prove non-existence?

## Comparing $\mathbb{N}$ and set of all subsets of $\mathbb{N}$

### Theorem (Cantor, 1891)

There is no bijection between  $\mathbb{N}$  and the set of all subsets of  $\mathbb{N}$ .

- ▶ Proving existence just needs one to exhibit a function
- ▶ But how do we prove non-existence? **Try contradiction.**

## Comparing $\mathbb{N}$ and set of all subsets of $\mathbb{N}$

### Theorem (Cantor, 1891)

There is no bijection between  $\mathbb{N}$  and the set of all subsets of  $\mathbb{N}$ .

**Proof by contradiction:** Suppose there is such a bijection, say  $f$ . This would imply that each  $i \in \mathbb{N}$  maps to some set  $f(i) \subseteq \mathbb{N}$ .

	0	1	2	3	...
$f(0)$	✓	×	×	×	...
$f(1)$	✓	×	✓	✓	...
$f(2)$	×	×	×	×	...
$f(3)$	×	✓	×	✓	...

## Comparing $\mathbb{N}$ and set of all subsets of $\mathbb{N}$

### Theorem (Cantor, 1891)

There is no bijection between  $\mathbb{N}$  and the set of all subsets of  $\mathbb{N}$ .

**Proof by contradiction:** Suppose there is such a bijection, say  $f$ . This would imply that each  $i \in \mathbb{N}$  maps to some set  $f(i) \subseteq \mathbb{N}$ .

	0	1	2	3	...
$f(0)$	<del>✓</del> ×	×	×	×	...
$f(1)$	✓	<del>×</del> ✓	✓	✓	...
$f(2)$	×	×	<del>×</del> ✓	×	...
$f(3)$	×	✓	×	<del>✓</del> ×	...

- Consider the set  $S \subseteq \mathbb{N}$  obtained by switching the diagonal elements, i.e.,  $S = \{i \in \mathbb{N} \mid i \notin f(i)\}$ .



# Comparing $\mathbb{N}$ and set of all subsets of $\mathbb{N}$

## Theorem (Cantor, 1891)

There is no bijection between  $\mathbb{N}$  and the set of all subsets of  $\mathbb{N}$ .

**Proof by contradiction:** Suppose there is such a bijection, say  $f$ . This would imply that each  $i \in \mathbb{N}$  maps to some set  $f(i) \subseteq \mathbb{N}$ .

	0	1	2	3	...
$f(0)$	<del>✓</del> ×	×	×	×	...
$f(1)$	✓	<del>×</del> ✓	✓	✓	...
$f(2)$	×	×	<del>×</del> ✓	×	...
$f(3)$	×	✓	×	<del>✓</del> ×	...

- ▶ Consider the set  $S \subseteq \mathbb{N}$  obtained by switching the diagonal elements, i.e.,  $S = \{i \in \mathbb{N} \mid i \notin f(i)\}$ .
- ▶ As  $f$  is bij,  $\exists j \in \mathbb{N}, f(j) = S$ .

# Comparing $\mathbb{N}$ and set of all subsets of $\mathbb{N}$

## Theorem (Cantor, 1891)

There is no bijection between  $\mathbb{N}$  and the set of all subsets of  $\mathbb{N}$ .

**Proof by contradiction:** Suppose there is such a bijection, say  $f$ . This would imply that each  $i \in \mathbb{N}$  maps to some set  $f(i) \subseteq \mathbb{N}$ .

	0	1	2	3	...
$f(0)$	<del>✓</del> ×	×	×	×	...
$f(1)$	✓	<del>×</del> ✓	✓	✓	...
$f(2)$	×	×	<del>×</del> ✓	×	...
$f(3)$	×	✓	×	<del>✓</del> ×	...

- ▶ Consider the set  $S \subseteq \mathbb{N}$  obtained by switching the diagonal elements, i.e.,  $S = \{i \in \mathbb{N} \mid i \notin f(i)\}$ .
- ▶ As  $f$  is bij,  $\exists j \in \mathbb{N}, f(j) = S$ .
- ▶  $S$  and  $f(j)$  differ at position  $j$ , for any  $j$ .

# Comparing $\mathbb{N}$ and set of all subsets of $\mathbb{N}$

## Theorem (Cantor, 1891)

There is no bijection between  $\mathbb{N}$  and the set of all subsets of  $\mathbb{N}$ .

**Proof by contradiction:** Suppose there is such a bijection, say  $f$ . This would imply that each  $i \in \mathbb{N}$  maps to some set  $f(i) \subseteq \mathbb{N}$ .

	0	1	2	3	...
$f(0)$	✓✗	×	×	×	...
$f(1)$	✓	✗✓	✓	✓	...
$f(2)$	×	×	✗✓	×	...
$f(3)$	×	✓	×	✓✗	...

- ▶ Consider the set  $S \subseteq \mathbb{N}$  obtained by switching the diagonal elements, i.e.,  $S = \{i \in \mathbb{N} \mid i \notin f(i)\}$ .
- ▶ As  $f$  is bij,  $\exists j \in \mathbb{N}, f(j) = S$ .
- ▶  $S$  and  $f(j)$  differ at position  $j$ , for any  $j$ .
- ▶ Thus,  $S \neq f(j)$  for all  $j \in \mathbb{N}$ , which is a contradiction!  $\square$

# Cantor's diagonalization

Does this proof look familiar??

# Cantor's diagonalization

Does this proof look familiar??




Figure: Cantor and Russell

# Cantor's diagonalization

Does this proof look familiar??




Figure: Cantor and Russell

- ▶  $S = \{i \in \mathbb{N} \mid i \notin f(i)\}$  is like the one from Russell's paradox.

# Cantor's diagonalization

Does this proof look familiar??




Figure: Cantor and Russell

- ▶  $S = \{i \in \mathbb{N} \mid i \notin f(i)\}$  is like the one from Russell's paradox.
- ▶ If  $\exists j \in \mathbb{N}$  such that  $f(j) = S$ , then we have a contradiction.
  - ▶ If  $j \in S$ , then  $j \notin f(j) = S$ .
  - ▶ If  $j \notin S$ , then  $j \notin f(j)$ , which implies  $j \in S$ .

# Cantor's diagonalization

Does this proof look familiar??




Figure: Cantor and Russell

In fact, using diagonalization Cantor showed that...

- ▶ There cannot be a bijection between **any** set and its power set (i.e., its set of subsets). **(H.W)**
- ▶ So there is an infinite hierarchy of “larger” infinities...



# Cantor's diagonalization

Does this proof look familiar??




Figure: Cantor and Russell

In fact, using diagonalization Cantor showed that...

- ▶ There cannot be a bijection between **any** set and its power set (i.e., its set of subsets). **(H.W)**
- ▶ So there is an infinite hierarchy of “larger” infinities...
- ▶ There is no bijection from  $\mathbb{R}$  to  $\mathbb{N}$  **(H.W)**. Moreover, there is a bijection from  $\mathbb{R}$  to set of subsets of  $\mathbb{N}$ .