SAMPLE

Worlds Beyond Earth: Triton

Neptune's Captured Moon

Aira Astralis

Free sample: Introduction & Chapter 1

putrabooks.github.io

Introduction

Triton is the largest natural satellite of the planet Neptune. It is the only Neptunian moon massive enough for its own gravity to have rounded it, and it keeps a thin, hazy atmosphere. Triton circles its planet against the spin, the one large moon in the Solar System that travels its orbit backwards. Triton is thought to have once been a dwarf planet from the Kuiper belt, captured into Neptune's orbit by the latter's gravity.

At 2,710 kilometers (1,680 mi) in diameter, Triton is the seventh-largest moon in the Solar System, the second-largest planetary moon in relation to its primary (after Earth's Moon), and larger than all of the known dwarf planets. The mean density is 2.061 g/cm3, reflecting a composition of approximately 30-45% water ice by mass, with the rest being mostly rock and metal.

Triton is differentiated, with a crust of primarily ice atop a probable subsurface ocean of liquid water and a solid rocky-metallic core at its center. Although Triton's orbit is nearly circular with a very low orbital eccentricity of 0.000016, its interior may still experience tidal heating through obliquity tides. Triton is one of the most geologically active worlds in the Solar System, with an estimated average surface age of less than 100 million years old.

Its surface is covered by frozen nitrogen and is geologically young, with very few impact craters. Young, intricate cryovolcanic and tectonic terrains suggest a complex geological history. The atmosphere of Triton is composed primarily of nitrogen, with minor components of methane and carbon monoxide. Triton's atmosphere is relatively thin and strongly variable, with its atmospheric surface pressure varying by up to a factor of three within the past 30 years.

Triton's atmosphere supports clouds of nitrogen ice crystals and a layer of organic atmospheric haze.

English astronomer William Lassell found Triton on the tenth of October, 1846, a mere seventeen days after the planet itself was confirmed, and it was the first of Neptune's moons to be found. Voyager 2's brief encounter in 1989 is still the only close-up look humanity has had, as of the mid-2020s.

The measurements of the late 1990s suggested a warming trend, the air measurably denser than at the moment of the flyby, a hint of the seasonal cycle at work.

Chapter 01: The Discovery of Triton

It is the only moon of Neptune massive enough to be rounded under its own gravity and hosts a thin, hazy atmosphere. Triton was the first Neptunian moon to be discovered, on October 10, 1846, by English astronomer William Lassell.

The 1989 flyby of Triton by the Voyager 2 spacecraft remains the only up-close visit to the moon as of 2026.

As the probe was able to study only about 40% of the moon's surface, multiple concept missions have been developed to revisit Triton. These include a Discovery-class Trident and New Frontiers-class Triton Ocean Worlds Surveyor and Nautilus. Triton was discovered by British astronomer William Lassell on October 10, 1846, just 17 days after the discovery of Neptune. When John Herschel received news of Neptune's discovery, he wrote to Lassell suggesting he search for possible moons. Lassell discovered Triton eight days later.

Lassell also claimed for a period to have discovered rings. Although Neptune was later confirmed to have rings, they are so faint and dark that it is not plausible he saw them. A brewer by trade, Lassell spotted Triton with his self-built 61 cm (24 in) aperture metal mirror reflecting telescope (also known as the "two-foot" reflector). This telescope was donated to the Royal Observatory, Greenwich in the 1880s, but was eventually dismantled.

Triton is named after the Greek sea god Triton, the son of Poseidon (the Greek god corresponding to the Roman Neptune). The name was first proposed by Camille Flammarion in his 1880 book Astronomie Populaire, and was officially adopted many decades later. Until the discovery of the second moon Nereid in 1949, Triton was commonly referred to as "the satellite of Neptune".

Lassell did not name his discovery; he later successfully suggested the name Hyperion for the eighth moon of Saturn when he discovered it. Planetary moons other than Earth's were never given symbols in the astronomical literature. Denis Moskowitz, a software engineer who designed most of the dwarf planet symbols, proposed a Greek tau (the initial of Triton) combined with Neptune's trident as the symbol of Triton (). This symbol is not widely used.

Triton is unique among all large moons in the Solar System for its retrograde orbit around its planet (i.e. Most of

the outer irregular moons of Jupiter and Saturn also have retrograde orbits, as do some of the irregular moons of Uranus and Neptune.

However, these moons are all much more distant from their primaries and are small in comparison, with the largest of them (Phoebe) having only 8% of the diameter (and 0.03% of the mass) of Triton, making Triton by far the largest moon with a retrograde orbit. Triton's orbit is associated with two tilts, the obliquity of Neptune's rotation to Neptune's orbit, 30 degrees, and the inclination of Triton's orbit to Neptune's rotation, 157 degrees (an inclination over 90 degrees indicates retrograde motion).

Triton's orbit precesses forward relative to Neptune's rotation with a period of about 678 Earth years (4.1 Neptunian years), making its Neptune-orbit-relative inclination vary between 127 degrees and 173 degrees. That inclination is currently 130 degrees; Triton's orbit is now near its maximum departure from coplanarity with Neptune's. Astronomers usually classify it as an irregular satellite, due to its ex situ origin and retrograde orbit.

However, like a regular satellite, its orbit is relatively close in and its precession is primarily controlled by Neptune rather than the Sun, conflicting with the usual definition of irregular satellites. As a result, Triton is often put in its own category. Triton's rotation is tidally locked to be synchronous with its orbit around Neptune: it keeps one face oriented towards the planet at all times.

Its equator is almost exactly aligned with its orbital plane. At present, Triton's rotational axis is about 40 degrees from Neptune's orbital plane, hence as Neptune orbits the Sun, Triton's polar regions take turns facing the Sun, resulting in seasonal changes as one pole, then the other moves into the sunlight. Such changes were observed in 2010. Triton's revolution around Neptune has become a nearly perfect circle with an eccentricity of almost zero.

Viscoelastic damping from tides alone is not thought to be capable of circularizing Triton's orbit in the time since the origin of the system, and gas drag from a prograde debris disc is likely to have played a substantial role. Tidal interactions also cause Triton's orbit, which is already closer to Neptune than the Moon is to Earth, to gradually decay further; predictions are that 3.6 billion years from now, Triton will pass within Neptune's Roche limit.

This will result in either a collision with Neptune's atmosphere or the breakup of Triton, forming a new ring system similar to that found around Saturn. However, a newer study done in 2025 has analyzed that Triton will reach Neptune's Roche limit in 28 billion years. This indicates a more stable orbital evolution of Triton over a long period of time.

The current understanding of moons in retrograde orbits means they cannot form in the same region of the solar nebula as the planets they orbit. Therefore, Triton must have been captured from elsewhere in the Solar System. Astrophysicists believe it might have originated in the Kuiper belt, a ring of small icy objects extending from just inside the orbit of Neptune to about 50 AU from the Sun.

Thought to be the point of origin for the majority of short-period comets observed from Earth, the belt is also home to several large dwarf planets

including Pluto, which is now recognized as the largest in a population of Kuiper belt objects (the plutinos) locked in resonant orbits with Neptune. Triton is only slightly larger than Pluto and is nearly identical in composition, which has led to the hypothesis that the two share a common origin.

This has been further supported in a 2024 study of the chemical composition of Pluto and Triton which suggests they originated in the same region of the outer Solar System before the latter was pulled into Neptune's orbit.

Studying prior data on the two bodies, the team found that both have a large amount of nitrogen and trace amounts of methane and carbon monoxide, which could have accumulated in the outer regions of the young nebula "For some reason, Triton was then ejected from this region and ensnared by Neptune".

The proposed capture of Triton may explain several features of the Neptunian system, including the extremely eccentric orbit of Neptune's moon Nereid and the scarcity of moons as compared to the other giant planets. Triton's initially eccentric orbit would have intersected the orbits of irregular moons and disrupted those of smaller regular moons, dispersing them through gravitational interactions.

Triton's eccentric post-capture orbit would have also resulted in tidal heating of its interior, which could have kept Triton fluid for a billion years; this inference is supported by evidence of differentiation in Triton's interior. This source of internal heat disappeared following tidal locking and circularization of the orbit. Two types of mechanisms have been proposed for Triton's capture.

To be gravitationally captured by a planet, a passing body must lose sufficient energy to be slowed down to a speed less than that required to escape. An early model of how Triton may have been slowed was by collision with

another object, either one that happened to be passing by Neptune (which is unlikely), or a moon or proto-moon in orbit around Neptune (which is more likely).

A more recent hypothesis suggests that, before its capture, Triton was part of a binary system.

When this binary encountered Neptune, it interacted in such a way that the binary dissociated, with one portion of the binary expelled, and the other, Triton, becoming bound to Neptune. This event is more likely for more massive companions. This hypothesis is supported by several lines of evidence, including binaries being very common among the large Kuiper belt objects. The event was brief but gentle, saving Triton from collisional disruption.

However, simulations in 2017 showed that after Triton's capture, and before its orbital eccentricity decreased, it probably did collide with at least one other moon, and caused collisions between other moons. Triton is the seventh-largest moon and sixteenth-largest object in the Solar System and is modestly larger than the dwarf planets Pluto and Eris.

It is the largest known object believed to have originated in the Kuiper Belt. It is also the largest retrograde moon in the Solar System. It accounts for more than 99.5% of all the mass known to orbit Neptune, including the planet's rings and fifteen other known moons, and is also more massive than all known moons in the Solar System smaller than itself combined.

Also, with a diameter 5.5% that of Neptune, it is the largest moon of a gas giant relative to its planet in terms of diameter, although Titan is bigger relative to Saturn in terms of mass (the ratio of Triton's mass to that of Neptune is approximately 1:4788). It has a radius, density (2.061 g/cm3), temperature, and chemical composition similar to that of Pluto. Triton's surface is covered with a transparent layer of annealed frozen nitrogen.

Only 40% of Triton's surface has been observed and studied, but it may be entirely covered in such a thin sheet of nitrogen ice. Triton's surface consists of 55% nitrogen ice with other ices mixed in. Water ice comprises 15-35% and frozen carbon dioxide (dry ice) the remaining 10-20%. Trace ices include 0.1% methane and 0.05% carbon monoxide. There could also be ammonia ice on the surface, as there are indications of ammonia dihydrate in the lithosphere.

Triton's mean density implies that it probably consists of about 30-45% water ice (including relatively small amounts of volatile ices), with the remainder being rocky material. Triton's surface area is 23 million km2, which is 4.5% of Earth, or 15.5% of Earth's land area. Triton has an unusually high albedo, reflecting 60-95% of the sunlight that reaches it, and it has changed only slightly since the first observations. By comparison, the Moon reflects only 11%.

This high albedo causes Triton to reflect a lot of whatever little sunlight there is instead of absorbing it, causing it to have the coldest recorded temperature in the Solar System at 38 K (-235 degreesC). Triton's reddish color is thought to be the result of methane ice, which is converted to tholins under exposure to ultraviolet radiation.

Because Triton's surface indicates a long history of melting, models of its interior posit that Triton is differentiated, like Earth, into a solid core, a mantle and a crust. Water, the most abundant volatile in the Solar System, comprises Triton's mantle, enveloping a core of rock and metal.

The discovery's immediate aftermath is a study in how slow the outer Solar System was to give up its facts. Lassell's first observations were confirmed by other telescopes within months, but the moon's orbit resisted precise determination for years, because the object was faint, the planet had barely completed a fraction of a revolution, and every measurement carried the errors of the pre-photographic age; by mid-century the orbit was good enough to show the retrograde direction, and that single result, published quietly in the ephemeris tables, waited a hundred years to be understood as the moon's true introduction. Lassell himself moved on to bigger mirrors and other discoveries, and his accounts of the Triton observations, kept in the notebooks of a working brewer-astronomer, make clear that he regarded the companion of the new planet as a sidelight to the mirror-making he considered his real trade.

A paragraph belongs here about the moon the nineteenth century saw, because the contrast with what the modern reader sees is itself instructive. To Lassell and his contemporaries Triton was a star of the thirteenth magnitude, a point of light inseparable from the glare of its planet except in the largest instruments, and the whole apparatus of nineteenth-century astronomy, the micrometers, the visual estimates, the long nights at the eyepiece, went into measuring that point's position to a fraction of an arcsecond. The campaign succeeded: the orbit was mapped, the retrograde direction established, the mass bounded from the perturbations, all from a light that a modern pair of binoculars could not find. The discovery chapter's second lesson, after the luck of timing, is the economy of the old

astronomy, which squeezed a moon's whole biography out of the position of a single dot, and the modern reader who knows the geysers and the pink frost should spare a thought for the century in which the dot was all there was.

End of free sample. The full book is available at putrabooks.github.io