SAMPLE

Cosmos Unveiled: Ultra-Diffuse Galaxies

Aira Astralis

Free sample: Introduction & Chapter 1
putrabooks.github.io

Introduction

In 2015, two teams pointed new kinds of telescopes at the Coma cluster and found something nobody expected: a population of galaxies as large as the Milky Way, shining with only a hundredth of its light. They were not new objects in any physical sense. They had been sitting in survey images for decades, invisible in plain sight, spread across the sky like smoke where everyone had been looking for campfires. Almost overnight, ultra-diffuse galaxies became one of the liveliest problems in astronomy, and they have kept that title ever since.

The surprise was not that faint galaxies exist. Astronomers have catalogued dwarfs for a century, and they have known since the 1980s that some galaxies spread their light so thin that they nearly vanish against the sky. The surprise was the combination of size and dimness. A galaxy with the footprint of a giant and the starlight of a dwarf did not fit the tidy ladder of galaxy types that a century of classification had built. Either such galaxies were freaks, rare accidents of cosmic history, or the ladder itself was missing a rung. That fork, freaks or missing rung, is worth holding through the whole book, because it is the standard shape of a discovery in astronomy, and the ultra-diffuses display it in an unusually clean form. When a new class of object appears, the first question is always how much of the sky it represents, and the answer arrived within a few years of the first discoveries: once the surveys knew what to look for, the ultra-diffuses turned up everywhere, in clusters, in groups, in the field, dozens then hundreds then catalogued by the thousand, and the freak hypothesis quietly dissolved. What remained was the more interesting verdict, the one this book spends its chapters on: the galaxy population had a whole faint wing the classification had never modeled, and the missing rung had been missing for a century.

The answer has turned out to be richer than either choice. Ultra-diffuse galaxies are common, they live everywhere from dense clusters to empty field, and they refuse to settle into a single story. Some appear to be almost nothing but dark matter, ghosts whose few stars ride in enormous invisible halos. A handful appear to have no dark matter at all, and those objects have forced a fight over the meaning of galaxy mass itself. Between the extremes, most seem to be ordinary dwarfs that were puffed up by spin, by stellar feedback, or by the rough handling of their neighborhoods.

This book is an account of that problem: what ultra-diffuse galaxies are, how they were found, what their existence does to theories of galaxy formation, and what remains unresolved. It is written for the curious reader who wants the structure of the argument rather than a list of facts, and it holds to a few commitments. It names no products, cites no citations, and leans on no statistics that expire. The physics here is old and stable, and the story is current enough to be genuinely unfinished, which is the best kind of story science has to offer.

The chapters that follow take the story in the order the field lived it: the long blindness and the instruments that ended it, the discovery census, the anatomy of the class, the formation theories, and the two dark matter controversies that put the ghosts on the front pages of astronomy. Later chapters widen out to the environments, the heritage, the simulations, and the craft of measurement, and the book closes with the open questions, which in this subject are not a formality but the actual point. The reader needs no mathematics beyond a willingness to picture sizes and brightnesses, and no background beyond a general acquaintance with the night sky's inventory.

Chapter 01: A Problem Hiding in Plain Sight

To understand why ultra-diffuse galaxies went unnoticed for so long, it helps to know what a telescope image actually records. A galaxy's light arrives spread across a patch of sky, and the eye or the camera sums up brightness per unit area, which astronomers call surface brightness. A large galaxy that is intrinsically dim can spread its starlight across so much sky that the surface brightness falls below the glow of the night sky itself, and once that happens, the galaxy does not look like an object at all. It looks like a slight brightening of the background, no more conspicuous than a breath on a window.

Telescopes do not fix this the way intuition suggests. A larger mirror gathers more light, but for an extended object it also gathers more sky, and the surface-brightness limit barely improves. The classic detectors made the problem worse. Photographic plates responded to light in a way that suppressed faint signals, a property that made the plates forgiving of overexposure but blind to the very faintest, largest structures. Galaxies spread across many square minutes of arc simply did not register.

The first cracks in this blindness appeared in the 1970s and 1980s. Careful photographic work, and in particular a technique that coaxed faint detail out of photographic plates, revealed galaxies of astonishing faintness. The most famous of them, later called Malin 1, turned out to be a spiral as wide as several Milky Ways, with a disk so faint

that it had been catalogued as a minor smudge for years. Its discovery told astronomers that the galaxy population had an entire faint wing, and it opened the study that became known as low surface brightness astronomy.

For two decades, that wing belonged mostly to the spirals. The compact, starry universe of clusters was thought to be mapped, because cluster surveys were tuned to find dense, bright things: ellipticals, lenticulars, ordinary dwarfs. What went unmeasured was the middle ground, objects both large and dim, especially in crowded cluster fields where the light of bright galaxies washes out everything around them.

Then instrument design changed the game. In the early 2010s, a small group of astronomers built a telescope out of commercial telephoto lenses, a design chosen deliberately for the property that mattered most: the lenses scatter very little stray light, so the instrument sees faint, extended structures without being blinded by nearby bright sources. The array, named for the insect whose vision it imitates, combined many such lenses into one camera with an enormous field of view. Its first deep images of nearby clusters, taken with long integrations and careful stacking, showed large, ghostly galaxies that standard surveys had missed, not because they were far away, but because they were large and nearly featureless.

The mathematics of the blindness is worth one more look, because it explains an old embarrassment. A galaxy the size of the Milky Way, with a hundredth of its stars, placed at the distance of the Coma cluster, has a central surface brightness more than ten times fainter than the typical night sky background in a good survey image. Human vision, which is good at edges and contrast, simply reports nothing there. Algorithms can do better, but only if someone tells them to look, and for decades, nobody had thought to ask how many enormous, barely luminous galaxies the universe might hold.

The lesson is a standing warning for all of observational astronomy: telescopes do not see the universe, they see the fraction of the universe that survives the instrument. Every telescope is a filter, and the filters of the late twentieth century were built to find compact, bright galaxies, so compact, bright galaxies were found in abundance. The ultra-diffuses did not hide. They were simply transparent to the questions the instruments were asking. Sit with that sentence a moment, because it is the book's methodological heart and its warning against the confidence of any catalog. An instrument is not a neutral window; it is a question repeated mechanically across the sky, and it answers that question and no other, so the galaxy catalogs of the twentieth century were, in strict honesty, catalogs of galaxies bright enough in their centers to trip the detection software, a selection nobody chose and everybody inherited. The ultra-diffuses had been in the plates and the frames all along, sitting unremarked in the backgrounds of famous cluster photographs, and it took a survey built for another purpose entirely, counting galaxy clusters by their faint smudges, to trip over them. The lesson generalizes to every science that surveys: what you find is shaped by what you ask, and the deepest discoveries are the ones waiting behind an unasked question.

A second reason deserves equal weight: the culture of classification itself. The great atlases of galaxies were built to answer the questions of their era, which concerned the bright, structured, surveyable objects, and a classification scheme is a kind of law about what features matter. Galaxies that presented no features, that arrived as smooth monotonous fields, fell outside the scheme's vocabulary. Astronomers are trained to describe what their categories can name, and an object with no category, no name, and no theory demanding it tends to be filed as a defect: a plate flaw, a reflection, a cloud. The ultra-diffuses spent decades being classified as errors, and the classification was not careless, it was simply the taxonomy working as built, on objects the taxonomy never anticipated.

The blindness also had a numerical form that is easy to state. Survey depth is usually quoted in magnitudes, which measure total brightness, and total brightness is a quantity that favors compact objects. A galaxy's total light is its surface brightness multiplied by its area, so a large, faint galaxy can have a respectable total magnitude while remaining undetectable in practice, because its light is spread across a field where the sky itself is brighter than the signal everywhere you look. Two objects with the same total magnitude can differ by a factor of a hundred in detectability. The survey limit that mattered was never magnitude. It was surface brightness, and surface brightness was a quantity the classic catalogues rarely printed.

The professional habits of the era reinforced the blindness in a quieter way. Observing time on large telescopes was scarce, and allocation committees judged proposals by expected results, which meant that the safest use of a precious night was a target guaranteed to produce a spectrum or a detection. A proposal to stare at blank sky for hours, looking for objects that might not exist, was not a proposal a committee could fund in good conscience, and so the deep, patient imaging that would have found the ghosts decades early was simply never done. Discoveries are often attributed to serendipity, but the missing ingredient is usually permission, and the ultra-diffuses waited, in part, for an era when a small team could build its own instrument and answer to no committee for how it spent its

darkness.

The visibility-paradox arithmetic gets a line. The objects were invisible for a working reason, a galaxy spread over the area of the moon shining at a hundredth the sky's glow per area, the account keeping the arithmetic's lesson, the brightest examples being detectable in century-old plates and still unreported for seventy years, the hiding file closing with the standing explanation, the ghosts were not hidden by distance but by their own manners, spread thin enough to match the sky's own noise.

The plate-archive amnesty gets a line. Some ghosts appear in the oldest plates on re-inspection, the century-old exposures having caught examples nobody catalogued, the account keeping the amnesty's generosity, the class having been photographable all along and merely unexamined, the hiding file finishing with the standing hindsight, the ghosts were photographed before they were discovered, and the photographs waited a working lifetime for a reader.