SAMPLE

Cosmos Unveiled: White Dwarfs

Aira Astralis

Free sample: Introduction & Chapter 1
putrabooks.github.io

Introduction

A white dwarf is the stellar core remnant left behind after a low- or intermediate-mass star has exhausted its nuclear fuel and gently shed its outer layers. Roughly the size of Earth yet typically carrying more than half the mass of the Sun, a white dwarf packs matter into densities around a million times that of water: a single teaspoon of its material would weigh about a tonne on Earth. The star no longer burns hydrogen or anything else. Instead, it shines for billions of years by slowly leaking away the enormous store of residual heat trapped inside it.

This radiation, which initially has a high color temperature, lessens and reddens over time. As a white dwarf ages, it cools and dims, marching down the color spectrum from blue-white through white and yellow to a dull red. Eventually, white dwarfs may become cool enough that their material begins to crystallize, turning the carbon-and-oxygen core into something like a vast cosmic diamond. Because they cool so slowly, the oldest white dwarfs in the galaxy are expected to persist for an extraordinarily long time - on the order of ten to the thirty-eighth power years - before fading into the theoretical state known as a black dwarf.

White dwarfs are not rare curiosities. They are the fate awaiting the overwhelming majority of stars: anything from roughly one-tenth to about eight times the mass of the Sun ends its life this way, which means the vast majority of the galaxy's stars will one day be white dwarfs. Our own Sun will join their number in roughly five billion years, first swelling into a red giant, casting off its envelope as a planetary nebula, and settling down as a slowly cooling ember about the size of Earth.

The story of how astronomers came to understand these strange objects is one of the great detective tales in the history of science. For decades, the companion of Sirius sat in the sky as an embarrassment - a star too faint to be real, radiating from a body so small and so dense that known physics could not explain it. Resolving that paradox required the birth of quantum mechanics, the willingness of a few astronomers to take impossible data seriously, and the slow accumulation of evidence from spectroscopy, stellar statistics, and eventually space observatories.

This book traces that story. It begins with the discovery of the first white dwarfs and the puzzle they posed, explains the extraordinary physics of degenerate matter that makes them possible, and walks through their key characteristics: their masses, sizes, compositions, atmospheres, and the strange crystalline futures that await them. Later chapters place white dwarfs in the broader context of stellar evolution, describe how astronomers observe and measure them, and explore the role they play in some of the most energetic events in the universe - including the Type Ia supernovae used to measure the expansion of the cosmos itself.

One way to grasp the scale of the white dwarf phenomenon is to count. Within a sphere of a few dozen light-years around the Sun - a region containing perhaps a thousand ordinary stars - astronomers already know of several dozen white dwarfs, and the tally keeps growing with each deep survey. Extrapolated across the galaxy, the count of white dwarfs reaches into the tens of billions. Nearly every one is invisible to the naked eye, and yet together they represent the most common endpoint of stellar life. Learning to read them - their spectra, their colors, their slow fading - is learning to read the future of nearly every star.

A note on how to read this book: the chapters are ordered so that each builds on the last, from the historical discovery through the physics to the open research frontier, but each is also written to stand alone. Readers new to the subject may wish to read Chapters 1 and 2 together as a single story; readers with more background may prefer to jump directly to the chapters on modern research, mysteries, and future directions. Throughout, the emphasis falls on understanding rather than memorizing: the goal is that by the final page, the reader can explain not just what a white dwarf is, but why it must be.

Chapter 01: The Discovery of White Dwarfs

The discovery of white dwarfs is a story about taking anomalies seriously. In the nineteenth century, positional measurements of some stars became precise enough to detect small changes in their location across the sky. Friedrich Bessel used such measurements to determine that the stars Sirius, the brightest star in Canis Major, and Procyon, the brightest in Canis Minor, were changing their positions periodically. In 1844 he predicted that both stars had unseen companions, their slow wobbles betraying the gravitational tug of dark partners. Bessel roughly estimated the period of the companion of Sirius to be about half a century, and in 1851 the astronomer C. A. F. Peters computed a full orbit for it.

It was not until 31 January 1862 that the predicted companion revealed itself. Alvan Graham Clark, testing a new

telescope lens, observed a previously unseen star close to Sirius - later identified as the predicted companion, now known as Sirius B. The discovery was a triumph of celestial mechanics: a star found not by being seen, but by being computed. Sirius B, however, immediately created a problem. Given its distance and its faintness, it had to be extraordinarily dim for its color - and if it was a normal star at that luminosity, its temperature should have made it much brighter than it appeared.

The second white dwarf was found hiding in the triple star system 40 Eridani. Its companion, 40 Eridani B, was plainly hot and small, yet so intrinsically faint that its existence defied the stellar physics of the day. The third came in 1917, when Adriaan van Maanen discovered an isolated faint white star with a large proper motion, now called van Maanen's Star. These three - Sirius B, 40 Eridani B, and van Maanen's Star - are remembered as the classical white dwarfs, the founding members of the class.

Spectroscopy deepened the mystery. In 1915, Walter Adams announced that he had found the spectrum of Sirius B to be similar to that of Sirius itself, meaning the tiny companion was as hot as its brilliant primary - as hot as a star that outshone it by a factor of ten thousand. In 1922, Willem Luyten published a series of three papers on faint white stars that had high proper motion, meaning they were low-luminosity stars close to the Earth. In his third paper he used the term "dwarf" for these objects, and the term was later popularized by Arthur Eddington as "white dwarf."

The decisive confrontation came from a simple estimate. If a star is in a binary system, as is the case for Sirius B or 40 Eridani B, it is possible to estimate its mass from orbital mechanics. Since hotter bodies radiate more energy than colder ones, a star's surface brightness can be estimated from its effective temperature, which the spectrum provides. If the star's distance is also known, its absolute luminosity and radius follow. When Ernst Opik estimated the densities of visual binary stars in 1916, he found that 40 Eridani B had a density of over 25,000 times that of the Sun - a figure so absurd that many astronomers simply set the data aside.

Eddington did not set it aside. As he put it in 1927: "We learn about the stars by receiving and interpreting the messages which their light brings to us. The message of the companion of Sirius when it was decoded ran: 'I am composed of matter 3,000 times denser than anything you have ever come across; a tonne of my substance would be a little nugget that you could put in a matchbox.' What reply can one make to such a message? The reply which most of us made in 1914 was: 'Shut up. Don't talk nonsense.'" In 1939, Henry Norris Russell looked back on the discovery and noted that Pickering had suggested that such exceptions lead to breakthroughs - and in this case the exception led to the discovery of white dwarfs, and to a new understanding of matter itself.

The observational limits of the era merit emphasis. In Bessel's time, detecting the wobble of Sirius meant measuring its position against the sky to a precision of a fraction of an arcsecond, night after night, year after year, with instruments whose lenses were ground by hand. The companion he inferred was never seen in his lifetime; Bessel died in 1846, sixteen years before Clark's discovery, still confident that the unseen stars were real. When Clark finally found Sirius B, he was not even looking for it - he was testing the optical quality of an eighteen-and-a-half-inch refractor, the largest in America at the time, by observing the brightest star in the sky as a difficult target.

Clark's father and brother ran the firm of Alvan Clark & Sons, the great American telescope makers, and the sighting caused an immediate sensation. Here, at last, was Bessel's invisible star, dimmer than ten thousandths of its brilliant companion, and yet unmistakably present. Astronomers then made the fateful assumption that the companion was a normal, cool, red star - an assumption that, once its spectrum was finally read in 1915, collapsed spectacularly. Sirius B was not cool at all. It was one of the hottest objects ever measured, radiating from a surface far hotter than Sirius itself, and its faintness could only mean one thing: it was astonishingly small.

Van Maanen's discovery in 1917 followed a different logic. His star had no brilliant companion to hide beside; it betrayed itself by moving. Among the faint stars of the sky, a large proper motion is the signature of proximity, and a faint, white-colored star moving quickly across the sky had to be both nearby and intrinsically dim. Van Maanen's Star turned out to be only about fourteen light-years away, making it one of the nearest stars of any kind - and, at the time of its discovery, the most perplexing.

The naming of the class settled slowly. The first spectral type designations treated these stars as oddities among the main sequence. It was Luyten's statistical work on high-proper-motion white stars that established them as a group worth a name, and Eddington's brilliant popularization in "The Internal Constitution of the Stars" that fixed "white dwarf" in the vocabulary of astronomy. The name is not perfectly accurate - hot young members are bluish, old cool members are yellowish or reddish, and none is a dwarf in the stellar sense of a main-sequence star - but it

captured exactly the two things that startled the 1920s: their color, and their size.

The three classical white dwarfs also framed the puzzle in three different ways, and the differences mattered. Sirius B posed a dynamical puzzle: a star whose mass could be weighed from its orbit. 40 Eridani B posed a photometric puzzle: a star too hot for its brightness, in a system whose distance was well known from its bright companions. Van Maanen's Star posed a statistical puzzle: a high-proper-motion star whose dimness implied an absolute luminosity far below the main sequence. Three independent lines of evidence, each pointing to the same impossible conclusion - that a class of stars existed with densities no known physics could support.

It is worth noting what did not happen next. No single dramatic experiment settled the matter. Instead, the case was built the way most of science is actually built: a decimal place added to a parallax here, a cleaner spectrum there, a careful tally of the faint blue stars by Luyten through the 1920s and 1930s. By the time Fowler applied the new quantum mechanics to the problem in 1926, the observational ground had been prepared for decades. The theory did not conquer the data; it was handed to them.