SAMPLE

Cosmos Unveiled: Active Galactic Nuclei

The Engines Inside Quasars

Aira Astralis

Free sample: Introduction & Chapter 1

putrabooks.github.io

Contents

Introduction

Chapter 01: A Light That Should Not Exist

Chapter 02: The Engine at the Center

Chapter 03: The Whirlpool Before the Fall

Chapter 04: Jets of Fire Across a Million Years

Chapter 05: The Zoo and Its Single Animal

Chapter 06: Reading the Light

Chapter 07: Weighing What Cannot Be Seen

Chapter 08: Hosts and Partners

Chapter 09: The Blinking Beacon

Chapter 10: Monsters Through Cosmic Time

Chapter 11: The Thermostat of the Galaxies

Chapter 12: Watching the Watchers

Chapter 13: The Unsolved Core

Chapter 14: Why the Beacons Matter

Conclusion

This free sample contains the Introduction and Chapter 1. The complete book contains the full contents above.

Introduction

There are places in the universe where a single region the size of our solar system pours out more light than an entire galaxy of stars. Astronomers spent decades puzzling over them before anyone believed the answer, because the answer seemed absurd: matter falling into a black hole, and shining brilliantly in the act of being devoured.

These objects are called active galactic nuclei, usually shortened to AGN, and this book is their story. It is a story that runs from a puzzled optician's notebook in 1943 to the largest surveys of the modern sky, and from the strangest-looking photographs in early radio astronomy to the confident picture we now hold of disks of doomed gas whirling around black holes heavier than a billion suns. It is also a story about how science works when the evidence points somewhere nobody wanted to look.

The phrase itself deserves a careful reading. A galactic nucleus is simply the heart of a galaxy, the crowded central bulge where stars pack most densely and where the deepest gravitational well is found. Nearly every large galaxy has a supermassive black hole waiting in that well. In most galaxies, including our own Milky Way, the black hole sits in near darkness, starved of fuel, detectable only by the sway of nearby stars. But in a small fraction of galaxies, gas and dust drift too close, the feeding begins, and the nucleus lights up in every band of the electromagnetic spectrum at once. That lit-up nucleus is what the word active means.

The numbers deserve a moment of stillness. A bright AGN can outshine its host galaxy a hundred or even a thousand times over. The region doing the shining is comparable in size to our solar system, while the galaxy around it spans a hundred thousand light-years. If you scaled the galaxy down to the size of a city, the engine would be a coin on a downtown sidewalk, and the coin would be brighter than the city.

The puzzle was so extreme that astronomers repeatedly proposed and abandoned explanations before accepting the black hole answer. Stars could not do it; no cluster of stars could pack tightly enough without collapsing. Nuclear reactions in ordinary gas could not do it; the fuel would run out too quickly. Only gravity offered a furnace efficient enough, and the black hole, long treated as a place where physics goes silent, turned out to be the most efficient power plant the universe has ever built. Matter that spirals inward before crossing the horizon can release several to a dozen percent of its rest-mass energy on the way down, far beyond the fraction released by nuclear fusion.

The chapters that follow build the picture one piece at a time. We begin with the discovery, because the discovery itself is a lesson in how a scientific community argues its way to an uncomfortable truth. Then we descend through the machinery: the black hole, the whirling disk, the swirling clouds of gas, the twin jets that some of these objects hurl into intergalactic space. We meet the menagerie of names, quasars and Seyferts and blazars, and see how a seemingly chaotic zoo organized itself into a single unified model viewed from different angles. We learn how astronomers weigh the invisible monster at the center, how these objects flicker and what their flickering reveals, and how the population of active nuclei has changed across the thirteen-odd billion years of cosmic history. Finally we turn to the deepest questions still standing, because for all the progress of eight decades, the cores of active galaxies remain one of astronomy's most productive frontiers.

One warning is worth giving at the start. The mind's image of a black hole as a vacuum cleaner, sucking in everything nearby, is wrong, and the wrongness matters. A black hole only captures what wanders so close that it cannot escape, and the disk around it shines precisely because most of the infalling matter fights its fate all the way down, colliding, heating, and radiating. The monster is bright because its meals are violent, not because it is greedy. Keep that correction in mind and the whole subject opens up; forget it and every explanation that follows will seem paradoxical.

This book is written for the reader with curiosity but no special training. Every idea is introduced before it is used, every term defined when it first appears, and the mathematics that professionals use has been translated into plain language and proportion. The aim is that by the final page, the reader will not merely know the facts about active galactic nuclei but will be able to reconstruct them, to see why the evidence forces the conclusions, and to feel the particular satisfaction that comes from understanding one of the universe's genuine extremes from the inside out.

Chapter 01: A Light That Should Not Exist

The story begins with a pattern rather than a bang. In 1943, Carl Seyfert, a young astronomer working at Mount

Wilson Observatory in California, published a short study of a dozen unusual galaxies. Most were spiral galaxies that looked ordinary at first glance, but their cores were strangely intense, small points of light far brighter than the center of a normal galaxy. Stranger still, their spectra were unlike anything stellar. Ordinary starlight shows a smooth rainbow crossed by dark absorption lines, the fingerprints of atoms in the star's cool outer layers. The spectra of

Seyfert's galaxies were crossed instead by bright emission lines, glowing bands of color that a star never produces, and some of those lines were enormously broad, smeared across hundreds of angstroms, as if the glowing gas were being flung about at thousands of kilometers per second.

Two conclusions hid in that data, and both were unsettling. Bright emission lines mean hot, thin gas, excited by something, and there simply is no known way for an ordinary galaxy's heart to fill its core with such gas at such temperatures. And the breadth of the lines meant violent motion in a tiny space, because the cores were unresolved points even in the best photographs. Small and bright and fast: whatever lived in those galactic hearts was no ordinary collection of stars.

The astronomy of that era had no framework for such objects, and Seyfert's galaxies were filed away as curiosities, noted and largely ignored for two decades. Textbooks mentioned them in passing. Few researchers suspected they were the visible edge of a far larger phenomenon, because the most extreme members of the family had not yet been recognized as galaxies at all.

The next clue came from a completely different corner of science: radio. In the years after the Second World War, the radar technology developed for battle was turned toward the sky, and radio astronomy was born. Early radio surveys found the sky surprisingly full of sources, and by the late 1950s the Third Cambridge Catalogue had listed hundreds of them. A few stood out as exquisitely compact, and when optical astronomers looked at their positions, they found strange blue points of light that looked like stars but refused to behave like them.

The most famous of these became known as 3C 273, the two hundred and seventy-third source in the Cambridge catalogue. Its optical counterpart was a starlike point with a faint jet of material sticking out to one side, and its spectrum was a mess of broad emission lines that matched nothing in any

laboratory. Through the early 1960s, this was one of astronomy's standing embarrassments: bright objects visible on photographic plates, with spectra no one could interpret.

The breakthrough came in February 1963, and the way it came is worth telling because it captures the mood of the field. Maarten Schmidt, a Dutch-born astronomer at Caltech, had been staring at the spectrum of 3C 273, and on a day of frustration he tried something radical. Hydrogen has a famous line, a red line at 6563 angstroms, the unmistakable signature of glowing hydrogen in the laboratory and in stars. Nothing in 3C 273's spectrum sat there. But Schmidt realized that if the hydrogen line were shifted far toward the red, moved by an enormous velocity, then other hydrogen lines would land exactly where the mystery lines sat. The required shift was about sixteen percent toward the red, a velocity of nearly a tenth the speed of light.

To see why this was shocking, picture the scale. In 1963, the known universe of galaxies stretched out with redshifts of a few percent at most. A sixteen percent redshift meant 3C 273 was not a star in our galaxy but an object roughly two billion light-years away. And it was not a galaxy of a hundred billion stars. It looked like a single point. A single point, at two billion light-years, visible from Earth with ordinary telescopes, competing in brightness with the light of entire nearby galaxies.

The word for this object already existed, coined with some embarrassment by Hong-Yee Chiu in 1964, mashing together the phrase quasi-stellar radio source: quasar. Within months, astronomers re-examined other blue starlike points and found more redshifted monsters. Within a few years, quasars were being discovered at redshifts beyond two, pushing outward through the early universe, and the argument began about what could possibly power them.

The candidates were canvassed and rejected one by one, and each rejection teaches something. A dense cluster of stars was proposed, but the brightness demanded so many stars packed into so small a region that collisions and collapse seemed unavoidable, and the spectra of starlight never produce those furious emission lines. A chain of supernova explosions was proposed, hundreds of them per year in the same core, but supernovae brighten and fade on schedule and quasars shine steadily for centuries at least; no chain could be sustained that long at the required rate without consuming a galaxy's worth of stars. Exotic nuclear reactions in super-dense gas were proposed, and failed on efficiency and stability. Every conventional furnace was too weak, too short-lived, or too unstable.

Gravity offered the way out. Two researchers in particular, Edwin Salpeter in the United States and Yakov

Zeldovich in the Soviet Union, argued in 1964 that matter falling toward a black hole could release energy with an efficiency no star could match. At first the idea was met with skepticism by some of the field's most respected figures, including Fred Hoyle, whose objection is worth remembering: a black hole with no magnetic field or internal structure seemed too simple a machine to radiate anything. The answer, worked out over the following decade, is that the black hole does not radiate at all. The disk of matter spiraling into it does, heated by friction and turbulence as it converts its orbital motion into light before the final plunge. The hole's job is to make the whirlpool, and the whirlpool makes the shine.

By the early 1970s, the theory had a name, accretion, and a champion, Donald

Lynden-Bell, who laid out in detail how a black hole of millions of solar masses, fed at the rate of a few suns' worth of gas per year, could power the observed quasars. Radio observations soon found the expected signature of violence: galaxies with lobes of radio-emitting plasma flung hundreds of thousands of light-years into space, obviously powered by something at the dead center. And when the Seyfert galaxies of 1943 were re-examined, astronomers realized that the same engine was running in them, in a smaller, closer, easier-to-study version.

The discovery chapter of this subject closes with a satisfying unification: the starlike quasars, the strange Seyfert cores, and the roaring radio galaxies were all placed on one axis. Their common property, and the property that defines the entire subject of this book, is a galactic nucleus radiating far more power than stars can account for, from a region far too small to contain enough stars. Once the community accepted that single statement, everything else in the following chapters becomes an engineering problem: what is the engine, what does the fuel line look like, and what happens to the exhaust.

End of free sample. The full book is available at putrabooks.github.io