

SMAC Results

Yuri Lavinas

4/16/2017

Contents


| | |
|---------------------------------|-----------|
| Summary | 1 |
| Convergency plots | 2 |
| ANOVA test and HSD Tukey | 14 |
| All regions | 14 |

Summary


In this document we show the convergency plots for the GAModel and ReducedGAmodeL with tournize size (k) 3 and tournsize (k) 2. The value of k being 2 was select by SMAC after it was executed for 2 days for both models.

Convergency plots


Kanto 2005 GAModel (Orange, k=3, Blue k=2)




Kanto 2005 ReducedGAModel (Orange, k=3, Blue k=2)







Kanto 2006 GAModel (Orange, k=3, Blue k=2)




Kanto 2006 ReducedGAModel (Orange, k=3, Blue k=2)







Kanto 2009 GAModel (Orange, k=3, Blue k=2)




Kanto 2009 ReducedGAModel (Orange, k=3, Blue k=2)




Kanto 2010 GAModel (Orange, k=3, Blue k=2)




Kanto 2010 ReducedGAModel (Orange, k=3, Blue k=2)




Kansai 2005 GAModel (Orange, k=3, Blue k=2)




Kansai 2005 ReducedGAModel (Orange, k=3, Blue k=2)




Kansai 2006 GAModel (Orange, k=3, Blue k=2)




Kansai 2006 ReducedGAModel (Orange, k=3, Blue k=2)




Kansai 2007 GAModel (Orange, k=3, Blue k=2)




Kansai 2007 ReducedGAModel (Orange, k=3, Blue k=2)




Kansai 2008 GAModel (Orange, k=3, Blue k=2)




Kansai 2008 ReducedGAModel (Orange, k=3, Blue k=2)




Kansai 2009 GAModel (Orange, k=3, Blue k=2)




Kansai 2009 ReducedGAModel (Orange, k=3, Blue k=2)




Kansai 2010 GAModel (Orange, k=3, Blue k=2)




Kansai 2010 ReducedGAModel (Orange, k=3, Blue k=2)




Tohoku 2005 GAModel (Orange, k=3, Blue k=2)




Tohoku 2005 ReducedGAModel (Orange, k=3, Blue k=2)




Tohoku 2006 GAModel (Orange, k=3, Blue k=2)




Tohoku 2006 ReducedGAModel (Orange, k=3, Blue k=2)




Tohoku 2007 GAModel (Orange, k=3, Blue k=2)




Tohoku 2007 ReducedGAModel (Orange, k=3, Blue k=2)




Tohoku 2008 GAModel (Orange, k=3, Blue k=2)




Tohoku 2008 ReducedGAModel (Orange, k=3, Blue k=2)




Tohoku 2009 GAModel (Orange, k=3, Blue k=2)




Tohoku 2009 ReducedGAModel (Orange, k=3, Blue k=2)











Tohoku 2010 GAModel (Orange, k=3, Blue k=2)







Tohoku 2010 ReducedGAModel (Orange, k=3, Blue k=2)









ANOVA test and HSD Tukey


All regions

```
resultANOVA = aov(loglikeValues~model+years+regions, data = data)
summary(resultANOVA)
```


```
##           Df      Sum Sq  Mean Sq  F value  Pr(>F)
## model      1       3409     3409     1.972 0.16090
## years      5      26894     5379     3.111 0.00896 **
## regions    3 216542658 72180886 41751.046 < 2e-16 ***
## Residuals 470     812555     1729
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

```
tuk = TukeyHSD(resultANOVA)
op <- par(mar = c(5,14,4,2)+0.1)
plot(tuk,las=0)
```


95% family-wise confidence level



95% family-wise confidence level



95% family-wise confidence level



KANTO

```
subTabela = data[data$regions=="Kanto",]
print("In Kanto")


## [1] "In Kanto"

resultANOVA = aov(loglikeValues~model+years, data = subTabela)
summary(resultANOVA)


##              Df Sum Sq Mean Sq F value    Pr(>F)
## model          1   4397    4397   12.85 0.000499 ***
## years          5 139054   27811   81.29 < 2e-16 ***
## Residuals     113   38661     342
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

tuk = TukeyHSD(resultANOVA)
op <- par(mar = c(5,14,4,2)+0.1)
plot(tuk,las=0)
```

95% family-wise confidence level



95% family-wise confidence level



EASTJAPAN

```
subTabela2 = data[data$regions=="EastJapan",]
print("In EastJapan")


## [1] "In EastJapan"

resultANOVA = aov(loglikeValues~model+years, data = subTabela2)
summary(resultANOVA)

##           Df Sum Sq Mean Sq F value    Pr(>F)
## model         1  10522    10522     2.91 0.090773 .
## years         5   97986    19597     5.42 0.000166 ***
## Residuals    113 408578      3616
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1


tuk = TukeyHSD(resultANOVA)
op <- par(mar = c(5,14,4,2)+0.1)
plot(tuk,las=0)
```


95% family-wise confidence level



Differences in mean levels of model

95% family-wise confidence level



Differences in mean levels of years

```
subTabela3 = data[data$regions=="Kansai",]
print("In Kansai")
```

```
## [1] "In Kansai"
```


```
resultANOVA = aov(loglikeValues~model+years, data = subTabela3)
summary(resultANOVA)
```

```
##           Df Sum Sq Mean Sq  F value Pr(>F)
## model      1     15      15      1.617  0.206
## years      5  89623  17925  1877.966 <2e-16 ***
## Residuals 113   1079      10
## ---
```


```
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

```
tuk = TukeyHSD(resultANOVA)
op <- par(mar = c(5,14,4,2)+0.1)
plot(tuk,las=0)
```

95% family-wise confidence level



95% family-wise confidence level



TOHOKU

```
subTabela3 = data[data$regions=="Tohoku",]
print("In Tohoku")


## [1] "In Tohoku"

resultANOVA = aov(loglikeValues~model+years, data = subTabela3)
summary(resultANOVA)

##           Df Sum Sq Mean Sq F value    Pr(>F)
## model      1   2321    2321   39.02 7.59e-09 ***
## years      5  43900    8780  147.60 < 2e-16 ***
## Residuals 113   6722     59
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

tuk = TukeyHSD(resultANOVA)
op <- par(mar = c(5,14,4,2)+0.1)
plot(tuk,las=0)
```

95% family-wise confidence level



95% family-wise confidence level

