

Carbohydrates: Structure, Reactivity, and Enzymatic Mechanisms in Biochemical Systems

Wei Qi Zhang

August 29, 2026

Abstract

Biochemistry provides the mechanistic foundation for cellular structure and function. The first paper of the biochemistry trilogy established the chemical logic of fatty acids through esterification and transesterification mechanisms, emphasizing how simple organic reactions generate biologically essential lipids. This second paper extends that mechanistic framework to carbohydrates, presenting the structural and reactive foundations of monosaccharides, followed by a detailed study of carbohydrate enzymology and glycation chemistry through concrete reaction mechanisms. An appendix then introduces nucleic-acid chemistry, illustrating ribonucleotide reduction and $\text{dNTP} \longrightarrow \text{dNTP}$ phosphorylation as worked examples and concluding with the Bloomfield-Alberty theorem. Together, these elements provide the conceptual bridge to the third paper, which will examine generic structure, nucleic-acid reactivity, and the biochemical mechanisms underlying medication-genome interactions.

1 Introduction

Biochemical systems rely on a small set of molecular classes: lipids, carbohydrates, nucleic acids, and proteins. Their reactivity defines the architecture and function of living cells. The first paper of this trilogy established the mechanistic foundations of fatty-acid chemistry, emphasizing esterification and transesterification as the core transformations that generate and remodel lipid structures. These reactions illustrated how simple organic mechanisms, when embedded in biological contexts, give rise to complex macromolecular assemblies and dynamic metabolic pathways.

This second paper extends that mechanistic framework to carbohydrates, a class of biomolecules whose structural diversity and reactivity underpin energy metabolism, cellular recognition, and enzymatic analysis. Carbohydrates exhibit a rich interplay between stereochemistry, conformational dynamics, and electronic effects, making them uniquely suited for mechanistic analysis. Their chemistry is governed by the behavior of the anomeric center, oxocarbenium ion character, and the dissociative pathways. As such, carbohydrate reactions

provide a natural platform for examining S_N1 , S_N2 , E1 and E2 mechanisms in a biologically relevant context.

Beyond intrinsic reactivity, carbohydrate enzymology reveals how biological catalysts manipulate these mechanisms. Building on Sinnott’s framework of intrinsic binding energy and transition-state stabilization, this paper analyzes how glycosidases, transferases, and related enzymes achieve rate acceleration and stereochemical control. These enzymatic pathways are contrasted with non-enzymatic glycation chemistry, where reducing sugars undergo Schiff-base formation, Amadori rearrangements, and enediol-mediated transformations that contribute to cellular aging and metabolic pathology.


To support the conceptual bridge toward the third paper in the trilogy, an appendix provides a concise introduction to nucleic-acid chemistry, including worked examples such as ribonucleotide reductase (RNR) and the phosphorylation of dNDPs to dNTPs. These mechanisms illustrate how the same principles of stereoelectronics, catalysis, and transition-state theory extend from carbohydrate chemistry into nucleotide metabolism. The appendix concludes with the Bloomfield-Alberty theorem, whose mathematical treatment reinforces the energetic and mechanistic concepts developed throughout this paper. Together, these elements position carbohydrates as a central mechanistic pillar of biochemistry, linking the lipid-chemistry of the first paper to the nucleic-acid reactivity and genomic mechanisms that will be explored in the third paper.

2 Reaction Mechanisms in Carbohydrate Chemistry: S_N1 , S_N2 , E1 and E2

Carbohydrate reactivity is governed by the interplay between stereochemistry, conformational dynamics, and the electronic structure of the anomeric center. Although classical organic mechanisms - S_N1 , S_N2 , E1 and E2 - remain the conceptual backbone, their behavior in carbohydrate systems differs substantially from simple alkyl substrates. This section establishes the mechanistic foundations by first recalling the canonical reactions in non-carbohydrate systems, then extending these principles to monosaccharides and glycosidic chemistry.

2.1 Canonical S_N2 Reactivity: A Reference Mechanism

Before analyzing carbohydrate S_N2 constraints it is useful to anchor the discussion with a standard alkyl-halide substitution. Consider the reaction:



This transformation proceeds through a concerted S_N2 mechanism, characterized by:

- **Backside attack** - nucleophile approaches opposite the C–Br bond.
- **Walden inversion** - stereochemical inversion at the reacting carbon.
- **Second order kinetics** - rate $\propto$ [substrate]*[nucleophile].
- **Transition-state geometry** - pentacoordinate carbon with partial bonds.

This reaction is fast because the reacting carbon is primary-like (sterically accessible) and the leaving group (Br^-) is excellent. The nucleophile (NH_2^-) is strong, small, and unhindered.

This example provides the mechanistic baseline against which carbohydrate S_N2 behavior can be contrasted.

2.2 S_N2 Reactivity at the Anomeric Center

Carbohydrates rarely undergo classical S_N2 reactions at the anomeric carbon because:

- **Steric congestion** - ring structure restricts backside attack.
- **Orbital alignment constraints** - axial/equatorial orientation affects accessibility.
- **Poor leaving groups** - glycosidic oxygen is not a halide.
- **Solvent effects** - aqueous environments disfavor strong nucleophiles.

As a result, true S_N2 displacement at $C1$ is rare, except in specially activated systems (e.g., glycosyl fluorides, triflates). Even then, such reactions are highly stereospecific and requires precise orbital alignment.

2.3 S_N1 Reactivity and Oxocarbenium-Ion Character

Most biologically relevant carbohydrate substitutions proceed through S_N1 -like pathways involving an oxocarbenium ion or a hemiacetal-derived cationic transition state.

- **Oxocarbenium stabilization** - resonance between $C1$ and ring oxygen.
- **Dissociative mechanism** - leaving group departure precedes nucleophilic attack.
- **Stereochemical scrambling** - mixture of α/β products unless constrained.
- **Solvent stabilization** - water stabilizes cationic intermediates.

The S_N1 -like pathway is central to glycosylation chemistry, enzymatic catalysis, and glycation reactions.

2.4 E1 and E2 Pathways in Carbohydrate Chemistry

Elimination reactions arise in carbohydrates typically arise from:

- **Dehydration** - acid-catalyzed loss of water from hemiacetals.
- **Enolization** - formation of enediols during glycation.
- **β -elimination** - base-induced elimination in activated systems.

E1 pathways often accompany S_N1 -like mechanisms, while *E2* pathways require strong bases and activated leaving groups. These conditions are uncommon in biological environments but relevant in synthetic carbohydrate chemistry.

2.5 Mechanistic Consequences for Biological Systems

The constraints described above lead to several important biochemical consequences:

- **Enzymes favor dissociative pathways** - glycosidases stabilize oxocarbenium-like transition states.
- **Neighboring-group participation** - 2-OH or acyl groups assist leaving-group departure.
- **Stereochemical control** - retaining vs inverting mechanisms.
- **Glycation chemistry parallels $S_N1/E1$** - open-chain aldehyde reacts through cationic intermediates.

3 Carbohydrate Enzymology: Intrinsic Binding Energy, Transition-State Stabilization, and Catalytic Strategy

Carbohydrate-active enzymes operate in a mechanistic landscape dominated by stereoelectronic constraints, oxocarbenium-ion character, and the need for precise control over reaction pathways. Sinnott's framework provides the most coherent theoretical foundation for understanding these systems: enzymes accelerate reactions not primarily by stabilizing substrates, but by selectively stabilizing the transition state and use intrinsic binding energy to enforce the geometry required for catalysis.

This section develops the mechanistic logic of carbohydrate enzymology through Sinnott's principles, focusing on glycosidases, glycosyltransferases, and related catalytic systems.

3.1 Intrinsic Binding Energy and the Enforcement of Catalytic Geometry

Sinnott's central argument is that enzymes use intrinsic binding energy - the portion of binding energy that cannot be attributed to ground-state stabilization - to distort substrates into conformations resembling the transition state.

- **Intrinsic binding energy** - energy available only when the substrate is in the catalytic pose.
- **Transition-state enforcement** - enzyme active sites pre-organize the substrate.
- **Ground-state destabilization** - substrate is forced into a higher-energy conformation.
- **Stereoelectronic alignment** - orbitals are oriented for oxocarbenium-like transition state formation.

In carbohydrate chemistry, the relevant transition state resembles an oxocarbenium ion, with partial positive charge at $C1$ and a flattened, sp^2 -like geometry. Enzymes bind substrates in conformations that mimic this geometry, lowering the activation barrier for glycosidic bond cleavage or formation.

3.2 Dissociative S_N1 -like Pathways in Glycosidases

Most glycosidases operate through dissociative S_N1 -like mechanisms rather than classical S_N2 . The leaving group departs first, generating an oxocarbenium-like transition state stabilized by the enzyme.

- **Oxocarbenium-ion stabilization** - resonance between $C1$ and ring oxygen.
- **General acid catalysis** - protonation of the leaving group.
- **Electronic stabilization** - active-site residues stabilize positive charge.
- **Conformational selection** - enzyme selects a half-chair or boat conformation.

The dissociative pathway is favored because:

- backside attack is sterically impossible at $C1$.
- glycosidic oxygen is a poor leaving group without protonation.
- aqueous environments stabilize cationic intermediates.

Such mechanistic preference explains the divergence of carbohydrate chemistry from classical alkyl-halide S_N2 behavior.

3.3 Retaining and Inverting Glycosidases

Glycosidases fall into two mechanistic classes:

3.3.1 Retaining Glycosidases

They proceed through a **double-displacement mechanism**:

- Covalent glycosyl-enzyme intermediate.
- General acid-base catalysis.
- Two sequential S_N1 -like steps.

Stereochemistry at $C1$ is retained because the intermediate is attacked from the same face twice.

3.3.2 Inverting Glycosidases

They proceed through a **single-displacement mechanism** involving:

- One general-acid and one general-base.
- Direct nucleophilic attack on oxocarbenium-like transition state.
- Stereochemical inversion.

Although often described as " S_N2 -like", the mechanism is not classical S_N2 ; it is a **concerted but dissociative** pathway with oxocarbenium character.

3.4 Neighboring-Group Participation (NGP) in Carbohydrate Enzymology

Carbohydrates possess multiple hydroxyl groups capable of neighboring-group participation, dramatically altering reaction pathways.

- **2-OH participation** - formation of a transient bicyclic intermediate.
- **Acyl-group participation** - anchimeric assistance from protecting groups.
- **Rate acceleration** - stabilization of cationic transition state.
- **Stereochemical control** - NGP enforces specific attack trajectories.

NGP is one of the most powerful stereoelectronic effects in carbohydrate chemistry and is widely exploited in both enzymology and synthetic chemistry.

3.5 Associative Pathways in Glycosyltransferases

While glycosidases favor dissociative pathways, glycosyltransferases often operate through associative S_N2 -like mechanisms due to activated leaving groups such as UDP, GDP or CMP.

- **Activated leaving groups** - nucleotide diphosphates.

- **Concerted attack** - nucleophile approaches as leaving group departs.
- **Stereospecificity** - inversion at $C1$.
- **Metal-ion interactions** - Mg^{2+} stabilizes negative charge.

These pathways resemble classical S_N2 more closely than glycosidase mechanisms, but still exhibit oxocarbenium-like transition-state character.

3.6 Enzymatic Enforcement of Oxocarbenium-Like Transition States

Across all carbohydrate-active enzymes, the unifying mechanistic theme is transition-state enforcement. Enzymes stabilize:

- **Flattened $C1$ geometry** - partial sp^2 character.
- **Positive charge delocalization** - resonance with ring oxygen.
- **Conformational distortion** - half-chair, boat or envelope conformations.
- **Leaving-group activation** - protonation or coordination.

Such stabilization is the mechanistic core carbohydrate enzymology and key to glycation chemistry.

4 Glycation Chemistry: Non-Enzymatic Reactivity of Reducing Sugars

Glycation chemistry represents the non-enzymatic counterpart to the carbohydrate mechanisms and enzymatic strategies developed in previous sections of this paper. While enzymatic glycosidic reactions rely on transition-state enforcement and precise stereoelectronic control, glycation proceeds through open-chain aldehyde chemistry, imine formation, rearrangements, and oxidative pathways that collectively generate a diverse set of products known as **advanced glycation end-products** (AGEs).

This section develops the mechanistic foundations of glycation, beginning with reducing-sugar reactivity and progressing through Schiff-base formation, Amadori rearrangement, enediol chemistry, and downstream oxidative pathways.

4.1 Reducing Sugars and Open-Chain Aldehyde Reactivity

Carbohydrates exist in equilibrium between cyclic hemiacetal forms and a small but chemically significant open-chain aldehyde form. This aldehyde fraction is the mechanistic origin of glycation.

- **Reducing sugars** - monosaccharides capable of adopting an open-chain aldehyde form.
- **Aldehyde electrophilicity** - carbonyl carbon susceptible to nucleophilic attack.
- **Ring-chain tautomerization** - equilibrium provides a continuous supply of reactive aldehyde.
- **Biological relevance** - glucose, ribose, and fructose participate readily.

Even though the open-chain form is typically ~ 0.01 , its reactivity dominates glycation chemistry.

4.2 Schiff-Base Formation: The Initial Glycation Step

The first step in glycation is nucleophilic attack by an amine (typically the ϵ -amine of lysine or the N-terminus of a protein) on the aldehyde carbonyl of a reducing sugar.

- **Schiff base formation** - nucleophilic attack $\rightarrow$ carbinolamine $\rightarrow$ dehydration $\rightarrow$ imine.
- **Reversibility** - imine formation is reversible and pH-dependent.
- **Protein modification** - forms early glycation adducts.
- **Stereoelectronic considerations** - imine formation alters local electronic structure.

The Schiff base is not the final product; it is a gateway intermediate that undergoes rearrangement.

4.3 Amadori Rearrangement: 1,2-Enolization and Ketosamine Formation

The Schiff base undergoes a 1,2-enolization that converts the aldimine into a more stable ketoamine, known as the Amadori product.

- **Amadori rearrangement** - proton transfer $\rightarrow$ enolization $\rightarrow$ rearrangement.
- **Ketosamine stability** - more stable than the initial imine.
- **Biological significance** - precursor to AGEs.
- **Mechanistic analogy** - rearrangement resembles cationic shifts seen in S_N1 chemistry.

The Amadori product is the central branching point of glycation pathways.

4.4 Enediol Chemistry: Oxidation, Fragmentation, and Reactive Intermediates

Further transformations involve enediol formation, a key mechanistic step that opens oxidative and fragmentation pathways.

- **Enediol formation** - tautomerization of the ketoamine.
- **Oxidation Pathways** - formation of reactive dicarbonyls (glyoxal, methylglyoxal)
- **Fragmentation** - cleavage of carbon skeleton.
- **Radical Chemistry** - oxidative stress accelerates enediol reactions.

Enediols are highly reactive, enabling formation of multiple downstream products.

4.5 Advanced Glycation End-Products (AGEs)

Advanced glycation end-products arise from oxidative, condensative, and cross-linking reactions involving Amadori products, enediols, and reactive dicarbonyl intermediates. In contrast to the previous glycation steps, AGE formation is irreversible, structurally diverse, and mechanistically rich.

- **AGE formation** - condensation of dicarbonyls with lysine, arginine, and cysteine residues.
- **Protein cross-linking** - formation of intermolecular bridges (e.g., pentosidine, crossline).
- **Structural modification** - altered protein folding, increased rigidity, denaturation.
- **Oxidative pathways** - reactive oxygen species accelerate dicarbonyl formation.
- **Pathophysiological relevance** - aging, diabetes, neurodegeneration, vascular damage.

AGE chemistry is driven by reactive dicarbonyls such as glyoxal, methylglyoxal, and 3-deoxyglucosone, which arise from enediol oxidation and fragmentation. These electrophiles react rapidly with nucleophilic amino-acid side chains, generating stable adducts and cross-links that accumulate over time.

Mechanistically, AGE formation represents the non-enzymatic extreme of carbohydrate reactivity: uncontrolled, multi-pathway, and structurally heterogeneous.

4.6 Mechanistic Parallels Between Glycation and Enzymatic Carbohydrate Chemistry

Although glycation is non-enzymatic, its mechanisms mirror the enzymatic pathways described in the previous section.

- **Oxocarbenium-like intermediates** - cationic transition states appear in both enzymatic and non-enzymatic reactions.
- **General acid-base catalysis** - pH dependent imine formation parallels enzymatic proton transfers.
- **Conformational effects** - ring opening and tautomerization resemble enzyme-induced distortions.
- **Stereoelectronic constraints** - aldehyde reactivity and enediol formation follow the same orbital logic.
- **Dissociative pathways** - glycation rearrangements echo S_N1 -like carbohydrate chemistry.

These parallels reinforce the conceptual unity of carbohydrate reactivity: whether controlled by enzymes or occurring spontaneously, the underlying chemistry is governed by the same stereoelectronic principles.

5 Conclusion

This second paper developed the mechanistic foundations of carbohydrate reactivity by integrating classical organic reaction pathways, stereoelectronic constraints, and enzymatic transition-state stabilization. Beginning with the $S_N1/S_N2/E1/E2$ framework, we showed that carbohydrate chemistry is governed by steric crowding at the anomeric center, restricted approach trajectories, and the inherent tendency of cyclic sugars to form oxocarbenium-like intermediates. These constraints explain why dissociative pathways dominate and why enzymatic catalysis must enforce distorted ring conformations to achieve rate acceleration.

Building on this foundation, the paper examined Sinnott’s intrinsic binding-energy model, highlighting how glycosidases use precise electrostatic and conformational control to stabilize transition states that resemble oxocarbenium ions. This enzymatic logic provides a unifying explanation for the stereochemical outcomes and rate enhancements observed across retaining and inverting mechanisms.

The final section extended these principles to non-enzymatic glycation chemistry. Reducing-sugar aldehyde reactivity, Schiff-base formation, Amadori rearrangement, enediol chemistry, and AGE formation were presented as spontaneous analogues of the same stereoelectronic forces that govern enzymatic pathways. Although uncontrolled and structurally heterogeneous, glycation reactions follow the same underlying orbital and conformational logic as enzyme-mediated

carbohydrate transformations.

Together, these results establish a coherent mechanistic picture of carbohydrate reactivity across both enzymatic and non-enzymatic contexts. The appendix provides a brief bridge to nucleic-acid chemistry, setting the stage for the subsequent paper, where the mechanistic principles developed here will be extended to the structure, reactivity, and biosynthesis of nucleotides and nucleic acids.

The purpose of this appendix is to establish the mechanistic continuity between the carbohydrate chemistry developed in Paper 2 and the nucleotide chemistry that will form the core of Paper 3. Although nucleotides differ structurally from simple carbohydrates, their reactivity is governed by the same stereoelectronic principles: constrained sugar conformations, anomeric effects, oxocarbenium-like transition states, and enzyme-enforced orbital alignment. To illustrate these parallels, the appendix presents three mechanistic components of nucleotide biosynthesis: Class II ribonucleotide reduction, phosphoryl-transfer chemistry in dNDP $\rightarrow$ dNTP conversion, and the Bloomfield–Alberty catalytic framework used to analyze multi-intermediate enzymatic pathways.

- **Pentose stereoelectronics** - ribose/deoxyribose reactivity parallels carbohydrate mechanisms.
- **Radical sugar chemistry** - Class II RNR manipulates ribose using controlled radical intermediates.
- **Phosphoryl transfer** - associative/dissociative pathways governed by metal-ion catalysis.
- **catalytic thermodynamics** - Bloomfield–Alberty formalism connects to Sinnott’s intrinsic binding-energy model.

These topics provide the mechanistic bridge from this paper to the subsequent paper, allowing the trilogy to transition from carbohydrate reactivity to the biosynthesis and chemical logic of nucleic acids.

A Ribonucleotide Reductase (Class II mechanism)

Class II ribonucleotide reductases use **adenosylcobalamin (AdoCbl)** to generate the 5-deoxyadenosyl radical that initiates sugar-ring chemistry. Unlike class I and III, class II enzymes are oxygen and metal independent and self-contained, making them ideal for illustrating the core mechanistic logic of ribonucleotide reduction without additional structural detail. Mechanism:

1. **AdoCbl homolysis** - $\text{AdoCbl} \longrightarrow \text{Co(II)} + \text{5-deoxyadenosyl radical}$ catalyzed by enzyme.
2. **Hydrogen abstraction** - 5-dAdo abstracts H from C3 and producing a C3-radical on ribose.

3. **Formation of 5-dADOH** - 5-dAdo becomes 5-deoxyadenosine (5-dAdoH).
4. **activation and departure of OH** - radical rearrangement activates hydroxyl.
5. **Protonation of hydroxyl** - cysteine protonates OH $\rightarrow$ H₂O forming carbocation intermediate.
6. **Return of radical to AdoCbl** - ribose radical abstracts H from 5-dAdoH, regenerating 5-Ado, recombines with Co(II) forming AdoCbl.

Mechanistically, class II RNR demonstrates how enzymes manipulate ribose stereochemistry using radical hydrogen abstraction, C3 radical formation, 2-OH elimination and radical return to regenerate AdoCbl. This radical-based strategy parallels the stereoelectronic constraints discussed in this paper: conformational control, transition-state stabilization, and precise orbital alignment.

B dNDP $\longrightarrow$ dNTP Phosphorylation


Mechanism:

1. **Phosphate transfer** - non-substrate-specific transfer where NDPK works for all nucleoside diphosphates.
2. **Ping-Pong mechanism with phospho-histidine intermediate** - ATP donates a phosphate to a specific histidine residue on NDPK.
3. **Formation of phosphate linkage with histidine** - phospho-His with dNDP binding and phospho-his transfers phosphate yielding dNTP.
4. **Histidine dephosphorylation** - Histidine returns to its non-phosphorylated form.
5. **RNR-controlled reaction** - Reaction driven by ATP to ADP transfer with RNR ratio controls.

Phosphoryl-transfer chemistry is governed by associative / dissociative pathways, metal-ion catalysis, and transition-state stabilization. These principles mirror the stereoelectronic constraints observed in carbohydrate enzymology.

C Mathematical Proof of Bloomfield-Alberty Framework

Consider a general bi-bi reaction:



with enzyme complexes: $E, EA, EB, EAB, EP, EQ, EPQ$ written in the form valid for any number of enzyme complexes under steady-state conditions.

Assume for every intermediate X :

$$\frac{d[X]}{dt} \approx 0$$

yielding a system of linear equations relating the concentrations of all enzyme species that is collapsible into single expression:

$$v = \frac{V_{max}[A][B]}{D}$$

where D is a denominator containing all binding terms.

The denominator must have the form:

$$D = \sum_i c_i \Pi_j [X_j]$$

where:

- each c_i is an observable kinetic parameter;
- each term is a product of reactant / product concentrations;
- the structure is independent of the number of intermediates

For a simple ordered bi-bi mechanism, the denominator reduces to:

$$D = K_A K_B + K_B[A] + K_A[B] + [A][B]$$

Thus,

$$v = \frac{V_{max}[A][B]}{K_A K_B + K_B[A] + K_A[B] + [A][B]}$$

yielding the Bloomfield-Alberty bi-bi rate equation.

The Bloomfield-Alberty treatment provides the thermodynamic and kinetic foundation for analyzing multi-intermediate enzymatic pathways. Its emphasis on intrinsic binding energy and transition-state stabilization directly complements Sinnott's model from section 2.

D Summary

Together, these derivations establish the mechanistic continuity between carbohydrate chemistry and nucleotide biosynthesis. They form the conceptual bridge into the next paper, where nucleic-acid structure, reactivity, and enzymatic pathways will be developed in full detail.

References

- [1] SINNOTT, Michael L.; *Carbohydrate Chemistry and Biochemistry: Structure and Mechanism*; RSC Publishing (Cambridge); 2007; pp.299-356, 361-387, 408-423, 478-492.
- [2] FERSHT, Alan; *Enzyme Structure and Mechanism*; W.H. Freeman (New York); 1985; pp.47-120.
- [3] BLACKBURN, Michael G., GAIT, Michael J., LOACKES, David, WILLIAMS, David M.; *Nucleic Acids in Chemistry and Biology*; 3rd ed.; RSC Publishing; 2006; pp.13-136, 209-379.
- [4] BRUCE, Paula Y.; *Organic Chemistry*; 2nd ed.; Pearson ERPI; 2012; pp.520-549.

Additional References

- [1] ZHANG, Wei Qi; 2026. *Biochemistry Enzyme Catalysis*. notes on enzyme catalysis in biochemistry.
- [2] ZHANG, Wei Qi; 2026. *Biochemistry Nucleic Acid Notes*. notes on nucleic-acid biochemistry.
- [3] ZHANG, Wei Qi; 2026. *Biochemistry Glycation Chemistry Notes*. notes on glycation biochemistry.