


Chapter 20 - Pathogenic Gram-Negative Cocci and Bacilli

NIMESH PATEL | HLSC 2400

OCTOBER 26, 2017

MICROBIOLOGY

5th Edition



WITH
DISEASES BY
TAXONOMY

ROBERT W. BAUMAN

PowerPoint® Lecture
Presentations prepared by
Mindy Miller-Kittrell,
North Carolina State
University

CHAPTER 20

Pathogenic Gram- Negative Cocci and Bacilli

Pathogenic, Gram-Negative, Aerobic Bacilli

- ***Bartonella***

- Aerobic bacilli
- Found in animals but causes disease only in humans
- Three species are pathogenic:
 - *Bartonella bacilliformis*
 - **Bartonellosis**
 - Causes fever, anemia, headache, muscle and joint pain, and chronic skin infections
 - Often fatal
 - Transmitted by bloodsucking sand flies
 - Endemic in Peru, Ecuador, and Columbia

Pathogenic, Gram-Negative, Aerobic Bacilli

- ***Bartonella***

- *Bartonella quintana*

- Trench fever (**five-day fever**)
 - Common during World War I
 - Transmitted by human body lice
 - Causes headaches, fever, and pain in the long bones
 - Causes two diseases in immunocompromised patients
 - Bacillary angiomatosis
 - Fever, skin lesions, and proliferating blood vessels
 - Bacillary peliosis hepatis
 - Blood-filled pockets in the liver

Pathogenic, Gram-Negative, Aerobic Bacilli

- ***Bartonella***

- *Bartonella henselae*

- **Cat-scratch disease**

- Transmitted through cat scratches and bites
 - Common disease in children in the United States
 - Causes fever, malaise, and swelling at infection site

- *Bartonella* infections are treated with various antimicrobials

Figure 20.21 Cat scratch disease.



Pathogenic, Gram-Negative, Aerobic Bacilli

- ***Brucella***


- Small, nonmotile, aerobic coccobacilli
- Can infect animals or humans
- *Brucella melitensis* causes **brucellosis** (**Bang's disease and undulant fever**)
 - Often an asymptomatic or mild disease
 - Illness is characterized by a fluctuating fever
 - Human infection due to contact with contaminated dairy products or infected animal blood, urine, or placentas
 - Vaccine available for use in domesticated animals
 - Not used in humans since it can cause disease.

Pathogenic, Gram-Negative, Aerobic Bacilli

- ***Bordetella***

- Pathogenesis, Epidemiology, and Disease
 - Small, aerobic, nonmotile coccobacillus
 - ***B. pertussis* is the most important**
 - Causes pertussis (whooping cough)
 - Most cases of disease are in children - ciliated epithelial cells of the trachea are damaged
 - Adhesins and toxins mediate the disease
 - *Pertussis toxin*
 - *Adenylate cyclase toxin*
 - *Dermonecrotic toxin*
 - *Tracheal cytotoxin*
 - Bacteria inhaled in aerosols multiply in epithelial cells

Figure 20.25 The approximate time course for the progression of pertussis.



Pathogenic, Gram-Negative, Aerobic Bacilli

- ***Bordetella***

- Diagnosis, Treatment, and Prevention
 - Diagnosis
 - Pertussis symptoms usually diagnostic
 - Treatment
 - Primarily supportive
 - Prevention
 - Immunization with diphtheria, tetanus, attenuated pertussis (DTaP) or Tdap vaccine
 - Tdap - a reduced dose of the diphtheria and pertussis vaccines

Pathogenic, Gram-Negative, Aerobic Bacilli

- ***Burkholderia***

- *Burkholderia cepacia*

- Aerobic, flagellated betaproteobacterium
 - Can decompose a broad range of organic molecules
 - Assists in cleanup of contaminated environmental sites
 - Used by farmers to reduce fungal infection of plant crops
 - Opportunistic pathogen of cystic fibrosis patients
 - Resistant to many antimicrobial drugs

- *Burkholderia pseudomallei*

- Causes melioidosis (Whitmore's disease), an Asian and Australian tropical disease that is emerging as a threat in other locales
 - CDC has identified as a potential bioweapon

Pathogenic, Gram-Negative, Aerobic Bacilli

- ***Burkholderia: Bioterrorism***

Why might germs that cause melioidosis be used as a weapon?

- They are found in nature in certain parts of the world (<https://www.cdc.gov/melioidosis/risk.html>)
- They cause a disease that can make people very sick. Without prompt treatment with specific antibiotics, people sick with melioidosis can die
- In the past, some countries that have used bioweapons during war have used germs closely related to the one that causes melioidosis

Source: <https://www.cdc.gov/melioidosis/bioterrorism/threat.html>

Pathogenic, Gram-Negative, Aerobic Bacilli

- ***Burkholderia: Bioterrorism***

How dangerous is melioidosis?

- Naturally resistant to many commonly used antibiotics
- Without treatment, up to 9 out of every 10 people who get it die.
- When people with melioidosis get treatment with the correct antibiotics, fewer than 4 out of 10 people die.
- Medical treatment in an intensive care facility can decrease deaths even more. Only 2 out of 10 people die when they get this level of medical treatment.

Source: <https://www.cdc.gov/melioidosis/bioterrorism/threat.html>

Pathogenic, Gram-Negative, Aerobic Bacilli

- **Pseudomonads**

- Aerobic bacilli
- Ubiquitous in soil, decaying organic matter, and moist environments
- **Problematic in hospitals**
- Opportunistic pathogens

Pathogenic, Gram-Negative, Aerobic Bacilli

- **Pseudomonads**

- *Pseudomonas aeruginosa*
 - Rarely part of normal human microbiota
 - Rarely causes disease in healthy individuals
 - Despite producing various virulence factors
 - Fimbriae, adhesins, capsule, toxins, and enzymes
 - Opportunistic infections in immunocompromised patients
 - Can colonize almost any organ or system
 - Also infects the lungs of cystic fibrosis patients
 - Biofilm protects bacteria from phagocytosis
 - Treatment is difficult due to drug resistance

Figure 20.26 A *Pseudomonas aeruginosa* infection.



Pathogenic, Gram-Negative, Aerobic Bacilli

- **Pseudomonads**

- *Moraxella* and *Acinetobacter*

- Aerobic, short, plump bacilli
 - Rarely cause disease in humans

- ***Moraxella catarrhalis***

- Opportunistic infections of the sinuses, bronchi, ears, and lungs
 - Susceptible to most antibacterial drugs

- ***Acinetobacter***

- Grows in soil, water, and sewage
 - Opportunistic infections of the respiratory, urinary, and central nervous systems
 - Often resistant to antibiotics

Pathogenic, Gram-Negative, Aerobic Bacilli

- ***Francisella***

- *Francisella tularensis*

- Nonmotile, strictly aerobic coccobacillus
 - Intracellular parasite of animals and amoebae in water
 - Diverse range of hosts
 - Reservoirs in the United States include rabbits, muskrats, and ticks

Pathogenic, Gram-Negative, Aerobic Bacilli

- ***Francisella***

- *Francisella tularensis*

- **Tularemia** (*rabbit fever* or *tick fever*)

- Zoonotic disease
 - Spread through bite of an infected tick or contact with an infected animal
 - Also spread by bacteria in meat, water, and in aerosols
 - The bacteria is highly infectious
 - Tularemia may be misdiagnosed
 - Symptoms are similar to other diseases
 - Treated with antimicrobial drugs
 - Infection prevented by avoiding reservoirs of *Francisella*

Pathogenic, Gram-Negative, Aerobic Bacilli

- ***Legionella***
 - Pathogenesis, Epidemiology, and Disease
 - Aerobic, slender, pleomorphic bacteria
 - Universal inhabitants of water
 - **Humans inhale bacteria in aerosols from water sources**
 - Intracellular parasites
 - *L. pneumophila* causes most disease in humans

Pathogenic, Gram-Negative, Aerobic Bacilli

- ***Legionella***

- Pathogenesis, Epidemiology, and Disease
 - Legionnaires' disease (*Legionellosis*)
 - *Causative agent: Legionella pneumophila*
 - Transmitted in aerosols such as those produced by air-conditioning systems, vaporizers, and grocery store misters
 - Results in pneumonia
 - Can be fatal
 - Particularly immunocompromised individuals
 - Pontiac fever
 - Similar to Legionnaires' disease
 - Doesn't produce pneumonia and is not fatal

Pathogenic, Gram-Negative, Aerobic Bacilli

- ***Legionella***
 - Diagnosis, Treatment, and Prevention
 - Diagnosis
 - Identification of *Legionella* by antibody staining or other serological tests
 - Treatment
 - Pontiac fever is self-limiting
 - Fluoroquinolone and azithromycin used to treat Legionnaires' disease
 - Prevention
 - Elimination of the bacteria is not feasible

Pathogenic, Gram-Negative, Aerobic Bacilli

- ***Coxiella***

- *Coxiella burnetii*

- Extremely small, aerobic bacteria
 - Obligate intracellular parasite
 - Originally thought to be a virus
 - Infective body enables survival in harsh environment
 - Human disease associated with farm animals and pets
 - Transmitted mainly by inhalation of the infective bodies

Pathogenic, Gram-Negative, Aerobic Bacilli

- ***Coxiella***

- *Coxiella burnetii*

- **Q fever**

- Causes fever, headache, chills, muscle pain, and mild pneumonia
 - Chronic Q fever can cause endocarditis years later
 - Chronic Q fever treated with long-term antimicrobials
 - Vaccine developed but not available in the United States and Canada
 - Prevent by avoiding inhalation of contaminated dust

Pathogenic, Gram-Negative, Anaerobic Bacilli

- Predominant microbiota of the gastrointestinal, urinary, reproductive, and lower respiratory tracts
- Important for human health
 - Inhibit the growth of most pathogens
 - Synthesize vitamins and vitamin precursors
 - Aid in digestion of food
- Cause disease when introduced into other parts of the body

Pathogenic, Gram-Negative, Anaerobic Bacilli

- ***Bacteroides***

- Normal microbiota of the intestinal and upper respiratory tracts
- *Bacteroides fragilis*
 - Most important pathogen of this genus
 - Produces various virulence factors (e.g., lipid A)
 - Involved in a variety of conditions
 - Abdominal infections
 - Genital infections in women
 - Wound infections of the skin
 - Infections can be treated with metronidazole

Pathogenic, Gram-Negative, Anaerobic Bacilli

- ***Prevotella***

- Normal microbiota of urinary, genital, and upper respiratory tracts
- Differ from *Bacteroides* in their sensitivity to bile
- Produce several virulence factors (e.g., capsule)
- Involved in various conditions
 - **Sinus and ear infections**
 - **Almost all periodontal infections**
 - Gynecological infections
 - Brain abscesses
 - Abdominal infections
- Treat by surgical removal of infected tissue and carbapenem