

Section 1.5

Solution Sets of Linear Systems

Today we will learn to describe and draw the solution set of an arbitrary system of linear equations Ax = b, using spans.




Today we will learn to describe and draw the solution set of an arbitrary system of linear equations Ax = b, using spans.



Recall: the **solution set** is the collection of all vectors x such that Ax = b is true.


Today we will learn to describe and draw the solution set of an arbitrary system of linear equations Ax = b, using spans.



Recall: the **solution set** is the collection of all vectors x such that Ax = b is true.

Last time we discussed the set of vectors b for which Ax = b has a solution.

Today we will learn to describe and draw the solution set of an arbitrary system of linear equations Ax = b, using spans.



Recall: the **solution set** is the collection of all vectors x such that Ax = b is true.

Last time we discussed the set of vectors b for which Ax = b has a solution.

We also described this set using spans, but it was a different problem.

Everything is easier when b = 0, so we start with this case.

Everything is easier when b = 0, so we start with this case.

Definition

A system of linear equations of the form Ax = 0 is called **homogeneous**.

Everything is easier when b = 0, so we start with this case.

Definition

A system of linear equations of the form Ax = 0 is called **homogeneous.**

These are linear equations where everything to the right of the = is zero.

Everything is easier when b = 0, so we start with this case.

Definition

A system of linear equations of the form Ax = 0 is called **homogeneous.**

These are linear equations where everything to the right of the = is zero. The opposite is:

Definition

A system of linear equations of the form Ax = b with $b \neq 0$ is called **nonhomogeneous** or **inhomogeneous**.

Everything is easier when b = 0, so we start with this case.

Definition

A system of linear equations of the form Ax = 0 is called **homogeneous.**

These are linear equations where everything to the right of the = is zero. The opposite is:

Definition

A system of linear equations of the form Ax = b with $b \neq 0$ is called **nonhomogeneous** or **inhomogeneous**.

A homogeneous system always has the solution $x = _$.

Everything is easier when b = 0, so we start with this case.

Definition

A system of linear equations of the form Ax = 0 is called **homogeneous.**

These are linear equations where everything to the right of the = is zero. The opposite is:

Definition

A system of linear equations of the form Ax = b with $b \neq 0$ is called **nonhomogeneous** or **inhomogeneous**.

A homogeneous system always has the solution x=0. This is called the **trivial solution**.

Everything is easier when b = 0, so we start with this case.

Definition

A system of linear equations of the form Ax = 0 is called **homogeneous.**

These are linear equations where everything to the right of the = is zero. The opposite is:

Definition

A system of linear equations of the form Ax = b with $b \neq 0$ is called **nonhomogeneous** or **inhomogeneous**.

Everything is easier when b = 0, so we start with this case.


Definition

A system of linear equations of the form Ax = 0 is called **homogeneous.**

These are linear equations where everything to the right of the = is zero. The opposite is:

Definition

A system of linear equations of the form Ax = b with $b \neq 0$ is called **nonhomogeneous** or **inhomogeneous**.



Everything is easier when b = 0, so we start with this case.


Definition

A system of linear equations of the form Ax = 0 is called **homogeneous.**

These are linear equations where everything to the right of the = is zero. The opposite is:

Definition

A system of linear equations of the form Ax = b with $b \neq 0$ is called **nonhomogeneous** or **inhomogeneous**.



Everything is easier when b = 0, so we start with this case.


Definition

A system of linear equations of the form Ax = 0 is called **homogeneous.**

These are linear equations where everything to the right of the = is zero. The opposite is:

Definition

A system of linear equations of the form Ax = b with $b \neq 0$ is called **nonhomogeneous** or **inhomogeneous**.



Question

$$A = \begin{pmatrix} 1 & 3 & 4 \\ 2 & -1 & 2 \\ 1 & 0 & 1 \end{pmatrix}$$
?

Question

What is the solution set of Ax = 0, where

$$A = \begin{pmatrix} 1 & 3 & 4 \\ 2 & -1 & 2 \\ 1 & 0 & 1 \end{pmatrix}$$
?

We know how to do this: first form an augmented matrix and row reduce.

Question

What is the solution set of Ax = 0, where

$$A = \begin{pmatrix} 1 & 3 & 4 \\ 2 & -1 & 2 \\ 1 & 0 & 1 \end{pmatrix}$$
?

We know how to do this: first form an augmented matrix and row reduce.

The only solution is the trivial solution x = 0.

Question

What is the solution set of Ax = 0, where

$$A = \begin{pmatrix} 1 & 3 & 4 \\ 2 & -1 & 2 \\ 1 & 0 & 1 \end{pmatrix}$$
?

We know how to do this: first form an augmented matrix and row reduce.

The only solution is the trivial solution x = 0.

Observation

Since the last column (everything to the right of the =) was zero to begin, it will always stay zero!

Question

What is the solution set of Ax = 0, where

$$A = \begin{pmatrix} 1 & 3 & 4 \\ 2 & -1 & 2 \\ 1 & 0 & 1 \end{pmatrix}$$
?

We know how to do this: first form an augmented matrix and row reduce.

The only solution is the trivial solution x = 0.

Observation

Since the last column (everything to the right of the =) was zero to begin, it will always stay zero! So it's not really necessary to write augmented matrices in the homogeneous case.

Question

$$A = \begin{pmatrix} 1 & -3 \\ 2 & -6 \end{pmatrix}$$
?

Question

What is the solution set of Ax = 0, where

$$A = \begin{pmatrix} 1 & -3 \\ 2 & -6 \end{pmatrix}$$
?

This last equation is called the **parametric vector form** of the solution.

Question

What is the solution set of Ax = 0, where

$$A = \begin{pmatrix} 1 & -3 \\ 2 & -6 \end{pmatrix}$$
?

This last equation is called the **parametric vector form** of the solution.

It is obtained by listing equations for all the variables, in order, including the free ones, and making a vector equation.

Question

$$A = \begin{pmatrix} 1 & -3 \\ 2 & -6 \end{pmatrix}$$
?

Answer:
$$x = x_2 \begin{pmatrix} 3 \\ 1 \end{pmatrix}$$
 for any x_2 in **R**.

Question

$$A = \begin{pmatrix} 1 & -3 \\ 2 & -6 \end{pmatrix}$$
?

Answer:
$$x = x_2 \begin{pmatrix} 3 \\ 1 \end{pmatrix}$$
 for any x_2 in **R**. The solution set is Span $\left\{ \right.$

Question

$$A = \begin{pmatrix} 1 & -3 \\ 2 & -6 \end{pmatrix}$$
?


Answer:
$$x = x_2 \begin{pmatrix} 3 \\ 1 \end{pmatrix}$$
 for any x_2 in **R**. The solution set is Span $\left\{ \begin{pmatrix} 3 \\ 1 \end{pmatrix} \right\}$.

Question

What is the solution set of Ax = 0, where

$$A = \begin{pmatrix} 1 & -3 \\ 2 & -6 \end{pmatrix}$$
?

Answer: $x = x_2 \begin{pmatrix} 3 \\ 1 \end{pmatrix}$ for any x_2 in **R**. The solution set is Span $\left\{ \begin{pmatrix} 3 \\ 1 \end{pmatrix} \right\}$.




Question

What is the solution set of Ax = 0, where

$$A = \begin{pmatrix} 1 & -3 \\ 2 & -6 \end{pmatrix}$$
?

Answer: $x = x_2 \begin{pmatrix} 3 \\ 1 \end{pmatrix}$ for any x_2 in **R**. The solution set is Span $\left\{ \begin{pmatrix} 3 \\ 1 \end{pmatrix} \right\}$.



Note: one free variable means the solution set is a line in \mathbb{R}^2 (2 = # variables = # columns).

Question

$$A = \begin{pmatrix} 1 & 3 & 1 \\ 2 & -1 & -5 \\ 1 & 0 & -2 \end{pmatrix}$$
?

Question

$$A = \begin{pmatrix} 1 & 3 & 1 \\ 2 & -1 & -5 \\ 1 & 0 & -2 \end{pmatrix}$$
?

Example, continued

Question

$$A = \begin{pmatrix} 1 & 3 & 1 \\ 2 & -1 & -5 \\ 1 & 0 & -2 \end{pmatrix}$$
?


Answer: Span
$$\left\{ \begin{pmatrix} 2 \\ -1 \\ 1 \end{pmatrix} \right\}$$
.

Example, continued

Question

$$A = \begin{pmatrix} 1 & 3 & 1 \\ 2 & -1 & -5 \\ 1 & 0 & -2 \end{pmatrix}$$
?

Answer: Span
$$\left\{ \begin{pmatrix} 2 \\ -1 \\ 1 \end{pmatrix} \right\}$$
.




Question

What is the solution set of Ax = 0, where

$$A = \begin{pmatrix} 1 & 3 & 1 \\ 2 & -1 & -5 \\ 1 & 0 & -2 \end{pmatrix}$$
?

Answer: Span
$$\left\{ \begin{pmatrix} 2 \\ -1 \\ 1 \end{pmatrix} \right\}$$
.



Note: *one* free variable means the solution set is a *line* in \mathbb{R}^3 (3 = # variables = # columns).

Question

$$A = \begin{pmatrix} 1 & 2 & 0 & -1 \\ -2 & -3 & 4 & 5 \\ 2 & 4 & 0 & -2 \end{pmatrix}$$
?

Question

$$\begin{pmatrix}
1 & 2 & 0 & -1 \\
-2 & -3 & 4 & 5 \\
2 & 4 & 0 & -2
\end{pmatrix}$$

Question

$$\begin{pmatrix} 1 & 2 & 0 & -1 \\ -2 & -3 & 4 & 5 \\ 2 & 4 & 0 & -2 \end{pmatrix} \xrightarrow{\text{row reduce}} \begin{pmatrix} 1 & 0 & -8 & -7 \\ 0 & 1 & 4 & 3 \\ 0 & 0 & 0 & 0 \end{pmatrix}$$

Homogeneous Systems

Question

$$\begin{pmatrix} 1 & 2 & 0 & -1 \\ -2 & -3 & 4 & 5 \\ 2 & 4 & 0 & -2 \end{pmatrix} \quad \overset{\text{row reduce}}{\longrightarrow} \quad \begin{pmatrix} 1 & 0 & -8 & -7 \\ 0 & 1 & 4 & 3 \\ 0 & 0 & 0 & 0 \end{pmatrix}$$

$$\overset{\text{equations}}{\longrightarrow} \quad \begin{cases} x_1 & -8x_3 - 7x_4 = 0 \\ x_2 + 4x_3 + 3x_4 = 0 \end{cases}$$

Homogeneous Systems

Question

$$\begin{pmatrix} 1 & 2 & 0 & -1 \\ -2 & -3 & 4 & 5 \\ 2 & 4 & 0 & -2 \end{pmatrix} \xrightarrow{\text{row reduce}} \begin{pmatrix} 1 & 0 & -8 & -7 \\ 0 & 1 & 4 & 3 \\ 0 & 0 & 0 & 0 \end{pmatrix}$$

$$\stackrel{\text{equations}}{\longleftrightarrow} \begin{cases} x_1 & -8x_3 - 7x_4 = 0 \\ x_2 + 4x_3 + 3x_4 = 0 \end{cases}$$

$$\xrightarrow{\text{parametric form}} \begin{cases} x_1 = 8x_3 + 7x_4 \\ x_2 = -4x_3 - 3x_4 \\ x_3 = x_3 \\ x_4 = x_4 \end{cases}$$

$$\begin{pmatrix} 1 & 2 & 0 & -1 \\ -2 & -3 & 4 & 5 \\ 2 & 4 & 0 & -2 \end{pmatrix} \xrightarrow{\text{row reduce}} \begin{pmatrix} 1 & 0 & -8 & -7 \\ 0 & 1 & 4 & 3 \\ 0 & 0 & 0 & 0 \end{pmatrix}$$

$$\xrightarrow{\text{equations}} \begin{cases} x_1 & -8x_3 - 7x_4 = 0 \\ x_2 + 4x_3 + 3x_4 = 0 \end{cases}$$

$$\xrightarrow{\text{parametric form}} \begin{cases} x_1 = 8x_3 + 7x_4 \\ x_2 = -4x_3 - 3x_4 \\ x_3 = x_3 \\ x_4 = x_4 \end{cases}$$

$$\xrightarrow{\text{parametric vector form}} x = \begin{pmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \end{pmatrix} = \begin{pmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \end{pmatrix} = \begin{pmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \end{pmatrix} = \begin{pmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \end{pmatrix} = \begin{pmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \end{pmatrix}$$

$$\begin{pmatrix} 1 & 2 & 0 & -1 \\ -2 & -3 & 4 & 5 \\ 2 & 4 & 0 & -2 \end{pmatrix} \xrightarrow{\text{row reduce}} \begin{pmatrix} 1 & 0 & -8 & -7 \\ 0 & 1 & 4 & 3 \\ 0 & 0 & 0 & 0 \end{pmatrix}$$

$$\xrightarrow{\text{equations}} \begin{cases} x_1 & -8x_3 - 7x_4 = 0 \\ x_2 + 4x_3 + 3x_4 = 0 \end{cases}$$

$$\begin{cases} x_1 = 8x_3 + 7x_4 \\ x_2 = -4x_3 - 3x_4 \\ x_3 = x_3 \\ x_4 = x_4 \end{cases}$$

$$\xrightarrow{\text{parametric vector form}} \begin{cases} x = 8x_3 + 7x_4 \\ x_2 = -4x_3 - 3x_4 \\ x_3 = x_3 \\ x_4 = x_4 \end{cases}$$

$$\begin{pmatrix} 1 & 2 & 0 & -1 \\ -2 & -3 & 4 & 5 \\ 2 & 4 & 0 & -2 \end{pmatrix} \quad \text{row reduce} \quad \begin{pmatrix} 1 & 0 & -8 & -7 \\ 0 & 1 & 4 & 3 \\ 0 & 0 & 0 & 0 \end{pmatrix}$$

$$\stackrel{\text{equations}}{\longleftrightarrow} \quad \begin{cases} x_1 & -8x_3 - 7x_4 = 0 \\ x_2 + 4x_3 + 3x_4 = 0 \end{cases}$$

$$\stackrel{\text{parametric form}}{\longleftrightarrow} \quad \begin{cases} x_1 = 8x_3 + 7x_4 \\ x_2 = -4x_3 - 3x_4 \\ x_3 = x_3 \\ x_4 = x_4 \end{cases}$$

$$\begin{pmatrix} x_1 \end{pmatrix} \quad \begin{pmatrix} 8 \end{pmatrix}$$

parametric vector form
$$x = \begin{pmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \end{pmatrix} = x_3 \begin{pmatrix} 8 \\ -4 \\ 1 \\ 0 \end{pmatrix} + x_4 \begin{pmatrix} 7 \\ -3 \\ 0 \\ 1 \end{pmatrix}.$$

Homogeneous Systems Example, continued

Question

$$A = \begin{pmatrix} 1 & 2 & 0 & -1 \\ -2 & -3 & 4 & 5 \\ 2 & 4 & 0 & -2 \end{pmatrix}$$
?

Homogeneous Systems Example, continued

Question

$$A = \begin{pmatrix} 1 & 2 & 0 & -1 \\ -2 & -3 & 4 & 5 \\ 2 & 4 & 0 & -2 \end{pmatrix}$$
?

Answer: Span
$$\left\{ \begin{pmatrix} 8 \\ -4 \\ 1 \\ 0 \end{pmatrix}, \right.$$

Homogeneous Systems Example, continued

Question

$$A = \begin{pmatrix} 1 & 2 & 0 & -1 \\ -2 & -3 & 4 & 5 \\ 2 & 4 & 0 & -2 \end{pmatrix}$$
?

Answer: Span
$$\left\{ \begin{pmatrix} 8 \\ -4 \\ 1 \\ 0 \end{pmatrix}, \begin{pmatrix} 7 \\ -3 \\ 0 \\ 1 \end{pmatrix} \right\}$$
.

Homogeneous Systems

Example, continued

Question

What is the solution set of Ax = 0, where

$$A = \begin{pmatrix} 1 & 2 & 0 & -1 \\ -2 & -3 & 4 & 5 \\ 2 & 4 & 0 & -2 \end{pmatrix}$$
?

Answer: Span
$$\left\{ \begin{pmatrix} 8 \\ -4 \\ 1 \\ 0 \end{pmatrix}, \begin{pmatrix} 7 \\ -3 \\ 0 \\ 1 \end{pmatrix} \right\}$$
.

[not pictured here]

Homogeneous Systems

Example, continued

Question

What is the solution set of Ax = 0, where

$$A = \begin{pmatrix} 1 & 2 & 0 & -1 \\ -2 & -3 & 4 & 5 \\ 2 & 4 & 0 & -2 \end{pmatrix}?$$

Answer: Span
$$\left\{ \begin{pmatrix} 8 \\ -4 \\ 1 \\ 0 \end{pmatrix}, \begin{pmatrix} 7 \\ -3 \\ 0 \\ 1 \end{pmatrix} \right\}$$
.

[not pictured here]

Note: *two* free variables means the solution set is a *plane* in \mathbf{R}^4 (4 = # variables = # columns).

Homogeneous systems

Let A be an $m \times n$ matrix.

Homogeneous systems

Let A be an $m \times n$ matrix. Suppose that the free variables in the homogeneous equation Ax = 0 are x_i, x_j, x_k, \ldots

Homogeneous systems

Let A be an $m \times n$ matrix. Suppose that the free variables in the homogeneous equation Ax = 0 are x_i, x_j, x_k, \dots

Then the solutions to Ax = 0 can be written in the form

$$x = x_i v_i + x_j v_j + x_k v_k + \cdots$$

for some vectors v_i, v_j, v_k, \ldots in R-,

Homogeneous systems

Let A be an $m \times n$ matrix. Suppose that the free variables in the homogeneous equation Ax = 0 are x_i, x_j, x_k, \dots

Then the solutions to Ax = 0 can be written in the form

$$x = x_i v_i + x_j v_j + x_k v_k + \cdots$$

for some vectors v_i, v_j, v_k, \ldots in \mathbb{R}^n , and any scalars x_i, x_j, x_k, \ldots

Homogeneous systems

Let A be an $m \times n$ matrix. Suppose that the free variables in the homogeneous equation Ax = 0 are x_i, x_j, x_k, \dots

Then the solutions to Ax = 0 can be written in the form

$$x = x_i v_i + x_j v_j + x_k v_k + \cdots$$

for some vectors v_i, v_j, v_k, \ldots in \mathbf{R}^n , and any scalars x_i, x_j, x_k, \ldots

The solution set is

Span{

Homogeneous systems

Let A be an $m \times n$ matrix. Suppose that the free variables in the homogeneous equation Ax = 0 are x_i, x_j, x_k, \dots

Then the solutions to Ax = 0 can be written in the form

$$x = x_i v_i + x_j v_j + x_k v_k + \cdots$$

for some vectors v_i, v_j, v_k, \ldots in \mathbf{R}^n , and any scalars x_i, x_j, x_k, \ldots

The solution set is

$$\mathsf{Span}\big\{v_i,\ v_j,\ v_k,\ldots\big\}.$$

Homogeneous systems

Let A be an $m \times n$ matrix. Suppose that the free variables in the homogeneous equation Ax = 0 are x_i, x_j, x_k, \dots

Then the solutions to Ax = 0 can be written in the form

$$x = x_i v_i + x_i v_i + x_k v_k + \cdots$$

for some vectors v_i, v_j, v_k, \ldots in \mathbf{R}^n , and any scalars x_i, x_j, x_k, \ldots

The solution set is

$$Span\{v_i, v_j, v_k, \ldots\}.$$

The equation above is called the **parametric vector form** of the solution.

How many solutions can there be to a homogeneous system with more equations than variables?

- A. 0
- B. 1
- **C**. ∞

How many solutions can there be to a homogeneous system with more equations than variables?

- A. 0
- B. 1
- **C**. ∞

The trivial solution is always a solution to a homogeneous system, so answer A is impossible.

How many solutions can there be to a homogeneous system with more equations than variables?

- A. (
- B. 1
- **C**. ∞

The trivial solution is always a solution to a homogeneous system, so answer A is impossible.

This matrix has only one solution to Ax = 0:

$$A = \begin{pmatrix} 1 & 0 \\ 0 & 1 \\ 0 & 0 \end{pmatrix}$$

How many solutions can there be to a homogeneous system with more equations than variables?

- A. (
- B. 1
- **C**. ∞

The trivial solution is always a solution to a homogeneous system, so answer A is impossible.

This matrix has only one solution to Ax = 0:

$$A = \begin{pmatrix} 1 & 0 \\ 0 & 1 \\ 0 & 0 \end{pmatrix}$$

This matrix has infinitely many solutions to Ax = 0:

$$A = \begin{pmatrix} 1 & 1 \\ 0 & 0 \\ 0 & 0 \end{pmatrix}$$

Question

$$A = \begin{pmatrix} 1 & -3 \\ 2 & -6 \end{pmatrix} \quad \text{and} \quad b = \begin{pmatrix} -3 \\ -6 \end{pmatrix}?$$

Nonhomogeneous Systems Example

Question

What is the solution set of Ax = b, where

$$A = \begin{pmatrix} 1 & -3 \\ 2 & -6 \end{pmatrix} \quad \text{and} \quad b = \begin{pmatrix} -3 \\ -6 \end{pmatrix}?$$

The only difference from the homogeneous case is the constant vector $p = {-3 \choose 0}$.

Question

What is the solution set of Ax = b, where

$$A = \begin{pmatrix} 1 & -3 \\ 2 & -6 \end{pmatrix} \quad \text{and} \quad b = \begin{pmatrix} -3 \\ -6 \end{pmatrix}?$$

The only difference from the homogeneous case is the constant vector $p = \binom{-3}{0}$.

Note that p is itself a solution: take $x_2 = 0$.

Example, continued

Question

$$A = \begin{pmatrix} 1 & -3 \\ 2 & -6 \end{pmatrix} \quad \text{and} \quad b = \begin{pmatrix} -3 \\ -6 \end{pmatrix}?$$

Answer:
$$x = x_2 \begin{pmatrix} 3 \\ 1 \end{pmatrix} + \begin{pmatrix} -3 \\ 0 \end{pmatrix}$$
 for any x_2 in **R**.

Example, continued

Question

What is the solution set of Ax = b, where

$$A = \begin{pmatrix} 1 & -3 \\ 2 & -6 \end{pmatrix} \quad \text{and} \quad b = \begin{pmatrix} -3 \\ -6 \end{pmatrix}?$$

Answer:
$$x = x_2 \begin{pmatrix} 3 \\ 1 \end{pmatrix} + \begin{pmatrix} -3 \\ 0 \end{pmatrix}$$
 for any x_2 in **R**.


This is a *translate* of Span $\left\{ \begin{pmatrix} 3 \\ 1 \end{pmatrix} \right\}$: it is the parallel line through $p = \begin{pmatrix} -3 \\ 0 \end{pmatrix}$.

What is the solution set of Ax = b, where

$$A = \begin{pmatrix} 1 & -3 \\ 2 & -6 \end{pmatrix}$$
 and $b = \begin{pmatrix} -3 \\ -6 \end{pmatrix}$?

Answer:
$$x = x_2 \begin{pmatrix} 3 \\ 1 \end{pmatrix} + \begin{pmatrix} -3 \\ 0 \end{pmatrix}$$
 for any x_2 in **R**.

This is a *translate* of Span $\left\{ \begin{pmatrix} 3 \\ 1 \end{pmatrix} \right\}$: it is the parallel line through $p = \begin{pmatrix} -3 \\ 0 \end{pmatrix}$.




What is the solution set of Ax = b, where

$$A = \begin{pmatrix} 1 & -3 \\ 2 & -6 \end{pmatrix} \quad \text{and} \quad b = \begin{pmatrix} -3 \\ -6 \end{pmatrix}?$$

Answer:
$$x = x_2 \begin{pmatrix} 3 \\ 1 \end{pmatrix} + \begin{pmatrix} -3 \\ 0 \end{pmatrix}$$
 for any x_2 in **R**.

This is a *translate* of Span $\left\{ \begin{pmatrix} 3 \\ 1 \end{pmatrix} \right\}$: it is the parallel line through $p = \begin{pmatrix} -3 \\ 0 \end{pmatrix}$.



It can be written

$$\mathsf{Span}\!\left\{ \begin{pmatrix} \mathbf{3} \\ \mathbf{1} \end{pmatrix} \right\} + \begin{pmatrix} -3 \\ 0 \end{pmatrix}.$$

Nonhomogeneous Systems Example

Question

$$A = \begin{pmatrix} 1 & 3 & 1 \\ 2 & -1 & -5 \\ 1 & 0 & -2 \end{pmatrix} \quad \text{and} \quad b = \begin{pmatrix} -5 \\ -3 \\ -2 \end{pmatrix}?$$

Example, continued

Question

$$A = \begin{pmatrix} 1 & 3 & 1 \\ 2 & -1 & -5 \\ 1 & 0 & -2 \end{pmatrix} \quad \text{and} \quad b = \begin{pmatrix} -5 \\ -3 \\ -2 \end{pmatrix}?$$

Answer: Span
$$\left\{ \begin{pmatrix} 2 \\ -1 \\ 1 \end{pmatrix} \right\} +$$

Example, continued

Question

$$A = \begin{pmatrix} 1 & 3 & 1 \\ 2 & -1 & -5 \\ 1 & 0 & -2 \end{pmatrix} \quad \text{and} \quad b = \begin{pmatrix} -5 \\ -3 \\ -2 \end{pmatrix}?$$

Answer: Span
$$\left\{ \begin{pmatrix} 2 \\ -1 \\ 1 \end{pmatrix} \right\} + \begin{pmatrix} -2 \\ -1 \\ 0 \end{pmatrix}$$
.


Example, continued

Question

What is the solution set of Ax = b, where

$$A = \begin{pmatrix} 1 & 3 & 1 \\ 2 & -1 & -5 \\ 1 & 0 & -2 \end{pmatrix} \quad \text{and} \quad b = \begin{pmatrix} -5 \\ -3 \\ -2 \end{pmatrix}?$$

Answer: Span
$$\left\{ \begin{pmatrix} 2 \\ -1 \\ 1 \end{pmatrix} \right\} + \begin{pmatrix} -2 \\ -1 \\ 0 \end{pmatrix}$$
.



The solution set is a translate of

Span
$$\left\{ \begin{pmatrix} 2 \\ -1 \\ 1 \end{pmatrix} \right\}$$
:

it is the parallel line through

$$p = \begin{pmatrix} -2 \\ -1 \\ 0 \end{pmatrix}.$$

Key Observation

The set of solutions to Ax = b, if it is nonempty, is obtained by taking one **specific** or **particular solution** p to Ax = b, and adding all solutions to Ax = 0.

Key Observation

The set of solutions to Ax = b, if it is nonempty, is obtained by taking one **specific** or **particular solution** p to Ax = b, and adding all solutions to Ax = 0.

Why? If
$$Ap = b$$
 and $Ax = 0$, then
$$A(p+x) = Ap + Ax = b + 0 = b,$$

Key Observation

The set of solutions to Ax=b, if it is nonempty, is obtained by taking one **specific** or **particular solution** p to Ax=b, and adding all solutions to Ax=0.

Why? If
$$Ap = b$$
 and $Ax = 0$, then

$$A(p+x) = Ap + Ax = b + 0 = b,$$

so p + x is also a solution to Ax = b.

Key Observation

The set of solutions to Ax = b, if it is nonempty, is obtained by taking one **specific** or **particular solution** p to Ax = b, and adding all solutions to Ax = 0.

Why? If Ap = b and Ax = 0, then

$$A(p+x) = Ap + Ax = b + 0 = b,$$

so p + x is also a solution to Ax = b.

We know the solution set of Ax = 0 is a span.

Key Observation

The set of solutions to Ax = b, if it is nonempty, is obtained by taking one **specific** or **particular solution** p to Ax = b, and adding all solutions to Ax = 0.

Why? If Ap = b and Ax = 0, then

$$A(p+x) = Ap + Ax = b + 0 = b,$$

so p + x is also a solution to Ax = b.

We know the solution set of Ax = 0 is a span. So the solution set of Ax = b is a *translate* of a span: it is *parallel* to a span.

Key Observation

The set of solutions to Ax = b, if it is nonempty, is obtained by taking one **specific** or **particular solution** p to Ax = b, and adding all solutions to Ax = 0.

Why? If Ap = b and Ax = 0, then

$$A(p+x) = Ap + Ax = b + 0 = b,$$

so p + x is also a solution to Ax = b.

We know the solution set of Ax = 0 is a span. So the solution set of Ax = b is a *translate* of a span: it is *parallel* to a span. (Or it is empty.)

Key Observation


The set of solutions to Ax = b, if it is nonempty, is obtained by taking one **specific** or **particular solution** p to Ax = b, and adding all solutions to Ax = 0.

Why? If Ap = b and Ax = 0, then

$$A(p+x) = Ap + Ax = b + 0 = b,$$

so p + x is also a solution to Ax = b.

We know the solution set of Ax = 0 is a span. So the solution set of Ax = b is a *translate* of a span: it is *parallel* to a span. (Or it is empty.)



This works for *any* specific solution p: it doesn't have to be the one produced by finding the parametric vector form and setting the free variables all to zero, as we did before.


Homogeneous vs. Nonhomogeneous Systems Varying *b*

If we understand the solution set of Ax = 0, then we understand the solution set of Ax = b for all b: they are all translates (or empty).

Homogeneous vs. Nonhomogeneous Systems Varying *b*


If we understand the solution set of Ax = 0, then we understand the solution set of Ax = b for all b: they are all translates (or empty).

For instance, if $A = \begin{pmatrix} 1 & -3 \\ 2 & -6 \end{pmatrix}$, then the solution sets for varying b look like this:



If we understand the solution set of Ax = 0, then we understand the solution set of Ax = b for all b: they are all translates (or empty).


For instance, if $A = \begin{pmatrix} 1 & -3 \\ 2 & -6 \end{pmatrix}$, then the solution sets for varying b look like this:



Which *b* gives the solution set Ax = b in red in the picture?

If we understand the solution set of Ax = 0, then we understand the solution set of Ax = b for all b: they are all translates (or empty).

For instance, if $A = \begin{pmatrix} 1 & -3 \\ 2 & -6 \end{pmatrix}$, then the solution sets for varying b look like this:




Which b gives the solution set Ax = b in red in the picture?

Choose p on the red line, and set b = Ap. Then p is a specific solution to Ax = b, so the solution set of Ax = b is the red line.

If we understand the solution set of Ax = 0, then we understand the solution set of Ax = b for all b: they are all translates (or empty).

For instance, if $A = \begin{pmatrix} 1 & -3 \\ 2 & -6 \end{pmatrix}$, then the solution sets for varying b look like this:




Which b gives the solution set Ax = b in red in the picture?

Choose p on the red line, and set b = Ap. Then p is a specific solution to Ax = b, so the solution set of Ax = b is the red line.

Note the cool optical illusion!

If we understand the solution set of Ax = 0, then we understand the solution set of Ax = b for all b: they are all translates (or empty).

For instance, if $A = \begin{pmatrix} 1 & -3 \\ 2 & -6 \end{pmatrix}$, then the solution sets for varying b look like this:



Which *b* gives the solution set Ax = b in red in the picture?

Choose p on the red line, and set b = Ap. Then p is a specific solution to Ax = b, so the solution set of Ax = b is the red line.

Note the cool optical illusion!

For a matrix equation Ax = b, you now know how to find which b's are possible, and what the solution set looks like for all b, both using spans.