

Section 1.4

The Matrix Equation $Ax = b$

Matrix \times Vector


Let A be an $m \times n$ matrix

Matrix \times Vector

the first number is
the number of rows

the second number is
the number of columns

Let A be an $m \times n$ matrix

A diagram consisting of two blue arrows. The first arrow starts from the text 'the first number is the number of rows' and points down to the variable 'm' in the expression 'm x n'. The second arrow starts from the text 'the second number is the number of columns' and points down to the variable 'n' in the expression 'm x n'.

Matrix \times Vector

the first number is
the number of rows

the second number is
the number of columns

Let A be an $m \times n$ matrix

$$A = \begin{pmatrix} | & | & \cdots & | \\ v_1 & v_2 & \cdots & v_n \\ | & | & \cdots & | \end{pmatrix}$$

with columns v_1, v_2, \dots, v_n

Matrix \times Vector

the first number is
the number of rows

the second number is
the number of columns

Let A be an $m \times n$ matrix

$$A = \begin{pmatrix} | & | & \cdots & | \\ v_1 & v_2 & \cdots & v_n \\ | & | & \cdots & | \end{pmatrix} \quad \text{with columns } v_1, v_2, \dots, v_n$$

Definition

The **product** of A with a vector x in \mathbf{R}^n is the linear combination

$$Ax = \begin{pmatrix} | & | & \cdots & | \\ v_1 & v_2 & \cdots & v_n \\ | & | & \cdots & | \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{pmatrix} \stackrel{\text{def}}{=} x_1 v_1 + x_2 v_2 + \cdots + x_n v_n.$$

Matrix \times Vector

the first number is
the number of rows

the second number is
the number of columns

Let A be an $m \times n$ matrix

$$A = \begin{pmatrix} | & | & \cdots & | \\ v_1 & v_2 & \cdots & v_n \\ | & | & \cdots & | \end{pmatrix} \quad \text{with columns } v_1, v_2, \dots, v_n$$

Definition

The **product** of A with a vector x in \mathbf{R}^n is the linear combination

$$Ax = \begin{pmatrix} | & | & \cdots & | \\ v_1 & v_2 & \cdots & v_n \\ | & | & \cdots & | \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{pmatrix} \stackrel{\text{def}}{=} x_1 v_1 + x_2 v_2 + \cdots + x_n v_n.$$

this means the equality
is a *definition*

Matrix \times Vector

the first number is
the number of rows

the second number is
the number of columns

Let A be an $m \times n$ matrix

$$A = \begin{pmatrix} | & | & \cdots & | \\ v_1 & v_2 & \cdots & v_n \\ | & | & \cdots & | \end{pmatrix} \quad \text{with columns } v_1, v_2, \dots, v_n$$

Definition

The **product** of A with a vector x in \mathbf{R}^n is the linear combination

$$Ax = \begin{pmatrix} | & | & \cdots & | \\ v_1 & v_2 & \cdots & v_n \\ | & | & \cdots & | \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{pmatrix} \stackrel{\text{def}}{=} x_1 v_1 + x_2 v_2 + \cdots + x_n v_n.$$

this means the equality
is a *definition*

The output is a vector in \mathbf{R}^m .

Matrix \times Vector

the first number is
the number of rows

the second number is
the number of columns

Let A be an $m \times n$ matrix

$$A = \begin{pmatrix} | & | & \cdots & | \\ v_1 & v_2 & \cdots & v_n \\ | & | & \cdots & | \end{pmatrix} \quad \text{with columns } v_1, v_2, \dots, v_n$$

Definition

The **product** of A with a vector x in \mathbf{R}^n is the linear combination

$$Ax = \begin{pmatrix} | & | & \cdots & | \\ v_1 & v_2 & \cdots & v_n \\ | & | & \cdots & | \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{pmatrix} \stackrel{\text{def}}{=} x_1 v_1 + x_2 v_2 + \cdots + x_n v_n.$$

this means the equality
is a *definition*

The output is a vector in \mathbf{R}^m .

Matrix \times Vector

the first number is
the number of rows

the second number is
the number of columns

Let A be an $m \times n$ matrix

$$A = \begin{pmatrix} | & | & \cdots & | \\ v_1 & v_2 & \cdots & v_n \\ | & | & \cdots & | \end{pmatrix} \quad \text{with columns } v_1, v_2, \dots, v_n$$

Definition

The **product** of A with a vector x in \mathbf{R}^n is the linear combination

$$Ax = \begin{pmatrix} | & | & \cdots & | \\ v_1 & v_2 & \cdots & v_n \\ | & | & \cdots & | \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{pmatrix} \stackrel{\text{def}}{=} x_1 v_1 + x_2 v_2 + \cdots + x_n v_n.$$

Annotations:
- Blue arrow from "this means the equality is a definition" points to the $\stackrel{\text{def}}{=}$ symbol.
- Red arrow from "these must be equal" points to the v_n in the matrix and the x_n in the vector.

The output is a vector in \mathbf{R}^m .

Note that the number of **columns** of A has to equal the number of **rows** of x .

Matrix \times Vector

the first number is
the number of rows

the second number is
the number of columns

Let A be an $m \times n$ matrix

$$A = \begin{pmatrix} | & | & \cdots & | \\ v_1 & v_2 & \cdots & v_n \\ | & | & \cdots & | \end{pmatrix} \quad \text{with columns } v_1, v_2, \dots, v_n$$

Definition

The **product** of A with a vector x in \mathbf{R}^n is the linear combination

$$Ax = \begin{pmatrix} | & | & \cdots & | \\ v_1 & v_2 & \cdots & v_n \\ | & | & \cdots & | \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{pmatrix} \stackrel{\text{def}}{=} x_1 v_1 + x_2 v_2 + \cdots + x_n v_n.$$

Annotations:
- Blue arrow from "this means the equality is a definition" points to the $\stackrel{\text{def}}{=}$ symbol.
- Red arrow from "these must be equal" points to the v_n in the matrix and the x_n in the vector.

The output is a vector in \mathbf{R}^m .

Note that the number of **columns** of A has to equal the number of **rows** of x .

Example

$$\begin{pmatrix} 4 & 5 & 6 \\ 7 & 8 & 9 \end{pmatrix} \begin{pmatrix} 1 \\ 2 \\ 3 \end{pmatrix} =$$

Matrix Equations

An example

Question

Let v_1, v_2, v_3 be vectors in \mathbf{R}^3 .

Matrix Equations

An example

Question

Let v_1, v_2, v_3 be vectors in \mathbf{R}^3 . How can you write the vector equation

$$2v_1 + 3v_2 - 4v_3 = \begin{pmatrix} 7 \\ 2 \\ 1 \end{pmatrix}$$

in terms of matrix multiplication?

Matrix Equations

In general

Let v_1, v_2, \dots, v_n , and b be vectors in \mathbf{R}^m .

Matrix Equations

In general

Let v_1, v_2, \dots, v_n , and b be vectors in \mathbf{R}^m . Consider the vector equation

$$x_1 v_1 + x_2 v_2 + \cdots + x_n v_n = b.$$

Matrix Equations

In general

Let v_1, v_2, \dots, v_n , and b be vectors in \mathbf{R}^m . Consider the vector equation

$$x_1 v_1 + x_2 v_2 + \cdots + x_n v_n = b.$$

It is equivalent to the **matrix equation**

$$Ax = b$$

where

Matrix Equations

In general

Let v_1, v_2, \dots, v_n , and b be vectors in \mathbf{R}^m . Consider the vector equation

$$x_1 v_1 + x_2 v_2 + \cdots + x_n v_n = b.$$

It is equivalent to the **matrix equation**

$$Ax = b$$

where

$$A = \begin{pmatrix} | & | & \cdots & | \\ v_1 & v_2 & \cdots & v_n \\ | & | & \cdots & | \end{pmatrix} \quad \text{and} \quad x = \begin{pmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{pmatrix}.$$

Matrix Equations

In general

Let v_1, v_2, \dots, v_n , and b be vectors in \mathbf{R}^m . Consider the vector equation

$$x_1 v_1 + x_2 v_2 + \cdots + x_n v_n = b.$$

It is equivalent to the **matrix equation**

$$Ax = b$$

where

$$A = \left(\begin{array}{c|c|c|c} | & | & & | \\ v_1 & v_2 & \cdots & v_n \\ | & | & & | \end{array} \right) \quad \text{and} \quad x = \begin{pmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{pmatrix}.$$

Conversely, if A is any $m \times n$ matrix, then

$Ax = b$ is equivalent to the
vector equation

Matrix Equations

In general

Let v_1, v_2, \dots, v_n , and b be vectors in \mathbf{R}^m . Consider the vector equation

$$x_1 v_1 + x_2 v_2 + \cdots + x_n v_n = b.$$

It is equivalent to the **matrix equation**

$$Ax = b$$

where

$$A = \left(\begin{array}{c|c|c|c} | & | & & | \\ v_1 & v_2 & \cdots & v_n \\ | & | & & | \end{array} \right) \quad \text{and} \quad x = \begin{pmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{pmatrix}.$$

Conversely, if A is any $m \times n$ matrix, then

$$Ax = b \quad \text{is equivalent to the} \quad x_1 v_1 + x_2 v_2 + \cdots + x_n v_n = b$$

vector equation

Matrix Equations

In general

Let v_1, v_2, \dots, v_n , and b be vectors in \mathbf{R}^m . Consider the vector equation

$$x_1 v_1 + x_2 v_2 + \cdots + x_n v_n = b.$$

It is equivalent to the **matrix equation**

$$Ax = b$$

where

$$A = \begin{pmatrix} | & | & \cdots & | \\ v_1 & v_2 & \cdots & v_n \\ | & | & \cdots & | \end{pmatrix} \quad \text{and} \quad x = \begin{pmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{pmatrix}.$$

Conversely, if A is any $m \times n$ matrix, then

$Ax = b$ is equivalent to the
vector equation

$$x_1 v_1 + x_2 v_2 + \cdots + x_n v_n = b$$

where v_1, \dots, v_n are

Matrix Equations

In general

Let v_1, v_2, \dots, v_n , and b be vectors in \mathbf{R}^m . Consider the vector equation

$$x_1 v_1 + x_2 v_2 + \cdots + x_n v_n = b.$$

It is equivalent to the **matrix equation**

$$Ax = b$$

where

$$A = \left(\begin{array}{c|c|c|c} | & | & & | \\ v_1 & v_2 & \cdots & v_n \\ | & | & & | \end{array} \right) \quad \text{and} \quad x = \begin{pmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{pmatrix}.$$

Conversely, if A is any $m \times n$ matrix, then

$$Ax = b \quad \text{is equivalent to the} \quad x_1 v_1 + x_2 v_2 + \cdots + x_n v_n = b$$

vector equation

where v_1, \dots, v_n are the columns of A , and x_1, \dots, x_n are

Matrix Equations

In general

Let v_1, v_2, \dots, v_n , and b be vectors in \mathbf{R}^m . Consider the vector equation

$$x_1 v_1 + x_2 v_2 + \cdots + x_n v_n = b.$$

It is equivalent to the **matrix equation**

$$Ax = b$$

where

$$A = \left(\begin{array}{c|c|c|c} | & | & & | \\ v_1 & v_2 & \cdots & v_n \\ | & | & & | \end{array} \right) \quad \text{and} \quad x = \begin{pmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{pmatrix}.$$

Conversely, if A is any $m \times n$ matrix, then

$$Ax = b \quad \text{is equivalent to the} \quad x_1 v_1 + x_2 v_2 + \cdots + x_n v_n = b$$

vector equation

where v_1, \dots, v_n are the columns of A , and x_1, \dots, x_n are the entries of x .

Linear Systems, Vector Equations, Matrix Equations, ...

We now have *four* equivalent ways of writing (and thinking about) linear systems:

We now have *four* equivalent ways of writing (and thinking about) linear systems:

1. As a system of equations:

$$2x_1 + 3x_2 = 7$$

$$x_1 - x_2 = 5$$

We now have *four* equivalent ways of writing (and thinking about) linear systems:

1. As a system of equations:

$$2x_1 + 3x_2 = 7$$

$$x_1 - x_2 = 5$$

2. As an augmented matrix:

$$\left(\begin{array}{cc|c} 2 & 3 & 7 \\ 1 & -1 & 5 \end{array} \right)$$

We now have *four* equivalent ways of writing (and thinking about) linear systems:

1. As a system of equations:

$$2x_1 + 3x_2 = 7$$

$$x_1 - x_2 = 5$$

2. As an augmented matrix:

$$\left(\begin{array}{cc|c} 2 & 3 & 7 \\ 1 & -1 & 5 \end{array} \right)$$

3. As a vector equation ($x_1 v_1 + \cdots + x_n v_n = b$):

$$x_1 \begin{pmatrix} 2 \\ 1 \end{pmatrix} + x_2 \begin{pmatrix} 3 \\ -1 \end{pmatrix} = \begin{pmatrix} 7 \\ 5 \end{pmatrix}$$

We now have *four* equivalent ways of writing (and thinking about) linear systems:

1. As a system of equations:

$$\begin{aligned}2x_1 + 3x_2 &= 7 \\ x_1 - x_2 &= 5\end{aligned}$$

2. As an augmented matrix:

$$\left(\begin{array}{cc|c} 2 & 3 & 7 \\ 1 & -1 & 5 \end{array} \right)$$

3. As a vector equation ($x_1 v_1 + \cdots + x_n v_n = b$):

$$x_1 \begin{pmatrix} 2 \\ 1 \end{pmatrix} + x_2 \begin{pmatrix} 3 \\ -1 \end{pmatrix} = \begin{pmatrix} 7 \\ 5 \end{pmatrix}$$

4. As a matrix equation ($Ax = b$):

$$\begin{pmatrix} 2 & 3 \\ 1 & -1 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \end{pmatrix} = \begin{pmatrix} 7 \\ 5 \end{pmatrix}$$

We now have *four* equivalent ways of writing (and thinking about) linear systems:

1. As a system of equations:

$$\begin{aligned}2x_1 + 3x_2 &= 7 \\ x_1 - x_2 &= 5\end{aligned}$$

2. As an augmented matrix:

$$\left(\begin{array}{cc|c} 2 & 3 & 7 \\ 1 & -1 & 5 \end{array} \right)$$

3. As a vector equation ($x_1 v_1 + \cdots + x_n v_n = b$):

$$x_1 \begin{pmatrix} 2 \\ 1 \end{pmatrix} + x_2 \begin{pmatrix} 3 \\ -1 \end{pmatrix} = \begin{pmatrix} 7 \\ 5 \end{pmatrix}$$

4. As a matrix equation ($Ax = b$):

$$\begin{pmatrix} 2 & 3 \\ 1 & -1 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \end{pmatrix} = \begin{pmatrix} 7 \\ 5 \end{pmatrix}$$

In particular, *all four have the same solution set.*

We now have *four* equivalent ways of writing (and thinking about) linear systems:

1. As a system of equations:

$$2x_1 + 3x_2 = 7$$

$$x_1 - x_2 = 5$$

2. As an augmented matrix:

$$\left(\begin{array}{cc|c} 2 & 3 & 7 \\ 1 & -1 & 5 \end{array} \right)$$

3. As a vector equation ($x_1 v_1 + \cdots + x_n v_n = b$):

$$x_1 \begin{pmatrix} 2 \\ 1 \end{pmatrix} + x_2 \begin{pmatrix} 3 \\ -1 \end{pmatrix} = \begin{pmatrix} 7 \\ 5 \end{pmatrix}$$

4. As a matrix equation ($Ax = b$):

$$\begin{pmatrix} 2 & 3 \\ 1 & -1 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \end{pmatrix} = \begin{pmatrix} 7 \\ 5 \end{pmatrix}$$

We will move back and forth freely between these over and over again, for the rest of the semester. Get comfortable with them now!

In particular, *all four have the same solution set.*

Matrix \times Vector

Another way

Definition

A **row vector** is a matrix with one row.

Matrix \times Vector

Another way

Definition

A **row vector** is a matrix with one row. The product of a row vector of length n and a (column) vector of length n is

$$\begin{pmatrix} a_1 & \cdots & a_n \end{pmatrix} \begin{pmatrix} x_1 \\ \vdots \\ x_n \end{pmatrix} \stackrel{\text{def}}{=} a_1 x_1 + \cdots + a_n x_n.$$

This is a _____.

Matrix \times Vector

Another way

Definition

A **row vector** is a matrix with one row. The product of a row vector of length n and a (column) vector of length n is

$$(a_1 \quad \cdots \quad a_n) \begin{pmatrix} x_1 \\ \vdots \\ x_n \end{pmatrix} \stackrel{\text{def}}{=} a_1 x_1 + \cdots + a_n x_n.$$

This is a scalar.

Matrix \times Vector

Another way

Definition

A **row vector** is a matrix with one row. The product of a row vector of length n and a (column) vector of length n is

$$(a_1 \quad \cdots \quad a_n) \begin{pmatrix} x_1 \\ \vdots \\ x_n \end{pmatrix} \stackrel{\text{def}}{=} a_1 x_1 + \cdots + a_n x_n.$$

This is a scalar.

If A is an $m \times n$ matrix with rows r_1, r_2, \dots, r_m ,

Matrix \times Vector

Another way

Definition

A **row vector** is a matrix with one row. The product of a row vector of length n and a (column) vector of length n is

$$(a_1 \quad \cdots \quad a_n) \begin{pmatrix} x_1 \\ \vdots \\ x_n \end{pmatrix} \stackrel{\text{def}}{=} a_1 x_1 + \cdots + a_n x_n.$$

This is a scalar.

If A is an $m \times n$ matrix with rows r_1, r_2, \dots, r_m , and x is a vector in \mathbf{R}^n , then

$$Ax = \begin{pmatrix} -r_1- \\ -r_2- \\ \vdots \\ -r_m- \end{pmatrix} x = \begin{pmatrix} r_1 x \\ r_2 x \\ \vdots \\ r_m x \end{pmatrix}$$

This is

Matrix \times Vector

Another way

Definition

A **row vector** is a matrix with one row. The product of a row vector of length n and a (column) vector of length n is

$$(a_1 \quad \cdots \quad a_n) \begin{pmatrix} x_1 \\ \vdots \\ x_n \end{pmatrix} \stackrel{\text{def}}{=} a_1 x_1 + \cdots + a_n x_n.$$

This is a scalar.

If A is an $m \times n$ matrix with rows r_1, r_2, \dots, r_m , and x is a vector in \mathbf{R}^n , then

$$Ax = \begin{pmatrix} -r_1- \\ -r_2- \\ \vdots \\ -r_m- \end{pmatrix} x = \begin{pmatrix} r_1 x \\ r_2 x \\ \vdots \\ r_m x \end{pmatrix}$$

This is a vector in \mathbf{R}^m (again).

Matrix \times Vector

Both ways

Example

$$\begin{pmatrix} 4 & 5 & 6 \\ 7 & 8 & 9 \end{pmatrix} \begin{pmatrix} 1 \\ 2 \\ 3 \end{pmatrix} =$$

Matrix \times Vector

Both ways

Example

$$\begin{pmatrix} 4 & 5 & 6 \\ 7 & 8 & 9 \end{pmatrix} \begin{pmatrix} 1 \\ 2 \\ 3 \end{pmatrix} =$$

Note this is the same as before:

Matrix \times Vector

Both ways

Example

$$\begin{pmatrix} 4 & 5 & 6 \\ 7 & 8 & 9 \end{pmatrix} \begin{pmatrix} 1 \\ 2 \\ 3 \end{pmatrix} =$$

Note this is the same as before:

Now you have *two* ways of computing Ax .

Matrix \times Vector

Both ways

Example

$$\begin{pmatrix} 4 & 5 & 6 \\ 7 & 8 & 9 \end{pmatrix} \begin{pmatrix} 1 \\ 2 \\ 3 \end{pmatrix} =$$

Note this is the same as before:

Now you have *two* ways of computing Ax .

In the second, you calculate Ax one entry at a time.

Matrix \times Vector

Both ways

Example

$$\begin{pmatrix} 4 & 5 & 6 \\ 7 & 8 & 9 \end{pmatrix} \begin{pmatrix} 1 \\ 2 \\ 3 \end{pmatrix} =$$

Note this is the same as before:

Now you have *two* ways of computing Ax .

In the second, you calculate Ax one entry at a time.

The second way is usually the most convenient, but we'll use both.

Spans and Solutions to Equations

Let A be a matrix with columns v_1, v_2, \dots, v_n :

$$A = \left(\begin{array}{c|c|c|c} | & | & & | \\ v_1 & v_2 & \cdots & v_n \\ | & | & & | \end{array} \right)$$

Spans and Solutions to Equations

Let A be a matrix with columns v_1, v_2, \dots, v_n :

$$A = \begin{pmatrix} | & | & \cdots & | \\ v_1 & v_2 & \cdots & v_n \\ | & | & \cdots & | \end{pmatrix}$$

Very Important Fact That Will Appear on Every Midterm and the Final

$Ax = b$ has a solution

Spans and Solutions to Equations

Let A be a matrix with columns v_1, v_2, \dots, v_n :

$$A = \begin{pmatrix} | & | & \cdots & | \\ v_1 & v_2 & \cdots & v_n \\ | & | & \cdots & | \end{pmatrix}$$

Very Important Fact That Will Appear on Every Midterm and the Final

$Ax = b$ has a solution

$$\iff \text{there exist } x_1, \dots, x_n \text{ such that } A \begin{pmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{pmatrix} = b$$


Spans and Solutions to Equations

Let A be a matrix with columns v_1, v_2, \dots, v_n :

$$A = \begin{pmatrix} | & | & \cdots & | \\ v_1 & v_2 & \cdots & v_n \\ | & | & \cdots & | \end{pmatrix}$$

Very Important Fact That Will Appear on Every Midterm and the Final

$Ax = b$ has a solution

 \iff there exist x_1, \dots, x_n such that $A \begin{pmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{pmatrix} = b$

"if and only if"


Spans and Solutions to Equations

Let A be a matrix with columns v_1, v_2, \dots, v_n :

$$A = \begin{pmatrix} | & | & \cdots & | \\ v_1 & v_2 & \cdots & v_n \\ | & | & \cdots & | \end{pmatrix}$$

Very Important Fact That Will Appear on Every Midterm and the Final

$Ax = b$ has a solution

 \iff there exist x_1, \dots, x_n such that $A \begin{pmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{pmatrix} = b$

"if and only if"

\iff there exist x_1, \dots, x_n such that $x_1 v_1 + \cdots + x_n v_n = b$


Spans and Solutions to Equations

Let A be a matrix with columns v_1, v_2, \dots, v_n :

$$A = \begin{pmatrix} | & | & \cdots & | \\ v_1 & v_2 & \cdots & v_n \\ | & | & \cdots & | \end{pmatrix}$$

Very Important Fact That Will Appear on Every Midterm and the Final

$Ax = b$ has a solution

 \iff there exist x_1, \dots, x_n such that $A \begin{pmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{pmatrix} = b$
"if and only if"

\iff there exist x_1, \dots, x_n such that $x_1 v_1 + \cdots + x_n v_n = b$

$\iff b$ is a linear combination of v_1, \dots, v_n

Spans and Solutions to Equations


Let A be a matrix with columns v_1, v_2, \dots, v_n :

$$A = \begin{pmatrix} | & | & \cdots & | \\ v_1 & v_2 & \cdots & v_n \\ | & | & \cdots & | \end{pmatrix}$$

Very Important Fact That Will Appear on Every Midterm and the Final

$Ax = b$ has a solution

“if and only if”

 \iff there exist x_1, \dots, x_n such that $A \begin{pmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{pmatrix} = b$

$$\iff \text{there exist } x_1, \dots, x_n \text{ such that } x_1 v_1 + \cdots + x_n v_n = b$$

$$\iff b \text{ is a linear combination of } v_1, \dots, v_n$$

$$\iff b \text{ is in the span of the columns of } A.$$

Spans and Solutions to Equations

Let A be a matrix with columns v_1, v_2, \dots, v_n :

$$A = \begin{pmatrix} | & | & \cdots & | \\ v_1 & v_2 & \cdots & v_n \\ | & | & \cdots & | \end{pmatrix}$$

Very Important Fact That Will Appear on Every Midterm and the Final

$Ax = b$ has a solution

“if and only if”



b is in the span of the columns of A .

Spans and Solutions to Equations


Let A be a matrix with columns v_1, v_2, \dots, v_n :

$$A = \begin{pmatrix} | & | & \cdots & | \\ v_1 & v_2 & \cdots & v_n \\ | & | & \cdots & | \end{pmatrix}$$

Very Important Fact That Will Appear on Every Midterm and the Final

$Ax = b$ has a solution

“if and only if”

 $\iff b$ is in the span of the columns of A .

The last condition is geometric.

Spans and Solutions to Equations

Example

Question

Let $A = \begin{pmatrix} 2 & 1 \\ -1 & 0 \\ 1 & -1 \end{pmatrix}$. Does the equation $Ax = \begin{pmatrix} 0 \\ 2 \\ 2 \end{pmatrix}$ have a solution?

Spans and Solutions to Equations


Example

Question

Let $A = \begin{pmatrix} 2 & 1 \\ -1 & 0 \\ 1 & -1 \end{pmatrix}$. Does the equation $Ax = \begin{pmatrix} 0 \\ 2 \\ 2 \end{pmatrix}$ have a solution?

Columns of A :

$$v = \begin{pmatrix} 2 \\ -1 \\ 1 \end{pmatrix} \quad w = \begin{pmatrix} 1 \\ 0 \\ -1 \end{pmatrix}$$



Spans and Solutions to Equations

Example

Question


Let $A = \begin{pmatrix} 2 & 1 \\ -1 & 0 \\ 1 & -1 \end{pmatrix}$. Does the equation $Ax = \begin{pmatrix} 0 \\ 2 \\ 2 \end{pmatrix}$ have a solution?

Columns of A :

$$v = \begin{pmatrix} 2 \\ -1 \\ 1 \end{pmatrix} \quad w = \begin{pmatrix} 1 \\ 0 \\ -1 \end{pmatrix}$$

Output vector:

$$b = \begin{pmatrix} 0 \\ 2 \\ 2 \end{pmatrix}$$




Spans and Solutions to Equations

Example

Question

Let $A = \begin{pmatrix} 2 & 1 \\ -1 & 0 \\ 1 & -1 \end{pmatrix}$. Does the equation $Ax = \begin{pmatrix} 0 \\ 2 \\ 2 \end{pmatrix}$ have a solution?



Columns of A :

$$v = \begin{pmatrix} 2 \\ -1 \\ 1 \end{pmatrix} \quad w = \begin{pmatrix} 1 \\ 0 \\ -1 \end{pmatrix}$$

Output vector:


$$b = \begin{pmatrix} 0 \\ 2 \\ 2 \end{pmatrix}$$

Spans and Solutions to Equations

Example

Question

Let $A = \begin{pmatrix} 2 & 1 \\ -1 & 0 \\ 1 & -1 \end{pmatrix}$. Does the equation $Ax = \begin{pmatrix} 0 \\ 2 \\ 2 \end{pmatrix}$ have a solution?



Columns of A :

$$v = \begin{pmatrix} 2 \\ -1 \\ 1 \end{pmatrix} \quad w = \begin{pmatrix} 1 \\ 0 \\ -1 \end{pmatrix}$$

Output vector:

$$b = \begin{pmatrix} 0 \\ 2 \\ 2 \end{pmatrix}$$


Is b contained in the span of the columns of A ?

Spans and Solutions to Equations

Example

Question

Let $A = \begin{pmatrix} 2 & 1 \\ -1 & 0 \\ 1 & -1 \end{pmatrix}$. Does the equation $Ax = \begin{pmatrix} 0 \\ 2 \\ 2 \end{pmatrix}$ have a solution?



Columns of A :

$$v = \begin{pmatrix} 2 \\ -1 \\ 1 \end{pmatrix} \quad w = \begin{pmatrix} 1 \\ 0 \\ -1 \end{pmatrix}$$

Output vector:

$$b = \begin{pmatrix} 0 \\ 2 \\ 2 \end{pmatrix}$$


Is b contained in the span of the columns of A ? It sure doesn't look like it.

Spans and Solutions to Equations

Example

Question

Let $A = \begin{pmatrix} 2 & 1 \\ -1 & 0 \\ 1 & -1 \end{pmatrix}$. Does the equation $Ax = \begin{pmatrix} 0 \\ 2 \\ 2 \end{pmatrix}$ have a solution?



Columns of A :

$$v = \begin{pmatrix} 2 \\ -1 \\ 1 \end{pmatrix} \quad w = \begin{pmatrix} 1 \\ 0 \\ -1 \end{pmatrix}$$

Output vector:

$$b = \begin{pmatrix} 0 \\ 2 \\ 2 \end{pmatrix}$$

Is b contained in the span of the columns of A ? It sure doesn't look like it.

Conclusion: $Ax = b$ is *inconsistent*.

Spans and Solutions to Equations

Example, continued

Question

Let $A = \begin{pmatrix} 2 & 1 \\ -1 & 0 \\ 1 & -1 \end{pmatrix}$. Does the equation $Ax = \begin{pmatrix} 0 \\ 2 \\ 2 \end{pmatrix}$ have a solution?

Answer: Let's check by solving the matrix equation using row reduction.

Spans and Solutions to Equations

Example, continued

Question

Let $A = \begin{pmatrix} 2 & 1 \\ -1 & 0 \\ 1 & -1 \end{pmatrix}$. Does the equation $Ax = \begin{pmatrix} 0 \\ 2 \\ 2 \end{pmatrix}$ have a solution?

Answer: Let's check by solving the matrix equation using row reduction.

In other words, the matrix equation

$$\begin{pmatrix} 2 & 1 \\ -1 & 0 \\ 1 & -1 \end{pmatrix} x = \begin{pmatrix} 0 \\ 2 \\ 2 \end{pmatrix}$$

has no solution, as the picture shows.

Spans and Solutions to Equations

Example

Question

Let $A = \begin{pmatrix} 2 & 1 \\ -1 & 0 \\ 1 & -1 \end{pmatrix}$. Does the equation $Ax = \begin{pmatrix} 1 \\ -1 \\ 2 \end{pmatrix}$ have a solution?

Spans and Solutions to Equations


Example

Question

Let $A = \begin{pmatrix} 2 & 1 \\ -1 & 0 \\ 1 & -1 \end{pmatrix}$. Does the equation $Ax = \begin{pmatrix} 1 \\ -1 \\ 2 \end{pmatrix}$ have a solution?

Columns of A :

$$v = \begin{pmatrix} 2 \\ -1 \\ 1 \end{pmatrix} \quad w = \begin{pmatrix} 1 \\ 0 \\ -1 \end{pmatrix}$$




Spans and Solutions to Equations

Example

Question

Let $A = \begin{pmatrix} 2 & 1 \\ -1 & 0 \\ 1 & -1 \end{pmatrix}$. Does the equation $Ax = \begin{pmatrix} 1 \\ -1 \\ 2 \end{pmatrix}$ have a solution?



Columns of A :

$$v = \begin{pmatrix} 2 \\ -1 \\ 1 \end{pmatrix} \quad w = \begin{pmatrix} 1 \\ 0 \\ -1 \end{pmatrix}$$

Solution vector:


$$b = \begin{pmatrix} 1 \\ -1 \\ 2 \end{pmatrix}$$

Spans and Solutions to Equations

Example

Question

Let $A = \begin{pmatrix} 2 & 1 \\ -1 & 0 \\ 1 & -1 \end{pmatrix}$. Does the equation $Ax = \begin{pmatrix} 1 \\ -1 \\ 2 \end{pmatrix}$ have a solution?



Columns of A :

$$v = \begin{pmatrix} 2 \\ -1 \\ 1 \end{pmatrix} \quad w = \begin{pmatrix} 1 \\ 0 \\ -1 \end{pmatrix}$$

Solution vector:


$$b = \begin{pmatrix} 1 \\ -1 \\ 2 \end{pmatrix}$$

Spans and Solutions to Equations

Example

Question

Let $A = \begin{pmatrix} 2 & 1 \\ -1 & 0 \\ 1 & -1 \end{pmatrix}$. Does the equation $Ax = \begin{pmatrix} 1 \\ -1 \\ 2 \end{pmatrix}$ have a solution?



Columns of A :

$$v = \begin{pmatrix} 2 \\ -1 \\ 1 \end{pmatrix} \quad w = \begin{pmatrix} 1 \\ 0 \\ -1 \end{pmatrix}$$

Solution vector:

$$b = \begin{pmatrix} 1 \\ -1 \\ 2 \end{pmatrix}$$


Is b contained in the span of the columns of A ?

Spans and Solutions to Equations

Example

Question

Let $A = \begin{pmatrix} 2 & 1 \\ -1 & 0 \\ 1 & -1 \end{pmatrix}$. Does the equation $Ax = \begin{pmatrix} 1 \\ -1 \\ 2 \end{pmatrix}$ have a solution?



Columns of A :

$$v = \begin{pmatrix} 2 \\ -1 \\ 1 \end{pmatrix} \quad w = \begin{pmatrix} 1 \\ 0 \\ -1 \end{pmatrix}$$

Solution vector:

$$b = \begin{pmatrix} 1 \\ -1 \\ 2 \end{pmatrix}$$

Is b contained in the span of the columns of A ? It looks like it: in fact,


$$b = \underline{\quad} v + \underline{\quad} w \implies x = \begin{pmatrix} \quad \\ \quad \end{pmatrix}.$$

Spans and Solutions to Equations

Example

Question

Let $A = \begin{pmatrix} 2 & 1 \\ -1 & 0 \\ 1 & -1 \end{pmatrix}$. Does the equation $Ax = \begin{pmatrix} 1 \\ -1 \\ 2 \end{pmatrix}$ have a solution?



Columns of A :

$$v = \begin{pmatrix} 2 \\ -1 \\ 1 \end{pmatrix} \quad w = \begin{pmatrix} 1 \\ 0 \\ -1 \end{pmatrix}$$

Solution vector:

$$b = \begin{pmatrix} 1 \\ -1 \\ 2 \end{pmatrix}$$

Is b contained in the span of the columns of A ? It looks like it: in fact,


$$b = 1v + (-1)w \implies x = \begin{pmatrix} 1 \\ -1 \end{pmatrix}.$$

Spans and Solutions to Equations

Example

Question

Let $A = \begin{pmatrix} 2 & 1 \\ -1 & 0 \\ 1 & -1 \end{pmatrix}$. Does the equation $Ax = \begin{pmatrix} 1 \\ -1 \\ 2 \end{pmatrix}$ have a solution?



Columns of A :

$$v = \begin{pmatrix} 2 \\ -1 \\ 1 \end{pmatrix} \quad w = \begin{pmatrix} 1 \\ 0 \\ -1 \end{pmatrix}$$

Solution vector:

$$b = \begin{pmatrix} 1 \\ -1 \\ 2 \end{pmatrix}$$

Is b contained in the span of the columns of A ? It looks like it: in fact,

$$b = 1v + (-1)w \implies x = \begin{pmatrix} 1 \\ -1 \end{pmatrix}.$$

Spans and Solutions to Equations

Example, continued

Question

Let $A = \begin{pmatrix} 2 & 1 \\ -1 & 0 \\ 1 & -1 \end{pmatrix}$. Does the equation $Ax = \begin{pmatrix} 1 \\ -1 \\ 2 \end{pmatrix}$ have a solution?

Answer: Let's do this systematically using row reduction.

Spans and Solutions to Equations

Example, continued

Question

Let $A = \begin{pmatrix} 2 & 1 \\ -1 & 0 \\ 1 & -1 \end{pmatrix}$. Does the equation $Ax = \begin{pmatrix} 1 \\ -1 \\ 2 \end{pmatrix}$ have a solution?

Answer: Let's do this systematically using row reduction.

This is consistent with the picture on the previous slide:

$$1 \begin{pmatrix} 2 \\ -1 \\ 1 \end{pmatrix} - 1 \begin{pmatrix} 1 \\ 0 \\ -1 \end{pmatrix} = \begin{pmatrix} 1 \\ -1 \\ 2 \end{pmatrix} \quad \text{or}$$

Spans and Solutions to Equations

Example, continued

Question

Let $A = \begin{pmatrix} 2 & 1 \\ -1 & 0 \\ 1 & -1 \end{pmatrix}$. Does the equation $Ax = \begin{pmatrix} 1 \\ -1 \\ 2 \end{pmatrix}$ have a solution?

Answer: Let's do this systematically using row reduction.

This is consistent with the picture on the previous slide:

$$1 \begin{pmatrix} 2 \\ -1 \\ 1 \end{pmatrix} - 1 \begin{pmatrix} 1 \\ 0 \\ -1 \end{pmatrix} = \begin{pmatrix} 1 \\ -1 \\ 2 \end{pmatrix} \quad \text{or} \quad A \begin{pmatrix} 1 \\ -1 \end{pmatrix} = \begin{pmatrix} 1 \\ -1 \\ 2 \end{pmatrix}.$$

Poll

Which of the following true statements can be checked by eyeballing them, *without* row reduction?

A. $\begin{pmatrix} 0 \\ 1 \\ 2 \end{pmatrix}$ is in the span of $\begin{pmatrix} 3 \\ 3 \\ 4 \end{pmatrix}$, $\begin{pmatrix} 0 \\ 10 \\ 20 \end{pmatrix}$, $\begin{pmatrix} 0 \\ -1 \\ -2 \end{pmatrix}$.

B. $\begin{pmatrix} 0 \\ 1 \\ 2 \end{pmatrix}$ is in the span of $\begin{pmatrix} 3 \\ 3 \\ 4 \end{pmatrix}$, $\begin{pmatrix} 0 \\ 5 \\ 7 \end{pmatrix}$, $\begin{pmatrix} 0 \\ 6 \\ 8 \end{pmatrix}$.

C. $\begin{pmatrix} 0 \\ 1 \\ 2 \end{pmatrix}$ is in the span of $\begin{pmatrix} 3 \\ 3 \\ 4 \end{pmatrix}$, $\begin{pmatrix} 0 \\ 1 \\ 0 \end{pmatrix}$, $\begin{pmatrix} 0 \\ 0 \\ \sqrt{2} \end{pmatrix}$.

D. $\begin{pmatrix} 0 \\ 1 \\ 2 \end{pmatrix}$ is in the span of $\begin{pmatrix} 5 \\ 7 \\ 0 \end{pmatrix}$, $\begin{pmatrix} 6 \\ 8 \\ 0 \end{pmatrix}$, $\begin{pmatrix} 3 \\ 3 \\ 4 \end{pmatrix}$.

When Solutions Always Exist

Here are criteria for a linear system to always have a solution.

Theorem

Let A be an $m \times n$ (non-augmented) matrix. The following are equivalent

When Solutions Always Exist

Here are criteria for a linear system to always have a solution.

Theorem

Let A be an $m \times n$ (non-augmented) matrix. The following are equivalent

↑
recall that this means
that for given A , either they're
all true, or they're all false

When Solutions Always Exist

Here are criteria for a linear system to always have a solution.

Theorem

Let A be an $m \times n$ (non-augmented) matrix. The following are equivalent

1. $Ax = b$ has a solution for all b in \mathbf{R}^m .

recall that this means
that for given A , either they're
all true, or they're all false

When Solutions Always Exist

Here are criteria for a linear system to always have a solution.

Theorem

Let A be an $m \times n$ (non-augmented) matrix. The following are equivalent

1. $Ax = b$ has a solution for all b in \mathbf{R}^m .

recall that this means
that for given A , either they're
all true, or they're all false

When Solutions Always Exist

Here are criteria for a linear system to always have a solution.

Theorem

Let A be an $m \times n$ (non-augmented) matrix. The following are equivalent

1. $Ax = b$ has a solution for all b in \mathbf{R}^m .
2. The span of the columns of A is all of \mathbf{R}^m .

recall that this means
that for given A , either they're
all true, or they're all false

When Solutions Always Exist

Here are criteria for a linear system to always have a solution.

Theorem

Let A be an $m \times n$ (non-augmented) matrix. The following are equivalent

1. $Ax = b$ has a solution for all b in \mathbf{R}^m .
2. The span of the columns of A is all of \mathbf{R}^m .
3. A has a pivot in each row.

recall that this means
that for given A , either they're
all true, or they're all false

When Solutions Always Exist

Here are criteria for a linear system to always have a solution.

Theorem

Let A be an $m \times n$ (non-augmented) matrix. The following are equivalent

1. $Ax = b$ has a solution for all b in \mathbf{R}^m .
2. The span of the columns of A is all of \mathbf{R}^m .
3. A has a pivot in each row.

recall that this means
that for given A , either they're
all true, or they're all false

Why is (1) the same as (2)?

When Solutions Always Exist

Here are criteria for a linear system to always have a solution.

Theorem

Let A be an $m \times n$ (non-augmented) matrix. The following are equivalent

1. $Ax = b$ has a solution for all b in \mathbf{R}^m .
2. The span of the columns of A is all of \mathbf{R}^m .
3. A has a pivot in each row.

recall that this means
that for given A , either they're
all true, or they're all false

Why is (1) the same as (2)? This was the Very Important box from before.

When Solutions Always Exist

Here are criteria for a linear system to always have a solution.

Theorem

Let A be an $m \times n$ (non-augmented) matrix. The following are equivalent

1. $Ax = b$ has a solution for all b in \mathbf{R}^m .
2. The span of the columns of A is all of \mathbf{R}^m .
3. A has a pivot in each row.

recall that this means
that for given A , either they're
all true, or they're all false

Why is (1) the same as (2)? This was the Very Important box from before.

Why is (1) the same as (3)?

When Solutions Always Exist

Here are criteria for a linear system to always have a solution.

Theorem

Let A be an $m \times n$ (non-augmented) matrix. The following are equivalent

1. $Ax = b$ has a solution for all b in \mathbf{R}^m .
2. The span of the columns of A is all of \mathbf{R}^m .
3. A has a pivot in each row.

recall that this means
that for given A , either they're
all true, or they're all false

Why is (1) the same as (2)? This was the Very Important box from before.

Why is (1) the same as (3)? If A has a pivot in each row then its reduced row echelon form looks like this:

$$\begin{pmatrix} 1 & 0 & \star & 0 & \star \\ 0 & 1 & \star & 0 & \star \\ 0 & 0 & 0 & 1 & \star \end{pmatrix}$$

When Solutions Always Exist

Here are criteria for a linear system to always have a solution.

Theorem

Let A be an $m \times n$ (non-augmented) matrix. The following are equivalent

1. $Ax = b$ has a solution for all b in \mathbf{R}^m .
2. The span of the columns of A is all of \mathbf{R}^m .
3. A has a pivot in each row.

recall that this means
that for given A , either they're
all true, or they're all false

Why is (1) the same as (2)? This was the **Very Important** box from before.

Why is (1) the same as (3)? If A has a pivot in each row then its reduced row echelon form looks like this:

$$\begin{pmatrix} 1 & 0 & \star & 0 & \star \\ 0 & 1 & \star & 0 & \star \\ 0 & 0 & 0 & 1 & \star \end{pmatrix} \quad \text{and } (A | b) \quad \begin{pmatrix} 1 & 0 & \star & 0 & \star & | & \star \\ 0 & 1 & \star & 0 & \star & | & \star \\ 0 & 0 & 0 & 1 & \star & | & \star \end{pmatrix}.$$

reduces to this:

When Solutions Always Exist

Here are criteria for a linear system to always have a solution.

Theorem

Let A be an $m \times n$ (non-augmented) matrix. The following are equivalent

1. $Ax = b$ has a solution for all b in \mathbf{R}^m .
2. The span of the columns of A is all of \mathbf{R}^m .
3. A has a pivot in each row.

recall that this means
that for given A , either they're
all true, or they're all false

Why is (1) the same as (2)? This was the **Very Important** box from before.

Why is (1) the same as (3)? If A has a pivot in each row then its reduced row echelon form looks like this:

$$\begin{pmatrix} 1 & 0 & \star & 0 & \star \\ 0 & 1 & \star & 0 & \star \\ 0 & 0 & 0 & 1 & \star \end{pmatrix} \quad \text{and } (A | b) \quad \begin{pmatrix} 1 & 0 & \star & 0 & \star & | & \star \\ 0 & 1 & \star & 0 & \star & | & \star \\ 0 & 0 & 0 & 1 & \star & | & \star \end{pmatrix}.$$

reduces to this:

There's no b that makes it inconsistent, so there's always a solution.

When Solutions Always Exist

Here are criteria for a linear system to always have a solution.

Theorem

Let A be an $m \times n$ (non-augmented) matrix. The following are equivalent

1. $Ax = b$ has a solution for all b in \mathbf{R}^m .
2. The span of the columns of A is all of \mathbf{R}^m .
3. A has a pivot in each row.

recall that this means
that for given A , either they're
all true, or they're all false

Why is (1) the same as (2)? This was the **Very Important** box from before.

Why is (1) the same as (3)? If A has a pivot in each row then its reduced row echelon form looks like this:

$$\begin{pmatrix} 1 & 0 & \star & 0 & \star \\ 0 & 1 & \star & 0 & \star \\ 0 & 0 & 0 & 1 & \star \end{pmatrix} \quad \text{and } (A | b) \quad \begin{pmatrix} 1 & 0 & \star & 0 & \star & | & \star \\ 0 & 1 & \star & 0 & \star & | & \star \\ 0 & 0 & 0 & 1 & \star & | & \star \end{pmatrix}.$$

reduces to this:

There's no b that makes it inconsistent, so there's always a solution. If A doesn't have a pivot in each row, then its reduced form looks like this:

When Solutions Always Exist

Here are criteria for a linear system to always have a solution.

Theorem

Let A be an $m \times n$ (non-augmented) matrix. The following are equivalent

1. $Ax = b$ has a solution for all b in \mathbf{R}^m .
2. The span of the columns of A is all of \mathbf{R}^m .
3. A has a pivot in each row.

recall that this means
that for given A , either they're
all true, or they're all false

Why is (1) the same as (2)? This was the **Very Important** box from before.

Why is (1) the same as (3)? If A has a pivot in each row then its reduced row echelon form looks like this:

$$\begin{pmatrix} 1 & 0 & \star & 0 & \star \\ 0 & 1 & \star & 0 & \star \\ 0 & 0 & 0 & 1 & \star \end{pmatrix} \quad \text{and } (A | b) \quad \begin{pmatrix} 1 & 0 & \star & 0 & \star & \star \\ 0 & 1 & \star & 0 & \star & \star \\ 0 & 0 & 0 & 1 & \star & \star \end{pmatrix}.$$

reduces to this:

There's no b that makes it inconsistent, so there's always a solution. If A doesn't have a pivot in each row, then its reduced form looks like this:

$$\begin{pmatrix} 1 & 0 & \star & 0 & \star \\ 0 & 1 & \star & 0 & \star \\ 0 & 0 & 0 & 0 & 0 \end{pmatrix}$$

When Solutions Always Exist

Here are criteria for a linear system to always have a solution.

Theorem

Let A be an $m \times n$ (non-augmented) matrix. The following are equivalent

1. $Ax = b$ has a solution for all b in \mathbf{R}^m .
2. The span of the columns of A is all of \mathbf{R}^m .
3. A has a pivot in each row.

recall that this means
that for given A , either they're
all true, or they're all false

Why is (1) the same as (2)? This was the Very Important box from before.

Why is (1) the same as (3)? If A has a pivot in each row then its reduced row echelon form looks like this:

$$\begin{pmatrix} 1 & 0 & \star & 0 & \star \\ 0 & 1 & \star & 0 & \star \\ 0 & 0 & 0 & 1 & \star \end{pmatrix} \quad \text{and } (A | b) \quad \begin{pmatrix} 1 & 0 & \star & 0 & \star & \star \\ 0 & 1 & \star & 0 & \star & \star \\ 0 & 0 & 0 & 1 & \star & \star \end{pmatrix}.$$

reduces to this:

There's no b that makes it inconsistent, so there's always a solution. If A doesn't have a pivot in each row, then its reduced form looks like this:

$$\begin{pmatrix} 1 & 0 & \star & 0 & \star \\ 0 & 1 & \star & 0 & \star \\ 0 & 0 & 0 & 0 & 0 \end{pmatrix} \quad \begin{array}{l} \text{and this can be} \\ \text{made} \\ \text{inconsistent:} \end{array} \quad \begin{pmatrix} 1 & 0 & \star & 0 & \star & 0 \\ 0 & 1 & \star & 0 & \star & 0 \\ 0 & 0 & 0 & 0 & 0 & 16 \end{pmatrix}.$$

Properties of the Matrix–Vector Product

Let c be a scalar, u, v be vectors, and A a matrix.

Properties of the Matrix–Vector Product

Let c be a scalar, u, v be vectors, and A a matrix.

- ▶ $A(u + v) = Au + Av$

Properties of the Matrix–Vector Product

Let c be a scalar, u, v be vectors, and A a matrix.

- ▶ $A(u + v) = Au + Av$
- ▶ $A(cv) = cAv$

Properties of the Matrix–Vector Product

Let c be a scalar, u, v be vectors, and A a matrix.

- ▶ $A(u + v) = Au + Av$
- ▶ $A(cv) = cAv$

See Lay, §1.4, Theorem 5.

Properties of the Matrix–Vector Product

Let c be a scalar, u, v be vectors, and A a matrix.

- ▶ $A(u + v) = Au + Av$
- ▶ $A(cv) = cAv$

See Lay, §1.4, Theorem 5.

Consequence: If u and v are solutions to $Ax = 0$, then so is every vector in $\text{Span}\{u, v\}$.

Properties of the Matrix–Vector Product

Let c be a scalar, u, v be vectors, and A a matrix.

- ▶ $A(u + v) = Au + Av$
- ▶ $A(cv) = cAv$

See Lay, §1.4, Theorem 5.

Consequence: If u and v are solutions to $Ax = 0$, then so is every vector in $\text{Span}\{u, v\}$. Why?

Properties of the Matrix–Vector Product

Let c be a scalar, u, v be vectors, and A a matrix.

- ▶ $A(u + v) = Au + Av$
- ▶ $A(cv) = cAv$

See Lay, §1.4, Theorem 5.

Consequence: If u and v are solutions to $Ax = 0$, then so is every vector in $\text{Span}\{u, v\}$. Why?

Important

The set of solutions to $Ax = 0$ is a span.