

## Section 6.4

### The Gram–Schmidt Process

# Motivation

All of the procedures we learned in §§6.2–6.3 require an *orthogonal* basis  $\{u_1, u_2, \dots, u_m\}$ .

# Motivation

All of the procedures we learned in §§6.2–6.3 require an *orthogonal* basis  $\{u_1, u_2, \dots, u_m\}$ .

- ▶ Finding the  $\mathcal{B}$ -coordinates of a vector  $x$  using dot products:

$$x = \sum_{i=1}^m \frac{x \cdot u_i}{u_i \cdot u_i} u_i$$

# Motivation

All of the procedures we learned in §§6.2–6.3 require an *orthogonal* basis  $\{u_1, u_2, \dots, u_m\}$ .

- ▶ Finding the  $\mathcal{B}$ -coordinates of a vector  $x$  using dot products:

$$x = \sum_{i=1}^m \frac{x \cdot u_i}{u_i \cdot u_i} u_i$$

- ▶ Finding the orthogonal projection of a vector  $x$  onto the span  $W$  of  $u_1, u_2, \dots, u_m$ :

$$\text{proj}_W(x) = \sum_{i=1}^m \frac{x \cdot u_i}{u_i \cdot u_i} u_i.$$

# Motivation

All of the procedures we learned in §§6.2–6.3 require an *orthogonal* basis  $\{u_1, u_2, \dots, u_m\}$ .

- Finding the  $\mathcal{B}$ -coordinates of a vector  $x$  using dot products:

$$x = \sum_{i=1}^m \frac{x \cdot u_i}{u_i \cdot u_i} u_i$$

- Finding the orthogonal projection of a vector  $x$  onto the span  $W$  of  $u_1, u_2, \dots, u_m$ :

$$\text{proj}_W(x) = \sum_{i=1}^m \frac{x \cdot u_i}{u_i \cdot u_i} u_i.$$

**Problem:** What if your basis isn't orthogonal?

# Motivation

All of the procedures we learned in §§6.2–6.3 require an *orthogonal* basis  $\{u_1, u_2, \dots, u_m\}$ .

- Finding the  $\mathcal{B}$ -coordinates of a vector  $x$  using dot products:

$$x = \sum_{i=1}^m \frac{x \cdot u_i}{u_i \cdot u_i} u_i$$

- Finding the orthogonal projection of a vector  $x$  onto the span  $W$  of  $u_1, u_2, \dots, u_m$ :

$$\text{proj}_W(x) = \sum_{i=1}^m \frac{x \cdot u_i}{u_i \cdot u_i} u_i.$$

**Problem:** What if your basis isn't orthogonal?

**Solution:** The Gram–Schmidt process: take any basis and make it orthogonal.

# The Gram–Schmidt Process

## Procedure

### The Gram–Schmidt Process

Let  $\{v_1, v_2, \dots, v_m\}$  be a basis for a subspace  $W$  of  $\mathbf{R}^n$ . Define:

1.  $u_1 = v_1$

# The Gram–Schmidt Process

## Procedure

### The Gram–Schmidt Process

Let  $\{v_1, v_2, \dots, v_m\}$  be a basis for a subspace  $W$  of  $\mathbf{R}^n$ . Define:

1.  $u_1 = v_1$

2.  $u_2 = v_2 - \text{proj}_{\text{Span}\{u_1\}}(v_2) \qquad = v_2 - \frac{v_2 \cdot u_1}{u_1 \cdot u_1} u_1$



# The Gram–Schmidt Process

## Procedure

### The Gram–Schmidt Process

Let  $\{v_1, v_2, \dots, v_m\}$  be a basis for a subspace  $W$  of  $\mathbf{R}^n$ . Define:

$$1. \ u_1 = v_1$$

$$2. \ u_2 = v_2 - \text{proj}_{\text{Span}\{u_1\}}(v_2) \qquad = v_2 - \frac{v_2 \cdot u_1}{u_1 \cdot u_1} u_1$$

$$3. \ u_3 = v_3 - \text{proj}_{\text{Span}\{u_1, u_2\}}(v_3) \qquad = v_3 - \frac{v_3 \cdot u_1}{u_1 \cdot u_1} u_1 - \frac{v_3 \cdot u_2}{u_2 \cdot u_2} u_2$$

# The Gram–Schmidt Process

## Procedure

### The Gram–Schmidt Process

Let  $\{v_1, v_2, \dots, v_m\}$  be a basis for a subspace  $W$  of  $\mathbf{R}^n$ . Define:

$$1. \quad u_1 = v_1$$

$$2. \quad u_2 = v_2 - \text{proj}_{\text{Span}\{u_1\}}(v_2) = v_2 - \frac{v_2 \cdot u_1}{u_1 \cdot u_1} u_1$$

$$3. \quad u_3 = v_3 - \text{proj}_{\text{Span}\{u_1, u_2\}}(v_3) = v_3 - \frac{v_3 \cdot u_1}{u_1 \cdot u_1} u_1 - \frac{v_3 \cdot u_2}{u_2 \cdot u_2} u_2$$

$$\vdots$$

$$m. \quad u_m = v_m - \text{proj}_{\text{Span}\{u_1, u_2, \dots, u_{m-1}\}}(v_m) = v_m - \sum_{i=1}^{m-1} \frac{v_m \cdot u_i}{u_i \cdot u_i} u_i$$

# The Gram–Schmidt Process

## Procedure

### The Gram–Schmidt Process

Let  $\{v_1, v_2, \dots, v_m\}$  be a basis for a subspace  $W$  of  $\mathbf{R}^n$ . Define:

$$1. \quad u_1 = v_1$$

$$2. \quad u_2 = v_2 - \text{proj}_{\text{Span}\{u_1\}}(v_2) = v_2 - \frac{v_2 \cdot u_1}{u_1 \cdot u_1} u_1$$

$$3. \quad u_3 = v_3 - \text{proj}_{\text{Span}\{u_1, u_2\}}(v_3) = v_3 - \frac{v_3 \cdot u_1}{u_1 \cdot u_1} u_1 - \frac{v_3 \cdot u_2}{u_2 \cdot u_2} u_2$$

$$\vdots$$

$$m. \quad u_m = v_m - \text{proj}_{\text{Span}\{u_1, u_2, \dots, u_{m-1}\}}(v_m) = v_m - \sum_{i=1}^{m-1} \frac{v_m \cdot u_i}{u_i \cdot u_i} u_i$$

Then  $\{u_1, u_2, \dots, u_m\}$  is an *orthogonal* basis for the same subspace  $W$ .

# The Gram–Schmidt Process

## Procedure

### The Gram–Schmidt Process

Let  $\{v_1, v_2, \dots, v_m\}$  be a basis for a subspace  $W$  of  $\mathbf{R}^n$ . Define:

$$1. \quad u_1 = v_1$$

$$2. \quad u_2 = v_2 - \text{proj}_{\text{Span}\{u_1\}}(v_2) = v_2 - \frac{v_2 \cdot u_1}{u_1 \cdot u_1} u_1$$

$$3. \quad u_3 = v_3 - \text{proj}_{\text{Span}\{u_1, u_2\}}(v_3) = v_3 - \frac{v_3 \cdot u_1}{u_1 \cdot u_1} u_1 - \frac{v_3 \cdot u_2}{u_2 \cdot u_2} u_2$$

$$\vdots$$

$$m. \quad u_m = v_m - \text{proj}_{\text{Span}\{u_1, u_2, \dots, u_{m-1}\}}(v_m) = v_m - \sum_{i=1}^{m-1} \frac{v_m \cdot u_i}{u_i \cdot u_i} u_i$$

Then  $\{u_1, u_2, \dots, u_m\}$  is an *orthogonal* basis for the same subspace  $W$ .

### Remark

In fact, for every  $i$  between 1 and  $n$ , the set  $\{u_1, u_2, \dots, u_i\}$  is an orthogonal basis for  $\text{Span}\{v_1, v_2, \dots, v_i\}$ .

# The Gram–Schmidt Process

Two vectors

Find an orthogonal basis  $\{u_1, u_2\}$  for  $W = \text{Span}\{v_1, v_2\}$ , where


$$v_1 = \begin{pmatrix} 1 \\ 1 \\ 0 \end{pmatrix} \quad \text{and} \quad v_2 = \begin{pmatrix} 1 \\ 1 \\ 1 \end{pmatrix}.$$

# The Gram–Schmidt Process

Two vectors

Find an orthogonal basis  $\{u_1, u_2\}$  for  $W = \text{Span}\{v_1, v_2\}$ , where

$$v_1 = \begin{pmatrix} 1 \\ 1 \\ 0 \end{pmatrix} \quad \text{and} \quad v_2 = \begin{pmatrix} 1 \\ 1 \\ 1 \end{pmatrix}.$$




# The Gram–Schmidt Process

Two vectors

Find an orthogonal basis  $\{u_1, u_2\}$  for  $W = \text{Span}\{v_1, v_2\}$ , where

$$v_1 = \begin{pmatrix} 1 \\ 1 \\ 0 \end{pmatrix} \quad \text{and} \quad v_2 = \begin{pmatrix} 1 \\ 1 \\ 1 \end{pmatrix}.$$



**Important:**  $\text{Span}\{u_1, u_2\} = \text{Span}\{v_1, v_2\} = W$ : this is an *orthogonal* basis for the *same* subspace.

# The Gram–Schmidt Process

Three vectors

Find an orthogonal basis  $\{u_1, u_2, u_3\}$  for  $W = \text{Span}\{v_1, v_2, v_3\} = \mathbf{R}^3$ , where

$$v_1 = \begin{pmatrix} 1 \\ 1 \\ 0 \end{pmatrix} \quad v_2 = \begin{pmatrix} 1 \\ 1 \\ 1 \end{pmatrix} \quad v_3 = \begin{pmatrix} 3 \\ 1 \\ 1 \end{pmatrix}.$$



# The Gram–Schmidt Process

Three vectors

Find an orthogonal basis  $\{u_1, u_2, u_3\}$  for  $W = \text{Span}\{v_1, v_2, v_3\} = \mathbf{R}^3$ , where

$$v_1 = \begin{pmatrix} 1 \\ 1 \\ 0 \end{pmatrix} \quad v_2 = \begin{pmatrix} 1 \\ 1 \\ 1 \end{pmatrix} \quad v_3 = \begin{pmatrix} 3 \\ 1 \\ 1 \end{pmatrix}.$$

**Important:**  $\text{Span}\{u_1, u_2, u_3\} = \text{Span}\{v_1, v_2, v_3\} = W$ : this is an *orthogonal* basis for the *same* subspace.

# The Gram–Schmidt Process


Three vectors, continued

$$v_1 = \begin{pmatrix} 1 \\ 1 \\ 0 \end{pmatrix}, \quad v_2 = \begin{pmatrix} 1 \\ 1 \\ 1 \end{pmatrix}, \quad v_3 = \begin{pmatrix} 3 \\ 1 \\ 1 \end{pmatrix} \xrightarrow{\text{G-S}} u_1 = \begin{pmatrix} 1 \\ 1 \\ 0 \end{pmatrix}, \quad u_2 = \begin{pmatrix} 0 \\ 0 \\ 1 \end{pmatrix}, \quad u_3 = \begin{pmatrix} 1 \\ -1 \\ 0 \end{pmatrix}$$

# The Gram–Schmidt Process

Three vectors, continued

$$v_1 = \begin{pmatrix} 1 \\ 1 \\ 0 \end{pmatrix}, \quad v_2 = \begin{pmatrix} 1 \\ 1 \\ 1 \end{pmatrix}, \quad v_3 = \begin{pmatrix} 3 \\ 1 \\ 1 \end{pmatrix} \xrightarrow{\text{G-S}} u_1 = \begin{pmatrix} 1 \\ 1 \\ 0 \end{pmatrix}, \quad u_2 = \begin{pmatrix} 0 \\ 0 \\ 1 \end{pmatrix}, \quad u_3 = \begin{pmatrix} 1 \\ -1 \\ 0 \end{pmatrix}$$



# The Gram–Schmidt Process

Three vectors in  $\mathbf{R}^4$

Find an orthogonal basis  $\{u_1, u_2, u_3\}$  for  $W = \text{Span}\{v_1, v_2, v_3\}$ , where

$$v_1 = \begin{pmatrix} 1 \\ 1 \\ 1 \\ 1 \end{pmatrix} \quad v_2 = \begin{pmatrix} -1 \\ 4 \\ 4 \\ -1 \end{pmatrix} \quad v_3 = \begin{pmatrix} 4 \\ -2 \\ -2 \\ 0 \end{pmatrix}.$$

## Poll

What happens if you try to run Gram–Schmidt on a linearly dependent set of vectors  $\{v_1, v_2, \dots, v_m\}$ ?

- A. You get an inconsistent equation.
- B. For some  $i$  you get  $u_i = u_{i-1}$ .
- C. For some  $i$  you get  $u_i = 0$ .
- D. You create a rift in the space-time continuum.

## Poll

What happens if you try to run Gram–Schmidt on a linearly dependent set of vectors  $\{v_1, v_2, \dots, v_m\}$ ?

- A. You get an inconsistent equation.
- B. For some  $i$  you get  $u_i = u_{i-1}$ .
- C. For some  $i$  you get  $u_i = 0$ .
- D. You create a rift in the space-time continuum.

If  $\{v_1, v_2, \dots, v_m\}$  is linearly dependent, then some  $v_i$  is in  $\text{Span}\{v_1, v_2, \dots, v_{i-1}\} = \text{Span}\{u_1, u_2, \dots, u_{i-1}\}$ .

## Poll

What happens if you try to run Gram–Schmidt on a linearly dependent set of vectors  $\{v_1, v_2, \dots, v_m\}$ ?

- A. You get an inconsistent equation.
- B. For some  $i$  you get  $u_i = u_{i-1}$ .
- C. For some  $i$  you get  $u_i = 0$ .
- D. You create a rift in the space-time continuum.

If  $\{v_1, v_2, \dots, v_m\}$  is linearly dependent, then some  $v_i$  is in  $\text{Span}\{v_1, v_2, \dots, v_{i-1}\} = \text{Span}\{u_1, u_2, \dots, u_{i-1}\}$ .

This means

$$\begin{aligned} v_i &= \text{proj}_{\text{Span}\{u_1, u_2, \dots, u_{i-1}\}}(v_i) \\ \implies u_i &= v_i - \text{proj}_{\text{Span}\{u_1, u_2, \dots, u_{i-1}\}}(v_i) = 0. \end{aligned}$$

## Poll

What happens if you try to run Gram–Schmidt on a linearly dependent set of vectors  $\{v_1, v_2, \dots, v_m\}$ ?

- A. You get an inconsistent equation.
- B. For some  $i$  you get  $u_i = u_{i-1}$ .
- C. For some  $i$  you get  $u_i = 0$ .
- D. You create a rift in the space-time continuum.

If  $\{v_1, v_2, \dots, v_m\}$  is linearly dependent, then some  $v_i$  is in  $\text{Span}\{v_1, v_2, \dots, v_{i-1}\} = \text{Span}\{u_1, u_2, \dots, u_{i-1}\}$ .

This means

$$\begin{aligned} v_i &= \text{proj}_{\text{Span}\{u_1, u_2, \dots, u_{i-1}\}}(v_i) \\ \implies u_i &= v_i - \text{proj}_{\text{Span}\{u_1, u_2, \dots, u_{i-1}\}}(v_i) = 0. \end{aligned}$$

In this case, you can simply discard  $u_i$  and  $v_i$  and continue: so Gram–Schmidt produces an orthogonal basis from any spanning set!



# QR Factorization

## QR Factorization Theorem

Let  $A$  be a matrix with linearly independent columns. Then

$$A = QR$$

where  $Q$  has orthonormal columns and  $R$  is upper-triangular with positive diagonal entries.

# QR Factorization

## QR Factorization Theorem

Let  $A$  be a matrix with linearly independent columns. Then

$$A = QR$$

where  $Q$  has orthonormal columns and  $R$  is upper-triangular with positive diagonal entries.

**Recall:** A set of vectors  $\{v_1, v_2, \dots, v_m\}$  is **orthonormal** if they are orthogonal unit vectors:  $v_i \cdot v_j = 0$  when  $i \neq j$ , and  $v_i \cdot v_i = 1$ .

# QR Factorization

## QR Factorization Theorem

Let  $A$  be a matrix with linearly independent columns. Then

$$A = QR$$

where  $Q$  has orthonormal columns and  $R$  is upper-triangular with positive diagonal entries.

**Recall:** A set of vectors  $\{v_1, v_2, \dots, v_m\}$  is **orthonormal** if they are orthogonal unit vectors:  $v_i \cdot v_j = 0$  when  $i \neq j$ , and  $v_i \cdot v_i = 1$ .

**Check:** A matrix  $Q$  has orthonormal columns if and only if  $Q^T Q = I$ .

# QR Factorization

## QR Factorization Theorem

Let  $A$  be a matrix with linearly independent columns. Then

$$A = QR$$

where  $Q$  has orthonormal columns and  $R$  is upper-triangular with positive diagonal entries.

**Recall:** A set of vectors  $\{v_1, v_2, \dots, v_m\}$  is **orthonormal** if they are orthogonal unit vectors:  $v_i \cdot v_j = 0$  when  $i \neq j$ , and  $v_i \cdot v_i = 1$ .

**Check:** A matrix  $Q$  has orthonormal columns if and only if  $Q^T Q = I$ .

The columns of  $A$  are a basis for  $W = \text{Col } A$ .

# QR Factorization

## QR Factorization Theorem

Let  $A$  be a matrix with linearly independent columns. Then

$$A = QR$$

where  $Q$  has orthonormal columns and  $R$  is upper-triangular with positive diagonal entries.

**Recall:** A set of vectors  $\{v_1, v_2, \dots, v_m\}$  is **orthonormal** if they are orthogonal unit vectors:  $v_i \cdot v_j = 0$  when  $i \neq j$ , and  $v_i \cdot v_i = 1$ .

**Check:** A matrix  $Q$  has orthonormal columns if and only if  $Q^T Q = I$ .

The columns of  $A$  are a basis for  $W = \text{Col } A$ . The columns of  $Q$  come from Gram-Schmidt as applied to the columns of  $A$ , after normalizing to unit vectors.

# QR Factorization

## QR Factorization Theorem

Let  $A$  be a matrix with linearly independent columns. Then

$$A = QR$$

where  $Q$  has orthonormal columns and  $R$  is upper-triangular with positive diagonal entries.

**Recall:** A set of vectors  $\{v_1, v_2, \dots, v_m\}$  is **orthonormal** if they are orthogonal unit vectors:  $v_i \cdot v_j = 0$  when  $i \neq j$ , and  $v_i \cdot v_i = 1$ .

**Check:** A matrix  $Q$  has orthonormal columns if and only if  $Q^T Q = I$ .

The columns of  $A$  are a basis for  $W = \text{Col } A$ . The columns of  $Q$  come from Gram–Schmidt as applied to the columns of  $A$ , after normalizing to unit vectors. The columns of  $R$  come from the steps in Gram–Schmidt.

# QR Factorization

## QR Factorization Theorem

Let  $A$  be a matrix with linearly independent columns. Then

$$A = QR$$

where  $Q$  has orthonormal columns and  $R$  is upper-triangular with positive diagonal entries.

**Recall:** A set of vectors  $\{v_1, v_2, \dots, v_m\}$  is **orthonormal** if they are orthogonal unit vectors:  $v_i \cdot v_j = 0$  when  $i \neq j$ , and  $v_i \cdot v_i = 1$ .

**Check:** A matrix  $Q$  has orthonormal columns if and only if  $Q^T Q = I$ .

The columns of  $A$  are a basis for  $W = \text{Col } A$ . The columns of  $Q$  come from Gram–Schmidt as applied to the columns of  $A$ , after normalizing to unit vectors. The columns of  $R$  come from the steps in Gram–Schmidt.

Here is the procedure for producing a  $QR$  factorization.

# QR Factorization

## Example

Find the  $QR$  factorization of  $A = \begin{pmatrix} 1 & 1 & 0 \\ 1 & 1 & 1 \\ 0 & 1 & 1 \end{pmatrix}$ .



# QR Factorization

## Example

Find the  $QR$  factorization of  $A = \begin{pmatrix} 1 & 1 & 0 \\ 1 & 1 & 1 \\ 0 & 1 & 1 \end{pmatrix}$ .

(The columns of  $A$  are the vectors  $v_1, v_2, v_3$  from a previous example.)

# QR Factorization

## Example

Find the QR factorization of  $A = \begin{pmatrix} 1 & 1 & 0 \\ 1 & 1 & 1 \\ 0 & 1 & 1 \end{pmatrix}$ .

(The columns of  $A$  are the vectors  $v_1, v_2, v_3$  from a previous example.)

Step 1: Run Gram-Schmidt

$$u_1 = v_1 = \begin{pmatrix} 1 \\ 1 \\ 0 \end{pmatrix}$$

$$u_2 = v_2 - \frac{v_2 \cdot u_1}{u_1 \cdot u_1} u_1 = v_2 - 1 u_1 = \begin{pmatrix} 0 \\ 0 \\ 1 \end{pmatrix}$$

$$\begin{aligned} u_3 &= v_3 - \frac{v_3 \cdot u_1}{u_1 \cdot u_1} u_1 - \frac{v_3 \cdot u_2}{u_2 \cdot u_2} u_2 \\ &= v_3 - 2 u_1 - 1 u_2 = \begin{pmatrix} 1 \\ -1 \\ 0 \end{pmatrix} \end{aligned}$$

# QR Factorization

## Example

Find the QR factorization of  $A = \begin{pmatrix} 1 & 1 & 0 \\ 1 & 1 & 1 \\ 0 & 1 & 1 \end{pmatrix}$ .

(The columns of  $A$  are the vectors  $v_1, v_2, v_3$  from a previous example.)

**Step 1:** Run Gram–Schmidt and solve for  $v_1, v_2, v_3$  in terms of  $u_1, u_2, u_3$ .

$$u_1 = v_1 = \begin{pmatrix} 1 \\ 1 \\ 0 \end{pmatrix} \qquad v_1 = u_1$$

$$u_2 = v_2 - \frac{v_2 \cdot u_1}{u_1 \cdot u_1} u_1 = v_2 - 1 u_1 = \begin{pmatrix} 0 \\ 0 \\ 1 \end{pmatrix} \qquad v_2 = u_1 + u_2$$

$$\begin{aligned} u_3 &= v_3 - \frac{v_3 \cdot u_1}{u_1 \cdot u_1} u_1 - \frac{v_3 \cdot u_2}{u_2 \cdot u_2} u_2 \\ &= v_3 - 2 u_1 - 1 u_2 = \begin{pmatrix} 1 \\ -1 \\ 0 \end{pmatrix} \qquad v_3 = 2u_1 + u_2 + u_3 \end{aligned}$$

# QR Factorization

Example, continued

$$v_1 = 1 u_1 \quad v_2 = 1 u_1 + 1 u_2 \quad v_3 = 2 u_1 + 1 u_2 + 1 u_3$$

Step 2: Write  $A = \hat{Q}\hat{R}$ , where  $\hat{Q}$  has *orthogonal* columns  $u_1, u_2, u_3$  and  $\hat{R}$  is upper-triangular with 1s on the diagonal.

# QR Factorization

Example, continued

$$v_1 = 1 u_1 \quad v_2 = 1 u_1 + 1 u_2 \quad v_3 = 2 u_1 + 1 u_2 + 1 u_3$$

**Step 2:** Write  $A = \hat{Q}\hat{R}$ , where  $\hat{Q}$  has *orthogonal* columns  $u_1, u_2, u_3$  and  $\hat{R}$  is upper-triangular with 1s on the diagonal.

Do this by putting the above equations in matrix form:

$$A \longrightarrow \begin{pmatrix} | & | & | \\ v_1 & v_2 & v_3 \\ | & | & | \end{pmatrix} = \begin{pmatrix} | & | & | \\ u_1 & u_2 & u_3 \\ | & | & | \end{pmatrix} \begin{pmatrix} 1 & 1 & 2 \\ 0 & 1 & 1 \\ 0 & 0 & 1 \end{pmatrix}$$

$\hat{Q}$   $\hat{R}$

# QR Factorization

Example, continued

$$v_1 = 1u_1 \quad v_2 = 1u_1 + 1u_2 \quad v_3 = 2u_1 + 1u_2 + 1u_3$$

**Step 2:** Write  $A = \hat{Q}\hat{R}$ , where  $\hat{Q}$  has *orthogonal* columns  $u_1, u_2, u_3$  and  $\hat{R}$  is upper-triangular with 1s on the diagonal.

Do this by putting the above equations in matrix form:

$$A \longrightarrow \begin{pmatrix} | & | & | \\ v_1 & v_2 & v_3 \\ | & | & | \end{pmatrix} = \begin{pmatrix} | & | & | \\ u_1 & u_2 & u_3 \\ | & | & | \end{pmatrix} \begin{pmatrix} 1 & 1 & 2 \\ 0 & 1 & 1 \\ 0 & 0 & 1 \end{pmatrix}$$

$\hat{Q}$   $\hat{R}$

first column of  $A = \begin{pmatrix} | & | & | \\ u_1 & u_2 & u_3 \\ | & | & | \end{pmatrix} \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix} = 1u_1 = v_1$

# QR Factorization

Example, continued

$$v_1 = 1u_1 \quad v_2 = 1u_1 + 1u_2 \quad v_3 = 2u_1 + 1u_2 + 1u_3$$

**Step 2:** Write  $A = \hat{Q}\hat{R}$ , where  $\hat{Q}$  has *orthogonal* columns  $u_1, u_2, u_3$  and  $\hat{R}$  is upper-triangular with 1s on the diagonal.

Do this by putting the above equations in matrix form:

$$A \longrightarrow \begin{pmatrix} | & | & | \\ v_1 & v_2 & v_3 \\ | & | & | \end{pmatrix} = \begin{pmatrix} | & | & | \\ u_1 & u_2 & u_3 \\ | & | & | \end{pmatrix} \begin{pmatrix} 1 & 1 & 2 \\ 0 & 1 & 1 \\ 0 & 0 & 1 \end{pmatrix}$$

$\hat{Q}$

$\hat{R}$

$$\text{first column of } A = \begin{pmatrix} | & | & | \\ u_1 & u_2 & u_3 \\ | & | & | \end{pmatrix} \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix} = 1u_1 = v_1$$

$$\text{second column of } A = \begin{pmatrix} | & | & | \\ u_1 & u_2 & u_3 \\ | & | & | \end{pmatrix} \begin{pmatrix} 1 \\ 1 \\ 0 \end{pmatrix} = 1u_1 + 1u_2 = v_2$$

# QR Factorization

Example, continued

$$v_1 = 1u_1 \quad v_2 = 1u_1 + 1u_2 \quad v_3 = 2u_1 + 1u_2 + 1u_3$$

**Step 2:** Write  $A = \hat{Q}\hat{R}$ , where  $\hat{Q}$  has *orthogonal* columns  $u_1, u_2, u_3$  and  $\hat{R}$  is upper-triangular with 1s on the diagonal.

Do this by putting the above equations in matrix form:

$$A \longrightarrow \begin{pmatrix} | & | & | \\ v_1 & v_2 & v_3 \\ | & | & | \end{pmatrix} = \begin{pmatrix} | & | & | \\ u_1 & u_2 & u_3 \\ | & | & | \end{pmatrix} \begin{pmatrix} 1 & 1 & 2 \\ 0 & 1 & 1 \\ 0 & 0 & 1 \end{pmatrix}$$

$\hat{Q}$

$\hat{R}$

$$\text{first column of } A = \begin{pmatrix} | & | & | \\ u_1 & u_2 & u_3 \\ | & | & | \end{pmatrix} \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix} = 1u_1 = v_1$$

$$\text{second column of } A = \begin{pmatrix} | & | & | \\ u_1 & u_2 & u_3 \\ | & | & | \end{pmatrix} \begin{pmatrix} 1 \\ 1 \\ 0 \end{pmatrix} = 1u_1 + 1u_2 = v_2$$

$$\text{third column of } A = \begin{pmatrix} | & | & | \\ u_1 & u_2 & u_3 \\ | & | & | \end{pmatrix} \begin{pmatrix} 2 \\ 1 \\ 1 \end{pmatrix} = 2u_1 + 1u_2 + 1u_3 = v_3$$



# QR Factorization

Example, continued

$$A = \hat{Q}\hat{R} \quad \begin{pmatrix} 1 & 1 & 0 \\ 1 & 1 & 1 \\ 0 & 1 & 1 \end{pmatrix} = \begin{pmatrix} 1 & 0 & 1 \\ 1 & 0 & -1 \\ 0 & 1 & 0 \end{pmatrix} \begin{pmatrix} 1 & 1 & 2 \\ 0 & 1 & 1 \\ 0 & 0 & 1 \end{pmatrix}$$

**Step 3:** Scale the columns of  $\hat{Q}$  to get unit vectors, and scale the rows of  $\hat{R}$  by the opposite factor, to get  $Q$  and  $R$ .

# QR Factorization

Example, continued

$$A = \hat{Q}\hat{R} \quad \begin{pmatrix} 1 & 1 & 0 \\ 1 & 1 & 1 \\ 0 & 1 & 1 \end{pmatrix} = \begin{pmatrix} 1 & 0 & 1 \\ 1 & 0 & -1 \\ 0 & 1 & 0 \end{pmatrix} \begin{pmatrix} 1 & 1 & 2 \\ 0 & 1 & 1 \\ 0 & 0 & 1 \end{pmatrix}$$

**Step 3:** Scale the columns of  $\hat{Q}$  to get unit vectors, and scale the rows of  $\hat{R}$  by the opposite factor, to get  $Q$  and  $R$ .

$$\begin{pmatrix} 1 & 1 & 0 \\ 1 & 1 & 1 \\ 0 & 1 & 1 \end{pmatrix} = \begin{pmatrix} 1 & 0 & 1 \\ 1 & 0 & -1 \\ 0 & 1 & 0 \end{pmatrix} \begin{pmatrix} 1 & 1 & 2 \\ 0 & 1 & 1 \\ 0 & 0 & 1 \end{pmatrix}.$$

# QR Factorization

Example, continued

$$A = \hat{Q}\hat{R} \quad \begin{pmatrix} 1 & 1 & 0 \\ 1 & 1 & 1 \\ 0 & 1 & 1 \end{pmatrix} = \begin{pmatrix} 1 & 0 & 1 \\ 1 & 0 & -1 \\ 0 & 1 & 0 \end{pmatrix} \begin{pmatrix} 1 & 1 & 2 \\ 0 & 1 & 1 \\ 0 & 0 & 1 \end{pmatrix}$$

**Step 3:** Scale the columns of  $\hat{Q}$  to get unit vectors, and scale the rows of  $\hat{R}$  by the opposite factor, to get  $Q$  and  $R$ .

$$\begin{pmatrix} 1 & 1 & 0 \\ 1 & 1 & 1 \\ 0 & 1 & 1 \end{pmatrix} = \begin{pmatrix} 1/\sqrt{2} & 0 & 1 \\ 1/\sqrt{2} & 0 & -1 \\ 0/\sqrt{2} & 1 & 0 \end{pmatrix} \begin{pmatrix} 1 \cdot \sqrt{2} & 1 \cdot \sqrt{2} & 2 \cdot \sqrt{2} \\ 0 & 1 & 1 \\ 0 & 0 & 1 \end{pmatrix}.$$

# QR Factorization

Example, continued

$$A = \hat{Q}\hat{R} \quad \begin{pmatrix} 1 & 1 & 0 \\ 1 & 1 & 1 \\ 0 & 1 & 1 \end{pmatrix} = \begin{pmatrix} 1 & 0 & 1 \\ 1 & 0 & -1 \\ 0 & 1 & 0 \end{pmatrix} \begin{pmatrix} 1 & 1 & 2 \\ 0 & 1 & 1 \\ 0 & 0 & 1 \end{pmatrix}$$

**Step 3:** Scale the columns of  $\hat{Q}$  to get unit vectors, and scale the rows of  $\hat{R}$  by the opposite factor, to get  $Q$  and  $R$ .

$$\begin{pmatrix} 1 & 1 & 0 \\ 1 & 1 & 1 \\ 0 & 1 & 1 \end{pmatrix} = \begin{pmatrix} 1/\sqrt{2} & 0/1 & 1 \\ 1/\sqrt{2} & 0/1 & -1 \\ 0/\sqrt{2} & 1/1 & 0 \end{pmatrix} \begin{pmatrix} 1 \cdot \sqrt{2} & 1 \cdot \sqrt{2} & 2 \cdot \sqrt{2} \\ 0 \cdot 1 & 1 \cdot 1 & 1 \cdot 1 \\ 0 & 0 & 1 \end{pmatrix}.$$

# QR Factorization

Example, continued

$$A = \hat{Q}\hat{R} \quad \begin{pmatrix} 1 & 1 & 0 \\ 1 & 1 & 1 \\ 0 & 1 & 1 \end{pmatrix} = \begin{pmatrix} 1 & 0 & 1 \\ 1 & 0 & -1 \\ 0 & 1 & 0 \end{pmatrix} \begin{pmatrix} 1 & 1 & 2 \\ 0 & 1 & 1 \\ 0 & 0 & 1 \end{pmatrix}$$

**Step 3:** Scale the columns of  $\hat{Q}$  to get unit vectors, and scale the rows of  $\hat{R}$  by the opposite factor, to get  $Q$  and  $R$ .

$$\begin{pmatrix} 1 & 1 & 0 \\ 1 & 1 & 1 \\ 0 & 1 & 1 \end{pmatrix} = \begin{pmatrix} 1/\sqrt{2} & 0/1 & 1/\sqrt{2} \\ 1/\sqrt{2} & 0/1 & -1/\sqrt{2} \\ 0/\sqrt{2} & 1/1 & 0/\sqrt{2} \end{pmatrix} \begin{pmatrix} 1 \cdot \sqrt{2} & 1 \cdot \sqrt{2} & 2 \cdot \sqrt{2} \\ 0 \cdot 1 & 1 \cdot 1 & 1 \cdot 1 \\ 0 \cdot \sqrt{2} & 0 \cdot \sqrt{2} & 1 \cdot \sqrt{2} \end{pmatrix}.$$

# QR Factorization

Example, continued

$$A = \hat{Q}\hat{R} \quad \begin{pmatrix} 1 & 1 & 0 \\ 1 & 1 & 1 \\ 0 & 1 & 1 \end{pmatrix} = \begin{pmatrix} 1 & 0 & 1 \\ 1 & 0 & -1 \\ 0 & 1 & 0 \end{pmatrix} \begin{pmatrix} 1 & 1 & 2 \\ 0 & 1 & 1 \\ 0 & 0 & 1 \end{pmatrix}$$

**Step 3:** Scale the columns of  $\hat{Q}$  to get unit vectors, and scale the rows of  $\hat{R}$  by the opposite factor, to get  $Q$  and  $R$ .

$$\begin{pmatrix} 1 & 1 & 0 \\ 1 & 1 & 1 \\ 0 & 1 & 1 \end{pmatrix} = \begin{pmatrix} 1/\sqrt{2} & 0/1 & 1/\sqrt{2} \\ 1/\sqrt{2} & 0/1 & -1/\sqrt{2} \\ 0/\sqrt{2} & 1/1 & 0/\sqrt{2} \end{pmatrix} \begin{pmatrix} 1 \cdot \sqrt{2} & 1 \cdot \sqrt{2} & 2 \cdot \sqrt{2} \\ 0 \cdot 1 & 1 \cdot 1 & 1 \cdot 1 \\ 0 \cdot \sqrt{2} & 0 \cdot \sqrt{2} & 1 \cdot \sqrt{2} \end{pmatrix}.$$

Note that the entries in the  $i$ th column of  $Q$  multiply by the entries in the  $i$ th row of  $R$ , so this doesn't change the product.

# QR Factorization

Example, continued

$$A = \hat{Q}\hat{R} \quad \begin{pmatrix} 1 & 1 & 0 \\ 1 & 1 & 1 \\ 0 & 1 & 1 \end{pmatrix} = \begin{pmatrix} 1 & 0 & 1 \\ 1 & 0 & -1 \\ 0 & 1 & 0 \end{pmatrix} \begin{pmatrix} 1 & 1 & 2 \\ 0 & 1 & 1 \\ 0 & 0 & 1 \end{pmatrix}$$

**Step 3:** Scale the columns of  $\hat{Q}$  to get unit vectors, and scale the rows of  $\hat{R}$  by the opposite factor, to get  $Q$  and  $R$ .

$$\begin{pmatrix} 1 & 1 & 0 \\ 1 & 1 & 1 \\ 0 & 1 & 1 \end{pmatrix} = \begin{pmatrix} 1/\sqrt{2} & 0/1 & 1/\sqrt{2} \\ 1/\sqrt{2} & 0/1 & -1/\sqrt{2} \\ 0/\sqrt{2} & 1/1 & 0/\sqrt{2} \end{pmatrix} \begin{pmatrix} 1 \cdot \sqrt{2} & 1 \cdot \sqrt{2} & 2 \cdot \sqrt{2} \\ 0 \cdot 1 & 1 \cdot 1 & 1 \cdot 1 \\ 0 \cdot \sqrt{2} & 0 \cdot \sqrt{2} & 1 \cdot \sqrt{2} \end{pmatrix}.$$

Note that the entries in the  $i$ th column of  $Q$  multiply by the entries in the  $i$ th row of  $R$ , so this doesn't change the product.

The final  $QR$  decomposition is:

$$A = QR \quad Q = \begin{pmatrix} 1/\sqrt{2} & 0 & 1/\sqrt{2} \\ 1/\sqrt{2} & 0 & -1/\sqrt{2} \\ 0 & 1 & 0 \end{pmatrix} \quad R = \begin{pmatrix} \sqrt{2} & \sqrt{2} & 2\sqrt{2} \\ 0 & 1 & 1 \\ 0 & 0 & \sqrt{2} \end{pmatrix}$$

# QR Factorization

Another example

Find the  $QR$  factorization of  $A = \begin{pmatrix} 1 & -1 & 4 \\ 1 & 4 & -2 \\ 1 & 4 & -2 \\ 1 & -1 & 0 \end{pmatrix}$ .



# QR Factorization

Another example

Find the  $QR$  factorization of  $A = \begin{pmatrix} 1 & -1 & 4 \\ 1 & 4 & -2 \\ 1 & 4 & -2 \\ 1 & -1 & 0 \end{pmatrix}$ .

(The columns are vectors from a previous example.)

**Step 1:** Run Gram–Schmidt and solve for  $v_1, v_2, v_3$  in terms of  $u_1, u_2, u_3$ :

## QR Factorization

Another example, continued

$$v_1 = 1 u_1 \quad v_2 = \frac{3}{2} u_1 + 1 u_2 \quad v_3 = 0 u_1 - \frac{4}{5} u_2 + 1 u_3$$

Step 2: Write  $A = \hat{Q}\hat{R}$ , where  $\hat{Q}$  has *orthogonal* columns  $u_1, u_2, u_3$  and  $\hat{R}$  is upper-triangular with 1s on the diagonal.

## QR Factorization

Another example, continued

$$A = \hat{Q}\hat{R} \quad \hat{Q} = \begin{pmatrix} 1 & -5/2 & 2 \\ 1 & 5/2 & 0 \\ 1 & 5/2 & 0 \\ 1 & -5/2 & -2 \end{pmatrix} \quad \hat{R} = \begin{pmatrix} 1 & 3/2 & 0 \\ 0 & 1 & -4/5 \\ 0 & 0 & 1 \end{pmatrix}$$

Step 3: Normalize the columns of  $\hat{Q}$  and the rows of  $\hat{R}$  to get  $Q$  and  $R$ :

## QR Factorization

Another example, continued

$$A = \hat{Q}\hat{R} \quad \hat{Q} = \begin{pmatrix} 1 & -5/2 & 2 \\ 1 & 5/2 & 0 \\ 1 & 5/2 & 0 \\ 1 & -5/2 & -2 \end{pmatrix} \quad \hat{R} = \begin{pmatrix} 1 & 3/2 & 0 \\ 0 & 1 & -4/5 \\ 0 & 0 & 1 \end{pmatrix}$$

Step 3: Normalize the columns of  $\hat{Q}$  and the rows of  $\hat{R}$  to get  $Q$  and  $R$ :

The final  $QR$  decomposition is

$$A = QR \quad Q = \begin{pmatrix} 1/2 & -1/2 & 1/\sqrt{2} \\ 1/2 & 1/2 & 0 \\ 1/2 & 1/2 & 0 \\ 1/2 & -1/2 & -1/\sqrt{2} \end{pmatrix} \quad R = \begin{pmatrix} 2 & 3 & 0 \\ 0 & 5 & -4 \\ 0 & 0 & 2\sqrt{2} \end{pmatrix}.$$

# QR Factorization

Application: computing determinants

Let  $A$  be an *invertible*  $n \times n$  matrix. Consider its  $QR$  factorization

$$A = QR.$$

# QR Factorization

Application: computing determinants

Let  $A$  be an *invertible*  $n \times n$  matrix. Consider its  $QR$  factorization

$$A = QR.$$

**Recall:** Since  $Q$  has orthonormal columns,  $Q^T Q = I_n$ , so  $Q^T = Q^{-1}$ .

# QR Factorization

Application: computing determinants

Let  $A$  be an *invertible*  $n \times n$  matrix. Consider its  $QR$  factorization

$$A = QR.$$

**Recall:** Since  $Q$  has orthonormal columns,  $Q^T Q = I_n$ , so  $Q^T = Q^{-1}$ .

But  $\det(Q^T) = \det(Q)$ , so

$$1 = \det(I_n) = \det(Q^T Q) = \det(Q^T) \det(Q) = \det(Q)^2.$$

# QR Factorization

Application: computing determinants

Let  $A$  be an *invertible*  $n \times n$  matrix. Consider its  $QR$  factorization

$$A = QR.$$

**Recall:** Since  $Q$  has orthonormal columns,  $Q^T Q = I_n$ , so  $Q^T = Q^{-1}$ .

But  $\det(Q^T) = \det(Q)$ , so

$$1 = \det(I_n) = \det(Q^T Q) = \det(Q^T) \det(Q) = \det(Q)^2.$$

It follows that  $\det(Q) = \pm 1$ .



# QR Factorization

Application: computing determinants

Let  $A$  be an *invertible*  $n \times n$  matrix. Consider its  $QR$  factorization

$$A = QR.$$

**Recall:** Since  $Q$  has orthonormal columns,  $Q^T Q = I_n$ , so  $Q^T = Q^{-1}$ .

But  $\det(Q^T) = \det(Q)$ , so

$$1 = \det(I_n) = \det(Q^T Q) = \det(Q^T) \det(Q) = \det(Q)^2.$$

It follows that  $\det(Q) = \pm 1$ .

(Since  $\det(R) > 0$ , in fact  $\det(Q)$  has the same sign as  $\det(A)$ .)

# QR Factorization

Application: computing determinants

Let  $A$  be an *invertible*  $n \times n$  matrix. Consider its  $QR$  factorization

$$A = QR.$$

**Recall:** Since  $Q$  has orthonormal columns,  $Q^T Q = I_n$ , so  $Q^T = Q^{-1}$ .

But  $\det(Q^T) = \det(Q)$ , so

$$1 = \det(I_n) = \det(Q^T Q) = \det(Q^T) \det(Q) = \det(Q)^2.$$

It follows that  $\det(Q) = \pm 1$ .

(Since  $\det(R) > 0$ , in fact  $\det(Q)$  has the same sign as  $\det(A)$ .)

Therefore,

$$\det(A) = \det(Q) \det(R) = \pm \det(R).$$

# QR Factorization

Application: computing determinants

Let  $A$  be an *invertible*  $n \times n$  matrix. Consider its  $QR$  factorization

$$A = QR.$$

**Recall:** Since  $Q$  has orthonormal columns,  $Q^T Q = I_n$ , so  $Q^T = Q^{-1}$ .

But  $\det(Q^T) = \det(Q)$ , so

$$1 = \det(I_n) = \det(Q^T Q) = \det(Q^T) \det(Q) = \det(Q)^2.$$

It follows that  $\det(Q) = \pm 1$ .

(Since  $\det(R) > 0$ , in fact  $\det(Q)$  has the same sign as  $\det(A)$ .)

Therefore,

$$\det(A) = \det(Q) \det(R) = \pm \det(R).$$

But  $R$  is upper-triangular, so it's easy to compute its determinant!

# QR Factorization

Application: computing determinants

Let  $A$  be an *invertible*  $n \times n$  matrix. Consider its QR factorization

$$A = QR.$$

**Recall:** Since  $Q$  has orthonormal columns,  $Q^T Q = I_n$ , so  $Q^T = Q^{-1}$ .

But  $\det(Q^T) = \det(Q)$ , so

$$1 = \det(I_n) = \det(Q^T Q) = \det(Q^T) \det(Q) = \det(Q)^2.$$

It follows that  $\det(Q) = \pm 1$ .

(Since  $\det(R) > 0$ , in fact  $\det(Q)$  has the same sign as  $\det(A)$ .)

Therefore,

$$\det(A) = \det(Q) \det(R) = \pm \det(R).$$

But  $R$  is upper-triangular, so it's easy to compute its determinant!

In fact, if  $v_1, v_2, \dots, v_n$  are the columns of  $A$ , and  $u_1, u_2, \dots, u_n$  are the vectors you obtain by applying Gram–Schmidt, then the  $(i, i)$  entry of  $R$  is  $\|u_i\|$ , so

$$\det(A) = \pm \|u_1\| \|u_2\| \cdots \|u_n\|.$$

# QR Factorization

Application: computing determinants

Let  $A$  be an *invertible*  $n \times n$  matrix. Consider its QR factorization

$$A = QR.$$

**Recall:** Since  $Q$  has orthonormal columns,  $Q^T Q = I_n$ , so  $Q^T = Q^{-1}$ .

But  $\det(Q^T) = \det(Q)$ , so

$$1 = \det(I_n) = \det(Q^T Q) = \det(Q^T) \det(Q) = \det(Q)^2.$$

It follows that  $\det(Q) = \pm 1$ .

(Since  $\det(R) > 0$ , in fact  $\det(Q)$  has the same sign as  $\det(A)$ .)

Therefore,

$$\det(A) = \det(Q) \det(R) = \pm \det(R).$$

But  $R$  is upper-triangular, so it's easy to compute its determinant!

In fact, if  $v_1, v_2, \dots, v_n$  are the columns of  $A$ , and  $u_1, u_2, \dots, u_n$  are the vectors you obtain by applying Gram–Schmidt, then the  $(i, i)$  entry of  $R$  is  $\|u_i\|$ , so

$$\det(A) = \pm \|u_1\| \|u_2\| \cdots \|u_n\|.$$

So you can use Gram–Schmidt to compute determinants (up to sign)!

# QR Factorization

Application: computing eigenvalues

Let  $A$  be an  $n \times n$  matrix with real eigenvalues. Here is an algorithm:

# QR Factorization

Application: computing eigenvalues

Let  $A$  be an  $n \times n$  matrix with real eigenvalues. Here is an algorithm:

$$A = Q_1 R_1 \quad QR \text{ factorization}$$

# QR Factorization

Application: computing eigenvalues

Let  $A$  be an  $n \times n$  matrix with real eigenvalues. Here is an algorithm:

$$A = Q_1 R_1 \quad QR \text{ factorization}$$

$$A_1 = R_1 Q_1 \quad \text{swap the } Q \text{ and } R$$



# QR Factorization

Application: computing eigenvalues

Let  $A$  be an  $n \times n$  matrix with real eigenvalues. Here is an algorithm:

$$A = Q_1 R_1 \quad QR \text{ factorization}$$

$$A_1 = R_1 Q_1 \quad \text{swap the } Q \text{ and } R$$

$$= Q_2 R_2 \quad \text{find its } QR \text{ factorization}$$

# QR Factorization

Application: computing eigenvalues

Let  $A$  be an  $n \times n$  matrix with real eigenvalues. Here is an algorithm:

$$A = Q_1 R_1 \quad \text{QR factorization}$$

$$A_1 = R_1 Q_1 \quad \text{swap the } Q \text{ and } R$$

$$= Q_2 R_2 \quad \text{find its QR factorization}$$

$$A_2 = R_2 Q_2 \quad \text{swap the } Q \text{ and } R$$

# QR Factorization

Application: computing eigenvalues

Let  $A$  be an  $n \times n$  matrix with real eigenvalues. Here is an algorithm:

$$A = Q_1 R_1 \quad \text{QR factorization}$$

$$A_1 = R_1 Q_1 \quad \text{swap the } Q \text{ and } R$$

$$= Q_2 R_2 \quad \text{find its QR factorization}$$

$$A_2 = R_2 Q_2 \quad \text{swap the } Q \text{ and } R$$

$$= Q_3 R_3 \quad \text{find its QR factorization}$$

# QR Factorization

Application: computing eigenvalues

Let  $A$  be an  $n \times n$  matrix with real eigenvalues. Here is an algorithm:

$$A = Q_1 R_1 \quad QR \text{ factorization}$$

$$A_1 = R_1 Q_1 \quad \text{swap the } Q \text{ and } R$$

$$= Q_2 R_2 \quad \text{find its } QR \text{ factorization}$$

$$A_2 = R_2 Q_2 \quad \text{swap the } Q \text{ and } R$$

$$= Q_3 R_3 \quad \text{find its } QR \text{ factorization}$$

et cetera

# QR Factorization

Application: computing eigenvalues

Let  $A$  be an  $n \times n$  matrix with real eigenvalues. Here is an algorithm:

$$\begin{aligned} A &= Q_1 R_1 && QR \text{ factorization} \\ A_1 &= R_1 Q_1 && \text{swap the } Q \text{ and } R \\ &= Q_2 R_2 && \text{find its } QR \text{ factorization} \\ A_2 &= R_2 Q_2 && \text{swap the } Q \text{ and } R \\ &= Q_3 R_3 && \text{find its } QR \text{ factorization} \\ &&& \text{et cetera} \end{aligned}$$

## Theorem

The matrices  $A_k$  converge to an upper triangular matrix, and the diagonal entries converge (quickly!) to the eigenvalues of  $A$ .

# QR Factorization

Application: computing eigenvalues

Let  $A$  be an  $n \times n$  matrix with real eigenvalues. Here is an algorithm:

$$\begin{aligned} A &= Q_1 R_1 && QR \text{ factorization} \\ A_1 &= R_1 Q_1 && \text{swap the } Q \text{ and } R \\ &= Q_2 R_2 && \text{find its } QR \text{ factorization} \\ A_2 &= R_2 Q_2 && \text{swap the } Q \text{ and } R \\ &= Q_3 R_3 && \text{find its } QR \text{ factorization} \\ &&& \text{et cetera} \end{aligned}$$

## Theorem

The matrices  $A_k$  converge to an upper triangular matrix, and the diagonal entries converge (quickly!) to the eigenvalues of  $A$ .

This gives a computationally efficient way (called the  $QR$  algorithm) to find the eigenvalues of a matrix.